



Australian Government

Great Barrier Reef  
Marine Park Authority

GREAT BARRIER REEF REGION STRATEGIC ASSESSMENT

# Strategic Assessment Report



The Great Barrier Reef Marine Park Authority acknowledges the continuing sea country management and custodianship of the Great Barrier Reef by Aboriginal and Torres Strait Islander Traditional Owners whose rich cultures, heritage values, enduring connections and shared efforts protect the Reef for future generations.

None of the recommendations in the report, if implemented, are intended to have the effect of extinguishing native title.





Australian Government

Great Barrier Reef  
Marine Park Authority

GREAT BARRIER REEF REGION STRATEGIC ASSESSMENT

# Strategic Assessment Report

© Commonwealth of Australia

Published by the Great Barrier Reef Marine Park Authority 2014

This work is copyright. Apart from any use as permitted under the *Copyright Act 1968*, no part may be reproduced by any process without the prior written permission of the Great Barrier Reef Marine Park Authority.

Requests and enquiries concerning reproduction and rights should be addressed to Communications,  
Great Barrier Reef Marine Park Authority, PO Box 1379, Townsville Queensland 4810

### Disclaimer

While reasonable efforts have been made to ensure that the contents of this publication are factually correct, the Great Barrier Reef Marine Park Authority does not make any representation or give any warranty regarding the accuracy, currency or suitability for any particular purpose of the source material contained in this publication. Readers should consult the source material referred to and, where necessary, seek appropriate advice about the suitability of the publication for their needs. To the extent permitted by law, the Great Barrier Reef Marine Park Authority shall not be liable for any loss, damage, cost or expense that may be occasioned directly or indirectly through the use of or reliance on the contents of this publication.

### National Library of Australia Cataloguing-in-Publication entry

Great Barrier Reef region strategic assessment : strategic assessment report / Great Barrier Reef Marine Park Authority.

ISBN 978 192212 649 8

Includes bibliographical references.

Environmental impact analysis—Queensland—Great Barrier Reef Region.  
Great Barrier Reef (Qld.)—Management.  
Great Barrier Reef (Qld.)—Environmental conditions.  
Great Barrier Reef (Qld.)—Economic conditions.  
Great Barrier Reef (Qld.)—Social conditions.

Great Barrier Reef Marine Park Authority.

333.91709943

### This publication should be cited as:

Great Barrier Reef Marine Park Authority 2014, *Great Barrier Reef Region Strategic Assessment: Strategic assessment report*, GBRMPA, Townsville.

Further information is available from:



**Australian Government**

**Great Barrier Reef  
Marine Park Authority**

2-68 Flinders Street East (PO Box 1379)

Townsville Queensland 4810

Telephone +617 4750 0700

Fax +617 4772 6093

Website [www.gbrmpa.gov.au](http://www.gbrmpa.gov.au)



# TABLE OF CONTENTS

	Glossary of commonly used terms and acronyms .....	xiii
<b>1</b>	<b>Introduction .....</b>	<b>1–1</b>
1.1	<b>The Great Barrier Reef .....</b>	<b>1–3</b>
1.1.1	Traditional Owner connections .....	1–3
1.1.2	International importance .....	1–3
1.1.3	National importance .....	1–6
1.1.4	Great Barrier Reef Marine Park Authority .....	1–6
1.1.5	Management approach .....	1–7
1.1.6	Evolving issues and management over time .....	1–7
1.2	<b>Comprehensive strategic assessment .....</b>	<b>1–8</b>
1.2.1	What is a strategic assessment? .....	1–8
1.2.2	Two complementary strategic assessments .....	1–8
1.2.3	Why undertake a strategic assessment? .....	1–9
1.2.4	Scope of the Great Barrier Reef Region strategic assessment .....	1–10
1.2.5	Matters of national environmental significance, including outstanding universal value .....	1–11
1.2.6	Strategic assessment process .....	1–12
1.2.7	Implications for the future .....	1–14
	<b>References .....</b>	<b>1–14</b>
<b>2</b>	<b>Assessment approach .....</b>	<b>2–1</b>
2.1	<b>Legislative basis .....</b>	<b>2–3</b>
2.2	<b>Matters of national environmental significance .....</b>	<b>2–3</b>
2.3	<b>Definitions and terminology .....</b>	<b>2–4</b>
2.4	<b>Principles .....</b>	<b>2–4</b>
2.5	<b>Complexity and scale .....</b>	<b>2–4</b>
2.6	<b>Steps in the assessment process .....</b>	<b>2–4</b>
2.7	<b>Assessment methods .....</b>	<b>2–5</b>
2.7.1	Assessment logic .....	2–5
2.7.2	Structured assessments .....	2–6
2.7.3	Assessing impacts .....	2–6
2.7.4	Assessing condition and trend .....	2–8
2.7.5	Benchmarking outstanding universal value .....	2–9
2.7.6	Assessing management effectiveness .....	2–9
2.7.7	Understanding resilience .....	2–9
2.7.8	Assessing future risk .....	2–9
2.7.9	Developing projected condition .....	2–9
2.7.10	Assessment structure .....	2–10
2.8	<b>Demonstration case studies .....</b>	<b>2–10</b>
2.9	<b>Recommended changes to management .....</b>	<b>2–11</b>
2.10	<b>Information used .....</b>	<b>2–11</b>
2.11	<b>Consultation and engagement .....</b>	<b>2–12</b>
2.11.1	Authority's advisory committees .....	2–12
2.11.2	Traditional Owner and stakeholder input .....	2–13
2.11.3	Input into assessment of management effectiveness .....	2–13
2.11.4	Presentations and meetings .....	2–14
2.12	<b>Independent peer review of the draft Strategic Assessment Report and draft Program Report .....</b>	<b>2–14</b>
2.13	<b>Public comment .....</b>	<b>2–14</b>
2.14	<b>Information gaps .....</b>	<b>2–14</b>
	<b>References .....</b>	<b>2–15</b>
<b>3</b>	<b>Current management .....</b>	<b>3–1</b>
3.1	<b>Context .....</b>	<b>3–3</b>
3.2	<b>Purpose .....</b>	<b>3–4</b>
3.3	<b>Strategic assessment area .....</b>	<b>3–5</b>
3.4	<b>Integrated management .....</b>	<b>3–5</b>
3.4.1	Jurisdiction .....	3–5
3.4.2	Great Barrier Reef Intergovernmental Agreement .....	3–6
3.4.3	Great Barrier Reef Ministerial Forum .....	3–7
3.5	<b>Legislation and conventions .....</b>	<b>3–7</b>
3.6	<b>Management tools .....</b>	<b>3–9</b>
3.6.1	Spatially explicit management tools .....	3–14
3.6.2	System-based regulatory tools .....	3–18
3.7	<b>Management activities .....</b>	<b>3–26</b>

<b>3.8</b>	<b>Partners in management .....</b>	<b>3–27</b>
<b>3.9</b>	<b>Management priorities .....</b>	<b>3–30</b>
3.9.1	Current priorities .....	3–30
<b>3.10</b>	<b>Applying the tools to achieve the priorities.....</b>	<b>3–30</b>
3.10.1	Addressing the key risks .....	3–30
3.10.2	Ensuring use is ecologically sustainable .....	3–32
3.10.3	Fostering stewardship .....	3–32
3.10.4	Effective and efficient organisation.....	3–33
3.10.5	Current resources for management .....	3–33
<b>3.11</b>	<b>Involvement in other planning and management frameworks .....</b>	<b>3–35</b>
3.11.1	National .....	3–35
3.11.2	State .....	3–35
3.11.3	Regional .....	3–36
3.11.4	Local.....	3–36
<b>3.12</b>	<b>Protecting matters of national environmental significance .....</b>	<b>3–36</b>
<b>3.13</b>	<b>Timeframe and review.....</b>	<b>3–38</b>
	<b>References .....</b>	<b>3–39</b>
<b>4</b>	<b>Matters of national environmental significance in the Great Barrier Reef Region .....</b>	<b>4–1</b>
<b>4.1</b>	<b>What are matters of national environmental significance? .....</b>	<b>4–3</b>
<b>4.2</b>	<b>World heritage properties.....</b>	<b>4–6</b>
4.2.1	Outstanding universal value .....	4–6
<b>4.3</b>	<b>The Great Barrier Reef Marine Park.....</b>	<b>4–8</b>
4.3.1	Biodiversity .....	4–9
4.3.2	Geomorphological features .....	4–14
4.3.3	Indigenous and historic heritage values .....	4–18
4.3.4	Community benefits of the environment.....	4–21
<b>4.4</b>	<b>National heritage places .....</b>	<b>4–23</b>
<b>4.5</b>	<b>Commonwealth marine areas .....</b>	<b>4–23</b>
<b>4.6</b>	<b>Listed migratory species .....</b>	<b>4–25</b>
4.6.1	Marine turtles.....	4–25
4.6.2	Estuarine crocodiles .....	4–28
4.6.3	Whales .....	4–28
4.6.4	Dolphins .....	4–28
4.6.5	Dugongs .....	4–29
4.6.6	Sharks .....	4–29
4.6.7	Seabirds .....	4–30
4.6.8	Shorebirds .....	4–32
<b>4.7</b>	<b>Listed threatened species and ecological communities .....</b>	<b>4–32</b>
4.7.1	Marine turtles.....	4–33
4.7.2	Marine mammals.....	4–33
4.7.3	Sharks and rays .....	4–33
4.7.4	Seabirds .....	4–34
<b>4.8</b>	<b>Wetlands of international importance .....</b>	<b>4–34</b>
4.8.1	Shoalwater and Corio Bays Area .....	4–34
4.8.2	Bowling Green Bay Area .....	4–35
<b>4.9</b>	<b>Environmental processes.....</b>	<b>4–36</b>
<b>4.10</b>	<b>Realising connectivity in the landscape .....</b>	<b>4–41</b>
<b>4.11</b>	<b>Connections between matters of national environmental significance.....</b>	<b>4–42</b>
<b>4.12</b>	<b>Summary of conclusions.....</b>	<b>4–46</b>
	<b>References .....</b>	<b>4–47</b>
<b>5</b>	<b>Drivers and activities .....</b>	<b>5–1</b>
<b>5.1</b>	<b>Background .....</b>	<b>5–3</b>
<b>5.2</b>	<b>Drivers of change .....</b>	<b>5–3</b>
5.2.1	Drivers relevant to the Region.....	5–3
5.2.2	Climate change .....	5–4
5.2.3	Economic growth.....	5–8
5.2.4	Population growth.....	5–11
5.2.5	Technological development.....	5–13
5.2.6	Societal attitudes .....	5–15
<b>5.3</b>	<b>Activities adjacent to the Region .....</b>	<b>5–16</b>
5.3.1	Agriculture .....	5–16
5.3.2	Aquaculture .....	5–19
5.3.3	Urban development.....	5–20
5.3.4	Industrial development and resource extraction.....	5–21
5.3.5	Port activities .....	5–22
<b>5.4</b>	<b>Activities within the Region .....</b>	<b>5–25</b>



5.4.1	Traditional use of marine resources.....	5–25
5.4.2	Tourism.....	5–26
5.4.3	Fishing — commercial.....	5–30
5.4.4	Fishing — recreational.....	5–35
5.4.5	Recreation.....	5–37
5.4.6	Shipping.....	5–41
5.4.7	Defence activities.....	5–46
5.4.8	Research activities.....	5–49
5.5	<b>Summary of conclusions</b> .....	5–49
	<b>References</b> .....	5–51
<b>6</b>	<b>Impacts on the values</b> .....	<b>6–1</b>
6.1	<b>Background</b> .....	<b>6–3</b>
6.1.1	Context.....	6–3
6.1.2	Challenges in assessing impacts.....	6–3
6.1.3	Terminology.....	6–4
6.2	<b>Impacts considered</b> .....	<b>6–4</b>
6.2.1	Identifying likely past and present impacts.....	6–4
6.3	<b>Legacy impacts</b> .....	<b>6–9</b>
6.4	<b>Impacts on biodiversity values</b> .....	<b>6–10</b>
6.4.1	Climate change impacts.....	6–10
6.4.2	Catchment run-off.....	6–15
6.4.3	Degradation of coastal ecosystems.....	6–26
6.4.4	Direct use.....	6–29
6.4.5	Assessment of impacts on biodiversity.....	6–46
6.5	<b>Impacts on geomorphological features</b> .....	<b>6–49</b>
6.6	<b>Impacts on Indigenous and historic heritage values</b> .....	<b>6–51</b>
6.6.1	Indigenous heritage values.....	6–51
6.6.2	Historic heritage values.....	6–54
6.7	<b>Impacts on community benefits of the environment</b> .....	<b>6–54</b>
6.8	<b>Cumulative impacts</b> .....	<b>6–55</b>
6.8.1	Qualitative models.....	6–55
6.8.2	Spatial approaches.....	6–58
6.8.3	Assessment of outcomes arising from cumulative impacts.....	6–66
6.9	<b>Effects on matters of national environmental significance</b> .....	<b>6–67</b>
6.9.1	World heritage properties.....	6–67
6.9.2	Great Barrier Reef Marine Park.....	6–68
6.9.3	National heritage places.....	6–68
6.9.4	Commonwealth marine area.....	6–69
6.9.5	Listed migratory and listed threatened species.....	6–69
6.9.6	Wetlands of international importance.....	6–69
6.10	<b>Key information gaps</b> .....	<b>6–69</b>
6.10.1	Biodiversity.....	6–70
6.10.2	Geomorphological features.....	6–71
6.10.3	Indigenous and historic heritage values.....	6–71
6.10.4	Community benefits of the environment.....	6–71
6.10.5	Cumulative impact assessment.....	6–71
6.10.6	Processes to address information gaps.....	6–72
6.11	<b>Summary of conclusions</b> .....	<b>6–73</b>
	<b>References</b> .....	<b>6–76</b>
<b>7</b>	<b>Current condition and trend</b> .....	<b>7–1</b>
7.1	<b>Biodiversity</b> .....	<b>7–3</b>
7.1.1	Summary of overall condition of biodiversity.....	7–13
7.1.2	Coral reefs.....	7–13
7.1.3	Seagrass meadows.....	7–15
7.1.4	Marine turtles.....	7–16
7.1.5	Inshore dolphins.....	7–17
7.1.6	Dugongs.....	7–19
7.1.7	Seabirds.....	7–19
7.1.8	Targeted bony fish species.....	7–19
7.1.9	Sharks and rays.....	7–19
7.2	<b>Geomorphological features</b> .....	<b>7–20</b>
7.3	<b>Indigenous and historic heritage values</b> .....	<b>7–22</b>
7.3.1	Traditional Owner connections.....	7–24
7.3.2	Cultural keystone species.....	7–24
7.3.3	Identifying Indigenous heritage values.....	7–25
7.3.4	Historic heritage values.....	7–25

<b>7.4</b>	<b>Community benefits of the environment .....</b>	<b>7–25</b>
7.4.1	Economic benefit.....	7–27
7.4.2	Visitation.....	7–28
7.4.3	Understanding and appreciation .....	7–28
7.4.4	Personal connections .....	7–29
7.4.5	Knowledge about community benefits.....	7–29
<b>7.5</b>	<b>Environmental processes.....</b>	<b>7–29</b>
<b>7.6</b>	<b>Condition and trend of matters of national environmental significance .....</b>	<b>7–33</b>
7.6.1	World heritage properties.....	7–33
7.6.2	Great Barrier Reef Marine Park.....	7–41
7.6.3	National heritage places.....	7–41
7.6.4	Commonwealth marine area .....	7–42
7.6.5	Listed migratory and listed threatened species .....	7–42
7.6.6	Wetlands of international importance .....	7–43
<b>7.7</b>	<b>Key information gaps.....</b>	<b>7–43</b>
7.7.1	Biodiversity — habitats and species.....	7–43
7.7.2	Geomorphological features .....	7–44
7.7.3	Indigenous and historic heritage values .....	7–44
7.7.4	Community benefits of the environment.....	7–45
7.7.5	Environmental processes.....	7–45
7.7.6	Processes to address information gaps .....	7–46
<b>7.8</b>	<b>Summary of conclusions.....</b>	<b>7–46</b>
	<b>References .....</b>	<b>7–47</b>

## **8 Management effectiveness — an independent assessment.....8–1**

<b>8.1</b>	<b>Background .....</b>	<b>8–3</b>
<b>8.2</b>	<b>Assessment method .....</b>	<b>8–4</b>
8.2.1	Impacts and values assessed .....	8–6
8.2.2	Calculation of grades.....	8–7
8.2.3	Traditional Owner and stakeholder input.....	8–9
<b>8.3</b>	<b>Protecting values .....</b>	<b>8–10</b>
8.3.1	Biodiversity protection .....	8–10
8.3.2	Indigenous and historic heritage values .....	8–11
8.3.3	Community benefits.....	8–14
<b>8.4</b>	<b>Assessing external impacts.....</b>	<b>8–15</b>
8.4.1	Climate change and extreme weather.....	8–15
8.4.2	Water quality protection (catchment run-off) .....	8–17
8.4.3	Coastal development (protection of coastal ecosystems) .....	8–18
<b>8.5</b>	<b>Managing direct use.....</b>	<b>8–19</b>
8.5.1	Tourism .....	8–19
8.5.2	Fishing — commercial.....	8–21
8.5.3	Fishing — recreational .....	8–22
8.5.4	Recreation .....	8–23
8.5.5	Port activities .....	8–24
8.5.6	Shipping .....	8–26
8.5.7	Defence activities .....	8–27
8.5.8	Research activities .....	8–28
<b>8.6</b>	<b>Effectiveness of specific measures to address impacts.....</b>	<b>8–29</b>
8.6.1	Understanding impacts and considering issues.....	8–30
8.6.2	Managing impacts .....	8–31
8.6.3	Providing certainty.....	8–32
8.6.4	Halt and reverse declines.....	8–44
8.6.5	Adapting to climate change .....	8–44
8.6.6	Integrating with government programs.....	8–45
8.6.7	Meeting Australia's international obligations .....	8–46
8.6.8	Monitoring, evaluating and reporting .....	8–47
<b>8.7</b>	<b>Effectiveness of specific measures to protect values.....</b>	<b>8–48</b>
8.7.1	Identifying values and determining their condition and trend .....	8–48
8.7.2	Monitoring, evaluating and reporting of values.....	8–49
<b>8.8</b>	<b>Effectiveness of specific measures to protect matters of national environmental significance.....</b>	<b>8–50</b>
<b>8.9</b>	<b>Summary of conclusions.....</b>	<b>8–55</b>
	<b>References .....</b>	<b>8–56</b>

## **9 Demonstration case studies .....**9–1

<b>9.1</b>	<b>Case studies selected .....</b>	<b>9–3</b>
9.1.1	Topics chosen .....	9–3
9.1.2	Meeting the selection criteria.....	9–4



<b>9.2</b>	<b>Assessing management effectiveness .....</b>	<b>9-5</b>
<b>9.3</b>	<b>Dugongs .....</b>	<b>9-6</b>
9.3.1	Significance.....	9-6
9.3.2	Key issues.....	9-6
9.3.3	Current management and its effectiveness .....	9-8
9.3.4	Outcomes and potential improvements .....	9-9
<b>9.4</b>	<b>Corals .....</b>	<b>9-10</b>
9.4.1	Significance.....	9-10
9.4.2	Key issues.....	9-10
9.4.3	Current management and its effectiveness .....	9-11
9.4.4	Outcomes and potential improvements .....	9-12
<b>9.5</b>	<b>Islands.....</b>	<b>9-13</b>
9.5.1	Significance.....	9-13
9.5.2	Responsibilities for islands.....	9-13
9.5.3	Key issues.....	9-14
9.5.4	Current management and its effectiveness .....	9-15
9.5.5	Outcomes and potential improvements .....	9-17
<b>9.6</b>	<b>Princess Charlotte Bay.....</b>	<b>9-18</b>
9.6.1	Significance.....	9-18
9.6.2	Key issues.....	9-20
9.6.3	Current management and its effectiveness .....	9-20
9.6.4	Outcomes and potential improvements .....	9-21
<b>9.7</b>	<b>Cairns Planning Area.....</b>	<b>9-21</b>
9.7.1	Significance.....	9-21
9.7.2	Key issues.....	9-22
9.7.3	Current management and its effectiveness .....	9-22
9.7.4	Outcomes and potential improvements .....	9-24
<b>9.8</b>	<b>Mackay Whitsunday — water quality improvement .....</b>	<b>9-25</b>
9.8.1	Significance.....	9-25
9.8.2	Key issues.....	9-25
9.8.3	Current management and its effectiveness .....	9-27
9.8.4	Outcomes and potential improvements .....	9-29
<b>9.9</b>	<b>Keppel Bay .....</b>	<b>9-29</b>
9.9.1	Significance.....	9-29
9.9.2	Key issues.....	9-30
9.9.3	Current management and its effectiveness .....	9-31
9.9.4	Outcomes and potential improvements .....	9-32
<b>9.10</b>	<b>East Coast Trawl Fishery .....</b>	<b>9-32</b>
9.10.1	Significance.....	9-32
9.10.2	Key issues.....	9-33
9.10.3	Current management and its effectiveness .....	9-36
9.10.4	Outcomes and potential improvements .....	9-37
<b>9.11</b>	<b>Summary of conclusions and guidance for future management.....</b>	<b>9-37</b>
	<b>References.....</b>	<b>9-38</b>
<b>10</b>	<b>Resilience and risk.....</b>	<b>10-1</b>
<b>10.1</b>	<b>Understanding ecosystem resilience .....</b>	<b>10-3</b>
10.1.1	Factors that affect ecosystem resilience.....	10-3
<b>10.2</b>	<b>Resilience of key values of biodiversity .....</b>	<b>10-4</b>
10.2.1	Coral reefs and corals.....	10-4
10.2.2	Seagrass meadows .....	10-4
10.2.3	Lagoonal habitats.....	10-4
10.2.4	Deep water and continental slope habitats .....	10-5
10.2.5	Inshore habitats .....	10-5
10.2.6	Bony fishes .....	10-5
10.2.7	Dugongs.....	10-5
10.2.8	Marine turtles .....	10-5
10.2.9	Sharks and rays .....	10-6
10.2.10	Seabirds.....	10-6
<b>10.3</b>	<b>Overall ecosystem resilience.....</b>	<b>10-6</b>
<b>10.4</b>	<b>Active restoration of ecosystem resilience.....</b>	<b>10-7</b>
<b>10.5</b>	<b>Social and cultural resilience .....</b>	<b>10-7</b>
<b>10.6</b>	<b>Assessing risks on the Region's values .....</b>	<b>10-8</b>
<b>10.7</b>	<b>Risks to biodiversity.....</b>	<b>10-9</b>
<b>10.8</b>	<b>Risks to geomorphological features .....</b>	<b>10-14</b>
<b>10.9</b>	<b>Risks to Indigenous heritage values.....</b>	<b>10-15</b>
<b>10.10</b>	<b>Risks to historic heritage values.....</b>	<b>10-19</b>
<b>10.11</b>	<b>Summary of risks to matters of national environmental significance.....</b>	<b>10-20</b>

10.12	Summary of conclusions.....	10-21
	References .....	10-25
11	Projected condition .....	11-1
11.1	Current condition and trend of values .....	11-3
11.2	Effectiveness of current management .....	11-3
11.2.1	Broadscale assessment .....	11-4
11.2.2	Outcomes of demonstration case studies .....	11-4
11.3	Ecosystem resilience.....	11-4
11.4	Overall risks to values .....	11-5
11.5	Projected condition.....	11-5
11.5.1	The future of the Great Barrier Reef.....	11-5
11.5.2	Projected condition of the Region's values .....	11-6
11.6	Future scenarios .....	11-15
11.6.1	Coral reef scenarios .....	11-16
11.6.2	Seagrass meadow scenarios .....	11-18
11.6.3	Dugong scenarios .....	11-18
	References .....	11-20
12	Recommended changes to management.....	12-1
12.1	How recommended improvements were identified .....	12-3
12.2	The need to strengthen the Authority's management .....	12-3
12.3	Recommended improvements .....	12-4
12.3.1	Recommended improvements to the Authority's management arrangements .....	12-4
12.3.2	Recommended improvements to related local, state and national government programs .....	12-9
12.4	Recommendations of the coastal zone assessment .....	12-12
13	Adaptive Management .....	13-1
13.1	Background .....	13-3
13.2	Comprehensive management program (Part B).....	13-4
13.3	The Permission System (Part C).....	13-6
13.3.1	Benefits of approving the Permission System.....	13-6
13.3.2	Activities under the program.....	13-6
13.3.3	Permission System objectives and outcomes .....	13-7
13.3.4	Permission System process .....	13-8
13.3.5	Summary of program commitments .....	13-20
13.3.6	Permission System alignment with the EPBC Act.....	13-21
13.4	Applying the principles of ecologically sustainable development.....	13-25
13.5	Applying adaptive management .....	13-29
13.5.1	Addressing risk.....	13-29
13.5.2	Addressing uncertainty.....	13-29
13.5.3	Implementing adaptive management to protect and manage matters of national environmental significance .....	13-30
13.5.4	Addressing World Heritage Committee recommendations .....	13-31
13.6	Monitoring and reporting.....	13-34
13.6.1	Monitoring.....	13-34
13.6.2	Review and public reporting .....	13-34
13.6.3	Adapting to new information.....	13-34
13.6.4	Parties responsible for monitoring and reporting.....	13-35
13.7	Addressing the endorsement criteria.....	13-35
13.8	Program review, modification and continuous improvement.....	13-42
13.8.1	Review and continuous improvement of Program Report Part B.....	13-42
13.8.2	Review and continuous improvement of Program Report Part C.....	13-42
	References .....	13-42
Appendix 1	Agreement with the then Minister for Sustainability, Environment, Water, Population and Communities	
Appendix 2	Great Barrier Reef Region Strategic Assessment Terms of Reference	
Appendix 3	Statement of Outstanding Universal Value, Great Barrier Reef – Property ID 154	
Appendix 4	Assessment of Management Effectiveness Terms of Reference	
Appendix 5	Traditional Owner and Stakeholder Engagement – Report on Workshops and Surveys	
Appendix 6	Matters of national environmental significance protected under the EPBC Act that could be impacted	
Appendix 7	Permission system case studies	
Appendix 8	Terms of Reference – concordance table	



# LIST OF FIGURES

Figure 1.1	Great Barrier Reef Region, World Heritage Area and Marine Park .....	1–4
Figure 1.2	Outstanding universal value of the Great Barrier Reef .....	1–5
Figure 1.3	Components of the comprehensive strategic assessment of the Great Barrier Reef World Heritage Area and adjacent coastal zone .....	1–9
Figure 1.4	An example of the different coastal boundaries for the Great Barrier Reef World Heritage Area, Region and Marine Park.....	1–11
Figure 1.5	Strategic assessment process .....	1–13
Figure 2.1	Steps in the assessment process .....	2–5
Figure 2.2	Assessment logic for the strategic assessment of the Great Barrier Reef Region.....	2–5
Figure 2.3	Indicative broad assessment areas for the Region and its catchment.....	2–8
Figure 3.1	Jurisdictional boundaries for the Great Barrier Reef Region.....	3–6
Figure 3.2	The Great Barrier Reef Zoning Plan .....	3–15
Figure 3.3	Administrative arrangements for the joint application and permit assessment process between the Authority and the Queensland Government .....	3–21
Figure 3.4	Administrative arrangements for environmental assessments under the GBRMP Act and EPBC Act.....	3–22
Figure 3.5	Bowling Green Bay Special Management Area.....	3–29
Figure 4.1	Timeline for the matters of national environmental significance relevant to the Great Barrier Reef Region.....	4–4
Figure 4.2	Matters of national environmental significance in the Great Barrier Reef Region.....	4–5
Figure 4.3	Spatial extent of the matters of national environmental significance for the Great Barrier Reef Region .....	4–6
Figure 4.4	Reef bioregions of the Great Barrier Reef .....	4–11
Figure 4.5	Non-reef bioregions of the Great Barrier Reef .....	4–12
Figure 4.6	Three-dimensional bathymetry of the Great Barrier Reef shelf.....	4–15
Figure 4.7	Three-dimensional bathymetry of submarine canyons along the Great Barrier Reef shelf edge....	4–17
Figure 4.8	Dominant values identified by Rockhampton Traditional Owner participants at a strategic assessment workshop .....	4–18
Figure 4.9	Boundaries of the Commonwealth marine area within and outside the Region.....	4–24
Figure 4.10	Identified important feeding and breeding areas for four species of marine turtles within the Region .....	4–27
Figure 4.11	Principal seabird breeding islands in the Region .....	4–30
Figure 4.12	Shoalwater and Corio Bays Area.....	4–35
Figure 4.13	Bowling Green Bay Area .....	4–36
Figure 4.14	Key environmental processes of the Great Barrier Reef Region .....	4–37
Figure 4.15	Crossing the blue highway.....	4–40
Figure 4.16	A method for identifying and prioritising coastal ecosystem functional connections to the Great Barrier Reef World Heritage Area and the Great Barrier Reef Marine Park.....	4–42
Figure 5.1	The influence of drivers and activities.....	5–3
Figure 5.2	Projected trends in climate change variables and the Region's biodiversity values .....	5–6
Figure 5.3	Mean atmospheric concentrations and the annual mean increase of carbon dioxide, 1958–2013.....	5–7
Figure 5.4	Major resource projects planned or underway in Queensland.....	5–10
Figure 5.5	Wells drilled in Queensland, 1954–2011 .....	5–11
Figure 5.6	Predicted population growth in regional centres of the catchment, 2006–2031 .....	5–12
Figure 5.7	Forecast average annual population growth by local government area, 2006–2031 .....	5–13
Figure 5.8	Activities within and adjacent to the Region.....	5–16
Figure 5.9	Land use in the Great Barrier Reef catchment, 2009 and 1999.....	5–17
Figure 5.10	Fertiliser use in the Great Barrier Reef catchment.....	5–19
Figure 5.11	Queensland trading ports in the Great Barrier Reef World Heritage Area, Region and Marine Park and their throughput, 2011–12 .....	5–23
Figure 5.12	Port infrastructure at Mackay and the Port of Hay Point.....	5–24
Figure 5.13	Distribution of tourism activity in the Great Barrier Reef Marine Park, 2012 .....	5–27
Figure 5.14	Number of tourism visitor days to the Great Barrier Reef Marine Park, 1994–2012 .....	5–28
Figure 5.15	International visitation to the Great Barrier Reef catchment, 2002–2011.....	5–28
Figure 5.16	Percentage of visitors to the Marine Park carried on high standard tourism operations, 2004–2012.....	5–29
Figure 5.17	Spatial distribution of catch by commercial line, net, pot and trawl fisheries, Great Barrier Reef Region, 2011 .....	5–32
Figure 5.18	Annual trawl fishing effort in the Great Barrier Reef Region, 1990–2011 .....	5–34
Figure 5.19	Number of recreational fishers by Queensland residential region in 2000 and 2010 .....	5–36
Figure 5.20	Recreational fishing compliance incidents by marine park zone, 2008–2012.....	5–37
Figure 5.21	Main activities of visitors to the Great Barrier Reef.....	5–38
Figure 5.22	Main destinations of recreational vessel-based trips, 2003 and 2008.....	5–38
Figure 5.23	Vessel registrations in the Great Barrier Reef catchment, 2011 .....	5–39
Figure 5.24	Number of recreational vessels registered in the Great Barrier Reef catchment, 2001–2011 .....	5–40
Figure 5.25	Total commercial vessel calls to ports adjacent to the Region, 2011–12.....	5–41
Figure 5.26	Designated Shipping Area and designated ship anchorages, Great Barrier Reef Region .....	5–42
Figure 5.27	Polled ship positions, Great Barrier Reef Region, 2000.....	5–44

Figure 5.28	Polled ship positions, Great Barrier Reef Region, 2010 .....	5–45
Figure 5.29	A projection of shipping increases at major ports adjacent to the Great Barrier Reef Region, 2012–2032 .....	5–46
Figure 5.30	Defence training areas relevant to the Region .....	5–47
Figure 6.1	Direct, indirect and consequential impacts .....	6–4
Figure 6.2	Cyclone winds (2005–2011) and flood plume exposure (1991–2011) in the Great Barrier Reef Region .....	6–11
Figure 6.3	Cyclone wave exposure in the Region, 2000–2011 .....	6–12
Figure 6.4	Summer sea surface temperature anomalies in the Region, 2001–2011 .....	6–13
Figure 6.5	Global average sea level, 1880–2011 .....	6–14
Figure 6.6	Rates of sea level rise in Australia .....	6–15
Figure 6.7	Annual freshwater discharge from the major rivers of the Great Barrier Reef catchment, 2002–2012 .....	6–15
Figure 6.8	Freshwater exposure in the Region, 2001–2011 .....	6–16
Figure 6.9	Years that chlorophyll concentrations exceeded the guidelines in the Region, 2003–2012 .....	6–18
Figure 6.10	Potential role of nutrients in the population dynamics of crown-of-thorns starfish .....	6–19
Figure 6.11	Extent of observed crown-of-thorns starfish outbreaks in the Region, 1986–2011 .....	6–20
Figure 6.12	Removal of crown-of-thorns starfish, offshore Cairns, August 2012 – September 2013 .....	6–21
Figure 6.13	Areas of exposure to PSII herbicide residues modelled in the Great Barrier Reef Region .....	6–23
Figure 6.14	Years that total suspended solids exceeded the guidelines in the Region, 2003–2012 .....	6–24
Figure 6.15	Water clarity at three tourism sites .....	6–25
Figure 6.16	Coastal ecosystems of the Great Barrier Reef catchment before European settlement .....	6–28
Figure 6.17	Coastal ecosystems of the Great Barrier Reef catchment as they are today .....	6–29
Figure 6.18	Dredge material disposed to date, approved for disposal or being assessed, by port .....	6–31
Figure 6.19	Surface large-scale currents for El Niño (2004), neutral (2007) and La Niña (2011) conditions at major ports .....	6–33
Figure 6.20	Retained and non-retained fisheries catch, Great Barrier Reef, 2007 .....	6–34
Figure 6.21	A breakdown of commercial fisheries non-retained catch, Great Barrier Reef, 2007 .....	6–35
Figure 6.22	Ecological groups retained by commercial fishing, Great Barrier Reef, 2007 .....	6–38
Figure 6.23	Man-made noise sources and the hearing ranges of marine mammals and fish .....	6–39
Figure 6.24	Reported vessel groundings in the Region, 1987–2012 .....	6–42
Figure 6.25	Reported spills and discharges in and adjacent to the Region, 1987–2012 .....	6–44
Figure 6.26	Qualitative model of how coral reefs are affected by drivers, activities and impacts .....	6–57
Figure 6.27	Qualitative model of how seagrass meadows are affected by drivers, activities and impacts .....	6–58
Figure 6.28	Exposure of coral reefs to key impacts, 2001–2011 .....	6–59
Figure 6.29	Cumulative exposure of coral reefs to key impacts, 2001–2011 .....	6–60
Figure 6.30	Key water quality exposures in the Region (part 1) .....	6–63
Figure 6.31	Key water quality exposures in the Region (part 2) .....	6–64
Figure 6.32	Combined key water quality exposures .....	6–65
Figure 6.33	Zones of influence of different activities .....	6–67
Figure 7.1	Shifting baselines on inshore coral reefs, Stone Island, offshore Bowen .....	7–5
Figure 7.2	Hard coral cover in the Great Barrier Reef, 1986–2012 .....	7–14
Figure 7.3	Average abundance of intertidal seagrass meadows, 1999–2012 .....	7–16
Figure 7.4	Cumulative monthly marine turtle strandings for the Queensland east coast, 1998–2012 .....	7–16
Figure 7.5	Overlap of protected areas and inshore dolphin populations .....	7–18
Figure 8.1	Framework for assessing management effectiveness of protected areas .....	8–4
Figure 8.2	Certainty provided for tourism activities by the Authority's management arrangements .....	8–34
Figure 8.3	Certainty provided for fishing and collecting activities by the Authority's management arrangements .....	8–36
Figure 8.4	Activities guide — an indication of what can be done in each zone .....	8–37
Figure 8.5	Certainty provided for research activities by the Authority's management arrangements .....	8–38
Figure 8.6	Certainty provided for recreation activities by the Authority's management arrangements .....	8–39
Figure 8.7	Certainty provided for traditional use by the Authority's management arrangements .....	8–40
Figure 8.8	Certainty provided for defence activities by the Authority's management arrangements .....	8–41
Figure 8.9	Certainty provided for shipping activities by the Authority's management arrangements .....	8–42
Figure 8.10	Certainty provided for commercial port operations by the Authority's management arrangements .....	8–43
Figure 9.1	Dugongs caught in the Queensland Shark Control Program, 1962–1999 .....	9–7
Figure 9.2	Relationships between the rainforest, islands and the migratory pied imperial pigeons .....	9–13
Figure 9.3	Princess Charlotte Bay .....	9–18
Figure 9.4	Life cycle of the largemouth sawfish .....	9–19
Figure 9.5	Cairns Planning Area .....	9–23
Figure 9.6	Coastal ecosystems in the Mackay Whitsunday natural resource management region, 2009 .....	9–26
Figure 9.7	Conceptual diagram of land use in the Mackay Whitsunday region .....	9–27
Figure 9.8	Key components of the <i>Reef Water Quality Protection Plan</i> .....	9–28
Figure 9.9	Keppel Bay and its islands .....	9–31
Figure 9.10	Chronology of the Queensland East Coast Trawl Fishery .....	9–35
Figure 11.1	Future modelled changes in the condition of coral reefs .....	11–17
Figure 11.2	Future modelled changes in the condition of seagrass meadows and dugongs .....	11–19
Figure 13.1	The four key steps in the permission assessment and decision process .....	13–9
Figure 13.2	Principles of ecologically sustainable use .....	13–25
Figure 13.3	Pathways of management intervention .....	13–31

# LIST OF TABLES

Table 1.1	Chronology of key management responses to emerging issues from the 1970s to the present.....	1–8
Table 1.2	Differences between the Great Barrier Reef World Heritage Area, Region and Marine Park .....	1–11
Table 2.1	Tools to improve understanding of cause-and-effect relationships.....	2–7
Table 3.1	Management tools employed by the Authority to protect and manage the Great Barrier Reef Region and relevant matters of national environmental significance .....	3–10
Table 3.2	Program guidance documents and <i>Environment Protection and Biodiversity Conservation Act 1999</i> Part 3 (matters of national environmental significance) .....	3–23
Table 3.3	Management tools used to address management topics in the Great Barrier Reef Region .....	3–26
Table 3.4	Current strategic priorities for the Authority .....	3–30
Table 3.5	Great Barrier Reef Marine Park Authority involvement in national planning and management frameworks .....	3–35
Table 3.6	Examples of Great Barrier Reef Marine Park Authority involvement in Queensland Government initiatives .....	3–36
Table 4.1	World heritage criteria relevant to the Great Barrier Reef.....	4–7
Table 4.2	Plants and animals of the Great Barrier Reef .....	4–14
Table 4.3	Values consistently identified in Traditional Owner and stakeholder workshops, 2012 .....	4–22
Table 4.4	Listed migratory species in the Great Barrier Reef Region .....	4–25
Table 4.5	Estimated population of listed migratory inshore and coastal foraging seabirds known to breed in the Great Barrier Reef Region .....	4–31
Table 4.6	Estimated population of listed migratory offshore and pelagic-foraging seabirds known to breed in the Region .....	4–32
Table 4.7	Listed threatened species in the Great Barrier Reef Region.....	4–33
Table 4.8	Key values and attributes of matters of national environmental significance .....	4–43
Table 4.9	Key environmental processes relevant to matters of national environmental significance.....	4–46
Table 5.1	Population, housing and income within natural resource management areas of the Great Barrier Reef catchment, 2011 .....	5–11
Table 5.2	Trading ports located within, or partly within, the Great Barrier Reef World Heritage Area, Region and Marine Park .....	5–22
Table 5.3	Commercial fisheries of the Great Barrier Reef Region, 2011 .....	5–33
Table 6.1	Impacts considered in the assessment.....	6–5
Table 6.2	Impacts and their sources.....	6–8
Table 6.3	Past activities resulting in direct legacy impacts on the Region's values .....	6–9
Table 6.4	Marine species captured in the Queensland Shark Control Program, 1993–2003 .....	6–36
Table 6.5	Large oils spills that have affected the values of the Great Barrier Reef .....	6–43
Table 6.6	Past and present effects of impacts on Great Barrier Reef habitats .....	6–47
Table 6.7	Past and present effects of impacts on Great Barrier Reef species and groups of species.....	6–48
Table 6.8	Past and present effects of impacts on Great Barrier Reef geomorphological features.....	6–50
Table 6.9	Past and present effect of impacts on Indigenous and historic heritage values.....	6–53
Table 6.10	Summary of impacts affecting outstanding universal value .....	6–68
Table 6.11	Summary of the past and present effects of impacts on the Region's values .....	6–75
Table 7.1	Current condition and trend of biodiversity values — Great Barrier Reef habitats .....	7–5
Table 7.2	Current condition and trend of biodiversity values — terrestrial habitats that support the Great Barrier Reef .....	7–8
Table 7.3	Current condition and trend of biodiversity values — species .....	7–10
Table 7.4	Summary of condition and trend for marine turtle species in the Region.....	7–17
Table 7.5	Current condition and trend of geomorphological features .....	7–20
Table 7.6	Current condition and trend of heritage values — Indigenous .....	7–22
Table 7.7	Current condition and trend of heritage values — historic heritage .....	7–23
Table 7.8	Current condition and trend of community benefits of the environment .....	7–26
Table 7.9	Economic contributions of Reef-related activities to Australia, 2012.....	7–28
Table 7.10	Current condition and trend of environmental processes.....	7–30
Table 7.11	Benchmarking the outstanding universal value of the Great Barrier Reef World Heritage Area ....	7–34
Table 7.12	Benchmarking the integrity of the Great Barrier Reef World Heritage Area.....	7–40
Table 7.13	Current condition and trend of the Great Barrier Reef national heritage place .....	7–41
Table 7.14	Current condition of key values .....	7–47
Table 8.1	Management tools considered in assessing management effectiveness .....	8–5
Table 8.2	A comparison of the scale and complexity of management topics assessed .....	8–7
Table 8.3	Assessment criteria to assess effectiveness of management.....	8–8
Table 8.4	Management effectiveness scores for biodiversity protection.....	8–10
Table 8.5	Management effectiveness scores for Indigenous heritage values.....	8–11
Table 8.6	Management effectiveness scores for historic heritage values .....	8–13
Table 8.7	Management effectiveness scores for community benefits .....	8–14
Table 8.8	Management effectiveness scores for climate change and extreme weather.....	8–15
Table 8.9	Management effectiveness scores for water quality protection (catchment run-off) .....	8–17
Table 8.10	Management effectiveness scores for coastal development (protection of coastal ecosystems) ...	8–18
Table 8.11	Management effectiveness scores for tourism .....	8–19
Table 8.12	Management effectiveness scores for fishing — commercial.....	8–21
Table 8.13	Management effectiveness scores for fishing — recreational.....	8–22
Table 8.14	Management effectiveness scores for recreation .....	8–23

Table 8.15	Management effectiveness scores for ports .....	8–24
Table 8.16	Management effectiveness scores for shipping .....	8–26
Table 8.17	Management effectiveness scores for defence activities .....	8–27
Table 8.18	Management effectiveness scores for research activities .....	8–28
Table 8.19	Effectiveness of the Authority's management arrangements to understand impacts .....	8–30
Table 8.20	Effectiveness of the Authority's management arrangements to consider issues .....	8–31
Table 8.21	Effectiveness of the Authority's management arrangements to manage impacts .....	8–32
Table 8.22	Effectiveness of the Authority's management arrangements to provide certainty .....	8–33
Table 8.23	Summary of management certainty provided for activities that occur in the Region .....	8–33
Table 8.24	Effectiveness of the Authority's management arrangements to halt and reverse declines .....	8–44
Table 8.25	Effectiveness of the Authority's management arrangements to adapt to climate change .....	8–45
Table 8.26	Effectiveness of the Authority's management arrangements to integrate with government programs .....	8–46
Table 8.27	Effectiveness of the Authority's management arrangements to meet Australia's international obligations .....	8–47
Table 8.28	Effectiveness of the Authority's management arrangements to monitor, evaluate and report .....	8–47
Table 8.29	Effectiveness of the Authority's management arrangements to identify values and determine their condition and trend .....	8–49
Table 8.30	Effectiveness of the Authority's management arrangements to monitor, evaluate and report on condition and trend .....	8–50
Table 8.31	Effectiveness of the Authority's management arrangements to protect each matter of national environmental significance .....	8–51
Table 8.32	Summary of management effectiveness results for management topics .....	8–56
Table 9.1	Meeting the selection criteria for demonstration case studies .....	9–4
Table 9.2	Matching demonstration case studies to management topics .....	9–5
Table 9.3	Matching demonstration case studies to management tools .....	9–6
Table 9.4	Tenure of islands in the World Heritage Area .....	9–14
Table 9.5	Mechanisms for identifying and protecting water quality values .....	9–27
Table 10.1	Gratings for likelihood and consequence .....	10–8
Table 10.2	Combining likelihood and consequence for overall risk .....	10–9
Table 10.3	Projected risks to biodiversity in the Region over the next 25 years .....	10–9
Table 10.4	Projected risks to geomorphological features in the Region over the next 25 years .....	10–14
Table 10.5	Projected risks to Indigenous heritage values in the Region over the next 25 years .....	10–15
Table 10.6	Projected risks to historic heritage values in the Region over the next 25 years .....	10–19
Table 10.7	Summary of projected risks to matters of national environmental significance in the Region .....	10–20
Table 10.8	Summary of projected risks to biodiversity values over the next 25 years .....	10–22
Table 10.9	Summary of projected risks to geomorphological features over the next 25 years .....	10–23
Table 10.10	Summary of projected risks to Indigenous heritage values over the next 25 years .....	10–24
Table 10.11	Summary of projected risks to historic heritage values over the next 25 years .....	10–25
Table 11.1	Projected condition of biodiversity values .....	11–7
Table 11.2	Projected condition of geomorphological features .....	11–11
Table 11.3	Projected condition of Indigenous and historic heritage values .....	11–12
Table 11.4	Projected condition of matters of national environmental significance .....	11–14
Table 12.1	Recommended improvements to the Authority's management arrangements .....	12–5
Table 12.2	Alignment between the recommended improvements of the Authority and the Queensland Government .....	12–12
Table 13.1	Implementation of measures to strengthen management and forward commitments .....	13–5
Table 13.2	The Authority's commitment to safeguard matters of national environmental significance .....	13–8
Table 13.3	Steps in the Authority's Permission System and EPBC Act assessment and decision-making process .....	13–10
Table 13.4	Content requirements for Permit Applications .....	13–17
Table 13.5	Comparison of the EPBC Act requirements and the Permission System .....	13–22
Table 13.6	Applying the principles of ecologically sustainable development .....	13–26
Table 13.7	Progress on the recommendations of the 2012 World Heritage Centre and International Union Conservation of Nature reactive monitoring mission .....	13–31
Table 13.8	Addressing endorsement criteria .....	13–36



## GLOSSARY OF COMMONLY USED TERMS AND ACRONYMS

**Action:** Includes a project, a development, an undertaking, an activity or series of activities, and an alteration to any of these things. (Adapted from EPBC Act)

**Adaptive management:** A systematic process for continually improving management practices through learning from the outcomes of previous management. It includes a monitoring, evaluation, reporting, and improvement cycle. (Adapted from *A guide to undertaking strategic assessments*)

**Attribute:** Those aspects which underpin and support matters of national environmental significance.

**Authority:** Great Barrier Reef Marine Park Authority.

**Avoiding impacts:** Measures taken so that actions have minimal negative effect on the environment. (Adapted from *A guide to undertaking strategic assessments*)

**Basin:** An area of land where surface water channels to a hydrological network and discharges into the sea (for example, a whole river system). Within the Great Barrier Reef catchment, 35 basins have been defined, based on the major river systems. A basin can include small creeks and streams that discharge directly to the sea.

**Benthic:** The bottom of the seafloor which includes the collection of organisms living on or in the bottom.

**Biodiversity:** The variability among living organisms from all sources (including terrestrial, marine and other aquatic ecosystems and the ecological complexes of which they are part). It includes diversity within species and between species, and diversity of ecosystems. (EPBC Act and GBRMP Act)

**Bycatch:** Any animal, plant or marine product that was unintentionally caught (that is, not targeted) during commercial, recreational or traditional fishing activities, including Queensland's Shark Control Program. Bycatch can either be retained (see by-product) or discarded.

**By-product:** Any animal, plant or marine product that was unintentionally caught but retained during a commercial, recreational or traditional fishing activity.

**Capital dredging:** Dredging for navigation, to create a new or enlarged channel, port, marina or boat harbour areas. Dredging for engineering purposes, to create trenches for pipes, cables, immersed tube tunnels, to remove material unsuitable for foundations and to remove overburden for aggregate.

**Carrying capacity:** The number of individuals an ecosystem can support without having any negative effects. It also includes a limit of resources and pollution levels that can be maintained without experiencing high levels of change.

**Coastal ecosystem:** Inshore, coastal and adjacent catchment ecosystems that connect the land and sea, and have the potential to influence the health and resilience of the Great Barrier Reef.

**Coastal reclamation:** The process of creating new land where there was ocean, wetlands, or other waterbodies by filling the area with 'land fill' or infrastructure such as groynes and jetties.

**Coastal zone:** The area of land and sea in or adjacent to the Great Barrier Reef containing Queensland waters, plus adjacent inland areas either within five kilometres of the coast or less than 10 metres above sea level (whichever is the further).

**Cultural keystone species:** The species that play special cultural roles for Indigenous and local peoples and are the ones they depend upon most extensively to meet their needs for food, clothing, shelter, fuel, medicine, and other necessities of life. These are the species that become embedded in a community's cultural traditions and narratives, their ceremonies, dances, songs, and discourse. Cultural keystone species can vary widely from one region to another and from one culture to another.

**Cumulative impact:** The impact on the environment resulting from the effects of one or more impacts, and the interactions between those impacts, added to other past, present, and reasonably foreseeable future pressures.

**Cumulative risk:** The combined risks to the environment by multiple impacts.

**Discarded catch:** See non-retained catch.

**Driver:** An overarching cause of change in the environment. (*Australia State of the Environment Report 2011*)

**Ecologically sustainable use:** Use of natural resources within their capacity to sustain natural processes while maintaining the life-support systems of nature and ensuring the benefit of their use to the present generation does not diminish the potential to meet the needs and aspirations of future generations. (EPBC Act)

**Ecosystem:** A dynamic complex of plant, animal and microorganism communities and their non-living environment interacting as a functional unit. (EPBC Act)

**Ecosystem-based management:** An integrated approach to managing an ecosystem and matters affecting that ecosystem, with the main object being to maintain ecological processes, biodiversity and functioning biological communities. (GBRMP Act)

**Ecosystem services:** Actions or attributes of ecosystems of benefit to humans, including regulation of the atmosphere, maintenance of soil fertility, food production, regulation of water flows, water filtration, pest control and waste disposal. It also includes social and cultural services, such as the opportunity for people to experience nature. (*Australia State of the Environment Report 2011*)

**Environment:** Ecosystems and their constituent parts, including people and communities; natural and physical resources; the qualities and characteristics of locations, places and areas heritage values of places; and the social, economic and cultural aspects of the above. (EPBC Act and GBRMP Act)

**Environmental Impact Statement:** An EIS is a statement of the likely impacts of an action on the environment and how they may be managed, mitigated or offset. Similar to a Public Environment Statement (PER) but broader in scope. (Part 8, Division 6, EPBC Act)

**EPBC Act:** *Environment Protection and Biodiversity Conservation Act 1999.*

**Extraction:** The removal of any animal, plant or marine product through legal commercial, recreational or traditional fishing activities including Queensland's Shark Control Program.

**Geomorphology:** Scientific study of landforms and the processes that shape them. (*Australia State of the Environment Report 2011*)

**GBRMP Act:** *Great Barrier Reef Marine Park Act 1975.*

**Great Barrier Reef catchment:** The area adjacent to the Great Barrier Reef Region which drains into the Region.

**Habitat:** The environment occupied by an organism or groups of organisms. (Adapted from EPBC Act)

**Halimeda:** Green macroalgae which is responsible for distinctive circular deposits on parts of the Great Barrier Reef.

**Heritage value:** A place's natural and cultural environment having aesthetic, historic, scientific or social significance, or other significance, for current and future generations of Australians. (EPBC Act and GBRMP Act)

**Impact:** An event or circumstance which has an effect, either positive or negative, on a value.

**Indigenous person:** A person who is a member of the Aboriginal race of Australia; or a descendant of an Indigenous inhabitant of the Torres Strait Islands. (GBRMP Act)

**Indigenous:** For the purposes of the strategic assessment, the term 'Indigenous' should be read to apply specifically to Traditional Owners and Traditional Owner groups.

**Indirect impact:** An impact that is not the direct result of a particular action but has been made possible by that action. These include downstream or upstream impacts, as well as facilitated or consequential impacts resulting from further actions (including actions by third parties). (Adapted from *A guide to undertaking strategic assessments*)

**Inshore:** Enclosed coastal and open coastal water bodies which extend from the mean low water mark out to approximately 20 kilometres, but also includes areas further offshore that are habitats for recognised inshore specialist species.

**Integrity:** A measure of the wholeness and intactness of the natural and/or cultural heritage and its attributes. (*Operational guidelines for the implementation of the World Heritage Convention* paragraphs no. 88–95)

**Listed migratory species:** A migratory species that is native or included under a relevant international convention, which has been placed by the Environment Minister on the published list of migratory species. (Adapted from EPBC Act)

**Listed threatened species:** A native species which is extinct, extinct in the wild, critically endangered, endangered, vulnerable or conservation dependent, as set out in the published list of threatened species established by the Environment Minister. (Adapted from EPBC Act)

**Maintenance dredging:** Dredging to ensure that previously dredged channels, berths or construction works are maintained at their designated dimensions.

**Marine Park:** Great Barrier Reef Marine Park.

**Matters of national environmental significance:** Those matters defined in the *Environment Protection and Biodiversity Conservation Act 1999*.

**Mesphotoc reefs:** Corals found at water depths where light penetration is low.

**Mitigating impacts:** Measures put in place to reduce the level of impact arising from an action, including indirect and cumulative impacts. (Adapted from *A guide to undertaking strategic assessments*)

**Morphology:** The form and structure of animals and plants, without regard to their functions.

**Net benefit:** Having an overall positive impact on the environment of the Great Barrier Reef.

**Non-retained (or discarded) catch:** Marine life caught by commercial, recreational or traditional fishers which has been discarded. This includes targeted species discarded due to size or catch restrictions, low market value, 'catch and release' practices, or bycatch that has been unintentionally captured.

**Outstanding universal value:** Cultural and/or natural heritage which is so exceptional as to transcend national boundaries and to be of such significance to humanity as a whole to make it worthy of special protection. (Adapted from *Operational guidelines for the implementation of the World Heritage Convention*)

**Offshore:** Offshore water bodies extend from approximately 20 kilometres out to the edge of the Great Barrier Reef Marine Park boundary.

**Offsetting impacts:** Measures intended to compensate for the residual adverse impacts of an action on the environment. (Adapted from *Environment Protection and Biodiversity Conservation Act 1999, Environmental Offsets Policy*, 2012)

**Palaeochannel:** An ancient stream or riverbed, cut into the rock or soil and overlaid by sediment after the stream has changed its course or dried up.

**Pelagic:** Relating to or living in or on oceanic waters. The pelagic zone of the ocean begins at the low tide mark and includes the entire oceanic water column and living organisms that inhabit this zone for all or part of their life (for example, plankton, pelagic fish).

**Poaching:** The illegal take of any animal, plant or marine product from land that is not one's own or is under official protection. It also includes the illegal take of a protected species.

**Protected species:** A species that is a cetacean; a listed marine species, a listed migratory species, a listed threatened ecological community, or a listed threatened species; a species of marine mammal, bird or reptile that is prescribed as endangered wildlife, vulnerable wildlife or rare wildlife under the *Nature Conservation Act 1992* (Qld); a species declared to be a protected species for the purposes of this definition; a species declared to be a strictly protected species for the purposes of this definition. (GBRMP Act)

**Precautionary principle:** The principle that lack of full scientific certainty should not be used as a reason for postponing a measure to prevent degradation of the environment where there are threats of serious or irreversible environmental damage. (EPBC Act and GBRMP Act)

**Pressure:** An activity or group of activities that cause an impact on a value.

**Program:** The Authority's management arrangements, including future commitments, as described in the Program Report.

**Public Environment Report:** A PER is a statement of the likely impacts on the environment and how they may be managed, mitigated or offset. Similar to an Environmental Impact Statement (EIS) but narrower in scope. (Part 8, Division 5, EPBC Act)

**Recreation:** An independent visit for enjoyment that is not part of a commercial operation. It is distinct from tourism where a visitor pays to use a commercial operation. (*Recreation Management Strategy for the Great Barrier Reef Marine Park*)

**Refugia:** An area where an organism can survive during a period of unfavourable conditions.

**Resilience:** The capacity of an ecosystem to recover from disturbance or withstand ongoing pressures.

**Region:** Great Barrier Reef Region as defined in the *Great Barrier Reef Marine Park Act 1975*.

**Retained catch:** Marine life caught and kept by commercial, recreational or traditional fishers including targeted and non-targeted species.

**Risk:** The possibility of something happening that impacts on objectives. It is the chance to either make a gain or a loss and is measured in terms of likelihood and consequence. (*Australian Standard for Risk Assessment* (AS/NZS ISO 31000:2009))

**Ship:** Vessels greater than 50 metres in overall length or carrying specialised product regardless of length (for example, oil tankers, chemical or liquefied gas carriers).

**SEWPaC:** The former Commonwealth Department of Sustainability, Environment, Water, Population and Communities, now the Department of the Environment.

**State Development Area:** Defined areas established to promote economic development for industry, infrastructure corridors and major public works.

**Take:** The act of or attempt to remove, gather, catch, kill, destroy, dredge for, raise, carry away, bring ashore, interfere with and obtain (by other means) an animal, plant or marine product.

**Taxa:** Groups of one or more populations of organisms.

**Threshold:** The breaking point above which an ecosystem or a component of an ecosystem can no longer sustain natural processes and remain in a healthy condition (for example, the point at which there is a phase shift from coral-dominated or algal-dominated reefs).

**Throughput:** The quantity of cargo that is passed through a port in a given period (exports and imports).

**Tourism:** Commercial activities that provide transport, accommodation or services to people who are visiting the Region principally for enjoyment. (Derived from GBRMP Act)

**Trading port:** A trading port refers to a port that has an associated pilotage area as defined in schedule 5 of the Transport Operations (Marine Safety) Regulation 2004. In these areas, the regional harbour master has the authority to direct the master of a ship to navigate or operate a ship in a prescribed way.

**Traditional Owner:** An Indigenous person recognised in the Indigenous community or by a relevant representative Aboriginal or Torres Strait Islander body as having spiritual or cultural affiliations with a site or area in the Marine Park, or as holding native title in relation to that site or area; and who is entitled to undertake activities under Aboriginal or Torres Strait Islander custom or tradition in that site or area.

**Traditional Owner group:** The group of Traditional Owners who, in accordance with Aboriginal or Torres Strait Islander custom, speak for a site or area.

**Trophic:** Of or relating to nutrition.

**Trigger value:** A point which, if exceeded, would mean there was a significant risk of adverse effects on an ecosystem or a component of an ecosystem. Exceeding a trigger value would 'trigger' action to address contributing impacts and/or review the trigger value.

**Turbidite:** A type of sedimentary rock composed of layered particles that grade upward from coarser to finer sizes and are thought to have originated from ancient turbidity currents in the oceans.

**Value:** Those aspects or attributes of an environment that make it of significance.

**Vulnerability:** The degree to which a system, organism or community is susceptible to, and unable to cope with, an impact.

**World Heritage Area:** Great Barrier Reef World Heritage Area.

**Wellbeing:** The combination of economic prosperity, community liveability and environmental integrity, which is determined by the quality, quantity, distribution, use and preservation of economic, human, social and natural capital. (Commonwealth of Australia (2012) Sustainability Framework (Version 0 – April 2012). Department of Sustainability, Environment, Water, Population and Communities, Canberra, Australia)

**Zones of influence:** Areas where impacts have detectable effects on values.



A vibrant underwater photograph of a coral reef. The scene is filled with various types of coral, including large, branching white corals and smaller, colorful patches of orange, purple, and blue. Numerous small, colorful fish, including yellow, orange, and blue ones, are swimming around the coral. The water is clear and blue, with sunlight filtering through from the surface, creating a bright and lively atmosphere.

## Chapter 1

## Introduction





## Table of contents

<b>1.1</b>	<b>The Great Barrier Reef .....</b>	<b>1–3</b>
1.1.1	Traditional Owner connections .....	1–3
1.1.2	International importance .....	1–3
1.1.3	National importance.....	1–6
1.1.4	Great Barrier Reef Marine Park Authority.....	1–6
1.1.5	Management approach.....	1–7
1.1.6	Evolving issues and management over time .....	1–7
<b>1.2</b>	<b>Comprehensive strategic assessment.....</b>	<b>1–8</b>
1.2.1	What is a strategic assessment? .....	1–8
1.2.2	Two complementary strategic assessments.....	1–8
1.2.3	Why undertake a strategic assessment?.....	1–9
1.2.4	Scope of the Great Barrier Reef Region strategic assessment .....	1–10
1.2.5	Matters of national environmental significance, including outstanding universal value.....	1–11
1.2.6	Strategic assessment process.....	1–12
1.2.7	Implications for the future .....	1–14
	<b>References .....</b>	<b>1–14</b>

**Cover page image:** Underwater reefscape of hard and soft corals and reef fish.

---

The Great Barrier Reef (the Reef) is an Australian icon and one of the most precious ecosystems on Earth. It is a world heritage property, recognised internationally for its outstanding universal value. Containing a maze of reefs and islands, it stretches more than 2300 kilometres along the Queensland coast (Figure 1.1). It is the world's largest coral reef ecosystem. It is rich in biodiversity, from mangroves and seagrasses to coral reefs and open waters.

Aboriginal and Torres Strait Islander peoples are the Traditional Owners of the Great Barrier Reef area and have a continuing connection to their land and sea country.

The Great Barrier Reef is critical to the cultural, economic and social wellbeing of the more than one million people who live in its catchment, and is valued by the national and international community. It is a marine protected area, conserving the Reef's environment and supporting a wide range of activities, including tourism, fishing, recreation, traditional use, research, defence, shipping and ports. The Reef's environment helps bring billions of dollars to Australia's economy each year and supports almost 70,000 jobs.<sup>1</sup>

## 1.1 The Great Barrier Reef

### 1.1.1 Traditional Owner connections

Traditional Owners have inherent rights and interests over their sea country. There are about 70 Aboriginal and Torres Strait Islander Traditional Owner clan groups whose customary estates include land and sea country within the Great Barrier Reef. The Great Barrier Reef Marine Park Authority (the Authority) acknowledges the Reef's Traditional Owners, past and present, and their unique and continuing association with the area.

For tens of thousands of years, much of what is now the Great Barrier Reef Region (the Region) was above sea level and occupied by past generations of Traditional Owners. Rising sea levels subsequently inundated the area and, as a result, many coastal, island and marine environments within and adjacent to the Region have only ever existed in the presence of Aboriginal and Torres Strait Islander peoples. At the same time, their continuous culture means Aboriginal and Torres Strait Islander connections to the marine environment retain the concept that the area was once a terrestrial landscape.

Indigenous heritage is dynamic and integrates nature, heritage and culture. Traditional Owners express their cultural heritage through their relationships with country, people, beliefs, knowledge, lore, language, symbols, ways of living, sea, land and objects — all of which arise from Indigenous spirituality. Many traditional cultural practices include plants, animals and the environment, making nature inseparable from cultural identity.

*“The sea, its natural resources and our identity as Traditional Owners, are inseparable... Our ancestors have hunted and fished in this sea country since time immemorial...”<sup>2</sup>*

Through the Convention on Biological Diversity<sup>3</sup>, the Australian Government has made a commitment to respect, preserve and maintain the knowledge, innovations and practices of Indigenous communities (Article 8(j)) and to protect and encourage customary use of biological resources in accordance with traditional cultural practices that are compatible with conservation or sustainable use requirements (Article 10(c)).

In all of its management arrangements, the Authority recognises the operation of section 211 of the *Native Title Act 1993* and the continued existence of native title rights and interests, and provides for management of traditional use of marine resources in accordance with Traditional Owner customs and traditions.

### 1.1.2 International importance

On 26 October 1981, the Great Barrier Reef was inscribed on the World Heritage List of the Convention Concerning the Protection of the World Cultural and Natural Heritage (the Convention)<sup>4</sup>, on the basis of its natural outstanding universal value (Figure 1.2).

The Great Barrier Reef was the first coral reef ecosystem in the world to be made a World Heritage Area, recognised for its natural beauty and natural phenomena; its representation of major stages in the Earth's evolutionary history, including man's interaction with the environment; its ecological and biological processes; and its habitats for the conservation of biodiversity (Figure 1.2). The Great Barrier Reef is one of only a small number of marine world heritage properties that have been inscribed by meeting all four 'natural' criteria.



Figure 1.1 Great Barrier Reef Region, World Heritage Area and Marine Park



# OUTSTANDING UNIVERSAL VALUE

The term outstanding universal value lies at the heart of the World Heritage Convention. It means a place is so exceptional that it transcends national boundaries and is of common importance for present and future generations. Places such as the Great Barrier Reef are on the World Heritage List because of their outstanding universal value, as judged against a standard set of criteria. They must also have integrity, a measure of wholeness and intactness, and an adequate system of protection and management.

## GLOBAL RECOGNITION




Figure 1.2 Outstanding universal value of the Great Barrier Reef

World heritage recognition highlights the continuing international significance of the Great Barrier Reef. When considering the Reef's nomination for world heritage listing, the then International Union for the Conservation of Nature and Natural Resources (IUCN) stated:

*'... if only one coral reef site in the world were to be chosen for the World Heritage List, the Great Barrier Reef is the site to be chosen.'*<sup>5</sup>

Articles 4 and 5 of the Convention set out the obligation and responsibility to protect, conserve and rehabilitate the Reef's values for all future generations and to present those values to the world.<sup>4</sup>

Notably, a criterion at the time of listing referred to the *'interaction between man and his natural environment'*. In the context of the Great Barrier Reef, this is recognition of the strong and continuing connections between Traditional Owners and their land and sea country — an aspect reflected in the consideration of values that underpin matters of national environmental significance in this assessment.

### 1.1.3 National importance

The Great Barrier Reef ecosystem has long been recognised for its remarkable biodiversity — from coastal estuarine systems, inshore fringing reefs and seagrass meadows to islands and spectacular outer barrier reefs. The Reef is an integral part of the Australian national identity. It is a subject of national pride, reflected in both the number of people who visit it and in the continuing public interest shown in its future health and management.

Early concerns for the Reef's future culminated in the Australian Government introducing the *Great Barrier Reef Marine Park Act 1975* (the Act). Today the main object of the Act is:

*'To provide for the long term protection and conservation of the environment, biodiversity and heritage values of the Great Barrier Reef Region.'*

The Act established the Authority and defined the Great Barrier Reef Region. It also enabled subsequent declaration of the Great Barrier Reef Marine Park (the Marine Park), introducing the concept of a multiple-use marine park where 'reasonable use' of natural resources could coexist with conservation.

Over subsequent years, sections of the Region were progressively declared as Marine Park. Today the Marine Park is one of the world's largest marine protected areas, covering approximately 344,000 square kilometres (Figure 1.1). It is complemented by the Great Barrier Reef Coast Marine Park, established in adjacent state waters under Queensland Government legislation. Both marine parks form part of the Great Barrier Reef World Heritage Area.


The Great Barrier Reef is also formally listed as a national heritage place under Australia's national environment legislation, the *Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act). Its premier importance to the Australian tourism industry is recognised through its inclusion as one of the nation's 16 National Landscapes, along with places such as the Red Centre, Kakadu and Sydney Harbour.

### 1.1.4 Great Barrier Reef Marine Park Authority

The Great Barrier Reef Marine Park Authority is an Australian Government statutory agency, established under the Great Barrier Reef Marine Park Act. The Authority reports to the Australian Government Minister responsible for the environment and advises the Minister on the control, care and development of the Marine Park.

In managing the Marine Park, the Authority must have regard to, and seek to act in a way that is consistent with, the objects of the Act, the protection of the world heritage values of the Great Barrier Reef World Heritage Area, and the principles of ecologically sustainable use.

The Authority is made up of a five person board of management, supported by a staff of about 200 people. It is based in Townsville, North Queensland, and maintains offices in Cairns, Mackay and Rockhampton along the Great Barrier Reef coast, plus a small office in Canberra.



*The Authority's Annual Report, provides an overview of its activities and is tabled in each House of the Parliament every financial year*



### 1.1.5 Management approach

The Great Barrier Reef Region is a multiple-use area, where biodiversity and heritage values are protected as well as the social and economic aspects of the environment.

Managing the Region is complex and involves a responsibility to maintain the area's natural and cultural integrity while allowing sustainable use. It is a challenging task because of the size and diversity of the Reef ecosystem; its economic importance; local, state, national and international interests; and jurisdictional, biophysical and social complexities.<sup>6</sup>

The Australian and Queensland governments work in a long-term partnership to protect and manage the Region. This cooperative approach was initially formalised by the Emerald Agreement in 1979 and reaffirmed in the Great Barrier Reef Intergovernmental Agreement signed in 2009. The Great Barrier Reef Ministerial Forum, consisting of two Australian and two Queensland government ministers, facilitates and oversees implementation and achievement of the objectives of the intergovernmental agreement.

The Field Management Program is a jointly-funded, cooperative partnership between the Australian Government and the Queensland Government. It undertakes operations and routine day-to-day management activities in the Marine Park, the adjacent Queensland Great Barrier Reef Coast Marine Park and on national park islands.

Many other Australian and Queensland government agencies have responsibilities within the Region, for example in relation to fisheries, tourism, science, natural resource management and shipping. The fundamental management partnership between the Australian and Queensland governments to protect the Great Barrier Reef is complemented by partnerships with these other agencies.

The Authority works in partnership with Traditional Owners on a range of sea country programs to conserve biodiversity and Indigenous heritage values, and promote sustainable use. Management is also enhanced through partnerships and stewardship arrangements with Great Barrier Reef stakeholders, such as industry associations, scientists, local government and community groups.

Through a network of Reef Advisory Committees and Local Marine Advisory Committees, the Authority receives technical and stakeholder advice on a range of issues associated with the health, use and management of the Great Barrier Reef.

Informed by the best available science, the Authority works to ensure long-term protection of the Reef environment, recognising the needs of communities and industries that rely on a healthy Marine Park for their economic, cultural and social wellbeing.

The Authority uses a number of tools to protect and manage the Marine Park, including zoning plans, plans of management, permits, policies and strategies, formal agreements and site management arrangements. Various management approaches are employed including education, planning, environmental impact assessment, monitoring, stewardship programs and enforcement.

Within the Marine Park, a number of activities are strictly prohibited by legislation (such as mining and oil drilling) and other activities are carefully managed according to the principles of ecologically sustainable use.

The Authority does not have direct management responsibilities for areas or activities outside the Marine Park, except in a few specific circumstances. However, it recognises the interconnectedness of the terrestrial and marine natural systems, and the significant effects that land-based activities can have on the Great Barrier Reef ecosystem. Consequently, the Authority actively works with those government departments, industries and communities whose actions have the potential to affect the marine ecosystem.

### 1.1.6 Evolving issues and management over time

The activities and impacts on the Reef that have required management over the past 50 years have changed substantially.

The prospect of drilling for oil and the mining of reef limestone created widespread public concern in the 1960s and were largely responsible for government initiatives to proactively manage the Great Barrier Reef in the 1970s. In the early 1980s, the Authority's priorities were to address the risks arising from the absence of a planning regime, the exponential growth of tourism, the lack of basic scientific knowledge to underpin management decisions and the lack of a comprehensive management framework. Also in the 1980s, crown-of-thorns starfish outbreaks were a focus for research and monitoring activities, and localised control at high value tourism sites was supported. By the late 1990s, management activities focused on the critical issues of biodiversity conservation, water quality, coastal development, fisheries, tourism and recreation. More recently, the impacts of climate change on the Reef, linkages between terrestrial and marine systems, improvements in catchment run-off and cumulative impacts of coastal development activities on ecosystem function have become key areas of additional management focus. Key management responses to these issues are listed in Table 1.1 and described in Chapter 3.

**Table 1.1 Chronology of key management responses to emerging issues from the 1970s to the present**

Since proclamation of the Great Barrier Reef Marine Park Act in 1975, the Authority has continually adapted its management arrangements to address the highest risks. *Adapted from the State Party Report on the state of conservation of the Great Barrier Reef World Heritage Area (Australia).*<sup>7</sup>

1970s	1980s	1990s	2000s	Present
<ul style="list-style-type: none"> <li>• <i>Great Barrier Reef Marine Park Act 1975</i></li> <li>• Great Barrier Reef Marine Park Authority established</li> <li>• Intergovernmental Agreement — Emerald Agreement</li> </ul>	<ul style="list-style-type: none"> <li>• Inscribed on World Heritage List</li> <li>• Marine Park sections proclaimed and zoning plans developed</li> <li>• Joint field management arrangements with Queensland Government established</li> <li>• Research and monitoring programs initiated</li> </ul>	<ul style="list-style-type: none"> <li>• <i>25-Year Strategic Plan</i></li> <li>• Plans of management for Cairns Area and Whitsundays</li> <li>• Compulsory pilotage</li> <li>• Dugong Protection Areas</li> <li>• Environmental management charge</li> <li>• Cooperative Research Centre</li> </ul>	<ul style="list-style-type: none"> <li>• Consolidated Great Barrier Reef Marine Park Zoning Plan (representative areas program)</li> <li>• Plans of management for Hinchinbrook and Shoalwater Bay</li> <li>• <i>Reef Water Quality Protection Plan</i></li> <li>• Aquaculture Regulations</li> <li>• Traditional Use of Marine Resources Agreements and sea country partnerships</li> <li>• Best practices and stewardship</li> <li>• Reef Guardian program</li> <li>• <i>Great Barrier Reef Climate Change Vulnerability Assessment and Action Plan</i></li> <li>• <i>Outlook Report 2009</i></li> <li>• Great Barrier Reef Intergovernmental Agreement</li> <li>• Research partnerships</li> </ul>	<ul style="list-style-type: none"> <li>• <i>Climate Change Adaptation Strategy</i></li> <li>• <i>Informing the outlook for Great Barrier Reef coastal ecosystems</i></li> <li>• <i>Biodiversity Conservation Strategy and vulnerability assessments</i></li> <li>• <i>Great Barrier Reef Region Strategic Assessment</i></li> </ul>

## 1.2 Comprehensive strategic assessment

### 1.2.1 What is a strategic assessment?

Under the Australian Government's national environmental law — the EPBC Act — a strategic assessment may be conducted as part of the environmental impact assessment process.

Strategic assessments differ greatly from assessments of individual projects which only look at the specific effects of activities at a local level. Instead, strategic assessments are conducted over much larger scales and timeframes, enabling consideration at the overall landscape level.<sup>8</sup>


The benefit of this approach is that it allows a broader set of issues to be investigated like the environmental, social, cultural and economic impacts of development and other activities. It also allows assessment of the direct, indirect and cumulative impacts on the environment from activities.

Importantly, this process provides the community, governments, businesses and industry with an opportunity to achieve both conservation and planning outcomes at a much larger scale than can be reached through project-by-project assessments.

### 1.2.2 Two complementary strategic assessments

The comprehensive strategic assessment of the Great Barrier Reef World Heritage Area and adjacent coastal zone is being undertaken by the Australian and Queensland governments. There are two components to the comprehensive strategic assessment — a marine component and a coastal component (Figure 1.3).

The comprehensive strategic assessment of the Great Barrier Reef analyses impacts affecting the Region, from activities on the land and on the water. The Authority is responsible for undertaking the marine component which is presented in this report. The Queensland Government is leading development of the strategic assessment of the Great Barrier Reef coastal zone.



**Figure 1.3 Components of the comprehensive strategic assessment of the Great Barrier Reef World Heritage Area and adjacent coastal zone**

There are areas of overlap and joint management responsibility across the two strategic assessments. These aspects will be considered and addressed in both assessments.

### 1.2.3 Why undertake a strategic assessment?

The Authority is undertaking this strategic assessment to evaluate and improve its effectiveness in managing existing and emerging risks to the Great Barrier Reef, focusing on the relevant matters of national environmental significance.

*The strategic assessment is part of the Australian Government's adaptive approach to managing the Great Barrier Reef.*

The findings of the strategic assessment have informed the recommendations of this report and a separate Program Report for the Great Barrier Reef Region. The Program Report is a detailed description of the Authority's management arrangements and future commitments to protect and manage matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area.

The comprehensive strategic assessment also forms part of the Australian Government's response to the World Heritage Committee's concerns regarding development impacts on the World Heritage Area, originally raised at its meeting in June 2011.

At that meeting, the World Heritage Committee requested that Australia undertake a comprehensive strategic assessment of the Great Barrier Reef World Heritage Area, identifying planned and potential developments that could impact on the outstanding universal value of the World Heritage Area, and develop a long-term plan for sustainable development. A World Heritage Centre/International Union for Conservation of Nature monitoring mission subsequently visited the area in March 2012. It concluded that the Great Barrier Reef continues to demonstrate outstanding universal value and proposed 14 recommendations.<sup>9</sup>

At its meeting in June 2012, the World Heritage Committee noted the findings of the monitoring mission and, amongst other things, requested that the recommendations in the mission report be addressed. Recommendation R5 relates to the conduct of the strategic assessment.

## World Heritage Centre/IUCN Monitoring Mission Report<sup>9</sup> — Recommendation R5

Complete the strategic assessment and resulting long-term plan for the sustainable development of the property for consideration by the World Heritage Committee at its 39<sup>th</sup> session in 2015. The assessment and long-term plan should be completed in a coordinated and fully consultative process, against a number of defined criteria for success, and considering the conclusions and recommendations of the mission as set out in this report. Expectations of the Strategic Assessment include that it will lead to:

- A long-term plan with agreed leadership at Federal and State levels, that addresses the entire property and the adjacent areas where activities can affect the outstanding universal value of the property, and ensures that any development that is approved results in an overall net-benefit for the property;
- Explicit incorporation of all elements that make up the outstanding universal value of the property, and in particular the long-term conservation of the integrity of the property, into the decision-making process regarding all development and use that may negatively impact the property, both within the boundaries of the World Heritage Area and in areas adjacent to the property;
- Improved effectiveness of the overall protection, planning and management of the outstanding universal value of the property as a whole, and the catchments, and coastal and marine areas that are intimately linked to it, including if necessary legal/statutory reforms to strengthen protection and management;
- A clear and target-driven framework to support planning and assessment of development proposals to protect outstanding universal value, and restore it where necessary, and to ensure resilience of the site, including the consideration of cumulative impacts;
- A clear analysis and related policies and strategies that will sustain long-term sustainable development, compatible with the protection of outstanding universal value, including consideration of all the economic sectors, including sustainable tourism and recreation and commercial fishing, as well as coastal development;
- Spatial policies that will identify appropriate and limited locations and standards for coastal development, and also identify areas that should not be subject to development, and which will provide greater business certainty regarding development proposals and community confidence and understanding of future development scenarios;
- Increased public confidence in their ability to engage with and influence policy and development decisions, including independent mechanisms to scrutinize and advise on the assessment of impacts of development;
- Support for new and enhanced policies and measures to regulate and manage shipping, and provide appropriate emergency planning and response;
- Appropriate systems to secure that, where development and use is permitted it will lead to net-benefits to the property as a whole, including from contributions from developers to mitigate impacts of development;
- Measures, such as legislative change to enhance compliance, that may increase the results achieved from the funding available for management, and to also increase overall levels of funding where required to provide for effective protection and management.

All management arrangements under the jurisdiction of the Authority have been considered in this strategic assessment, together with partnerships and collaborative arrangements with other Australian and Queensland government agencies, and partnerships with stakeholders and members of the Great Barrier Reef coastal community.

### 1.2.4 Scope of the Great Barrier Reef Region strategic assessment


The area considered in this strategic assessment is the Great Barrier Reef Region (Figure 1.1), as defined in the Great Barrier Reef Marine Park Act, and places outside that Region to the extent that the Region may be affected by actions in those places.

The Region extends from the tip of Cape York in the north to just past Lady Elliot Island in the south. Its western boundary is mean low water and it extends eastwards between 80 and 250 kilometres. Established in 1975, it covers 346,000 square kilometres and includes Commonwealth islands. It does not include the islands or internal waters that fall under Queensland Government jurisdiction.

There are geographically small but important differences between the boundaries of the Region, the Great Barrier Reef World Heritage Area and the Great Barrier Reef Marine Park (Table 1.2). These are illustrated spatially in Figure 1.4, using the area around the city of Cairns as an example.

At 348,000 square kilometres, the World Heritage Area is slightly larger than the Region because it includes all the islands and waters within its boundary, regardless of tenure.

Both the Region and the Marine Park include Commonwealth islands within their boundaries but do not include Queensland's islands and internal waters. In addition, there are 13 coastal areas excluded from the Marine Park. These areas were excluded to accommodate port interests, and contain trading ports and non-trading ports or marinas (see Section 5.3.5). In total, the Marine Park covers 344,400 square kilometres.



**Figure 1.4 An example of the different coastal boundaries for the Great Barrier Reef World Heritage Area, Region and Marine Park**

As illustrated in this map of the area around Cairns, the Great Barrier Reef World Heritage Area includes all internal waters and islands of Queensland (for example enclosed bays and estuaries, Green Island and Fitzroy Island). The Great Barrier Reef Region includes all waters seaward of low water, but not the internal waters and islands of Queensland (such as Green Island and Fitzroy Island). The Region includes Commonwealth islands. The Great Barrier Reef Marine Park has the same boundary as the Region, except that defined areas around most trading ports are excluded. The Marine Park boundary around ports is different from the port limit which is defined in relation to operation of the port itself.

**Table 1.2 Differences between the Great Barrier Reef World Heritage Area, Region and Marine Park**

The Great Barrier Reef Region is the subject of this strategic assessment.

Great Barrier Reef World Heritage Area	Great Barrier Reef Region	Great Barrier Reef Marine Park
348,000 km <sup>2</sup>	346,000 km <sup>2</sup>	344,400 km <sup>2</sup>
Inscribed 1981	Established 1975	Declared in sections between 1979 and 2001; amalgamated into one section in 2003
Includes: <ul style="list-style-type: none"> <li>all islands within outer boundary (about 1050)</li> <li>all waters seaward of low water mark (including internal waters of Queensland and port waters)</li> <li>all 12 trading ports</li> </ul>	Includes: <ul style="list-style-type: none"> <li>approximately 70 Commonwealth islands</li> <li>all waters seaward of low water mark (excluding Queensland internal waters)</li> </ul> Does <u>NOT</u> include: <ul style="list-style-type: none"> <li>internal waters of Queensland</li> <li>Queensland islands (about 980)</li> </ul>	Includes: <ul style="list-style-type: none"> <li>approximately 70 Commonwealth islands</li> <li>all waters seaward of low water mark (excluding Queensland internal waters)</li> </ul> Does <u>NOT</u> include: <ul style="list-style-type: none"> <li>internal waters of Queensland</li> <li>Queensland islands (about 980)</li> <li>13 coastal exclusion areas</li> </ul>

### 1.2.5 Matters of national environmental significance, including outstanding universal value

Matters of national environmental significance are Australia's national environmental assets as defined in the EPBC Act. The matters considered in this assessment are defined in the Authority's agreement with the then Minister for Sustainability, Environment, Water, Population and Communities in relation to this assessment (see Appendix 1) and the terms of reference (see Appendix 2).

This report is focused on assessing and recommending improvements to the current management arrangements to protect matters of national environmental significance relevant to the Great Barrier Reef Region.



These matters are:

- world heritage properties
- the Great Barrier Reef Marine Park
- national heritage places
- Commonwealth marine areas
- listed migratory species
- listed threatened species and ecological communities
- wetlands of international importance.

The report contains explicit consideration of matters relating to the outstanding universal value of the Great Barrier Reef World Heritage Area. Outstanding universal value is a concept central to world heritage properties (Figure 1.2). It relates to the exceptional qualities of global significance that make an area worthy of special protection and includes the concept of 'integrity' which is a measure of the wholeness or intactness of the property's natural heritage and its attributes.<sup>10</sup>

### 1.2.6 Strategic assessment process

In accordance with the EPBC Act, there is a set of legislative and non-legislative steps to be followed in undertaking a strategic assessment (Figure 1.5).

The Authority entered into an agreement with the then Minister for Sustainability, Environment, Water, Population and Communities in February 2012 (see Appendix 1). It subsequently developed terms of reference for the assessment in consultation with the then Department of Sustainability, Environment, Water, Population and Communities (see Appendix 2), and prepared drafts of the Strategic Assessment Report and the Program Report for public comment.

All responses received during the public comment period, along with advice from independent peer reviewers, have been taken into account in finalising the Strategic Assessment Report (this report) and the Program Report. These documents and a Supplementary Report explaining how public responses have been taken into account, along with the responses themselves, will be provided to the Minister for consideration.

The Minister may endorse the future management arrangements described in the Program Report if satisfied that the impacts relevant to the strategic assessment have been adequately addressed. The Minister may also recommend changes before endorsement. During these considerations, the Minister will have regard to the extent to which the endorsement criteria, as set out in the terms of reference, have been met.

#### Endorsement criteria for the strategic assessment


In determining whether or not to endorse the Program, the Minister will have regard to the extent to which the Program meets the objects of the *Environment Protection and Biodiversity Conservation Act 1999*. In particular, that it:

- a) protects the environment, especially those aspects of the environment that are matters of national environmental significance
- b) promotes ecologically sustainable development through the conservation and ecologically sustainable use of natural resources
- c) promotes the conservation of biodiversity
- d) provides for the protection and conservation of heritage
- e) promotes a cooperative approach to the protection and management of the environment
- f) assists in the cooperative implementation of Australia's international environmental responsibilities
- g) recognises the role of Indigenous people in the conservation and ecologically sustainable use of Australia's biodiversity
- h) promotes the use of Indigenous peoples' knowledge of biodiversity with the involvement of, and in co-operation with, the owners of the knowledge.

Without limiting the matters the Minister may consider when making the decision to endorse the Program, the Minister will consider the manner in which the Program:

- i) identifies direct, indirect and cumulative impacts on matters of national environmental significance
- j) avoids impacts on matters of national environmental significance
- k) mitigates the impacts on matters of national environmental significance
- l) offsets the impacts on matters of national environmental significance
- m) contributes to the enhancement of the existing environment and management of existing threats
- n) demonstrates adaption to reasonable climate change scenarios.

The Authority, the Department of the Environment and the Queensland Government have worked together throughout the process of developing this Strategic Assessment Report and the Program Report.



**Figure 1.5 Strategic assessment process**

The steps in the strategic assessment process are set out in the EPBC Act. This diagram has been adapted from *A guide to undertaking strategic assessments*.<sup>8</sup>

### 1.2.7 Implications for the future

The strategic assessment process aims to strengthen protection of matters of national environmental significance in the Region. It is intended to guide future management activities by the Authority and its partners.

This report provides the rationale for recommended improvements to management arrangements for the Region.

With a 25-year horizon, the accompanying Program Report provides the overarching strategic direction for the Authority's management of the Region. It defines the environmental outcomes required to protect the Great Barrier Reef and the management measures to achieve them.

---

## References

1. Deloitte Access Economics 2013, *Economic Contribution of the Great Barrier Reef 2011-2012*, Great Barrier Reef Marine Park Authority, Townsville.
2. Girringun Aboriginal Corporation 2005, *Girringun Region Sea Country statement; TUMRA accreditation manuscript for the Great Barrier Reef Marine Park Authority*, Girringun Aboriginal Corporation, Cardwell.
3. Secretariat of the Convention on Biological Diversity 1992, *Convention on Biological Diversity*, Secretariat of the Convention on Biological Diversity, Montreal, Canada.
4. UNESCO 1972, *Convention Concerning the Protection of the World Cultural and Natural Heritage*, UNESCO, Paris, France.
5. International Union for the Conservation of Nature 1981, *Great Barrier Reef World Heritage Nomination: IUCN Technical Report*, International Union for the Conservation of Nature, <[http://whc.unesco.org/archive/advisory\\_body\\_evaluation/154.pdf](http://whc.unesco.org/archive/advisory_body_evaluation/154.pdf)>.
6. Day, J.C. 2011, Great Barrier Reef Marine Park - Governance analysis, in *Governing Marine Protected Areas: getting the balance right, Vol 2, Technical Report*, eds P.J.S. Jones, W. Qiu and E.M. De Santo, United Nations Environment Program, Nairobi, Kenya.
7. Australian Government 2013, *State Party Report on the state of conservation of the Great Barrier Reef World Heritage Area (Australia) property ID N154 in response to World Heritage Committee decision WHC 35 COM 7B.10*, Department of Sustainability, Environment, Water, Population and Communities, Canberra.
8. Australian Government 2012, *A guide to undertaking strategic assessments*, Department of Sustainability, Environment, Water, Population and Communities, Canberra.
9. UNESCO 2012, *Mission Report Great Barrier Reef*, UNESCO, Paris, France.
10. UNESCO Intergovernmental Committee for the Protection of the World Cultural and Natural Heritage 2012, *Operational guidelines for the implementation of the World Heritage Convention*, UNESCO, Paris, France, <<http://whc.unesco.org/archive/opguide12-en.pdf>>.




A diver in a black wetsuit and mask is shown underwater, holding a clipboard and pen, positioned over a vibrant coral reef. The diver's wetsuit has "HYPERALINE" written on the sleeve. The coral reef is composed of various types of coral, including branching and table corals, in shades of yellow, orange, and pink. The water is clear and blue.

## Chapter 2

## Assessment approach





2.11.2	Traditional Owner and stakeholder input .....	2-13
2.11.3	Input into assessment of management effectiveness .....	2-13
2.11.4	Presentations and meetings.....	2-14
2.12	<b>Independent peer review of the draft Strategic Assessment Report and draft Program Report .....</b>	<b>2-14</b>
2.13	<b>Public comment .....</b>	<b>2-14</b>
2.14	<b>Information gaps .....</b>	<b>2-14</b>
	<b>References.....</b>	<b>2-15</b>

## Table of contents

2.1	<b>Legislative basis.....</b>	<b>2-3</b>
2.2	<b>Matters of national environmental significance.....</b>	<b>2-3</b>
2.3	<b>Definitions and terminology .....</b>	<b>2-4</b>
2.4	<b>Principles .....</b>	<b>2-4</b>
2.5	<b>Complexity and scale.....</b>	<b>2-4</b>
2.6	<b>Steps in the assessment process.....</b>	<b>2-4</b>
2.7	<b>Assessment methods .....</b>	<b>2-5</b>
2.7.1	Assessment logic.....	2-5
2.7.2	Structured assessments .....	2-6
2.7.3	Assessing impacts .....	2-6
2.7.4	Assessing condition and trend.....	2-8
2.7.5	Benchmarking outstanding universal value.....	2-9
2.7.6	Assessing management effectiveness.....	2-9
2.7.7	Understanding resilience .....	2-9
2.7.8	Assessing future risk.....	2-9
2.7.9	Developing projected condition .....	2-9
2.7.10	Assessment structure .....	2-10
2.8	<b>Demonstration case studies .....</b>	<b>2-10</b>
2.9	<b>Recommended changes to management .....</b>	<b>2-11</b>
2.10	<b>Information used .....</b>	<b>2-11</b>
2.11	<b>Consultation and engagement.....</b>	<b>2-12</b>
2.11.1	Authority's advisory committees .....	2-12

**Cover page image:** Diver undertaking a reef health impact survey.



**Extract from Great Barrier Reef Region Strategic Assessment terms of reference**

**7. Strategic assessment process**

**7.1 Collaboration with the Queensland Government and other Australian Government agencies**

- a) undertake the strategic assessment in consultation and collaboration with the Queensland Government, the Australian Government's Department of Sustainability, Environment, Water, Population and Communities and other relevant Australian Government agencies

**7.2 Community engagement**

- a) document how the community and stakeholders were engaged in the strategic assessment process and how views and comments were taken into account in the preparation of the Strategic Assessment Report and the Program Report

**7.4 Information and assessments**

- a) use the best available information to undertake the strategic assessment, including scientific data, expert opinion, and Traditional Owner and community knowledge
- b) document the methods used to undertake the strategic assessment
- c) for information used in the strategic assessment, indicate where possible:
  - i. the source of the information
  - ii. how recent the information is
  - iii. the reliability and limitations of the assessment.

The strategic assessments for the Great Barrier Reef Region (the Region) and coastal zone have been closely coordinated and undertaken using similar methods, where appropriate. The approaches used were aligned through the development of a joint technical framework. A range of relevant experts, including members of the Reef Water Quality Protection Plan Independent Science Panel, reviewed the joint technical framework to ensure the methods used were complementary, robust and effective. The aspects of the framework relevant to the strategic assessment for the Region are summarised in this chapter.

## 2.1 Legislative basis

The Great Barrier Reef Region strategic assessment is being carried out under Part 10 of the *Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act). It must meet the objects of that Act, which are:

- to provide for the protection of the environment, especially those aspects of the environment that are matters of national environmental significance
- to promote ecologically sustainable development through the conservation and ecologically sustainable use of natural resources
- to promote the conservation of biodiversity
- to provide for the protection and conservation of heritage
- to promote a cooperative approach to the protection and management of the environment involving governments, the community, landholders and Indigenous peoples
- to assist in the cooperative implementation of Australia's international environmental responsibilities
- to recognise the role of Indigenous people in the conservation and ecologically sustainable use of Australia's biodiversity
- to promote the use of Indigenous peoples' knowledge of biodiversity with the involvement of, and in cooperation with, the owners of the knowledge.

An agreement between the then Minister for Sustainability, Environment, Water, Population and Communities and the Great Barrier Reef Marine Park Authority (the Authority) signed on 16 February 2012 provides the basis for the assessment (see Appendix 1).

Terms of reference set out the requirements for the strategic assessment, including its contents and the development process (see Appendix 2). The Minister approved these on 30 August 2012.

## 2.2 Matters of national environmental significance

This report is focused on assessing and recommending improvements to management arrangements to protect the matters of national environmental significance, as set out in the EPBC Act, that are relevant to the Region.

Each of the matters of national environmental significance is described in detail in Chapter 4.

## 2.3 Definitions and terminology

A glossary of terms used is provided at the beginning of the report.

## 2.4 Principles

The following key principles informed development of the strategic assessment:

- establish a system-wide understanding of values and attributes, including ecosystem processes, which underpin matters of national environmental significance
- establish a system-wide understanding of pressures and impacts on matters of national environmental significance, recognising these are likely to vary at different spatial and temporal scales
- include spatial representations of values and attributes, impacts and risks wherever relevant and possible
- use demonstration cases to examine in finer detail how management systems identify, protect and manage impacts on matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area
- use information that is publicly available and, where available, information which has undergone peer review
- promote rigour and transparency through expert advice where published information is not available or sufficient
- acknowledge and report confidence, uncertainties and gaps in information
- present information in a simple and clear fashion suitable for the public.

## 2.5 Complexity and scale

The biophysical, social and jurisdictional complexity of the Region and its adjacent coastal zone, as well as the factors influencing them, means this strategic assessment is conducted on a number of scales.

At each step, a Reef-wide overview is presented and supplemented by more information on specific values, issues or areas on a range of scales. Where appropriate, and where the level of detail is available, spatial variation is described and, where possible, mapped.

Demonstration case studies are used to investigate values and impacts, and to illustrate management effectiveness in greater detail and at a finer scale.

## 2.6 Steps in the assessment process

The steps taken in this strategic assessment reflect its terms of reference. The following individual steps are presented as separate chapters in the report, with the outcomes of each chapter informing subsequent chapters:

- describing the Authority's current management arrangements (see Chapter 3)
- outlining the extent to which the matters of national environmental significance are relevant to the Region, and establishing the values or, in the case of the property's outstanding universal value the attributes, that are relevant to them (see Chapter 4)
- describing the drivers and activities relevant to the Region (see Chapter 5)
- describing and assessing the past and present impacts affecting the values and attributes, and summarising the implications for the matters of national environmental significance (see Chapter 6)
- assessing the current condition and trend of the values and attributes, and summarising the implications for the matters of national environmental significance, including benchmarking the outstanding universal value of the World Heritage Area (see Chapter 7)
- presenting an independent assessment of the effectiveness of the Authority's current management arrangements (see Chapter 8)
- illustrating in finer detail current management effectiveness through selected demonstration case studies (see Chapter 9)
- providing an understanding of ecosystem resilience, assessing the future risks to the values, and summarising the implications for the matters of national environmental significance (see Chapter 10)
- projecting the future condition for matters of national environmental significance (see Chapter 11)
- recommending changes to improve the effectiveness of management arrangements (see Chapter 12)
- describing how proposed future management arrangements will support adaptive management of matters of national environmental significance (see Chapter 13).

These steps are illustrated in Figure 2.1.

The outcomes of the strategic assessment are reflected in Chapter 12 and in the accompanying Program Report which sets out the Authority's 25-year program for the protection and management of the Great Barrier Reef.



Figure 2.1 Steps in the assessment process

## 2.7 Assessment methods

### 2.7.1 Assessment logic

The strategic assessment is based on the assessment logic encompassed in the DPSIR framework (**D**rivers, **P**ressures, **S**tate, **I**mpact, **R**esponse). This type of logic is widely adapted and applied in managing and reporting on the state of the environment (for example the *Australia State of the Environment Report 2011*<sup>1</sup>). The typical terms used in the DPSIR framework are adapted in this assessment to remain consistent with those in the terms of reference.

A summary diagram of the logic adopted in the strategic assessment is provided in Figure 2.2.




Figure 2.2 Assessment logic for the strategic assessment of the Great Barrier Reef Region

Based on the widely recognised Drivers, Pressures, State, Impact, Response (DPSIR) framework, the assessment examines the drivers and activities causing impacts or pressures that affect the state or condition of both the ecological and human systems of the Region. The benefits the community derives from the Region depends on the state of those systems. The management responses of the Authority and others can be directed at each component of the framework.

## 2.7.2 Structured assessments

Based on the terms of reference, this report provides a comprehensive assessment of drivers, activities and impacts that affect the Region's values, including relevant matters of national environmental significance of the Region, and their current and future management.

The assessments are based on **key values and attributes** of the matters of national environmental significance, as established in Chapter 4. They have been grouped into four broad categories: biodiversity, geomorphological features, Indigenous and historic values, and community benefits derived from the environment. The **environmental processes** critical to the functioning of the matters of national environmental significance are also considered. The key values and attributes were determined using a number of sources, primarily:

- *Great Barrier Reef Outlook Report 2009*<sup>2</sup>
- *Informing the outlook for Great Barrier Reef coastal ecosystems*<sup>3</sup>
- *Great Barrier Reef Biodiversity Conservation Strategy 2013*<sup>4</sup>
- *Great Barrier Reef Heritage Strategy 2005*<sup>5</sup>
- *Geological and geomorphological features of outstanding universal value in the Great Barrier Reef World Heritage Area*<sup>6</sup>
- Traditional Owner and stakeholder input, including through the Authority's Reef Advisory Committees and Local Marine Advisory Committees, and targeted workshops and meetings (Section 2.11).

An analysis of drivers and activities (see Chapter 5) is used to establish a set of **past and present impacts** and **future risks**. They are based on those outlined in the Outlook Report 2009 with some additions and amendments to reflect advances in understanding and the different focus of the strategic assessment.

**Structured assessments** applying standard criteria are undertaken on the following topics:

- effects of past and present impacts on the key values and attributes of matters of national environmental significance (see Chapter 6)
- current condition and trend of key values and attributes of the matters of national environmental significance (see Chapter 7)
- benchmarking outstanding universal value compared to its condition at the time of world heritage listing (see Chapter 7)
- effectiveness of current management (see Chapter 8)
- likely future risks to the Region's values including relevant matters of national environmental significance (see Chapter 10)
- projected condition of key values and attributes of the matters of national environmental significance (see Chapter 11).

## 2.7.3 Assessing impacts

The DPSIR framework is based on the cause-and-effect chain that links drivers, activities and impacts to the state of the values relevant to the matters of national environmental significance, and the benefits derived from the environment (Figure 2.2). Assessing many impacts in large, interconnected systems such as the Great Barrier Reef is complex. Any given activity may lead to multiple impacts. Similarly, any given impact may be driven by multiple activities and affect many matters of national environmental significance.

A progression of best practice approaches is used to assess past and present impacts on matters of national environmental significance, from simple lists to process models. How well each of these tools is likely to account for the causal links implied in the DPSIR framework is summarised in Table 2.1.

Previously, the Authority has undertaken assessments using some of the tools listed in Table 2.1, namely structured lists, value-impact matrices and influence diagrams (for example, in the *Outlook Report 2009*<sup>2</sup>), and conceptual diagrams (for example, in *Informing the outlook for Great Barrier Reef coastal ecosystems*<sup>3</sup>). In this assessment, most of the tools in the hierarchy are employed to progressively refine understanding of impacts and identify those of most concern.

**Structured lists** are used in Chapter 4 to identify key values and attributes of matters of national environmental significance (see Tables 4.8 and 4.9) and again in Chapter 6 to define the scope of the impacts to be considered and to connect the identified impacts to direct drivers and activities (see Table 6.2).






Based on a detailed analysis of each impact, **matrices** are used to assess the past and current effect of each impact on key values and attributes of biodiversity, geomorphological features, and Indigenous and historic heritage values (see Tables 6.6 to 6.9). The resulting impacts on community benefits are also described (see Section 6.7). While the matrices present a more complete understanding of the range of impacts and the effects of each impact acting on an individual value, they do not allow consideration of complex interactions and cumulative impacts.



**Conceptual diagrams** and influence diagrams are used to map relationships between different impacts, values and processes. Examples are presented throughout the report (see Figures 4.14 and 4.15). These types of diagrams are also employed during the process of building qualitative process models, the tenth tool in the hierarchy shown in Table 2.1.

**Table 2.1 Tools to improve understanding of cause-and-effect relationships**

The table was adapted from Table 2 of Hayes et al. (2012)<sup>9</sup>. Reproduced by permission of CSIRO © 2012

	Complexity of cause-effect relationship				
	None <sup>1</sup>	Simple <sup>2</sup>	Directed <sup>3</sup>	Diffuse <sup>4</sup>	Feedback <sup>5</sup>
<b>Tools</b>					
1. Unstructured list	●	●			
2. Objective-indicator matrix	●	●			
3. Structured list		●	●		
4. Value-impact matrix		●	●		
5. Conceptual diagram or cartoon		●	●		
6. Influence diagram		●	●	●	
7. Fuzzy cognitive map		●	●	●	
8. Statistical model		●	●	●	● <sup>6</sup>
9. Bayesian network			●	●	● <sup>7</sup>
10. Qualitative process model				●	●
11. Quantitative process model				●	●

<sup>1</sup> No cause-effect relationship.

<sup>2</sup> Pressure directly affects indicator variable.


<sup>3</sup> Pressure directly affects a variable that has knock-on effects to indicator variable.


<sup>4</sup> Pressure indirectly affects an indicator variable via multiple interaction pathways.


<sup>5</sup> Multiple pressures simultaneously affect complex system with feedbacks between variables.

<sup>6</sup> Explicit analysis of feedback not possible with classic statistical techniques. Incorporation of process models within statistical analyses of time series (e.g. state space modelling) can account for system feedbacks; such techniques, however, require extensive data, especially for large systems.

<sup>7</sup> With difficulty; standard Bayesian networks limited to acyclic graph structures. Dynamic Bayesian networks can account for feedbacks, but are difficult to parameterise and analyse, typically making them impractical for complex systems.

 pressure or impact

 system variable — an element of the ecological or human system or benefit derived from that system that forms part of the cause-and-effect relationship but is not measured

 indicator variable — a measurable indicator (it could be a specific ecosystem element (e.g. seagrass abundance) or benefit derived from the ecosystem (e.g. income) or a surrogate measure for the health of matters of national environmental significance)

**Qualitative models** are used in Chapter 6 to document how key impacts affect matters of national environmental significance associated with coral reefs and seagrass meadows (including dugong). These models can be used to assess the multiple drivers and activities that act simultaneously on complex ecological systems.<sup>7,8</sup> The models were developed in workshops with experts in these fields. An advantage of qualitative models is that they provide a relatively rapid and flexible means to understand system dynamics, predict cumulative impacts and consider potential management interventions. Because they can be constructed and analysed relatively quickly, they can be used to compare alternative models about how a system works.

**Bayesian networks**, based on qualitative models, are employed in Chapter 11 (see Figures 11.1 and 11.2) to model possible future scenarios for the condition of coral reefs, seagrass meadows and dugongs, given the predicted trends in some impacts. Bayesian networks are **statistical models** that represent variables within a system and their dependencies.

**Quantitative modelling** has been applied to generate a series of exposure maps of coral reefs to key impacts and of key water quality variables (see Section 6.8). These models are useful where management questions require definition of critical thresholds for limits to acceptable change to a value or attribute. Quantitative models can produce precise predictions for highly specified details of a system's biological and ecological components, processes and relationships; however, they typically require a large amount of data.

A more detailed explanation of the different types of modelling tools employed, together with a description of the modelling and spatial approaches adopted in the strategic assessment, is provided in Chapter 6 (see Section 6.8).

#### 2.7.4 Assessing condition and trend

Detailed consideration of the available evidence informs the assessment of the current condition and trend of the Region's key values and attributes including relevant matters of national environmental significance (see Chapter 7). The assessment of condition is graded relative to the best available knowledge about likely original condition. Where limited information is available, evidence of the condition of supporting ecosystems and processes was taken into consideration in assessing condition and trend. The outcomes are presented separately for key values and attributes, including more detailed analysis of the condition and trend of values and attributes of particular concern. Each assessment is comprehensively referenced.

To reflect differences within the Region, the current condition and trend of the biodiversity values and attributes are reported separately for four broad areas: northern inshore (N.I.), northern offshore (N.O.), southern inshore (S.I.) and southern offshore (S.O.) (Figure 2.3). While the boundaries are not precisely defined, the north-south dividing line is in the vicinity of Port Douglas, which marks the broad division between the more developed and less developed catchments adjacent to the Region. The inshore-offshore dividing line is generally about 20 kilometres offshore. It corresponds to enclosed coastal and open coastal water bodies described in the *Water Quality Guidelines for the Great Barrier Reef Marine Park*<sup>10</sup> but also includes areas further offshore that are habitats for recognised inshore specialist species such as dugongs.

The assessment of terrestrial habitats that support the Great Barrier Reef is also presented for four areas: northern inland (N.In.), northern coastal (N.C.), southern inland (S.In.) and southern coastal (S.C.) (Figure 2.3). Again, the north-south dividing line is around Port Douglas. The coastal areas are defined as being a maximum of five kilometres from the coastline or where land reaches the height of 10 metres Australian Height Datum, whichever is furthest from the coast. The inland area is the remainder of the Great Barrier Reef catchment.

An overview statement is also provided for each grouping of values and attributes. In assessing overall condition and trend, not all components were given equal weighting. For example, greater weight was given to habitats and species that are critical in supporting a range of other habitats and species.




Figure 2.3 Indicative broad assessment areas for the Region and its catchment

### 2.7.5 Benchmarking outstanding universal value

For world heritage values, the terms of reference require that the current condition of key indicators of outstanding universal value be benchmarked against the retrospective Statement of Outstanding Universal Value<sup>11</sup>. This is the official statement adopted by the World Heritage Committee for the property, and is the standard against which the property's state of conservation can be assessed. The statement is reproduced in full in Appendix 3.

The outstanding universal value is benchmarked by grading the current condition of the relevant attributes described in the statement, along with the trend of the value since 1981. The outcome is presented in Chapter 7.

### 2.7.6 Assessing management effectiveness

The Authority's current management arrangements for the Great Barrier Reef Region are summarised in Chapter 3.

A team of three independent assessors carried out an assessment of the effectiveness of these arrangements. The assessors have a comprehensive knowledge of protected area management, particularly in the Queensland marine environment. They reviewed documentation provided by the Authority, and consulted stakeholders through formal meetings and an online survey (Section 2.11).

The terms of reference for the assessment of management effectiveness are provided in Appendix 4. The outcomes are summarised in Chapter 8.

The assessment approach is based on the management effectiveness framework used in the Outlook Report 2009.<sup>2</sup> This framework, developed by the International Union for Conservation of Nature/World Commission on Protected Areas, was specifically designed to evaluate management of protected areas. Some amendments were made to the framework to meet the requirements of the terms of reference for the strategic assessment (see Appendix 2, Item 4.1.2) and to reflect advances in understanding and knowledge since 2009.

A similar assessment framework is used in each demonstration case study (Section 2.8), including a review of the outcomes by the independent assessors.

The outcomes of the management effectiveness assessments inform subsequent assessments of future risk (see Chapter 10) and projected condition (see Chapter 11), and recommendations to improve management (see Chapter 12 and the Program Report).

The full report titled the *Assessment of Management Effectiveness for the Strategic Assessment of the Great Barrier Reef Region* (dated 19 March 2013), and the Authority's response (the Supplementary Report) are available on the Authority's website at [www.gbrmpa.gov.au](http://www.gbrmpa.gov.au).

### 2.7.7 Understanding resilience

Assessment of the projected condition of matters of national environmental significance is informed, in part, by an understanding of ecosystem resilience — the capacity to recover from disturbance or withstand ongoing impacts. The description of ecosystem resilience considers factors such as: capacity to either resist (absorb) an impact, or to recover from that impact; capacity for recruitment; diversity; connectivity; scale and complexity (see Chapter 10).

### 2.7.8 Assessing future risk

To assess future risks to the Great Barrier Reef ecosystem posed by identified impacts, the Australian Standard for Risk Assessment (AS/NZS 31000:2009) was adopted. The assessment is based on the assessments of current trends in drivers and activities, past and present impacts, current state of the values, effectiveness of management and current ecosystem resilience.

Both the likelihood and consequence of each predicted impact are ranked on a five-point scale and an overall risk level for each threat is determined, based on a combination of these two factors.

Where relevant, an impact is individually assessed for its likely future effect on biodiversity, geomorphological features, Indigenous heritage values and historic heritage values. The flow-on implications for community benefits are outlined. The outcomes are presented in Chapter 10.

### 2.7.9 Developing projected condition

The projected condition of matters of national environmental significance (see Chapter 11) is based on:

- an understanding of drivers and activities and the past and present effects of impacts
- the evaluation of the current condition and trend of key values and attributes of the matters of national environmental significance, including consideration of integrity
- the effectiveness of current management arrangements to protect values and manage impacts, both on a broad scale and the demonstration case studies
- an understanding of ecosystem resilience
- the evaluation of overall risk, identifying the most serious future impacts
- a consideration of scenarios of the likely future projected condition of the Region, using the qualitative models outlined in Section 6.8.

The assessment of projected condition is presented for the key values and attributes and summarised for each relevant matter of national environmental significance.

## 2.7.10 Assessment structure

### Grading statements

Grading statements are used to standardise the assessments. For example, the condition of each biodiversity value or the effectiveness of each aspect of the Authority's management arrangements is graded by rating it against a set of grading statements — four grades for each assessment.

The grading statements are based on those used in the *Outlook Report 2009*<sup>2</sup> and adapted, where necessary, to align with the strategic assessment's terms of reference. Additional statements are introduced for the assessment of heritage values, based on grading statements used in the *Australia State of the Environment Report 2011*<sup>1</sup>.

The assessment of outstanding universal value is adapted from a grading system developed by the International Union for Conservation of Nature to assess natural world heritage sites.<sup>12</sup>

Based on the relevant grading statements, a grade is assigned for:

- the effect of impacts on values and attributes: very low, low, high or very high
- the current and projected condition of key values and attributes: very good, good, poor or very poor
- management effectiveness: very effective, effective, partially effective or ineffective
- future risks: low, medium, high or very high.

For each component, the grade is based on the best available information and is a 'grade of best fit'. It is a summary of all aspects and is not necessarily the appropriate grade for any individual area, species or impact. Where appropriate, specific information about individual elements of the component being assessed has been used to inform the assessment of the component as a whole (for example, knowledge of the condition of individual species of sharks and rays has informed assessment of the value 'sharks and rays').

### Trend

The approach to assessing values and impacts is further refined by including an indication of trend, similar to that used in the *Australia State of the Environment Report 2011*.<sup>1</sup>

In most cases, the information about the condition of values reported in the *Outlook Report 2009*<sup>2</sup> is used as the baseline for determining trend. Therefore, the trend reported relates to the past five years, conforming with the *Outlook Report* cycle. An indication of trend over this time period helps inform consideration of changes in the effectiveness of management arrangements since 2009.

In assessing outstanding universal value, a longer timeframe is adopted to enable this to be benchmarked with the world heritage property's condition at the time of listing in 1981.

Where relevant, and where the information is available, longer term trends are also described throughout the report.

There are four categories for trend: improving, stable, deteriorating or no clear trend. The category of 'no clear trend' is applied to a value when the available information is too variable to be able to establish a trend.

### Confidence

In line with the *Australia State of the Environment Report*<sup>1</sup>, the level of confidence in each assessment is rated as either:

- adequate high quality evidence and high level of consensus
- limited evidence or limited consensus
- very limited evidence, assessment based on anecdotal information.

Where there is limited publicly available information, it is often possible to provide an assessment based on knowledge from within managing agencies, and that of Traditional Owners, topic experts and informed stakeholders. When using this type of information, a confidence grade is given of 'very limited evidence, assessment based on anecdotal information'.

## 2.8 Demonstration case studies

Eight demonstration case studies are used to assess, in finer detail, the effectiveness of current management arrangements to protect and manage the Region's values and to guide improvements.

The case studies were chosen in consultation with the Queensland Government and the then Department of Sustainability, Environment, Water, Population and Communities, guided by criteria in the terms of reference (see Appendix 2, Item 4.2.2). Given the joint management arrangements between the Authority and the Queensland Government, some of the case studies were prepared together for both strategic assessment reports.



The case studies are:

- dugongs
- corals
- islands
- Princess Charlotte Bay
- Cairns Planning Area
- Mackay Whitsunday — water quality improvement
- Keppel Bay
- East Coast Trawl Fishery.

The case studies and the rationale for their selection are provided in Chapter 9.

## 2.9 Recommended changes to management

In accordance with the terms of reference (see Appendix 2, Item 6.1), this report includes recommended changes to the Authority's current management arrangements to improve its effectiveness to deliver on its objectives. Improvements to related local, state and national government programs are also recommended where relevant (see Chapter 12).

The outcomes of the assessment of current management effectiveness (see Chapter 8), the demonstration case studies (see Chapter 9) and the projections of future condition (see Chapter 11) helped inform these recommendations.

## 2.10 Information used

The strategic assessment is based on the best available information as at June 2013, including scientific data, expert opinion, and Traditional Owner and stakeholder knowledge.

All references are cited in the text and listed at the end of each chapter. The web addresses provided were correct at the time of printing.

In addition to peer-reviewed literature, much of the information and synthesis is drawn from the *Outlook Report 2009*, as well as more detailed subsequent work. Examples include *Improving the outlook for Great Barrier Reef coastal ecosystems*<sup>3</sup>, the *Great Barrier Reef Biodiversity Conservation Strategy 2013*<sup>4</sup> and its supporting vulnerability assessments<sup>13</sup>, the *Recreation Management Strategy for the Great Barrier Reef Marine Park*<sup>14</sup>, and the *2013 Scientific Consensus Statement Update* and supporting evidence.<sup>15,16,17,18,19</sup>

Information received from stakeholders through meetings, workshops and surveys in the latter half of 2012 contributed to the assessments of the Region's values, impacts occurring on those values, and management effectiveness, including potential areas for improvement.

Information relevant to Indigenous heritage values has been sourced from literature; meetings, workshops and surveys undertaken specifically for the strategic assessment in the latter half of 2012; and direct conversations with Traditional Owners. The Authority acknowledges and adopts the principle that Traditional Owners of the Region are the primary source of information on the value of their culture and heritage, and how it is best managed and conserved. The Authority has developed long and lasting relationships with the Region's Traditional Owners, and understands the importance of identifying Traditional Owners and other Indigenous peoples with rights and interests in the area, and of talking with the Traditional Owners who have the cultural authority to speak for their sea country. In addition, Authority staff have contributed their combined knowledge and experiences to the information provided. Some staff members are Traditional Owners of the Region or are Aboriginal or Torres Strait Islander people who have strong cultural connections with the Region. They have shared aspects of their cultural knowledge for the assessment.

Eight Sustainable Regional Development Program projects, funded by the Australian Government, were undertaken during 2012–13 to address a number of important knowledge gaps for matters relevant to the strategic assessment. Where available, relevant outcomes of these projects are incorporated into the report and the reports cited. These projects are:

- *Improved dredge material management for the Great Barrier Reef Region*<sup>20</sup>: To provide improved information on which to base dredge spoil management decisions for the five major ports in the Great Barrier Reef World Heritage Area.
- *Ship anchorage management in the Great Barrier Reef World Heritage Area*<sup>21</sup>: To identify environmental impacts of existing offshore anchoring for the five major Great Barrier Reef ports and the likely future impacts of increased shipping.
- *Great Barrier Reef coastal ecosystems assessment framework*<sup>22</sup>: To examine development impacts in selected basins within the Great Barrier Reef coastal zone to assess present and future development pressures and potential offset opportunities.

- *Great Barrier Reef resilience decision framework*<sup>23</sup>: To develop a resilience framework to inform decision making in the Great Barrier Reef coastal zone.
- *Economic contribution of the Great Barrier Reef*<sup>24</sup>: To update understanding of the Great Barrier Reef's economic contribution, including analysis of commercial and non-commercial uses and detailed regional-scale analysis.
- *Geological and geomorphological features of outstanding universal value in the Great Barrier Reef World Heritage Area*<sup>6</sup>: To identify geological and geomorphological features of outstanding universal value that may not have been previously identified and provide an overview of the pressures affecting values.
- *Defining the aesthetic values of the Great Barrier Reef World Heritage Area*<sup>25</sup>: To identify and map aesthetic values and analyse the sensitivity of those values to impacts.
- *Integrated monitoring framework for the Great Barrier Reef World Heritage Area*<sup>26</sup>: To establish a framework for a standardised and integrated ecological, social and economic monitoring program to address critical information needs, align existing monitoring programs and provide a baseline for assessing the condition of values and effects of pressures, as well as the impact on those values. This project was also funded through the National Environmental Research Program — Marine Biodiversity Hub.

## 2.11 Consultation and engagement

An engagement plan provided a structured approach to consultation and engagement, based around four objectives:

- to improve decision-making by gathering a range of information, views and experiences from Traditional Owners and stakeholders as part of the strategic assessment process
- to provide a transparent framework for the development of a common understanding of the broad range of issues, and to foster informed and engaged Traditional Owners and stakeholders as part of the process
- to seek feedback from Traditional Owners and stakeholders on the potential impact of different approaches, acknowledging that there may not be consensus
- to provide a mechanism to better understand implementation issues that may need to be considered.

### 2.11.1 Authority's advisory committees

Throughout the assessment process, the Authority consulted with its established expert and community advisory groups:

- Four expertise-based Reef Advisory Committees provided advice on catchment and coastal, ecosystem, Indigenous, and tourism and recreation issues.
- Twelve community-based Local Marine Advisory Committees provided advice on management issues across the Region. There are committees for Cape York, Douglas, Cairns, Cassowary Coast, Hinchinbrook, Townsville, Bowen–Burdekin, Whitsunday, Mackay, Capricorn Coast, Gladstone and Burnett.

The Reef Advisory Committees provided advice on elements of the strategic assessment as part of their normal meetings. Since June 2011, when development of a strategic assessment was first proposed, the matter has been an item for all of the 13 Reef Advisory Committee meetings held. Topics considered included the assessment's terms of reference, cumulative impact assessment approaches, the technical assessment framework, stakeholder engagement and the demonstration case studies. Out of session, members also participated in the online survey of management effectiveness (Section 2.11.3).

In addition, 39 Reef Advisory Committee members provided further advice at the combined Reef Advisory Committee workshop on the strategic assessment in February 2013.

Similarly, members of the Local Marine Advisory Committees provided input at their regular meetings and at 'mini-workshops' (Section 2.11.2). In addition, representatives of each committee contributed at the Local Marine Advisory Committee Chairs' meeting in October 2012, and at the workshop on the strategic assessment in February 2013.

Participants at the Local Marine Advisory Committee Chairs' meeting in October 2012 also provided input on a number of the demonstration case studies, particularly on those that are regionally specific, as well as feedback on the proposed approach for benchmarking outstanding universal value.

### 2.11.2 Traditional Owner and stakeholder input

Opportunities for broader Traditional Owner and stakeholder input were provided through a series of purpose-designed workshops and follow-up surveys.

This targeted consultation achieved representation from a diverse range of interests including Traditional Owners; local government; ports, shipping and related development sectors; mining and resource sectors; research organisations; tourism operators; commercial fishers; recreational users including fishers; natural resource managers; farmers; conservation groups, and the broader community.

There was a two-stage approach to seeking Traditional Owner and stakeholder input — an initial stage of understanding their values in relation to the Region, and the pressures and potential impacts on those values; followed by a second stage investigating their views of current management effectiveness and their aspirations for future management actions.

#### Stage one

An initial series of workshops explored Traditional Owner and stakeholder views about values and impacts. A total of 290 people attended the workshops between August and October 2012.

- Workshops were held in Cairns, Townsville and Rockhampton in August 2012.
- Twelve mini-workshops were held at the regular Local Marine Advisory Committee meetings in September 2012.
- Input from Traditional Owners was gathered at specific workshops in Cairns and Rockhampton in September 2012.

After the workshops, participants were invited to answer a short online survey to gain a deeper understanding of what is important to them and why. Of the 290 workshop participants, 135 people (47 per cent) responded to the survey. Results provided an indication of the relative importance that this group of people place, on Great Barrier Reef values, its threats and their aspirations.

#### Stage two

Traditional Owners and stakeholders who participated in the stage one workshops were invited to reconvene to provide further advice on elements of the strategic assessment. Advice was sought on the key findings to date, including any discrepancies between expert opinion and on-ground community knowledge, and on future management actions which would help protect the Region's values. In addition, the independent assessors of management effectiveness provided participants with their preliminary assessment of the Authority's management effectiveness, and received feedback.

The workshops were also an opportunity to update participants on the overall progress of the strategic assessment.

A total of 105 Traditional Owners and stakeholders attended workshops in Cairns, Townsville and Rockhampton in December 2012.

#### Outcomes

The outcomes of the Traditional Owner and stakeholder engagement were considered throughout the preparation of this report and are presented in relevant sections. The comments made during public consultation on the terms of reference were also taken into account in undertaking the strategic assessment.

The *Traditional Owner and Stakeholder Engagement – Report on Workshops and Surveys* (see Appendix 5) provides a description of the process followed and the advice received.

### 2.11.3 Input into assessment of management effectiveness

Gathering the views of Traditional Owners and stakeholders (including members of the Authority's advisory committees) was an important part of the independent review of management effectiveness. This recognised their valuable knowledge and practical experience of management issues and the tools used by the Authority. The assessors received the views of Traditional Owners and stakeholders through:

- the Local Marine Advisory Committee Chairs' meeting in October 2012, including testing an early version of the online survey for stakeholders
- the stage two Traditional Owner and stakeholder workshops
- an online survey specific to management effectiveness distributed to members of Local Marine Advisory Committees and Reef Advisory Committees, as well as participants of the regional workshops.

For the online survey, a total of 95 people made 172 assessments of 15 management topics. Participants without access to the internet were able to undertake the survey over the telephone or in writing. Face-to-face interviews were undertaken with Traditional Owners to ensure their views were documented. The number of people responding to some management topics was quite low, making statistical analysis difficult. However, some general patterns of effectiveness were evident for most of the tools, and were taken into account by the assessors.

#### 2.11.4 Presentations and meetings

In addition to structured Traditional Owner and stakeholder engagement, Authority representatives met with a range of interested organisations, groups and individuals while developing the strategic assessment to inform them of the process and receive their input.

### 2.12 Independent peer review of the draft Strategic Assessment Report and draft Program Report

In accordance with the terms of reference, the draft Strategic Assessment Report was peer reviewed. Consultants Sinclair Knight Merz (SKM) were commissioned to evaluate the draft Great Barrier Reef Region Strategic Assessment Report and draft Program Report against the following criteria:

- consistency with its terms of reference
- structure and cohesiveness of presentation
- breadth and depth of matters covered
- technical accuracy
- validity of conclusions drawn.

A comparative assessment of the Authority's strategic assessment (of marine regions and issues) and the Queensland strategic assessment (of the coastal zone) was also made, to identify any gaps or duplication in the management and protection of matters of national environmental significance.

The full report titled the *Great Barrier Reef Strategic Assessment, Independent Review Report* (dated 3 February 2014), and the Authority's response (the Supplementary Report) are available on the Authority's website at [www.gbrmpa.gov.au](http://www.gbrmpa.gov.au).

### 2.13 Public comment

The Authority and the Queensland Government implemented a joint process for public consultation on the Great Barrier Reef Region and Great Barrier Reef Coastal Zone draft Strategic Assessment Report and draft Program Report.

Consultation ran from 1 November 2013 to 31 January 2014. The period was lengthened from the minimum 28-day timeframe outlined in the terms of reference to take into account public interest, the length and complexity of the reports and the Christmas holiday period.

Consultants GHD were commissioned to facilitate a number of components of the public consultation process. This included the:

- development of a standalone website that gave the public access to draft strategic assessment reports and supporting information, in conjunction with an online/postal survey for feedback and a portal for the electronic lodgement of submissions.
- management of submissions received during public consultation, including hard copy lodgement of submissions via a nominated postal address, and the distribution of copies of submissions to the Authority, the Queensland Government and the Australian Department of the Environment
- analysis of submissions and preparation of a summary report synthesising information received during the public consultation process.

A total of 6616 submissions were received during the 13-week public consultation period.

The full report titled the *Great Barrier Reef Strategic Assessment – Public Consultation on Draft Reports* (dated April 2014), and the Authority's response (the Supplementary Report) are available on the Authority's website at [www.gbrmpa.gov.au](http://www.gbrmpa.gov.au).

The outcomes of the independent assessment of management effectiveness, the independent peer review, and the public consultation, together with the Authority's response (the Supplementary Report) and the final Program Report will be provided to the Minister for the Environment for endorsement consideration.

### 2.14 Information gaps

The chapters considering drivers and activities, impacts and the current condition and trend of values conclude with an outline of key information gaps. They include the outcomes of a review of the Authority's scientific information needs, which involved consideration of existing information and policy documents that guide the Authority's priorities for research and monitoring.




## References

1. Department of Sustainability, Environment, Water, Population and Communities 2011, *State of the Environment Report*, DSEWPaC, Canberra.
2. Great Barrier Reef Marine Park Authority 2009, *Great Barrier Reef Outlook Report 2009*, GBRMPA, Townsville.
3. Great Barrier Reef Marine Park Authority 2012, *Informing the outlook for Great Barrier Reef coastal ecosystems*, GBRMPA, Townsville.
4. Great Barrier Reef Marine Park Authority 2013, *Great Barrier Reef Biodiversity Conservation Strategy*, GBRMPA, Townsville.
5. Great Barrier Reef Marine Park Authority 2005, *Great Barrier Reef Marine Park Heritage Strategy*, GBRMPA, Townsville.
6. Whiteway, T., Smithers, S., Potter, A. and Brooke, B. 2013, *Geological and geomorphological features of Outstanding Universal Value in the Great Barrier Reef World Heritage Area: Technical report prepared for the Department of Sustainability, Environment, Water, Population and Communities*, Geoscience Australia and James Cook University, Townsville.
7. Dambacher, J.M., Li, H.W. and Rossignol, P.A. 2002, Relevance of community structure in assessing indeterminacy of ecological predictions, *Ecology* 83: 1372-1385.
8. Puccia, C.J. and Levins, R. 1985, *Qualitative modelling of complex systems: an introduction to loop analysis and time averaging*, Harvard University Press, Cambridge.
9. Hayes, K.R., Dambacher, J.M., Lyne, V., Sharples, R., Rochester, W.A., Dutra, L.X.C. and Smith, R. 2012, *Ecological indicators for Australia's Exclusive Economic Zone: Rationale and approach with application to the South West Marine Region*, Department of Sustainability, Environment, Water, Population and Communities, Hobart.
10. Great Barrier Reef Marine Park Authority 2010, *Water quality guidelines for the Great Barrier Reef Marine Park*, GBRMPA, Townsville.
11. Department of Sustainability, Environment, Water, Population and Communities 2012, *Statement of Outstanding Universal Value - Great Barrier Reef World Heritage Area*, DSEWPaC, Canberra.
12. International Union for the Conservation of Nature 2012, *IUCN conservation outlook assessments - Guidelines for their application to natural world heritage sites*, International Union for the Conservation of Nature, Gland, Switzerland.
13. Great Barrier Reef Marine Park Authority 2012, *Vulnerability assessments*, GBRMPA, Townsville, <<http://www.gbrmpa.gov.au/about-the-reef/biodiversity/draft-biodiversity-conservation-strategy/vulnerability-assessments>>.
14. Great Barrier Reef Marine Park Authority 2012, *Recreation management strategy for the Great Barrier Reef Marine Park*, GBRMPA, Townsville.
15. Brodie, J., Waterhouse, J., Schaffelke, B., Johnson, J., Kroon, F., Thorburn, P., Rolfe, J., Lewis, S., Warne, M., Fabricius, K., McKenzie, L. and Devlin, M. 2013, *Reef Water Quality Scientific Consensus Statement 2013*, Department of the Premier and Cabinet, Brisbane.
16. Schaffelke, B., Anthony, K., Blake, J., Brodie, J., Collier, C., Devlin, M., Fabricius, K., Martin, K., McKenzie, L., Negri, A.P., Ronan, M., Thompson, A. and Warne, M. 2013, Marine and coastal ecosystem impacts, in *Synthesis of evidence to support the Reef Water Quality Scientific Consensus Statement 2013*, Reef Water Quality Protection Plan Secretariat, Brisbane.
17. Johnson, J.E., Maynard, J.A., Devlin, M.J., Wilkinson, S., Anthony, K.R.N., Yorkston, H., Heron, S.F., Puotinen, M.L. and van Hoodonk, R. 2013, Resilience of Great Barrier Reef marine ecosystems and drivers of change, in *Synthesis of evidence to support the Reef Water Quality Scientific Consensus Statement 2013*, Department of the Premier and Cabinet, Brisbane.
18. Kroon, F., Turner, R., Smith, R., Warne, M., Hunter, H., Bartley, R., Wilkinson, S., Lewis, S., Waters, D. and Carroll, C. 2013, Sources of sediment, nutrients, pesticides and other pollutants in the Great Barrier Reef catchment, in *Synthesis of evidence to support the Reef Water Quality Scientific Consensus Statement 2013*, Department of the Premier and Cabinet, Brisbane.
19. Thorburn, P., Rolfe, J., Wilkinson, S., Silburn, M., Blake, J., Gongora, M., Windle, J., VanderGragt, M., Wegscheid, C., Ronan, M. and Carroll, C. 2013, The water quality and economic benefits of agricultural management practices, in *Synthesis of evidence to support the Reef Water Quality Scientific Consensus Statement 2013*, Department of the Premier and Cabinet, Brisbane.
20. Sinclair Knight Merz Pty Ltd and Asia-Pacific Applied Science Associates 2013, *Improved dredge material management for the Great Barrier Reef Region*, Great Barrier Reef Marine Park Authority, Townsville.
21. GHD Pty Ltd 2013, *Ship anchorage management in the Great Barrier Reef World Heritage Area*, Great Barrier Reef Marine Park Authority, Townsville.
22. Great Barrier Reef Marine Park Authority 2012, *Great Barrier Reef Coastal Ecosystems Assessment Framework*, GBRMPA, Townsville.
23. Anthony, K., Beeden, R., Dambacher, J.M. and Walsh, T. 'A draft qualitative framework for understanding cumulative impacts on values in the Great Barrier Reef World Heritage Area'.
24. Deloitte Access Economics 2013, *Economic Contribution of the Great Barrier Reef 2011-2012*, Great Barrier Reef Marine Park Authority, Townsville.
25. Context Pty Ltd 2013, *Defining the aesthetic values of the Great Barrier Reef*, Department of Sustainability, Environment, Water, Population and Communities, Canberra.
26. Hedge, P., Molloy, F., Sweatman, H., Hayes, K., Dambacher, J., Chandler, J., Gooch, M., Chin, A., Bax, N. and Walshe, T. 2013, *An integrated monitoring framework for the Great Barrier Reef World Heritage Area*, Great Barrier Reef Marine Park Authority, Townsville.







## Chapter 3

---

### Current management





3.11.3	Regional .....	3–36
3.11.4	Local.....	3–36
3.12	<b>Protecting matters of national environmental significance.....</b>	<b>3–36</b>
3.13	<b>Timeframe and review .....</b>	<b>3–38</b>
<b>References .....</b>		<b>3–39</b>

## Table of contents

3.1	<b>Context .....</b>	<b>3–3</b>
3.2	<b>Purpose .....</b>	<b>3–4</b>
3.3	<b>Strategic assessment area .....</b>	<b>3–5</b>
3.4	<b>Integrated management.....</b>	<b>3–5</b>
3.4.1	Jurisdiction.....	3–5
3.4.2	Great Barrier Reef Intergovernmental Agreement .....	3–6
3.4.3	Great Barrier Reef Ministerial Forum..	3–7
3.5	<b>Legislation and conventions.....</b>	<b>3–7</b>
3.6	<b>Management tools.....</b>	<b>3–9</b>
3.6.1	Spatially explicit management tools..	3–14
3.6.2	System-based regulatory tools .....	3–18
3.7	<b>Management activities .....</b>	<b>3–26</b>
3.8	<b>Partners in management .....</b>	<b>3–27</b>
3.9	<b>Management priorities .....</b>	<b>3–30</b>
3.9.1	Current priorities .....	3–30
3.10	<b>Applying the tools to achieve the priorities .....</b>	<b>3–30</b>
3.10.1	Addressing the key risks.....	3–30
3.10.2	Ensuring use is ecologically sustainable .....	3–32
3.10.3	Fostering stewardship.....	3–32
3.10.4	Effective and efficient organisation...	3–33
3.10.5	Current resources for management..	3–33
3.11	<b>Involvement in other planning and management frameworks .....</b>	<b>3–35</b>
3.11.1	National.....	3–35
3.11.2	State .....	3–35

Cover page image: Vessel anchored offshore from Ellison Reef.



**Extract from Great Barrier Reef Region Strategic Assessment terms of reference**

**1.1 Provide an overview of the current Program\*. For the purposes of the strategic assessment, the life of the Program is 25 years. The overview is to include a description of:**

- a. the purpose of the Program, including Program objectives
- b. the area to which the Program applies (the strategic assessment area)
- c. legislation, plans, policies and other mechanisms that make up the Program, including Program commitments
- d. relevant activities within the scope of the Program
- e. international, national, state and regional context (environmental, social, cultural and economic) in which the Program operates, including activities outside the strategic assessment area that may influence the Program
- f. relevant national, state, regional and local planning or management frameworks that affect the Program and contribute to protection and management of the matters of national environmental significance
- g. identification of how long the Program will be in effect and the process for review of the Program, including adaptive management
- h. identification of the relevant authorities responsible for the implementation of the Program.

\* Note: In this report 'current Program' is also referred to as 'current management arrangements'.

This chapter provides an overview of the current management arrangements of the Great Barrier Reef Marine Park Authority (the Authority) in relation to the Great Barrier Reef Region (the Region).

The Queensland Government's contribution to the management of the Great Barrier Reef World Heritage Area (World Heritage Area) and the protection of matters of national environmental significance in the Great Barrier Reef coastal zone is outlined in the complementary Great Barrier Reef Coastal Zone Strategic Assessment.

### 3.1 Context

Through an intergovernmental agreement, the Australian and Queensland governments have been working together for the long-term protection and conservation of the Great Barrier Reef Marine Park since its inception in 1975.

Management of the Region relies upon a number of Australian and Queensland government agencies to regulate access and to control or mitigate impacts associated with activities. These agencies use a combination of management tools, including zoning plans, plans of management, permits, policies and the *Reef Water Quality Protection Plan* (Reef Plan). They employ various management approaches including education, planning, environmental impact assessment, monitoring, stewardship and enforcement.

In this strategic assessment, the management arrangements under the jurisdiction of the Authority are considered. They include, but are not limited to:

- statutory instruments, including Regulations, zoning plans, plans of management and permits
- non-statutory mechanisms including policies, position statements and guidelines
- partnership and collaborative arrangements with other Australian and Queensland government agencies
- partnerships with Traditional Owners in the management of marine resources
- partnership and stewardship programs, including education programs and engagement with local governments, communities, Indigenous persons and industry
- research and monitoring programs.

Based on the Authority's statutory functions as set out in the *Great Barrier Reef Marine Park Act 1975* (the GBRMP Act), the Authority's management focus is the protection and conservation of the Great Barrier Reef Marine Park (the Marine Park). The Authority also assists in meeting Australia's international responsibilities in relation to the environment and protection of the world heritage properties of the Region.

As the strategic assessment is being carried out under the *Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act), this chapter concludes with a description of how the Authority's current management arrangements, established to protect the Marine Park and assist in the protection of the world heritage properties, also protect the other matters of national environmental significance relevant to the Region.

The description of the Authority's current management arrangements sets the basis for the assessments in later chapters, including the examination of the effectiveness of the Authority's current management.

## 3.2 Purpose

The purpose of the Authority's current management arrangements is to achieve:

*'The long-term protection, ecologically sustainable use, understanding and enjoyment of the Great Barrier Reef for all Australians and the international community through the care and development of the Marine Park.'*

This is derived from the objects of the GBRMP Act.

### **Objects of the Great Barrier Reef Marine Park Act 1975 — Section 2A**

- (1) The main object of this Act is to provide for the long-term protection and conservation of the environment, biodiversity and heritage values of the Great Barrier Reef Region.
- (2) The other objects of this Act are to do the following, so far as is consistent with the main object:
  - (a) allow ecologically sustainable use of the Great Barrier Reef Region for purposes including the following:
    - (i) public enjoyment and appreciation
    - (ii) public education about and understanding of the Region
    - (iii) recreational, economic and cultural activities
    - (iv) research in relation to the natural, social, economic and cultural systems and value of the Great Barrier Reef Region;
  - (b) encourage engagement in the protection and management of the Great Barrier Reef Region by interested persons and groups, including Queensland and local governments, communities, Indigenous persons, business and industry;
  - (c) assist in meeting Australia's international responsibilities in relation to the environment and protection of world heritage (especially Australia's responsibilities under the World Heritage Convention).
- (3) In order to achieve its objects, this Act:
  - (a) provides for the establishment, control, care and development of the Great Barrier Reef Marine Park; and
  - (b) establishes the Great Barrier Reef Marine Park Authority; and
  - (c) provides for zoning plans and plans of management; and
  - (d) regulates, including by a system of permissions, use of the Great Barrier Reef Marine Park in ways consistent with ecosystem-based management and the principles of ecologically sustainable use; and
  - (e) facilitates partnership with Traditional Owners in management of marine resources; and
  - (f) facilitates a collaborative approach to management of the Great Barrier Reef World Heritage Area with the Queensland Government.

In managing the Marine Park, the Authority must have regard to, and seek to act in a way that is consistent with, the objects of the Act, the protection of the world heritage values of the World Heritage Area, and the principles of ecologically sustainable use as set out in the Act, namely:

- (a) decision-making processes should effectively integrate both long-term and short-term environmental, economic, social and equitable considerations
- (b) the precautionary principle
- (c) the principle of intergenerational equity — that the present generation should ensure the health, diversity and productivity of the environment is maintained or enhanced for the benefit of future generations
- (d) the conservation of biodiversity and ecological integrity should be a fundamental consideration in decision making
- (e) improved valuation, pricing and incentive mechanisms should be promoted.

### 3.3 Strategic assessment area

In the Authority's agreement with the then Minister for Sustainability, Environment, Water, Population and Communities (see Appendix 1), the strategic assessment area is defined as: *the Great Barrier Reef Region and areas outside the Great Barrier Reef Region, to the extent that actions in those areas may affect the Great Barrier Reef Region.*

Through its management arrangements, the Authority generally seeks to directly address activities that occur in the Marine Park and to influence activities outside this area that affect or may affect the Marine Park. Section 7(1A) of the Act states that a matter shall be taken to relate to the Marine Park if it relates to either the use or management of an area that would or might affect the Marine Park or the use of a place outside the Marine Park for a purpose relating to the Marine Park.

Examples of how the Authority seeks to address activities that occur outside this area include:

- position statements to document the Authority's view on matters outside its jurisdiction but which have an effect on the Marine Park
- stewardship programs which seek to influence societal attitudes and behaviours that affect the Marine Park.

In regard to actions that may pollute water in a manner harmful to animals and plants in the Marine Park, Section 66(2)(e) of the GBRMP Act provides the capacity to regulate or prohibit these actions, whether they are within the Marine Park or elsewhere.

### 3.4 Integrated management

#### 3.4.1 Jurisdiction


Both the Australian and Queensland governments have direct legislative responsibilities within the World Heritage Area (Figure 3.1). Under Australia's constitution, regulation of natural resource management and environment protection on land are primarily the responsibility of state governments — in this case, Queensland. However, matters of national environmental significance, including world and national heritage properties and their values, are protected through national regulation.

The Australian and Queensland governments work in partnership to protect and manage the Region. The Queensland Government's role in the operation, management and regulation of the Region is based on:

- state title to Australia's territorial sea to a distance of three nautical miles from the territorial sea baseline (low water mark along the coast, including the coast of islands). However, Queensland's rights over its coastal waters are subject to the operation of the GBRMP Act, resulting in the Commonwealth having jurisdiction to regulate, through the Act, all waters within the Great Barrier Reef Marine Park in respect of matters relating to the Marine Park
- a role for the Queensland Government in management of the Marine Park, particularly permanent membership of the Marine Park Authority Board
- the ability for the Authority to delegate to Queensland Government officers or employees to act on the Authority's behalf where an arrangement is in force with the Queensland Government to enable this to occur
- the Queensland Government being responsible for the management of fisheries in the waters adjacent to the Queensland coast (including within the Marine Park). The Authority has a regulatory role in the management of fishing through its Zoning Plan and Regulations, as well as permit requirements for a limited number of commercial and developmental fisheries. The Authority has an advisory role to other agencies in relation to the management of fisheries in the Region. The Commonwealth Department of the Environment also has responsibilities relating to fisheries in the World Heritage Area through implementation of the EPBC Act. This requires the Australian Government to assess the environmental performance of those fisheries with an export component, and promote ecologically sustainable management.

The Queensland Government manages the Great Barrier Reef Coast Marine Park, established under the *Marine Parks Act 2004* (Qld). This adjoins the Great Barrier Reef Marine Park and covers the area between low water mark and highest astronomical tide, and Queensland's internal waters.

The State of Queensland has jurisdiction over approximately 980 islands. About 400 of these contain national parks under the *Nature Conservation Act 1992* (Qld) and are managed as part of the joint Field Management Program (Section 3.8). The remaining Queensland islands have a variety of tenures (freehold, leasehold and unallocated state land) and are not part of the Field Management Program. They are managed by a range of other Queensland Government agencies and local governments.



**Figure 3.1 Jurisdictional boundaries for the Great Barrier Reef Region**

The Great Barrier Reef encompasses both Commonwealth and Queensland jurisdictions. An agreement to jointly manage marine parks ensures integrated field management of both the Great Barrier Reef Marine Park and the adjacent Great Barrier Reef Coast Marine Park. Queensland territory extends from the land to the three nautical mile limit. Fisheries management, within the Region and beyond, is a Queensland Government responsibility. The assessment and approval provisions of the *Environment Protection and Biodiversity Conservation Act 1999* apply throughout the Region; however, its Commonwealth marine area provisions apply only in Commonwealth Territory. Adapted from the *Great Barrier Reef Outlook Report 2009*<sup>1</sup>.

Approximately 70 Commonwealth islands are part of the Great Barrier Reef Marine Park. Twenty-one of these include navigational lights or lightstations and, in 1988, responsibility for management of those islands was transferred from the Australian Maritime Safety Authority to the Great Barrier Reef Marine Park Authority. The Australian Maritime Safety Authority remains responsible for the operation of the navigation lights, and leases back this portion of land. The Authority manages Kent Island, Pilon Island, Unnamed Island (Pilon Island Group), Russell Island, North Reef Island, Low Isles, Pine Islet, Lady Elliot Island, Albany Rock, Coppersmith Island, Hannah Island, Bailey Islet, Clerke Island, Coquet Island, Eshelby Island, Hannibal Island, High Peak Islet, Rocky Island and South Brook Island, together with parts of Dent and Penrith islands.

The Department of Defence is responsible for all the remaining Commonwealth islands in the Region, except for Little Fitzroy Island which is owned by the Department of Finance and Deregulation.

Development and land use activities in adjacent coastal and water catchment areas play a critical role in the health and resilience of the Great Barrier Reef. The Queensland Government is responsible for natural resource management and land use planning on Queensland islands, and the coast and hinterland adjacent to the Region, including through the *Sustainable Planning Act 2009* and the *Coastal Protection and Management Act 1995*.

Under the EPBC Act, the Australian Government is responsible for regulating activities having or likely to have a significant impact on matters of national environmental significance, whether they are undertaken in or outside of the Region.

### 3.4.2 Great Barrier Reef Intergovernmental Agreement

The Australian and Queensland governments' cooperative approach to management of the Great Barrier Reef was initially formalised by the Emerald Agreement in 1979, and was updated by the Great Barrier Reef Intergovernmental Agreement in 2009.

The objective of this latest agreement is to ensure an integrated and collaborative approach is taken by the Australian and Queensland governments to manage marine and land environments within and adjacent to the World Heritage Area so as to:

- provide for the long-term protection and conservation of the environment and biodiversity of the Great Barrier Reef ecosystem, as encompassed by the World Heritage Area, and its transmission in good condition to future generations



- allow ecologically sustainable use of the Great Barrier Reef ecosystem subject to the overarching objective of long-term protection and conservation
- provide for meeting Australia's international responsibilities for the World Heritage Area under the World Heritage Convention.

To achieve these objectives, the agreement includes each government's ongoing commitment to:

- prohibit activities for the exploration and recovery of minerals or petroleum, and any drilling and mining within the World Heritage Area, including for the purposes of depositing materials
- maintain complementarity of relevant Commonwealth and Queensland management arrangements, in particular: marine park legislation and associated Regulations; zoning plans and plans of management; planning and development arrangements; environmental assessment and permit requirements; management of fishing activities
- continue a Commonwealth–Queensland Ministerial Council to facilitate implementation and achievement of the objectives of this agreement
- continue a joint program of field management, with shared funding on a 50:50 basis, for the Great Barrier Reef Marine Park and Queensland marine and national parks within the World Heritage Area
- continue joint action to halt and reverse the decline in quality of water entering the Reef
- continue joint action to maximise the resilience of the Reef to climate change
- address significant threats to the health and biodiversity of the Reef ecosystem, including pollution from the land and sea, the impacts of climate change, ecologically unsustainable fishing activities and other resource extraction activities
- periodically review the condition of the Reef ecosystem and any need for further action
- ensure that Indigenous traditional cultural practices continue to be recognised in the conservation and management of the Reef.

### 3.4.3 Great Barrier Reef Ministerial Forum

The Great Barrier Reef Ministerial Council was established under the 1979 Emerald Agreement, and arrangements for the Council were updated through the Great Barrier Reef Intergovernmental Agreement in 2009. From 1 July 2011, the Council has been called the Great Barrier Reef Ministerial Forum.

The Ministerial Forum is comprised of two ministers each from the Australian and Queensland governments, with responsibility for matters relating to the environment and marine parks, science, tourism and/or natural resource management. Ministers responsible for mining may not be members.

The Ministerial Forum's role is strategic in that it facilitates and oversees implementation and achievement of the objectives of the Intergovernmental Agreement. The Ministerial Council met 32 times under the 1979 Emerald Agreement. Four meetings have been held under the 2009 Intergovernmental Agreement.

Key strategic initiatives of the Great Barrier Reef Ministerial Forum and Ministerial Council include establishment of the joint Field Management Program in 1979, implementation of zoning plans, the establishment of Dugong Protection Areas, endorsement of the Reef Water Quality Protection Plan, and agreement to conduct a comprehensive strategic assessment of planned and potential coastal development affecting the Great Barrier Reef.

Since the integrated governance and management model for the Great Barrier Reef has been in place, it has facilitated development of complementary Commonwealth and state legislation, integrated management with relevant Commonwealth and state agencies, and guided the application of ecosystem-based management principles both inside and outside marine park boundaries. Indeed, it is considered by many to be the 'gold standard' for large-scale marine protected area management.<sup>2</sup>

## 3.5 Legislation and conventions

The principal legislation relating to protection and management of the Region is the Commonwealth GBRMP Act and its supporting Great Barrier Reef Marine Park Regulations 1983 (the GBRMP Regulations).

In addition, there is a range of other Commonwealth and Queensland legislation relevant to management of the Region. Management is also guided by Australia's obligations under relevant international conventions.

The legislation and conventions relevant to the Region are listed below:

### Great Barrier Reef Marine Park legislation

- *Great Barrier Reef Marine Park Act 1975* is the primary Act in respect to the Great Barrier Reef Marine Park.
- Great Barrier Reef Marine Park Regulations 1983 are the primary Regulations in force under the *Great Barrier Reef Marine Park Act 1975*.

- Great Barrier Reef Marine Park (Aquaculture) Regulations 2000 regulate the discharge of waste from aquaculture operations outside the Marine Park, which may affect animals and plants within the Marine Park. (Note: It is the Great Barrier Reef Marine Park Authority's intention to rescind the Great Barrier Reef Marine Park (Aquaculture) Regulations 2000, as equivalent levels of protection for the plants and animals in the Marine Park are provided through the *Environment Protection and Biodiversity Conservation Act 1999*).
- *Great Barrier Reef Marine Park (Environmental Management Charge—Excise) Act 1993* and *Great Barrier Reef Marine Park (Environmental Management Charge—General) Act 1999* govern operation of the environmental management charge.
- Great Barrier Reef Marine Park Zoning Plan 2003 is the primary planning instrument for the conservation and management of the Marine Park.
- Cairns Area Plan of Management 1998, Whitsundays Plan of Management 1998, Hinchinbrook Plan of Management 2004 and Shoalwater Bay (Dugong) Plan of Management 1997 establish more detailed management arrangements for specific areas of the Marine Park.

#### **Other Commonwealth legislation**

- *Environment Protection and Biodiversity Conservation Act 1999* regulates actions that have, will have or are likely to have, a significant impact on matters of national environmental significance.
- *Environment Protection (Sea Dumping) Act 1981* prohibits dumping of waste or other matter from any vessel, aircraft or platform in Australian waters, unless a permit has been issued.
- *Historic Shipwrecks Act 1976* prohibits certain activities in relation to historic shipwrecks and relics, and requires discoveries to be notified.
- *Native Title Act 1993* recognises and protects native title and includes a mechanism for determining claims to native title.
- *Protection of the Sea (Prevention of Pollution from Ships) Act 1983* gives effect to Australia's commitments under the International Convention for the Prevention of Pollution from Ships.
- *Sea Installations Act 1987* regulates the installation of structures including tourism pontoons and power cables.

#### **Queensland legislation**

- *Aboriginal Cultural Heritage Act 2003*
- *Coastal Protection and Management Act 1995*
- *Environmental Protection Act 1994*
- *Fisheries Act 1994*
- *Local Government Act 1993*
- *Marine Parks Act 2004*
- *Land Act 1994*
- *Mineral Resources Act 1989*
- *Marine Parks (Great Barrier Reef Coast) Zoning Plan 2004*
- *Marine Parks Regulation 2006*
- *Mineral Resources Act 1989*
- *Native Title (Queensland) Act 1993*
- *Nature Conservation Act 1992*
- *Queensland Heritage Act 1992*
- *State Development and Public Works Organisation Act 1971*
- *Sustainable Planning Act 2009*
- *Torres Strait Islander Cultural Heritage Act 2003*
- *Transport Operations (Marine Pollution) Act 1995*
- *Transport Operations (Marine Safety) Act 1994*
- *Transport Infrastructure Act 1994*
- *Vegetation Management Act 1999*
- *Water Act 2000*
- *Workplace Health and Safety Act 1995*.

#### **International agreements**

- Convention concerning the Protection of the World Cultural and Natural Heritage, 1972
- Convention on Biological Diversity, 1992
- Convention on International Trade in Endangered Species of Wild Fauna and Flora, 1973

- Convention on the Conservation of Migratory Species of Wild Animals, 1979
- Convention on Wetlands of International Importance Especially as Waterfowl Habitats, 1971
- China–Australia Migratory Bird Agreement, 1986
- International Convention for the Prevention of Pollution from Ships, 1973
- Japan–Australia Migratory Bird Agreement, 1974
- Republic of Korea–Australia Migratory Bird Agreement, 2007
- United Nations Convention on the Law of the Sea, 1982
- United Nations Framework Convention on Climate Change, 1992.


### 3.6 Management tools

The Authority uses a wide range of tools to manage the Region:

- Acts and regulations
- Zoning Plan
- plans of management
- permits (including environmental impact assessment)
- fees and charges
- Traditional Owner agreements
- compliance
- site infrastructure
- policy (including strategies, policies, position statements, site management arrangements and guidelines)
- partnerships
- stewardship and best practice
- education and community awareness
- research and monitoring
- reporting.

A summary description of each of these management tools is described in detail in Table 3.1, and outlined further in Section 3.7 and Section 3.8.

An emerging management tool is offsets. These are measures that compensate for the residual impacts of an action on the environment, after avoidance and mitigation measures are taken. Where appropriate, offsets are considered during the assessment phase of an environmental impact assessment under the EPBC Act. The GBRMP Regulations provide for permissions to be granted, subject to the provision of a security or financial guarantee for the recovery, restoration or removal of a structure and/or requiring the permission holder to make a financial contribution to undertake specified activities to protect the environment of the Marine Park.



*Strategies are specific arrangements that guide decision makers and the public*

Table 3.1 Management tools employed by the Authority to protect and manage the Great Barrier Reef Region and relevant matters of national environmental significance

Management tool	Purpose	Current components and activities
<b>Acts and Regulations</b>	The GBRMP Act and GBRMP Regulations govern the protection and management of the Great Barrier Reef Marine Park. They provide for the Zoning Plan and plans of management, and govern permitting decisions. The provisions are matched in areas of Queensland jurisdiction by the <i>Marine Parks Act 2004</i> and Regulations. Other Commonwealth and Queensland legislation also applies in the Region; for example, the <i>Environment Protection and Biodiversity Conservation Act 1999</i> .	<ul style="list-style-type: none"> <li>• <i>Great Barrier Reef Marine Park Act 1975</i></li> <li>• Great Barrier Reef Marine Park Regulations 1993</li> <li>• Providing advice; for example, on projects assessed under the <i>Environment Protection and Biodiversity Conservation Act 1999</i></li> <li>• Coordinating application of the Queensland <i>Marine Parks Act 2004</i> and Regulations; for example, in relation to joint marine parks permits</li> </ul>
<b>Zoning Plan</b>	Provides spatial control of use and, to a lesser extent, access within the Great Barrier Reef Marine Park. Establishes the framework for extractive use and the need for permits for some uses, such as tourism, infrastructure and research. Zoning plans are developed under Part 5 Division 2 of the <i>Great Barrier Reef Marine Park Act 1975</i> . Complementary arrangements are in place in adjacent areas under Queensland jurisdiction.	<ul style="list-style-type: none"> <li>• Great Barrier Reef Marine Park Zoning Plan 2003</li> </ul>
<b>Plans of management</b>	Set out specific arrangements for activities, areas, species or ecological communities. They complement zoning and permitting arrangements. Some components are legally binding. Plans of management are developed under Part VB of the <i>Great Barrier Reef Marine Park Act 1975</i> . There is the capacity for the Authority to enter into agreements or arrangements for management of an area, species or ecological community with a community group having a special interest in an area, including some form of native title.	<ul style="list-style-type: none"> <li>• Cairns Area Plan of Management 1998</li> <li>• Hinchinbrook Area Plan of Management 2004</li> <li>• Whitsundays Plan of Management 1998</li> <li>• Shoalwater Bay (Dugong) Plan of Management 1997</li> </ul>
<b>The permission system</b>	Facilitate opportunities for sustainable use of the Marine Park. Permits are issued mainly for tourism, research, harvest fisheries, dredging and infrastructure (for example, jetties and marinas) and include detailed risk-based environmental impact assessment. Matched in adjacent areas of Queensland jurisdiction, generally providing a joint permit. Fisheries licences are issued and managed by the Queensland Government.	<ul style="list-style-type: none"> <li>• Permits granted under Part 2A of the Great Barrier Reef Marine Park Regulations 1983 and under Queensland Marine Parks Regulations 2006</li> </ul>
<b>Fees and charges</b>	Three main fees and charges apply in the Marine Park: <ul style="list-style-type: none"> <li>• The cost of assessing an application for a permit for commercial activities is partly recovered through payment of a permit application assessment fee.</li> <li>• The environmental management charge applies to some commercial activities operating under a permit issued by the Authority. The revenue is applied to Marine Park management.</li> <li>• Bonds (usually as a bank guarantee) may be held by the Authority to cover the risks associated with a proposed activity.</li> </ul>	<ul style="list-style-type: none"> <li>• Permit application assessment fees are currently charged for activities of a commercial nature including tourist programs; vessel chartering; construction or maintenance of a facility; operation of a land-based sewage outfall; and the construction and operation of a mooring.</li> <li>• Most tourism visitors to the Marine Park pay the environmental management charge. For operations involving the hire of equipment, installation and operation of tourist facilities, and sewage outfalls, quarterly charges are paid by the operator.</li> <li>• Bonds are generally secured as part of a deed of agreement between the permittee and the Authority.</li> </ul>



Management tool	Purpose	Current components and activities
<b>Policy</b>	<p>Developed by the Authority, under Section 7(4) of the <i>Great Barrier Reef Marine Park Act 1975</i>, detailing the way in which the Authority intends to manage the Marine Park or perform its other functions.</p> <p>Policy documents are not legislative instruments. They are specific arrangements that guide decision makers and the public. Types of policy documents include: strategies, policies, site management arrangements, position statements and guidelines.</p>	<p><b>Strategies</b></p> <ul style="list-style-type: none"> <li>• <i>Great Barrier Reef Biodiversity Conservation Strategy 2013</i></li> <li>• <i>Great Barrier Reef Climate Change Adaptation Strategy and Action Plan 2012–2017</i></li> <li>• <i>Great Barrier Reef Heritage Strategy 2005</i></li> <li>• <i>Recreation Management Strategy for the Great Barrier Reef Marine Park</i></li> </ul> <p><b>Policies</b></p> <ul style="list-style-type: none"> <li>• Cruise shipping policy for the Great Barrier Reef Marine Park</li> <li>• Dredging and spoil disposal</li> <li>• Environmental impact management</li> <li>• Managing activities that include the direct take of a protected species from the Great Barrier Reef Marine Park</li> <li>• Managing bareboat operations in the Great Barrier Reef Marine Park</li> <li>• Managing scientific research in the Great Barrier Reef Marine Park</li> <li>• Managing tourism permissions to operate in the Great Barrier Reef Marine Park (including allocation, latency and tenure)</li> <li>• Marine tourism contingency plan for the Great Barrier Reef Marine Park</li> <li>• Moorings in the Great Barrier Reef Marine Park</li> <li>• Operational policy on whale and dolphin conservation in the Great Barrier Reef Marine Park</li> <li>• Sewage discharges from marine outfalls to the Great Barrier Reef Marine Park</li> <li>• Structures</li> </ul> <p><b>Site management arrangements</b></p> <ul style="list-style-type: none"> <li>• Site plans for Raine Island, Moulter Cay and MacLennan Cay; Clump Point, Mission Beach; Low Isles, offshore from Port Douglas; Michaelmas Cay locality; Upolu Cay Reef; Bauer Bay; South Molle Island; Blue Pearl Bay, Hayman Island; Whitsundays Plan of Management setting 5 site plans; Tongue Bay; Hill Inlet and Whitehaven Beach; Fitzroy Reef; Keppel Bay and islands; Lady Elliot Island Reef; Lady Musgrave Island Reef</li> </ul> <p><b>Position statements</b></p> <ul style="list-style-type: none"> <li>• Aquaculture within the Great Barrier Reef Marine Park</li> <li>• Conservation of dugongs in the Great Barrier Reef Marine Park</li> <li>• Indigenous participation in tourism and its management</li> <li>• Management of commercial jet ski operations around Magnetic Island</li> <li>• Management of tourist flights in the vicinity of Magnetic Island</li> <li>• Management of memorials within the Great Barrier Reef Marine Park</li> <li>• Managing access to the Restricted Access Special Management Areas surrounding Raine Island, Moulter Cay and MacLennan Cay</li> <li>• Marine tourism contingency plan for the Great Barrier Reef Marine Park</li> <li>• No structures sub-zones</li> <li>• Translocation of species in the Great Barrier Reef Marine Park</li> </ul> <p><b>Guidelines</b></p> <ul style="list-style-type: none"> <li>• Coral transplantation</li> <li>• Emergency disposal of foreign fishing vessels</li> <li>• Management of artificial reefs in the Great Barrier Reef Marine Park</li> </ul>

Management tool	Purpose	Current components and activities
		<ul style="list-style-type: none"> <li>Managing visitation to seabird breeding islands</li> <li>Permits Information Bulletin — no structure sub-zones</li> <li>Use of hydrodynamic numerical modelling for dredging projects in the Great Barrier Reef Marine Park</li> <li>EPBC Act referral guidelines for the outstanding universal value of the Great Barrier Reef World Heritage Area</li> </ul>
<b>Traditional Owner agreements</b>	<p>Traditional Use of Marine Resources Agreements are formal agreements describing how Traditional Owner groups work with Australian and Queensland governments to manage traditional activities in sea country. They are made in accordance with Part 2B of the Regulations. They do not affect the operation of Section 211 of the <i>Native Title Act 1993</i> and are not intended to extinguish native title rights and interests.</p> <p>Indigenous Land Use Agreements are between one or more native title groups and other people or parties about the use and management of land and waters.</p>	<ul style="list-style-type: none"> <li>Kuuku Ya'u People's Indigenous Land Use Agreement</li> <li>Traditional Use of Marine Resources Agreements for Wuthathi; Lama Lama; Yuku-Baja-Muliku; Yirrganydji; Giringun; Woppaburra; Port Curtis Coral Coast</li> </ul>
<b>Site infrastructure</b>	On-ground infrastructure is installed to manage use and protect the values of individual sites. Implemented and maintained by the Authority and the Queensland Government through the Field Management Program.	<ul style="list-style-type: none"> <li>No-anchoring areas</li> <li>Public moorings</li> <li>Reef protection markers</li> <li>Signs</li> <li>Transit lanes</li> </ul>
<b>Partnerships</b>	Formal arrangements, often executed through a memorandum of understanding or an agreement, to enable a partnership approach to management of the Marine Park.	<ul style="list-style-type: none"> <li>Great Barrier Reef Intergovernmental Agreement 2009 between the Australian and Queensland governments</li> <li>Fisheries Queensland undertakes much of the fisheries management within the Marine Park under an agreement with the Australian Government</li> <li>The Paddock to Reef monitoring program which is a commitment under the Reef Water Quality Protection Plan (Reef Plan)</li> <li>High Standard Tourism program with Ecotourism Australia</li> <li>Management agreement with the Department of Defence on the implementation of the strategic environmental assessment of defence activities in the Marine Park</li> <li>Marine Strandings Hotline</li> <li>Memorandum of understanding with the Department of the Environment, Water, Heritage and the Arts relating to the integration and application of the <i>Environment Protection and Biodiversity Conservation Act 1999</i> and the <i>Great Barrier Reef Marine Park Act 1975</i></li> <li>Memorandum of understanding with Queensland ports on port activities in or adjacent to the Great Barrier Reef Marine Park (2009)</li> <li>Local Marine Advisory Committees</li> <li>Reef Advisory Committees</li> </ul>
<b>Field management and compliance</b>	Programs and activities that encourage adherence with legal requirements, both through education and enforcement.	<ul style="list-style-type: none"> <li>Eyes and Ears Incident Reporting program</li> <li>Field Management Program jointly undertaken with the Queensland Government</li> </ul>

Management tool	Purpose	Current components and activities
<b>Stewardship and best practice</b>	Voluntary arrangements with stakeholders that provide the opportunity for contributions to protection and management. Provision of expertise and advice to stakeholders and natural resource management bodies.	<ul style="list-style-type: none"> <li>• Eyes and Ears Incident Reporting program</li> <li>• Eye on the Reef monitoring program</li> <li>• Low Isles Preservation Society</li> <li>• Marine Contingency Coordination Framework for Environmental Incidents</li> <li>• Marine monitoring program water quality monitoring volunteers</li> <li>• Pro-vision Reef Stewardship Action Plan</li> <li>• Reef Guardian program, including schools, councils, fishers, farmers and graziers, and tourism (in development)</li> <li>• Responsible Reef Practices (for tourism and recreational users)</li> </ul>
<b>Education and community awareness</b>	Programs to inform and motivate members of the community about the Great Barrier Reef and its protection and management, including ways they can contribute.	<ul style="list-style-type: none"> <li>• Community Access Points which distribute zoning maps and educational material</li> <li>• Onboard website for tourism operators</li> <li>• Reef Guardian Schools</li> <li>• Reef HQ Aquarium</li> <li>• The Authority's publications including Reef in Brief and fact sheets</li> <li>• The Authority's website and social media channels</li> </ul>
<b>Research and monitoring</b>	Undertaken, commissioned or partnered by the Authority to better inform decisions on protection and management of the Great Barrier Reef, guided by the Authority's <i>Scientific information needs for the management of the Great Barrier Reef Marine Park 2009–2014</i> .	<ul style="list-style-type: none"> <li>• Commissioned research projects to address specific management issues</li> <li>• Eye on the Reef monitoring program</li> <li>• Independent and partnered research by research institutions and the Great Barrier Reef Foundation</li> <li>• Marine monitoring program</li> <li>• National Environmental Research Program, Tropical Ecosystems Hub, 2011–2015</li> </ul>
<b>Reporting</b>	Undertaken by the Authority to meet statutory, national and international obligations, and to provide direction for strategic planning within the Authority.	<ul style="list-style-type: none"> <li>• Field Management business strategy (annual)</li> <li>• Great Barrier Reef Marine Park Authority annual report</li> <li>• <i>Great Barrier Reef Outlook Report</i> (five-yearly)</li> <li>• World Heritage periodic reporting (six-yearly)</li> </ul>

### 3.6.1 Spatially explicit management tools

The Authority utilises a range of spatially-explicit planning instruments. These include:

- at a Reef-wide scale, the Great Barrier Reef Marine Park Zoning Plan, including designated Shipping Areas
- at a regional scale, plans of management and Traditional Owner Agreements
- at a local scale, designated Special Management Areas, site specific management arrangements and site infrastructure.

#### Zoning Plan

The *Great Barrier Reef Marine Park Zoning Plan 2003* is the primary planning instrument for the conservation and management of the Marine Park.

The Zoning Plan aims, in conjunction with other management mechanisms, to protect and conserve the biodiversity of the Great Barrier Reef ecosystem, while providing opportunities for the ecologically sustainable use of, and access to, the Great Barrier Reef Region by current and future generations.

#### The objects of the Zoning Plan are:

- (a) to regulate the use of the Marine Park so as to:
  - (i) protect the ecosystem within the Great Barrier Reef Region; and
  - (ii) ensure the use is ecologically sustainable use; and
  - (iii) manage competing usage demands; and
- (b) to protect areas in the Marine Park that are of high conservation value; and
- (c) to protect and conserve the biodiversity of the Marine Park, including ecosystems, habitats, populations and genes; and
- (d) to regulate activities that exploit the resources of the Great Barrier Reef Region so as to:
  - (i) minimise the adverse effect of those activities on the Great Barrier Reef; and
  - (ii) ensure the ecologically sustainable use of the resources; and
- (e) to protect the world heritage values of the Great Barrier Reef World Heritage Area; and
- (f) to provide for the ecologically sustainable use of marine resources by Traditional Owners consistent with their traditional practices; and
- (g) to reserve some areas of the Great Barrier Reef Region for public enjoyment and appreciation; and
- (h) to preserve some areas of the Great Barrier Reef Region in a natural state, undisturbed except for the purposes of scientific research that cannot be undertaken elsewhere in the Marine Park.


The Zoning Plan divides the Great Barrier Reef Marine Park into eight zones (Figure 3.2). It sets out the purposes for which each zone may be used or entered without permission, and the purposes for which each zone may be used or entered only with the written permission of the Authority. The General Use Zone provides for the widest range of activities, while the Preservation Zone is the most restricted. The Commonwealth Islands Zone provides for the use or entry of areas of the Marine Park above mean low water on Commonwealth Islands. There are complementary zoning arrangements in adjacent areas under Queensland jurisdiction, within the Great Barrier Reef Coast Marine Park.

The Zoning Plan:

- provides for the protection of representative areas of biodiversity and other areas of high conservation value by assigning protective zoning to a range of habitats and other special or unique sites. It also provides for a range of commercial, recreational and research opportunities, and for the continuation of traditional activities
- expressly acknowledges the rights and interests of Indigenous Australians by providing for the management of the traditional use of marine resources, including traditional hunting, in accordance with Aboriginal and Torres Strait Islander custom and tradition
- acknowledges the contributions of scientific research to the management and understanding of the Great Barrier Reef, through the establishment of a system of Scientific Research Zones to facilitate research around scientific research stations
- provides for the management of remote natural areas of the Marine Park, and the designation of special management areas and shipping areas, as well as additional purposes, such as Commonwealth activities, for which zones may be used or entered. It also provides for emergency responses in relation to incidents such as risks to human life, pollution threats or vessel groundings in combination with other management arrangements, provides a high level of certainty about where and how extractive use, including fishing, may be undertaken.

In accordance with section 37 of the GBRMP Act, the Authority may amend a Zoning Plan if it (or its most recent amendment) has been in operation for at least seven years.





**Figure 3.2 The Great Barrier Reef Zoning Plan**

The Reef-wide Zoning Plan protects biodiversity values through a network of no-take zones for 33 per cent of its area and for at least 20 per cent of each bioregion. It provides for the spatial control of use, and sets out specific arrangements for activities consistent with the principles of ecologically sustainable use.

### Plans of management

Plans of management complement the Zoning Plan and impose controls on the granting of permissions. They address issues specific to an area, species or community in greater detail than can be accomplished by the broader Reef-wide zoning plans.

Plans of management are generally prepared for intensively used areas, particularly 'at-risk' groups of islands and reefs, and for the protection of species of conservation concern or ecological communities.

There are four plans of management for areas in the Marine Park, covering about eight per cent of the Marine Park. The plans for the Cairns Area, the Whitsundays and Hinchinbrook set out detailed management arrangements applying to all users of these areas, with a focus on protecting key natural values and allowing a range of uses. The Shoalwater Bay (Dugong) Plan of Management aims to manage activities in the bay that threaten its dugong population or impact on its seagrass meadows. The objects of plans of management are:

#### Objects of the plans of management — GBRMP Act Section 39Y

- (a) to ensure, for particular areas of the Marine Park in which the Authority considers that nature conservation values, cultural and heritage values, or scientific values, are, or may be, threatened, that appropriate proposals are developed to reduce or eliminate the threats;
- (b) to ensure management for the recovery and continued protection and conservation of species and ecological communities that are, or may become:
  - (i) extinct; or
  - (ii) extinct in the wild; or
  - (iii) critically endangered; or
  - (iv) endangered; or
  - (v) vulnerable; or
  - (vi) conservation dependent;
- (c) to ensure that activities within areas of the Marine Park are managed on the basis of ecologically sustainable use;
- (d) to provide a basis for managing the uses of a particular area of the Marine Park that may conflict with other uses of the area or with the values of the area;
- (e) to provide for the management of areas of the Marine Park in conjunction with community groups in circumstances where those groups have a special interest in the areas concerned;
- (f) to enable people using the Marine Park to participate in a range of recreational activities.

Plans of management are not subject to the sunset provisions of subsection 54(2) of the *Legislative Instruments Act 2003*, and may be amended from time to time.

### Traditional Owner agreements

Formal Traditional Use of Marine Resources Agreements (TUMRAs) describe how Traditional Owner groups work with Australian and Queensland governments to manage traditional use activities in sea country. Indigenous Land Use Agreements (ILUAs) are agreements between one or more native title groups and other people or parties about the use and management of land and waters.

The agreements address issues such as the sustainable take of culturally significant species, and supporting Traditional Owner cultural practice in the conservation and management of the Great Barrier Reef. The agreements incorporate traditional and contemporary scientific knowledge and environmental management for the ongoing protection of the Great Barrier Reef. TUMRAs are used by Traditional Owners as a formal tool to conserve and protect species and ecosystems critical to the health of people, culture and country.

Agreements can incorporate specific management strategies for the conservation and sustainable use of key species, as well as habitats. Other management activities that Traditional Owners may identify in their agreement implementation plan include restoring and maintaining waterways and coastal ecosystems; maintaining and protecting significant heritage values including sites; sharing and documenting traditional ecological knowledge, culture and language; and research and monitoring of sea country, including through partnerships with managing agencies and leading scientific institutions.

There are currently seven TUMRAs covering about 13 per cent of the Region. They include coverage over traditional sea country estates of Wuthathi; Lama Lama; Yuku-Baja-Muliku; Yirrganydji; Giringun; Woppaburra; and Port Curtis Coral Coast.

TUMRAs and ILUAs operate for a set time, after which the agreements and their timeframes are renegotiated.

### Supporting community-based management of sea country — Lama Lama people

Traditional Use of Marine Resources Agreements — a management tool available to Traditional Owners — recognise traditional lore, customary rights and traditional ecological knowledge in managing sea country through a formal partnership arrangement with the Authority and the Queensland Government. The agreements are developed by Traditional Owners to conserve biodiversity and protect their cultural and natural heritage values. The process of developing a marine resources agreement is managed by Traditional Owners, with support from the Authority, involving extensive consultation and negotiation within the Traditional Owner group. Through the structured planning process, the group documents its structure, governance, history, lore, custom and aspirations for managing sea country.

The Authority supports implementation of Traditional Use of Marine Resources Agreements by helping communities to develop implementation and compliance plans. In some areas, such as the Girringun region and Port Curtis Coral Coast, these plans have been incorporated into and implemented on country through Commonwealth-funded Indigenous ranger programs.

In 2013 — after a three year development phase — Lama Lama Traditional Owners had a Traditional Use of Marine Resources Agreement accredited with the Authority and the Queensland Government. The group looks forward to putting many of its plans into action through the partnership arrangement with the Authority.

*“This is an exciting time as we work to implement our ideas and thoughts to manage our sea country resources in the best possible way, not only for current Lama Lama people but for our future generations as well.”*



*Lama Lama Traditional Owners, at Maarpa Island*

During the development of their agreement, the Lama Lama people outlined how they plan to manage sea country resources such as turtle and dugong, and identified and documented culturally important areas of their sea country. Community meetings provided opportunities for Lama Lama Traditional Owners to come together, reach consensus and discuss aspirations for management of their sea country including hunting, conservation measures, science partnerships, future priorities and economic opportunities.

Meetings on sea country provided an important opportunity for the group, as elders and children have been able to visit various culturally significant sites within Princess Charlotte Bay. Children were shown the Marpa (Cliff) Islands and rock art on Ronganhu. They also explored the islands and talked about custom and culture.

The Authority supported the Lama Lama Traditional Owners' recent successful application for a Fisheries

Research and Development Corporation Indigenous Development Scholarship. The scholarship will sponsor a Lama Lama representative to visit a Pacific Island community to exchange information and learn about its ways of sea country management; traditional ecological knowledge; partnerships other communities have established with groups such as conservation volunteers; and how communities have established turtle and dugong ecotourism. Recognising the value of the knowledge exchange trip, the Authority is sponsoring an additional two Lama Lama Traditional Owners to join the trip. The information can then be shared with the broader Lama Lama community. The Authority will be the host organisation and will provide mentoring as well as technical and financial support. The scholarship will also sponsor five Lama Lama land and sea rangers to visit the Torres Strait Regional Authority's Land and Sea Management Unit to complete their coxswain training, with technical and mentoring support from the Authority.

Establishing relationships with James Cook University researchers has also been identified as a priority for the Lama Lama Traditional Owners. The possibility of researching inshore dolphins has already been investigated and resulted in the university providing dolphin identification training to the group. It has also led to a plan to bring a research vessel to Port Stewart to work with the rangers to undertake research on Lama Lama sea country.



*Lama Lama children playing at Maarpa Island*



### **Site specific management arrangements**

Site specific management arrangements are localised plans for use of a particular site. They guide decision makers and the public. They are a form of policy, developed by the Authority under section 7(4) of the GBRMP Act. They are not legislative instruments.

Site specific management arrangements identify significant values of the specific site and describe current management arrangements for these sites, concentrating on managing specific use issues and cumulative impacts at the site.

Review of site specific management arrangements falls within the formal control process undertaken by the Authority (see Section 6.2.13).

### **Site infrastructure**

On-ground infrastructure is installed to manage use and protect the values of individual sites throughout the Marine Park. For example, facilities such as moorings, no anchoring areas and transit lanes support sustainable use of popular sites within planning areas and elsewhere. Site infrastructure is installed and maintained by the Authority and the Queensland Government through the joint Field Management Program.

## **3.6.2 System-based regulatory tools**

The Authority utilises a range of system-based regulatory tools in its management of the Great Barrier Reef. These include its permission system, fees and charges, audit and compliance, and statutory monitoring, evaluation and reporting programs.

### **Permissions**

Permissions facilitate opportunities for use of the Marine Park, and are a tool to reduce impacts, separate potentially conflicting activities, collect data for planning and monitor potentially damaging activities. They are issued mainly for tourism activities, dredging and dredge material disposal, and infrastructure (for example, jetties and marinas). Harvest (dive-based) fishery operations within the Marine Park require a Marine Park permit, as do netting activities (other than bait netting) in the Special Management Area within Princess Charlotte Bay. Any developmental fishery program would also require a permit from the Authority.

As part of the permitting process, there is formal environmental impact assessment under the Act for evaluating the likely possible risks or impacts to the environment from a proposed activity or development. Common activities where a detailed assessment process is undertaken include, but are not limited to, the construction and operation of pontoons, jetties, moorings, pipelines and marinas, as well as dredging. The assessment criteria include ecological, social and economic considerations, Traditional Owner interests, as well as current and future use of the proposed location. These assessments will continue to be informed by the best available information using contemporary assessment methods including, where appropriate, modelling and mapping to understand the cause-and-effect relationships of impacts on values. A detailed description of the Authority's permission system is provided in Part C of the Program Report.

Joint Queensland Government–Authority permits are issued for activities which operate across jurisdictions. Fisheries licences are issued by the Queensland Government.

New permissions are generally granted for one year. This allows time to establish the operation, and provides the applicant and the Authority an opportunity to review the operation.

Existing permission holders who apply to continue the same operation and satisfy the assessment requirements are generally granted a permission for six to 15 years, depending on the activities to be conducted and the level of environmental certification they have achieved from a third party certification body.

### **Permission assessment and decision process**

The Great Barrier Reef Marine Park Regulations 1983 (the GBRMP Regulations) describe how permissions apply to the conduct of activities in the Great Barrier Reef Marine Park, and set out the statutory requirements for permissions including:

- applications for a permission, including advertising of applications
- consideration of applications, including mandatory and discretionary considerations when deciding whether or not to grant a permission
- granting and refusing permissions
- form, term and condition of permissions
- transfer, modification, suspension and revocation of permissions
- offence provisions.

For each permit application made, the Authority undertakes an assessment of the nature and scale of activities and makes a determination based on the acceptability of those activities on the environment.

There are four key phases in the permission assessment and decision process: scoping, assessment, decision, auditing and compliance. A brief description of each of these steps is provided below, and in more detail within Chapter 13.



The purpose of the **scoping phase** is to proactively discuss the initial concepts of a proposed action in order to explain regulatory arrangements and identify potential mitigations. For actions which may have a significant impact on the Marine Park, including matters of national environmental significance, reference is made to relevant GBRMP Act and EPBC Act guidance material. A permit application assessment fee and accompanying permit application form are provided, and must contain sufficient information to enable a clear understanding of the intended action.

The purpose of the **assessment phase** is to enable an informed decision on whether or not a permit will be granted to undertake an action within the Great Barrier Reef Marine Park. During the assessment phase, the Authority considers criteria outlined in the GBRMP Regulations, and relevant GBRMP Act and EPBC Act guidance material.

The purpose of the **decision phase** is to grant or refuse a permission based on information provided within the assessment phase. A permission may be granted with conditions consistent with the objects of the GBRMP Act, including the principles of ecologically sustainable development and the protection of matters of national environmental significance.

The purpose of the **audit and compliance phase** is to monitor the environmental performance of permitted activities against permit conditions. Where the results of performance monitoring conditions have not been met, compliance action may be undertaken.

### **The Environment Protection and Biodiversity Conservation Act 1999**

Certain activities also require assessment and approval under the EPBC Act. The EPBC Act provides the legal framework for the protection and management of nationally and internationally important flora, fauna, ecological communities, and heritage places. These are defined under the EPBC Act as matters of national environmental significance. The EPBC Act also protects, among other matters, the environment on areas of Commonwealth land and marine environment. Both of these categories of 'protected matters' are the subject of this strategic assessment (protected matters are defined in Part 3, the EPBC Act).

An action — including some actions for which permits are sought under the GBRMP Act — will require approval from the Minister for the Environment if the action has, will have, or is likely to have, a significant impact on matters protected under Part 3 of the EPBC Act. There is an onus on the proponent (person responsible for the action) of an activity to refer the action to the Minister for the Environment if it is likely that there will be a significant impact on one or more of these matters. This is typically done on a project-by-project basis.

The Department of the Environment's EPBC Act Significant Impact Guidelines 1.1<sup>3</sup> describe the meaning of 'significant impact'. Under the guidelines, a significant impact is one which could be considered to be important, notable or of consequence, in terms of the context or intensity. Whether or not an action is likely to have a significant impact depends upon the sensitivity, value and quality of the environment which is impacted, and upon the intensity, duration, magnitude and geographic extent of the impacts. To be likely, it is sufficient that a significant impact on the environment is a real and not a remote chance or possibility.

The EPBC Act Significant Impact Guidelines 1.1 identify the key considerations that should inform whether or not a proposed action should be referred. These considerations include whether protected matters are present in the area of interest, whether they may be impacted (directly or indirectly), and whether there are measures that could be taken to avoid or mitigate any impacts to reduce those impacts to below the significance threshold. Consideration should also be given to the sensitivity, value and quality of the environment, as well as the intensity, duration, magnitude and geographic extent of the impacts. There are a range of assessment processes which must then be followed for actions determined as having significant impacts.

Public comment is invited on both initial referrals as well as draft environmental impact assessments. In addition, proponents must publish their referrals and assessments, including their responses to public comments, and the Minister for the Environment must publish decisions and provide statements of reasons, if requested.

The EPBC Act also provides for compliance and enforcement of requirements. The Department of the Environment undertakes positive enforcement mechanisms through information sharing and promotion, and also conducts investigations and pursues enforcement where breaches of the legislative requirements may occur. The Minister for the Environment may revoke approvals or issue additional conditions to ensure there are not unacceptable impacts on matters protected under Part 3 of the EPBC Act.

### Inter-Government administrative arrangements for permits

The Authority and the **Queensland Parks and Wildlife Service** (QPWS) operate a joint application and permit assessment process for the determination of permissions under the GBRMP Act and *Marine Parks Act 2004* (Qld). Responsibilities at various stages are shared between the Authority and QPWS, however, the Authority takes the lead role in processing applications and preparing decision documentation. The flow chart shown in Figure 3.3 depicts the respective responsibilities and timeframes.

The Authority and the Queensland Government may elect to conduct separate marine park permit processes for development proposals where, for example, a project is declared a 'coordinated project' under the *Queensland State Development and Public Works Organisation Act 1971*. This generally applies to complex or significant infrastructure development proposals involving multiple approvals and potentially significant environmental effects.

The Authority provides the **Department of the Environment** with advice on referrals for actions outside the Great Barrier Reef catchment to determine if the action is likely to have a significant impact on relevant matters of national environmental significance in or adjacent to the World Heritage Area.

If an action is deemed to require approval under the EPBC Act (that is, deemed to be a 'controlled' action) the Authority provides advice to the Department throughout the assessment process.

Since 2001, the Authority has provided advice on over 250 EPBC Act referrals and assessments.

In November 2009, the Great Barrier Reef Marine Park became a Matter of National Environmental Significance under the EPBC Act. To facilitate an integrated and streamlined application and assessment process under the EPBC Act and the GBRMP Act, a Memorandum of Understanding between the Department of the Environment and the Great Barrier Reef Marine Park Authority was established to enable:

- the integration of environmental impact assessments and approval requirements under both the *Great Barrier Reef Marine Park Act 1975* and the *Environment Protection and Biodiversity Conservation Act 1999*
- consistent and legally effective decision making
- compliance to the statutory responsibilities of the Minister, the Department of the Environment and the Authority
- efficient and cost effective legislative administration.

Administrative processes for permit application and assessment between the Department of the Environment and the Authority are outlined in Figure 3.4.

### Policy

Developed by the Authority under section 7(4) of the GBRMP Act, policies guide decision makers and the public. They are not legislative instruments.

They detail the way in which the Authority intends to manage the Marine Park or perform its other functions, and the way in which the Authority considers that the Act or other legislative instruments apply.

There are three broad categories of guidance documents:

- **Policies:** give effect to the Authority's responsibilities, functions and powers outlined in the GBRMP Act, for example, managing environmental impacts.
- **Position Statements:** outline the Authority's position on issues where the Authority has a strong interest, but does not have legislative powers; for example, emerging risks from shipping.
- **Guidelines:** generally support the Authority's policies or position statements, and outline the Authority's process on certain issues, for example, managing visitation to seabird breeding islands.

A comprehensive list of policies, position statements and guidelines are presented in Table 3.1. Guidance documents are reviewed every four years and, if needed, updated more regularly to reflect the best available science and management, consistent with maintenance of high environmental standards.

A formal control process within the Authority regulates the development, amendment, revocation and approval of external policies and plans. Where policies or plans are likely to have a regulatory impact on business or the not-for-profit sector (such as changes to service charges, subsidies, compliance costs, restriction on competition), a 'Regulatory Impact Statement' is developed in consultation with the Office of Best Practice Regulation. Prior to the preparation of final drafts, every new or significantly reviewed external policy or plan is additionally required to be made available for public comment for a minimum period of 28 days.

This formal control process enables the review of external policies and plans every four years or sooner, as prompted by changes in legislation, standards, whole of government policy, advancements in scientific knowledge and/or advancements in better environmental management practices.

Table 3.2 outlines key policies relevant to the assessment of environmental impacts of activities on the Region's values, including matters of national environmental significance.




Figure 3.3 Administrative arrangements for the joint application and permit assessment process between the Authority and the Queensland Government




Figure 3.4 Administrative arrangements for environmental assessments under the GBRMP Act and EPBC Act



**Table 3.2 Program guidance documents and *Environment Protection and Biodiversity Conservation Act 1999* Part 3 (matters of national environmental significance)**

DOCUMENT	DESCRIPTION OF CONTENT
<b>POLICIES</b>	
<b>Environmental Impact Management Policy</b>	<p>Sets the framework for assessment, mitigation and management of environmental impacts associated with development activities in the Marine Park and the World Heritage Area.</p> <p>It provides a transparent, consistent and contemporary approach to environmental impact management of major developments in the Great Barrier Reef Marine Park. The Authority has developed the policy based on historical Marine Park Authority decisions, best available scientific and management information, and community consultation.</p>
<b>Environmental Assessment and Management Risk Assessment Guidelines</b>	<p>Provides guidance on the systematic application of risk assessment to inform decisions about the impacts human activities may have on the environment.</p> <p>The Authority has developed this guidance based on the requirements of the Australian and international risk management standard.</p>
<b>Dredging and Spoil Disposal Policy</b>	<p>Provides a transparent, consistent and contemporary approach to environmental impact management of dredging and spoil disposal in the Great Barrier Reef Marine Park. The Authority has developed the policy based on best available scientific and management information, and community consultation.</p> <p>Guides assessment of the impact of dredging and spoil disposal in accordance with the Environmental Impact Management Policy.</p>
<b>Structures Policy</b>	<p>Provides a transparent, consistent and contemporary approach to environmental impact management of structures in the Great Barrier Reef Marine Park. The Authority has developed the policy based on historical Marine Park Authority decisions, best available scientific and management information, and community consultation.</p> <p>Guides assessment of the impact of structures in accordance with the Environmental Impact Management Policy.</p>
<b>Sewage Discharges from Marine Outfalls to the Great Barrier Reef Marine Park Policy</b>	<p>Considered when assessing the impact of sewage discharges in accordance with the Environmental Impact Management Policy.</p> <p>Purpose is to minimise the potential impact of treated sewage discharges from marine outfalls in the Great Barrier Reef Marine Park. The Policy has been developed in consultation with the Queensland Government, and is complementary with the state policy where it applies to the discharge of treated sewage via marine outfalls to the Marine Park.</p>
<b>Moorings in the Marine Park</b>	<p>Considered when assessing the impact of moorings in accordance with the Environmental Impact Management Policy.</p> <p>The purpose of this policy is to detail the Authority's framework for the consistent and effective use and management of moorings in the Marine Park.</p> <p>The issues covered by the policy include availability of moorings, site planning requirements, standard mooring buoys, design, installation, maintenance, relocation and removal of moorings, moorings best practice and permit related matters.</p>
<b>Managing Scientific Research in the Marine Park</b>	<p>Considered when assessing the impact of scientific research in accordance with the Environmental Impact Management Policy.</p> <p>The issues covered by this policy include: umbrella and institutional permits; permit duration; native title; permit fees; scientific research in the Scientific Research Zone; the Buffer Zone; the Marine National Park Zone; and the Preservation Zone.</p>
<b>Managing Activities that include the direct take of a Protected Species from the Marine Park</b>	<p>Considered when assessing the impact of collection of protected species in accordance with the Environmental Impact Management Policy.</p> <p>The issues covered by this policy include: assessment considerations such as the conservation status of the species; zoning of the area of take; relevance to the conservation management of the Marine Park; permission information; provision for sick, injured or at risk animals; guidelines; photography; filming or sound recording; and public display for education; tourism interpretation; research and broodstock collection for aquaculture.</p> <p>The direct take of a Protected Species for the purpose of traditional use of marine resources is not addressed in this policy.</p>

DOCUMENT	DESCRIPTION OF CONTENT
<b>POSITION STATEMENTS</b>	
<b>Translocation of Species in the Great Barrier Reef Marine Park</b>	<p>Considered when assessing activities under the Environmental Impact Management Policy that may involve translocation of species.</p> <p>The desired outcome of this position statement is to reduce the risks associated with translocation-related activities in the Marine Park through a process of risk management, including a case-by-case activity risk assessment, and requiring the development of a translocation proposal for each relevant permit application.</p>
<b>Aquaculture</b>	<p>Considered when assessing the impact of aquaculture activities in accordance with the Environmental Impact Management Policy.</p> <p>The position statement provides a brief background on actual and potential impacts of aquaculture operations in the Marine Park, and a guiding statement of the approach that the Authority will take in assessing applications for aquaculture operations.</p>
<b>GUIDELINES</b>	
<b>Permits Information Bulletin — No Structures Sub-zones</b>	<p>Considered when assessing the impact of structures in accordance with the Environmental Impact Management Policy.</p> <p>The objective of No Structure Sub-zones is to ensure that some areas of the Great Barrier Reef Marine Park remain:</p> <ol style="list-style-type: none"> <li>a) in a natural state, largely unaltered by human works</li> <li>b) free from structures and permanently-moored facilities, except for approved vessel moorings, approved management, research and monitoring facilities and approved navigational markers which are essential for the protection, wise use, understanding and enjoyment of the Marine Park.</li> </ol>
<b>Water Quality Guidelines for the Great Barrier Reef Marine Park</b>	<p>Considered when assessing activities under the Environmental Impact Management Policy that may impact water quality.</p> <p>These guidelines have been established to support initiatives to halt and reverse the decline in the quality of water entering the Great Barrier Reef. The guidelines confirm the values of the World Heritage Area and recognise the protection of the ecological systems of the Great Barrier Reef World Heritage Area from water-borne contaminants as one of the critical issues for management.</p>
<b>The use of Hydrodynamic Numerical Modelling for Dredging Projects in the Great Barrier Reef Marine Park</b>	<p>Provides guidance on the use of three-dimensional hydrodynamic numerical models in the Great Barrier Reef Marine Park.</p> <p>These guidelines have been established to inform proponents about the specific hydrodynamic modelling and dredge plume modelling expected to be undertaken in order to predict the extent, intensity and persistence of dredge-generated sediment plumes, and the extent, severity and duration of the resultant impacts to benthic habitats.</p>

### Fees and charges

Three main fees and charges apply in the Marine Park:

- The cost of assessing an application for a permit for commercial activities is partly recovered through payment of a permit application assessment fee.
- The environmental management charge applies to some commercial activities operating under a permit issued by the Authority. The revenue is applied to Marine Park management.
- Bonds (usually as a bank guarantee) may be held by the Authority to cover the risks associated with a proposed activity. It primarily relates to recovery, restoration and/or removal of an activity/structure, and is designed to act as a financial incentive for a permission holder to meet required performance levels.

### Compliance

Compliance activities — both through education and enforcement — encourage adherence to legal requirements such as those contained in permits, plans of management and zoning, helping to ensure balanced and sustainable use of the Marine Park. The Field Management Program, jointly undertaken by the Authority and the Queensland Government, plays a key role in compliance activities in the Region, including in relation to fishing. A cooperative multi-agency approach allows a broad range of legislation and compliance tools to be used, with vessel and aerial surveillance activities by all agencies coordinated by the Field Management Compliance Coordination Unit. The Authority works closely with Traditional Owners in relation to traditional use of marine resources and in managing any risks associated with poaching of species of conservation concern (for example, dugongs and marine turtles), including through Indigenous community compliance liaison officers<sup>4</sup>. Informal activities such as under the community-based Eyes and Ears Incident Reporting program also help inform compliance actions.

### Field management operations

Field management activities, as part of the joint Australian and Queensland government Field Management Program, underpin regulatory approaches. They are a significant proportion of the Authority's work program and are at the frontline of efforts to protect the Great Barrier Reef and ensure use remains ecologically sustainable. Activities that support environmental regulation include: on-ground visitor management; visitor infrastructure and facilities (for example, moorings and reef protection markers); compliance; monitoring and assessment of habitats and vulnerable species; and incident response (for example, extreme weather events and shipping incidents).

### Statutory reporting

The Authority has a number of statutory, national and international obligations to report on the Great Barrier Reef, its condition and management arrangements. Analysis and reporting also provides direction for strategic planning within the Authority. Examples of reporting include:

- an annual report to the Parliament of Australia
- five-yearly reporting, including on reef condition, resilience and management effectiveness, through the *Outlook Report* (see below)
- responses to government priorities for example, this strategic assessment
- international reporting obligations such as contributing to world heritage reporting.

#### ***Great Barrier Reef Outlook Report — GBRMP Act, Section 54***

- (1) The Authority must prepare and give to the Minister a report in relation to the Great Barrier Reef Region every five years. The first report must be given to the Minister by 30 June 2009.
- (2) The report must be prepared in accordance with the regulations (if any).

##### *Content of report*

- (3) The report must contain the following matters:
  - (a) an assessment of the current health of the ecosystem within the Great Barrier Reef Region and of the ecosystem outside that region to the extent it affects that region;
  - (b) an assessment of the current biodiversity within that region;
  - (c) an assessment of the commercial and non-commercial use of that region;
  - (d) an assessment of the risks to the ecosystem within that region;
  - (e) an assessment of the current resilience of the ecosystem within that region;
  - (f) an assessment of the existing measures to protect and manage the ecosystem within that region;
  - (g) an assessment of the factors influencing the current and projected future environmental, economic and social values of that region;
  - (h) an assessment of the long-term outlook for the ecosystem within that region;
  - (i) any other matter prescribed by the regulations for the purposes of this paragraph.

#### ***Great Barrier Reef Outlook Report — GBRMP Regulations, Section 116A***

- (1) For paragraph 54(3)(i) of the Act, an assessment of the heritage values of the Great Barrier Reef Region is prescribed as a matter that must be contained in the Great Barrier Reef Outlook Report.
- (2) An ***assessment of the heritage values***, of the Great Barrier Reef Region, includes the following:
  - (a) an assessment of the current heritage values of the region;
  - (b) an assessment of the risks to the heritage values of the region;
  - (c) an assessment of the current resilience of the heritage values of the region;
  - (d) an assessment of the existing measures to protect and manage the heritage values of the region;
  - (e) an assessment of the factors influencing the current and projected future heritage values of the region;
  - (f) an assessment of the long-term outlook for the heritage values of the region.
- (3) In this regulation:  
***heritage values***, of the Great Barrier Reef Region, include the following values for the region:
  - (a) the Commonwealth heritage values;
  - (b) the heritage values;
  - (c) the Indigenous heritage values;
  - (d) the national heritage values;
  - (e) the world heritage values.

### 3.7 Management activities

The Authority's management tools are designed to protect the Region's values and allow ecological sustainable use. The current management activities of the Authority are grouped into 15 topics covering the major uses and the most serious areas of risk to the Region. The groupings are based on the management topics considered in the *Great Barrier Reef Outlook Report 2009*<sup>1</sup>, with amendments to reflect emerging issues.

The management topics addressed by the Authority are:

#### Values

- biodiversity protection
- heritage (Indigenous and historic)
- community benefits

#### External impacts on values

- climate change and extreme weather
- water quality protection (catchment run-off)
- coastal development (protection of coastal ecosystems)

#### Direct use

- tourism (marine-based)
- fishing — commercial
- fishing — recreational
- recreation
- port activities
- shipping
- defence activities
- research activities

Management tools are applied as appropriate to each management topic. The tools applied to each topic are summarised in Table 3.3.

**Table 3.3 Management tools used to address management topics in the Great Barrier Reef Region**

		Management topic														
		Biodiversity protection	Indigenous heritage	Historic heritage	Community benefits	Climate change and extreme weather	Water quality protection	Coastal development	Tourism	Fishing — commercial	Fishing — recreational	Recreation	Port activities	Shipping	Defence activities	Research activities
Regulatory	Management tool															
	Acts and Regulations	•	•	•	•		•	•	•	•	•	•	•	•	•	•
	Zoning Plan	•	•	•	•		•	•	•	•	•	•	•	•	•	•
	Plans of management	•	•	•	•				•	•	•	•			•	•
	The permission system	•	•	•	•		•	•	•	•*		•	•	•		•
	Traditional Owner agreements	•	•	•	•											
	Field management and compliance	•	•	•	•		•		•	•	•	•	•	•	•	•
	Site infrastructure	•	•	•	•				•		•	•				
Non-regulatory	Fees and charges						•		•	•*			•			
	Policy	•	•	•		•	•	•	•			•	•			•
	Partnerships	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
	Stewardship and best practice	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
	Education and community awareness	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
	Research and monitoring	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
	Reporting	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•

\*commercial harvest fisheries only



### 3.8 Partners in management

In addition to the Australian and Queensland governments' partnership established through the Intergovernmental Agreement, the Authority has partnership arrangements or stewardship programs with Traditional Owners, industry sectors, community groups and individuals who directly participate in protection and management of the Great Barrier Reef. The main contributors to protection and management are listed below:

#### Australian Government

- **Great Barrier Reef Marine Park Authority**  
Protects and manages the Great Barrier Reef Marine Park.
- **Department of the Environment**  
Develops and implements national policies, programs and legislation to protect and conserve Australia's natural environment and heritage. Responsible for implementing the Environment Protection and Biodiversity Conservation Act.
- **Department of Climate Change and Energy Efficiency**  
Leads the development and coordination of Australia's climate change policies and programs.
- **Australian Customs Service**  
Enforces a range of Commonwealth laws at sea and at various international entry points, including ports. Coastwatch provides aerial surveillance of Australian coastal waters.
- **Australian Maritime Safety Authority**  
Manages shipping activities throughout the Great Barrier Reef Region. Coordinates emergency responses to marine emergencies and marine pollution; for example, oil spills.
- **Australian Quarantine Inspection Service**  
Responsible for quarantine inspection services for all vessels entering Australian waters.
- **Department of Agriculture, Fisheries and Forestry**  
Develops and implements policies and programs to ensure Australia's agricultural, fisheries, food and forestry industries remain competitive, profitable and sustainable.
- **Department of Defence**  
Responsible for all defence activities within the Great Barrier Reef Region, including the management of defence activities in designated defence training areas.
- **Department of Resources, Energy and Tourism**  
Develops and delivers policies to increase Australia's international competitiveness in relation to resources, energy and tourism, consistent with the principles of environmental responsibility and sustainable development.
- **Australian Institute of Marine Science**  
Undertakes research that supports management of tropical marine environments — the World Heritage Area is a primary focus.
- **CSIRO**  
Australia's national science agency.

#### Queensland Government

- **Department of National Parks, Recreation, Sport and Racing**  
The **Queensland Parks and Wildlife Service** is responsible for day-to-day field management in the Great Barrier Reef Marine Park and the adjacent Great Barrier Reef Coast Marine Park, as well as island and mainland national parks.
- **Department of Environment and Heritage Protection**  
Responsible for management of Queensland's environment, including water, salinity, native title and threatened species. It is the Queensland Government's lead agency on environmental management matters, including the assessment and approval of works in intertidal areas, internal waters and the Great Barrier Reef catchment.
- **Department of Agriculture, Fisheries and Forestry**  
Responsible for management and research on fisheries and fisheries habitat in the Great Barrier Reef Region. The Queensland Boating and Fisheries Patrol enforces fisheries regulations, and marine parks and transport legislation.
- **Department of the Premier and Cabinet**  
Provides overall coordination and direction for Queensland Government involvement in Great Barrier Reef matters. Within the department, the Reef Secretariat provides leadership for the *Reef Water Quality Protection Plan*.
- **Department of State Development, Infrastructure and Planning**  
Delivers major infrastructure, economic development and planning services, including developing regional plans along the Great Barrier Reef coast.

- **Department of Transport and Main Roads**  
Provides policy and strategic advice relating to Queensland's ports system. Within the department, **Maritime Safety Queensland** is responsible for licencing, registration and the safe navigation of vessels. It is the lead response agency for oil and chemical spills.
- **Queensland Water Police**  
Enforces marine parks legislation and investigates crimes on the water.

#### Other partners

- **Traditional Owners**  
Traditional Owners have inherent rights and interests over their sea country in the Region. They work with the Authority to protect cultural and heritage values, conserve biodiversity and enhance the resilience of the Great Barrier Reef.
- **Native title bodies**  
Assist in the permit referral process for applications for activities in the marine parks.
- **Local government**  
Responsible for local planning and development decisions and providing public services such as local roads, waste removal and water treatment within the Great Barrier Reef catchment.
- **Natural resource management bodies**  
Support natural resource management activities in the Great Barrier Reef catchment.
- **Research institutions**  
Provide improved knowledge about the Great Barrier Reef and advice on its implications for management.
- **Industry groups and individual operators**  
Organise and participate in programs that contribute to protection and management (for example, tourism operators, commercial fishers, farmers, graziers, Association of Marine Park Tourism Operators and other tourism associations, Queensland Seafood Industry Association, Growcom, Queensland Cane Growers Association, and ports corporations).
- **Community groups**  
Organise and participate in community activities that help people to understand and protect the Great Barrier Reef.
- **Environmental non-government organisations**  
Raise public awareness about the state of the Great Barrier Reef and advocate for its increased conservation and protection.
- **Schools**  
Educate and engage students in marine activities and conservation.

### Fishers work with government to protect dugongs in Bowling Green Bay

Between July 2010 and September 2011, there were seven dugong deaths reported in Bowling Green Bay. This was of particular concern as it is considered that recovery of the severely depleted dugong population south of Cooktown requires mortality from human causes to be as close to zero as possible. The deaths were understood to be associated with incidental capture in fishing nets. Fishing representatives from the Burdekin Sustainable Fisheries Alliance — a local stewardship initiative — worked to address this issue in partnership with the Authority, Queensland Government agencies and the Queensland Seafood Industry Association as part of the Burdekin Regional Management Project.

Alliance members developed substantial modifications to the type of large mesh net used in the high-risk areas, making the nets shorter with a shallower drop and lead core rope weighting to further reduce the risk of catching dugongs. The group also suggested that an adjacent high-risk area where dugongs were known to inhabit should be closed to mesh netting.

In 2011, following public consultation, the Authority introduced amendments to the Great Barrier Reef Marine Park Regulations 1983 in relation to commercial netting rules, designating:

- the No Netting Area where no netting activities (other than small mesh bait netting) are allowed
- the Restricted Netting Area where limited lower-risk netting activities are allowed.

The large mesh nets now allowed are restricted to the type developed by the Burdekin Sustainable Fisheries Alliance. The areas within Bowling Green Bay where the rule changes apply are shown in Figure 3.5.

Since the changes were introduced, no dugong mortalities have been reported from the area.




Figure 3.5 Bowling Green Bay Special Management Area

A group of net fishers is now working with researchers to trial a new net design that may further reduce the likelihood of fatal interactions with species of conservation concern, while improving safety and maintaining fishing efficiency for target species such as barramundi. The modified net has a collapsible panel that allows large animals to push through the net rather than becoming entangled. The panels are readily refitted, therefore, minimising any impact on fishing efficiency. A preliminary study, conducted by the Moreton Bay Seafood Industry Association, provided 'proof of concept'. The net is now being tested through a research project funded by the Fisheries Research and Development Corporation.

## 3.9 Management priorities

### 3.9.1 Current priorities

To address the challenges the Reef is facing while achieving the greatest value for the available resources, the Authority's management must be well-targeted, knowledge-based, scientifically robust and measurable.

The Authority regularly reviews its management priorities and arrangements to ensure its resources are applied most effectively to achieve the long-term protection, ecologically sustainable use, understanding and enjoyment of the Great Barrier Reef. The four current priorities, and their intended effect, are listed in the *Great Barrier Reef Marine Park Authority Strategic Plan 2012–2016* (Table 3.4).

**Table 3.4 Current strategic priorities for the Authority**

The current priorities for the Authority are set out in its *Strategic Plan 2012–2016*

Strategic priorities	Intended effect
<b>Address the key risks affecting the outlook for the Great Barrier Reef</b>	<ul style="list-style-type: none"><li>• The declines in coastal ecosystem health, particularly in water quality and biodiversity, are halted and reversed; and, where time scales for this change are long, then plans are in place to achieve it as soon as possible.</li><li>• Coastal communities and Reef users understand and have capacity to adapt to pressures of climate change and prevent damaging practices threatening Reef health.</li></ul>
<b>Ensure management of the Marine Park supports ecologically sustainable use</b>	<ul style="list-style-type: none"><li>• The natural functions and resilience of the Reef's ecosystems are maintained, or if necessary restored, through effective decisions governing natural resource management and environmental protection, and working in partnership with Traditional Owners, coastal communities, Reef users and local government.</li><li>• The protective legal measures built into the <i>Great Barrier Reef Marine Park Act 1975</i> and related laws are effective, and supported by a comprehensive field management program delivered jointly with the Queensland Government.</li></ul>
<b>Encourage stewardship of the Great Barrier Reef by educating others, drawing on the best available scientific information</b>	<ul style="list-style-type: none"><li>• Management and all activities contributing to the health and resilience of the Reef are underpinned by the best available science.</li><li>• Communities and Reef users demonstrate a high level of participation in activities that contribute to the health and resilience of the Marine Park.</li></ul>
<b>Maintain a high performing, effective and efficient organisation</b>	<ul style="list-style-type: none"><li>• The Authority's workplaces are safe, positive and supportive of its staff.</li><li>• The Authority is an accountable, responsive and adaptive organisation that complies with all obligations to the government and the Australian public.</li></ul>

## 3.10 Applying the tools to achieve the priorities

The following is a summary of how the Authority applies its management tools to achieve its four current management priorities.

### 3.10.1 Addressing the key risks

The *Outlook Report 2009* identified climate change, continued declining water quality from catchment run-off, loss of coastal habitats from coastal development, and remaining impacts from fishing and illegal fishing and poaching as the priority issues reducing the resilience of the Great Barrier Reef. The Authority is working in partnership with other agencies and stakeholders to reduce the magnitude of these risks and build the resilience of the Reef ecosystem to withstand and recover from their effects.

The identified key risks principally arise beyond the boundaries of the Region, and their management is jurisdictionally complex. As a result, **partnership** arrangements (such as through the *Reef Water Quality Protection Plan*), **stewardship programs** (such as Reef Guardian programs for councils, farmers and fishers, and tourism and fishing industry actions on climate change), and **education** (such as through the Reef Guardian Schools program, Eyes and Ears Indigenous Compliance Training, Reef HQ Aquarium and the Authority's communications program) are key management tools.

The Authority also works in collaboration and plays a key advisory role in **environmental impact assessment** processes — both Commonwealth and state — for proposed developments that have the potential to affect the Great Barrier Reef environment (for example major port expansions). This work includes processes under the EPBC Act and the *State Development and Public Works Organisation Act 1971* (Qld). The Authority also provides comment and advice on catchment planning and management relevant to protecting the health of the Reef environment.

The Authority works in close collaboration with the Queensland Government on fisheries management. Of particular importance are strategies to address the remaining impacts from fishing, with an emphasis on illegal fishing and poaching, reducing incidental catch of species of conservation concern, reducing death of discarded



species, addressing issues associated with any unsustainable extraction of top predators, and fishing of fish spawning aggregations.

The **Zoning Plan** plays a key role in managing the location of fishing activities in the Region, and building the resilience of the Reef ecosystem. In addition, fishing activities must comply with Marine Park Regulations and Queensland fisheries legislation. Harvest (dive-based) fishery operations within the Marine Park require a Marine Park **permit**, as do netting activities (other than bait netting) in the Special Management Area within Princess Charlotte Bay. Any developmental fishery program will also require a permit from the Authority. The joint Field Management Program plays a key role in **compliance** activities in the Region, including in relation to fishing.

### Benefits of zoning

The Authority's Great Barrier Reef Zoning Plan 2003 substantially increased protection for biodiversity in the Marine Park; in particular, the increase in no-take zones from less than five per cent to 33 per cent of the Marine Park. The Zoning Plan set a global benchmark for marine conservation, providing protection from extractive uses for at least 20 per cent of every bioregion, while allowing for a range of ecologically sustainable uses and the continuation of traditional activities.<sup>1,5,6,7,8,9,10,11,12,13,14,15</sup>

The benefits of the Great Barrier Reef zoning were synthesised by a group of 21 leading scientists in 2010, in the international journal *Proceedings of the National Academy of Sciences*.<sup>16</sup> Their conclusions are summarised below.

*'Overall, the available evidence suggests that the large-scale network of marine reserves on the Great Barrier Reef is proving to be an excellent investment in social, economic, and environmental terms.'*<sup>16</sup>

**More, bigger fish:** The network of no-take marine reserves is providing significant, long-term benefits for the species of coral reef fish that are targeted by fishers (especially coral trout), with not only more fish, but bigger fish in reserves — some reserves having around twice as much fish biomass.<sup>16,17</sup>

**Improved fish recruitment:** Larger fish contribute disproportionately more larvae than smaller fish. Recent research in the Keppel Islands suggests increased reproduction by the more abundant, bigger fish in reserves is not just benefiting populations within those reserves, but the benefits are 'spilling over' to other reefs, including reserve and fished reefs.<sup>18</sup> Therefore, the reserve areas are having flow-on benefits to the entire ecosystem, not just the reefs within the reserve network.<sup>18</sup>

**Improved resilience:** The spillover effects also mean the connectivity between reserve reefs is intact.<sup>18</sup> Spatial analysis shows most reserve reefs are within the dispersal range of other reserve reefs, so they are able to function as a network.<sup>16</sup> Networks are more resilient than isolated components, meaning the Zoning Plan is contributing to the resilience of the overall system.

**Sharks, dugongs and turtles:** These species are harder to protect because they are slow growing and slow breeding. They are also highly mobile, moving in and out of protected zones. Despite this, available evidence shows zoning is benefiting these species, although additional protection is still needed.<sup>14, 21</sup> For effective management of mobile species, zoning needs to be complemented by measures to protect populations outside the reserve zones.<sup>16,19,20,21</sup>

**Reduced crown-of-thorns starfish outbreaks:** Outbreaks of crown-of-thorns starfish appear to be less frequent on reserve reefs than fished reefs.<sup>16,22</sup> This is particularly important as crown-of-thorns starfish have been the greatest cause of coral mortality in the Region in recent decades.<sup>23</sup>

**Zoning benefits for seabed habitats:** Zoning has improved protection of seabed habitats, with at least 20 per cent of most habitat types protected from trawling.<sup>16,24</sup>

**The benefits of zoning depend on compliance:** A relatively small amount of illegal fishing can rapidly reduce the benefits of marine reserves.<sup>16</sup> It is, therefore, important to have good enforcement and compliance to ensure the ecosystem benefits of zoning are maintained.

Partnerships with Traditional Owners — formalised in **Traditional Use of Marine Resources Agreements** — address issues such as the sustainable take of culturally significant species, and supporting Traditional Owner cultural practice in the conservation and management of the Great Barrier Reef. The agreements incorporate traditional and contemporary scientific knowledge and environmental management for the ongoing protection of the Great Barrier Reef, and are used by Traditional Owners as a formal tool to conserve and protect species and ecosystems critical to the health of people, culture and country. They can incorporate specific management strategies for the conservation and sustainable use of key species, as well as habitats. Other management activities that Traditional Owners may identify in their agreement implementation plan include restoring and maintaining waterways and coastal ecosystems; maintaining and protecting significant heritage values including sites; sharing and documenting traditional ecological knowledge, culture and language; and research and monitoring of sea country including through partnerships with managing agencies and leading scientific institutions. By accrediting Traditional Owner agreements, governments have a clear pathway towards understanding important Traditional Owner issues, and to address shared matters.

**Research** to improve understanding of the risks facing the Reef and their likely effects is supported in a range of research programs, including in the National Environmental Research Program Tropical Ecosystems Hub, the

Australian Research Council Centre of Excellence for Coral Reef Studies, the Australian Institute of Marine Science (AIMS), CSIRO, the Fisheries Research Development Corporation, the Australian Marine Mammal Centre, James Cook University, University of Queensland and the Great Barrier Reef Foundation. **Monitoring** of impacts (such as the marine monitoring program which samples water quality) and the state of the Reef ecosystem itself (such as field management monitoring surveys, the Eye on the Reef program and the AIMS long-term monitoring program) provide scientific evidence to support assessments of management effectiveness. Monitoring of social and economic drivers and activities is being developed.

**Policy** arrangements such as the *Great Barrier Reef Climate Change Adaptation Strategy and Action Plan 2012–2017* and the *Great Barrier Reef Biodiversity Conservation Strategy 2013* explain the way in which the Authority intends to manage the Marine Park or perform its other functions, and the way in which the Authority considers the Act or other legislative instruments apply. Policies are not legislative instruments.

Regular **reporting** on the condition of the Great Barrier Reef and on the factors affecting it helps track the progress being made towards reducing risks. This reporting includes five-yearly *Outlook Reports*, annual Paddock to Reef reporting by Reef Plan partners and the one-off report *Informing the outlook for Great Barrier Reef coastal ecosystems*.

### 3.10.2 Ensuring use is ecologically sustainable

The Authority has a direct management responsibility to ensure use of the Marine Park is ecologically sustainable.

The **Zoning Plan** defines the activities that can occur in various locations and provides the framework for extractive use within the Marine Park. Generally, applications are assessed and permits issued jointly with the Queensland Government. A range of uses of the Marine Park, such as tourism, infrastructure, some fisheries and some research activities, require a **permit** from the Authority. Permits enable the Australian and Queensland governments to reduce impacts on high use and sensitive areas, separate potentially conflicting activities, collect data for planning of Marine Parks and monitor potentially damaging activities. During 2011–2012, 625 permissions were granted, 36 permissions were varied and 70 were transferred to another permit holder. A permit application assessment **fee** generally applies for all commercial activities.

There is a formal **environmental impact assessment** process under the Act for evaluating the likely possible risks or impacts to the environment from a proposed activity or development. Common activities where a detailed assessment process is undertaken include, but are not limited to, the construction and operation of pontoons, jetties, moorings, pipelines and marinas as well as dredging. The assessment criteria include ecological, social and economic considerations, Traditional Owner interests, as well as current and future use of the proposed location.

As most ports in the Region are located outside the Marine Park, their activities are mostly managed by the Queensland Government, unless those activities are deemed to affect matters of national environmental significance under the EPBC Act, for which the Department of the Environment has responsibility. As part of environmental assessments under the EPBC Act, the Authority provides **specialist advice** on the impacts of development in, and adjacent to, the Marine Park to ensure consistency of decision making and environmentally sustainable outcomes. It also provides advice on assessments of fisheries under the EPBC Act.

The Zoning Plan establishes a **designated shipping area** for the Marine Park so coastal and international shipping traffic follow lower risk routes through the Reef. The Authority is a member of the North-East Shipping Management Group, which addresses the risks shipping poses to the Great Barrier Reef, Torres Strait and the Coral Sea region.

**Plans of management** for the Cairns Area, the Whitsundays and Hinchinbrook set out detailed management arrangements applying to all users of these areas. The plans focus on protecting key natural values and allowing a range of uses. **Site infrastructure** such as moorings, no-anchoring areas and transit lanes support sustainable use of popular sites within planning areas and elsewhere.

The Authority's activities are guided by a series of **policy documents** supporting sustainable use. Examples include a recreation management strategy, and policies on the management of tourism permits and on moorings.

Tourism visitors and some other users pay the **environmental management charge**. All funds received are applied directly to management including education, research, ranger patrols and policy development.

All these formal management arrangements are supported by a **compliance program** conducted under the joint Field Management Program. The Field Management Compliance Coordination Unit coordinates vessel and aerial surveillance activities across the Marine Park. A cooperative multi-agency approach allows a broad range of legislation and compliance tools to be used.

### 3.10.3 Fostering stewardship

The Authority recognises the Marine Park plays an important role in the lifestyles and livelihoods of Queenslanders. Engaging Traditional Owners, stakeholders and local communities through positive and constructive relationships is essential in successfully managing activities within the Region.

With its main office in Townsville, and other offices in Cairns, Mackay and Rockhampton, the Authority is able to closely connect with the Region's coastal communities.

The Authority places a strong emphasis on fostering **stewardship** with Marine Park users by encouraging and supporting them to play a role in protecting the area, and to achieve best practices in all their activities. Examples include the High Standard Tourism program which is a formal **partnership** with Ecotourism Australia and the tourism industry; the Eye on the Reef monitoring program predominantly with tourism operators; crown-of-thorns starfish control undertaken by the Association of Marine Park Tourism Operators; climate-related initiatives with the tourism and fishing industries; the Pro-vision Reef Stewardship Action Plan for coral and aquarium fishers, and the Reef Guardian program.

The Authority's Reef Guardian program recognises the good environmental work undertaken by communities and industries to protect the Reef. It involves a hands-on, community-based approach to make the Reef healthier and more resilient by working closely with those who rely on the Reef or live in its catchment.

A number of projects are underway to trial regional stewardship approaches; for example, a National Environmental Research Program funded project focused on the Mackay and Bowen–Burdekin Local Marine Advisory Committees.

**Traditional Use of Marine Resources Agreements** provide a mechanism for close collaboration with Traditional Owners.

A series of voluntary Responsible Reef Practices provide a **best practice** guide to recreational and tourism users on sustainable practices that can be voluntarily adopted when visiting the Marine Park, in the office and at home.

The Authority has four **Reef Advisory Committees**: Catchment and Coastal; Ecosystem; Indigenous; and Tourism and Recreation. One of their key roles is to advise the Authority on actions that can be taken to address the risks to the Marine Park identified in the *Outlook Report 2009*. They are competency-based committees comprising a cross-section of stakeholder interests.

The Authority is advised at a local level by voluntary community-based **Local Marine Advisory Committees**. Established in 1999, there are now 12 committees from Cooktown to Bundaberg. Members are appointed by the Authority and offer a wide range of perspectives. The committees enable local communities to have effective input into managing the Marine Park, while providing a forum for interest groups, government and the community to discuss issues around marine resources. They also help in communications between the public and the Authority.

**Education and communication** are recognised as essential to managing the Reef and preserving it for future generations. The Authority communicates through its website, traditional media outlets, social media, its regional offices, awareness campaigns, community events and a range of other avenues. Through Reef HQ Aquarium — the Australian Government's national education centre for the Great Barrier Reef — thousands of people each month receive messages about the Reef, risks to its resilience and what can be done to protect it. The Reef HQ Aquarium, located in Townsville and managed by the Authority, provides people of all ages and physical abilities the chance to see and gain an appreciation for a living coral reef ecosystem, what makes it special and why it needs to be protected. More than 3.5 million local, national and international visitors have visited the facility since it opened in 1987, and visitation continues to increase. Since 2009, the facility's reef videoconferencing program has showcased the living reef to more than 700 locations nationally and internationally, directly engaging more than 55,000 people.

The Authority's website is a key communication tool for the Authority — it conveys information about our work, the marine environment, health and threats to the Reef; provides news and education resources; and also conveys information that is available for consultation. On average, there are over 500,000 visits to the site each year from around the world. While online traffic slightly varies each year, about 30–40 per cent of users are return visitors. Mobile browsing, using smartphones and tablets, is increasing rapidly.

### 3.10.4 Effective and efficient organisation

The Authority recognises that being a high performing, effective and efficient organisation is central to achieving its goals and objectives. The Authority places strong emphasis on maintaining and developing effective knowledge management systems, services and processes.

The Authority's governance framework is based on the legislative requirements of the Great Barrier Reef Marine Park Act, as well as the *Financial Management and Accountability Act 1997*, the *Public Service Act 1999* and a focus on engaging the community in management of the Marine Park.

An Audit Committee has been established in compliance with Section 46 of the Financial Management and Accountability Act and Regulation 22C of the Financial Management and Accountability Regulations 1997. This committee provides independent assurance and assistance to the Chief Executive Officer and the Marine Park Authority Board on the Authority's risk, control and compliance framework, and its financial statement responsibilities.

### 3.10.5 Current resources for management

The Authority's budget for 2012–2013 was \$53.96 million. This included a direct appropriation from the Australian Government; the Australian and Queensland governments' contribution to field management; revenue from the environmental management charge; and targeted funding from the Caring for our Country program for the Reef Rescue initiative and the Ensuring the Resilience of our Reef initiative. In the past, a significant part of the Authority's budget (about 20 per cent) has come from the environmental management charge, principally from

visitors participating in a tourist activity. A downward trend in visitation to the Marine Park has resulted in a decline in revenue from the charge in recent years.

In addition to the Authority's resources, a range of Australian and Queensland government investments contribute substantially in monitoring and protecting the Great Barrier Reef, and increasing its long-term resilience.

### **Protecting the Great Barrier Reef — a priority for the nation**

A number of Australian and Queensland government investments contribute substantially to protection and management of the Region. For example, in addition to the Authority's resources, the Australian Government has provided the following resources over recent years:

- more than \$7.53 million for a new initiative to control crown-of-thorns starfish in the Great Barrier Reef Marine Park
- more than \$200 million through the Reef Rescue initiative over the five years to June 2013 to reduce the discharge of sediments, nutrients and pesticides from agricultural lands to the Great Barrier Reef lagoon. There is a government commitment to invest a further \$200 million in the Reef Rescue program to 2018.
- \$52.7 million in regional base funding to the six reef catchment natural resource management regional bodies (2008–2009 to 2012–2013)
- \$12.5 million over four years from 2013 to the Great Barrier Reef Foundation to coordinate research in areas such as reef resilience and climate change
- \$2.8 million for projects to inform the comprehensive strategic assessment and long-term sustainable development planning for the World Heritage Area and adjacent coastal zone
- \$12.4 million over four years for reef ecosystem research through the National Environmental Research Program, including \$3.68 million over four years for water quality research aimed at better understanding the drivers and impacts of water quality on the biodiversity of the Great Barrier Reef
- more than \$11.25 million for 14 projects in catchments that flow into the Great Barrier Reef lagoon for activities which restore, manage and better protect biodiversity
- \$61.9 million for the four years to December 2014 for research relevant to the Great Barrier Reef through the National Environmental Research Program Tropical Ecosystems Hub. This aims to address issues of concern for the management, conservation and sustainable use of the Great Barrier Reef, tropical rainforests including the Wet Tropics World Heritage Area, and the terrestrial and marine assets underpinning resilient communities in the Torres Strait.
- science and monitoring associated with the Australian Institute of Marine Science and the CSIRO.

The Queensland Government invests in a range of activities designed to protect the Reef. This includes:

- \$35 million a year spent on Reef water quality initiatives
- more than \$8 million a year for joint field management for the Marine Park which includes compliance, management of visitor facilities and education
- more than \$21 million a year for fisheries management, as well as an additional \$9 million for a buy out of the net fishery on the east coast of Queensland which is expected to have profitability benefits for the fishery, as well as conservation benefits
- \$12 million (state-wide) over three years for Everyone's Environment Grants which will provide funding to community-based groups to tackle environmental degradation. A number of the successful grants will directly contribute to improvements in Reef health; for example, through community Coastcare and waterway clean ups.
- \$500,000 a year to the Queensland Wetlands Program which provides a number of tools to help restore the function of important wetlands adjacent to the Reef
- about \$3 million a year to support on-ground activities through its natural resource management program in Reef catchments
- an investment of \$1 million to control crown-of-thorns starfish through the Skilling Queensland program. This program provides a range of activities including job preparation, work placement and accredited training to assist local long-term unemployed jobseekers. Work placement participants assist in small scale control of the crown-of-thorns starfish infestation on selected and popular coral reefs.



## 3.11 Involvement in other planning and management frameworks

### 3.11.1 National

A further layer to the Authority's operating environment is the broad range of national policies and programs that intersect with regulation and management of the Marine Park. The key policy areas relevant to the Marine Park relate to its major uses and pressures. These include environment protection, biodiversity conservation, fisheries management, natural resource management, coastal development, tourism and climate change. Other policy areas of relevance include customs, maritime safety, Indigenous affairs, resources and energy. Table 3.5 provides examples of how the Authority is involved in implementing or contributing to a range of national planning and management frameworks.

**Table 3.5 Great Barrier Reef Marine Park Authority involvement in national planning and management frameworks**

Portfolio area	Activities
<b>Environment</b>	<ul style="list-style-type: none"> <li><i>Environment Protection and Biodiversity Conservation Act 1999</i> — as set out in a memorandum of understanding with the then Department of Sustainability, Environment, Water, Population and Communities</li> <li><i>Reef Water Quality Protection Plan</i> — a collaborative program to ensure that run-off from broadscale land use has no detrimental impact on the health and resilience of the Great Barrier Reef by 2020</li> <li>Caring for our Country: crown-of-thorns starfish control; Ensuring the Resilience of our Reef</li> <li>Caring for our Country — Reef Rescue: Land and Sea Country Indigenous Partnerships program; marine monitoring program</li> <li>National Environmental Research Program</li> <li>National Representative System for Marine Protected Areas</li> <li>National working groups: marine biodiversity decline; national partnerships approach to the sustainable harvest of marine turtles and dugongs; national plan of action for the conservation and management of sharks; Oceans Policy Science Advisory Group</li> </ul>
<b>Tourism</b>	<ul style="list-style-type: none"> <li>National Landscapes</li> <li>National Long-term Tourism Strategy</li> </ul>
<b>Science</b>	<ul style="list-style-type: none"> <li>Australian Research Council — Linkage programs</li> <li>Australian Research Council — Centre of Excellence for Coral Reef Studies</li> <li>Integrated Marine Observing System</li> <li>Fisheries Research and Development Corporation</li> </ul>
<b>Indigenous — Closing the Gap</b>	<ul style="list-style-type: none"> <li>Caring for our Country — Reef Rescue Land and Sea Country Indigenous Partnerships Program</li> </ul>
<b>Education</b>	<ul style="list-style-type: none"> <li>Development of teaching units that cover key issues for the Reef and which meet the Australian Curriculum national requirements</li> </ul>
<b>Foreign Affairs</b>	<ul style="list-style-type: none"> <li>AusAID funding — managing Caribbean coral reefs in a changing climate; co-hosting the International Coral Reef Initiative with Belize</li> </ul>
<b>Climate Change</b>	<ul style="list-style-type: none"> <li>National Climate Change Adaptation Framework — a Council of Australian Governments initiative</li> <li>Participation in the National Climate Change Adaptation Research Facility's Marine Biodiversity and Resources Adaptation Research Network</li> </ul>
<b>Transport</b>	<ul style="list-style-type: none"> <li>Chairman of the Authority is on the board of the Australian Maritime Safety Authority</li> <li>North-East Shipping Management Group</li> </ul>

### 3.11.2 State

The Authority is involved in a range of Queensland Government initiatives relevant to the Great Barrier Reef and its catchment. Table 3.6 provides examples of how the Authority is involved in implementing or contributing to a range of Queensland Government initiatives.

**Table 3.6 Examples of Great Barrier Reef Marine Park Authority involvement in Queensland Government initiatives**

Portfolio	Activities
Environment	<ul style="list-style-type: none"> <li>Gladstone Healthy Harbour Partnership for the ongoing monitoring and improvement of Gladstone Harbour and surrounding catchments</li> <li>Queensland Wetlands Program governance group</li> <li>Reef Protection Package research and development program</li> </ul>
Fisheries	<ul style="list-style-type: none"> <li>Review of trawl management arrangements (Trawl Plan Review)</li> </ul>
Planning	<ul style="list-style-type: none"> <li>Development and implementation of state management plans — for example, Green Island, Michaelmas Cay</li> </ul>
Tourism	<ul style="list-style-type: none"> <li>Draft Queensland Ecotourism Plan</li> <li>Relevant national park and coastal marine park management plans</li> <li>Great Barrier Reef experience audits and product development</li> <li>Superyacht itineraries and information about marine park management</li> </ul>

### 3.11.3 Regional

The Authority partners with natural resource management bodies to achieve planning and management outcomes within the Great Barrier Reef catchment. Examples include the Reef Catchments' Urban Think Tank and Healthy Waterways Alliance.

### 3.11.4 Local

The Authority works collaboratively at a local level with councils, schools, researchers, Traditional Owners, stakeholders and other government agencies to achieve a wide range of management outcomes. It is represented on a wide range of formal committees such as local government groups and informal, issues-based working groups. The Authority's Local Marine Advisory Committees also play a role in information exchange between the community and the Authority.

## 3.12 Protecting matters of national environmental significance

The current management program contributes to the protection of each matter of national environmental significance. By virtue of protecting the values of the Marine Park, the associated matters of national environmental significance are also afforded protection.

A number of management tools, including the GBRMP Act and Regulations, the Zoning Plan, plans of management and the *Reef Water Quality Protection Plan 2009* afford protection to all matters of national environmental significance. Other tools, such as Dugong Protection Areas, are designed to protect specific matters of national environmental significance.

The Great Barrier Reef Intergovernmental Agreement underpins collaborative joint management arrangements between the Authority and the Queensland Government, relating to the protection and management of the Great Barrier Reef. This includes joint arrangements for field management, climate change adaptation and management of fisheries resources.

The values of the **World Heritage Area** are protected and conserved by the Authority under the GBRMP Act and the EPBC Act. In addition to its obligations to protect and conserve the environment, biodiversity and heritage values of the Region, the Authority has an obligation under Section 2A(2)(c) of the Act to 'assist in meeting Australia's international responsibilities in relation to the environment and protection of world heritage (especially Australia's responsibilities under the World Heritage Convention)'. It fulfils this obligation by providing advice to the Department of the Environment on activities (within or outside the property) likely to have a significant impact on the world heritage values of the property. The Authority's advisory role is critical, as many of the threats to the Great Barrier Reef ecosystem (such as coastal development and catchment land use practices) are the result of activities occurring outside the Region, and for which the Authority has no jurisdictional control.

Additionally, more than 99 per cent of the World Heritage Area is located within the Great Barrier Reef Marine Park for which the Authority has direct management responsibility, in collaboration with the Queensland Government and its partner agencies. Impacts to the property's outstanding universal value arising from activities within the Great Barrier Reef Region are managed through the Authority's suite of regulatory and non-regulatory management tools. The Authority seeks to reduce impacts on the property's outstanding universal value from activities occurring outside the Region through its partnership, stewardship and education programs, and in its advisory capacity as described above.

Obligations under the Convention Concerning the Protection of the World Cultural and Natural Heritage also include a duty of presentation and giving the heritage values of the property 'a function in the life of the

*community* (Articles 4 and 5). The Authority assists in meeting these obligations by working closely with the tourism industry to promote the presentation of the property's outstanding universal value. This includes the development of interpretive material, and the High Standard Tourism program associated with marine parks permits. The Authority's Reef Guardian program and its multiple-use management approach are central to ensuring the property's values are given a function in the life of the community.

The Authority also contributes to periodic reporting to the World Heritage Committee on Australia's international responsibilities for the environment and protection of the World Heritage Area.

The Great Barrier Reef was listed as a **national heritage place** in 2007. The place has the same boundary as the World Heritage Area, and its listing is based on the world heritage values identified by the World Heritage Committee. The Authority contributes to the protection and management of the national heritage place using the same tools and approaches as described above for the World Heritage Area.

The **Great Barrier Reef Marine Park** is a matter of national environmental significance, and the Authority is responsible for its protection and management under the GBRMP Act. Legislative management tools, such as the Regulations, the Zoning Plan, plans of management and permits, set out requirements for the protection of biodiversity and management of ecological sustainable use. The objects of the Act encourage community engagement in the protection and management of the Region, and there is strong focus on engagement with Traditional Owners, primarily through the Authority's Traditional Use of Marine Resources Agreement program, and stakeholders more broadly through the Authority's partnerships, education and stewardship activities, including its Reef Guardian program. These tools are important as they have the capacity to influence actions outside the Marine Park but which affect the Marine Park (such as catchment management).

The **Commonwealth marine area** includes all parts of the Great Barrier Reef Region beyond Queensland state waters (that is, greater than three nautical miles from high water) extending beyond the Region into the Torres Strait, Coral Sea and to the south of the Region. Within the Region, the Commonwealth marine area is a subset of the Marine Park. Management tools that relate to the Marine Park deliver protection to the Commonwealth marine area and its environment. Other initiatives to which the Authority is a party, such as the North-East Shipping Management Plan and Reef Plan, will provide benefits to species and habitats within the Commonwealth marine area.

All **listed migratory species** are protected under Regulations, which means they cannot be taken or interfered with unless a permit has been obtained. The policy *Managing activities that include the direct take of a protected species from the Great Barrier Reef Marine Park* informs the permitting process. In addition, the following management measures are in place to protect and manage listed migratory species:

#### **Sharks and rays**

- shark and rays spatial protection via inshore habitat conservation areas such as dugong protection areas
- vulnerability assessment for sharks and rays under the *Biodiversity Conservation Strategy 2013*.<sup>25,26</sup>

#### **Marine turtles**

- position statement on managing access to the Restricted Access Special Management Areas surrounding Raine Island, Moulter Cay and MacLennan Cay<sup>27</sup>
- *Raine Island Climate Change Adaptation Plan 2010–2070*<sup>28</sup>
- Indigenous Land and Sea Country Partnerships Program, including Traditional Use of Marine Resources Agreements and Indigenous Land Use Agreements, that can include provisions for the traditional use of marine resources
- code of practice for the sustainable management of dugong and marine turtle tourism in Australia<sup>29</sup>
- population model for the southern Great Barrier Reef green turtle stock<sup>30</sup>
- monitoring of foraging green turtles in Shoalwater Bay
- spatial and temporal closures of some islands and/or surrounding waters to protect nesting activities
- rehabilitation facilities for sick and injured marine turtles (Reef HQ Aquarium)
- vulnerability assessment for marine turtles under the *Biodiversity Conservation Strategy 2013*.<sup>26</sup>

#### **Estuarine crocodiles**

- signage warning the public of estuarine crocodile sightings, for example Low Isles.

#### **Seabirds and shorebirds**

- position statement on managing access to the Restricted Access Special Management Areas surrounding Raine Island, Moulter Cay and MacLennan Cay<sup>27</sup>
- guidelines for visitation to seabird breeding islands<sup>27</sup>
- coastal bird monitoring and information strategy
- prohibition on fishing with more than six hooks in the Great Barrier Reef Marine Park
- seasonal closures of islands and intertidal areas to protect important breeding and feeding areas
- vulnerability assessment for seabirds and shorebirds under the *Biodiversity Conservation Strategy 2013*.<sup>26</sup>

## Whales

- operational policy on whale and dolphin conservation in the Great Barrier Reef Marine Park<sup>27</sup>
- Regulations for managing vessel and aircraft interactions with whales and dolphins (Part 4A of the Regulations)
- Whale Protection Area in the Whitsundays to protect calving grounds
- vulnerability assessment for dwarf minke whales under the *Biodiversity Conservation Strategy*.

## Australian snubfin dolphin and the Indo-Pacific humpback dolphin

- operational policy on whale and dolphin conservation in the Great Barrier Reef Marine Park
- Regulations for managing vessel and aircraft interactions with whales and dolphins (Part 4A of the Regulations)
- Dugong Protection Areas provide some protection to inshore dolphin habitats
- vulnerability assessment for inshore dolphins under the *Biodiversity Conservation Strategy 2013*

## Dugongs

- position statement on the conservation of dugongs in the Great Barrier Reef Marine Park<sup>27</sup>
- Great Barrier Reef Marine Park Zoning Plan 2003 (dugongs were explicitly considered in its development)<sup>31</sup>
- Indigenous Land and Sea Country Partnerships Program — Traditional Use of Marine Resources Agreements
- Indigenous Land Use Agreements (these can include provisions for the traditional use of marine resources)
- code of practice for the sustainable management of dugong and marine turtle tourism in Australia<sup>27</sup>
- vulnerability assessment for dugongs under the *Biodiversity Conservation Strategy 2013*.<sup>26</sup>

All **listed threatened species** are protected under Great Barrier Reef Marine Park Regulations, which means they cannot be taken or interfered with, unless a permit has been obtained. Permits are not required for traditional use. The policy *Managing activities that include the direct take of a protected species from the Great Barrier Reef Marine Park* informs the permitting process. Specific management tools for listed threatened species are the same as those specified above for relevant listed migratory species.

The Shoalwater and Corio Bays Area and Bowling Green Bay are **listed wetlands of international importance** and are partly within the Region. The Authority, through its Field Management Program, undertakes dedicated management activities including patrols, engagement activities with Traditional Owners, supporting marine turtle and dugong monitoring, and monitoring shorebird populations. The Shoalwater Bay (Dugong) Plan of Management and Dugong Protection Area were designated to support the most important dugong seagrass habitat in the southern region of the Marine Park. Other management initiatives, such as Reef Plan, also indirectly address broader impacts that may affect the area.

## 3.13 Timeframe and review

The Authority has been responsible for the protection and management of the Great Barrier Reef since the introduction of the Great Barrier Reef Marine Park Act in 1975.

In recognition of the need to continually adapt its work to emerging issues, the Authority regularly reviews its management programs and priorities. An important part of the Authority's management approach is ensuring that work programs reflect current priorities, and that decision making is transparent and accountable. A number of reviews of various aspects of the Authority's business have been conducted over the past 20 years resulting in changes to organisational structure, legislative arrangements and work programs.<sup>32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50</sup>

The five-yearly production of an *Outlook Report*, including an assessment of management effectiveness, provides the foundation for cyclical business planning and decision making by the Authority.<sup>51</sup> The *Outlook Report 2014* will use the best available science and build upon the information presented in the *Outlook Report 2009* and the strategic assessment, to assess the current state of the Reef, commercial and non-commercial use, factors influencing its condition, management measures, risk, current resilience and the long-term outlook. It will form an important part of the ongoing adaptive management cycle, by reviewing and evaluating the information and outcomes of both reports and resultant management changes.



## References

1. Great Barrier Reef Marine Park Authority 2009, *Great Barrier Reef Outlook Report 2009*, GBRMPA, Townsville.
2. UNESCO 2012, *Mission report Great Barrier Reef*, UNESCO, Paris.
3. Department of the Environment 2013, *Significant impact guidelines 1.1: Matters of national environmental significance*, Commonwealth of Australia, <<http://www.environment.gov.au/resource/significant-impact-guidelines-11-matters-national-environmental-significance>>.
4. Great Barrier Reef Marine Park Authority 2011, *Traditional Owner connections to sea country*, GBRMPA, Townsville, <<http://www.gbrmpa.gov.au/our-partners/traditional-owners>>.
5. Hughes, T.P., Gunderson, L.H., Folke, C., Baird, A.H., Bellwood, D.R., Berkes, F., Crona, B., Helfgott, A., Leslie, H., Norberg, J., Nyström, M., Olsson, P., Osterblom, H., Scheffer, M., Schuttenberg, H., Steneck, R.S., Tengö, M., Troell, M., Walker, B., Wilson, J. and Worm, B. 2007, Adaptive management of the Great Barrier Reef and the Grand Canyon World Heritage Areas, *Ambio* 36(7): 586-592.
6. Fernandes, L., Day, J., Lewis, A., Slegers, S., Kerrigan, B., Breen, D., Cameron, D., Jago, B., Hall, J., Lowe, D., Innes, J., Tanzer, J., Chadwick, V., Thompson, L., Gorman, K., Simmons, M., Barnett, B., Sampson, K., De'ath, G., Mapstone, B., Marsh, H., Possingham, H., Ball, I., Ward, T., Dobbs, K., Aumend, J., Slater, D. and Stapleton, L. 2005, Establishing representative no take areas in the Great Barrier Reef: Large scale implementation of theory on marine protected areas, *Conservation Biology* 19(6): 1733-1744.
7. Forrester, T.L. 2013, *Personal communication: Lama Lama Traditional Owner clan group*.
8. McPhee, D. and Stone, S. Unpublished report, 'Assessment of by-catch reduction devices in Moreton Bay trawl and net fisheries'.
9. Marsh, H. and Lawler, I.R. 2006, *Dugong distribution and abundance on the urban coast of Queensland: A basis for management: Final report to Marine and Tropical Research Sciences Facility interim projects 2005-6*, James Cook University, Townsville.
10. Ruckelshaus, M., Klinger, T., Knowlton, N. and DeMaster, D.P. 2008, Marine ecosystem-based management in practice: Scientific and governance challenges, *Bioscience* 58(1): 53-63.
11. Pandolfi, J.M., Bradbury, R.H., Sala, E., Hughes, T.P., Bjorndal, K.A., Cooke, R.G., McArdle, D., McClenachan, L., Newman, M.J.H., Paredes, G., Warner, R.R. and Jackson, J.B.C. 2003, Global trajectories of the long-term decline of coral reef ecosystems, *Science* 301(5635): 955-958.
12. Day, J.C. 2011, Great Barrier Reef Marine Park - Governance analysis, in *Governing Marine Protected Areas: getting the balance right, Vol 2, Technical Report*, eds. P.J.S. Jones, W. Qiu and E.M. De Santo, United Nations Environment Program, Nairobi, Kenya.
13. Day, J.C. and Dobbs, K. 2013, Effective governance of a large and complex cross-jurisdictional marine protected area: Australia's Great Barrier Reef, *Marine Policy* 41: 14-24.
14. Department of Sustainability, Environment, Water, Population and Communities 2013, *State Party Report on the state of conservation of the Great Barrier Reef World Heritage Area (Australia) property ID N154 in response to World Heritage Committee decision WHC 35 COM 7B.10*, DSEWPoC, Canberra.
15. Day, J.C. 2002, Zoning - lessons from the Great Barrier Reef Marine Park, *Ocean and Coastal Management* 45(2-3): 139-156.
16. McCook, L.J., Ayling, T., Cappel, M., Choat, J.H., Evans, R.D., DeFreitas, D.M., Heupel, M., Hughes, T.P., Jones, G.P., Mapstone, B., Marsh, H., Mills, M., Molloy, F.J., Pitcher, C.R., Pressey, R.L., Russ, G.R., Sutton, S., Sweatman, H., Tobin, R., Wachenfeld, D. and Williamson, D. 2010, Adaptive management of the Great Barrier Reef: A globally significant demonstration of the benefits of networks of marine reserves, *Proceedings of the National Academy of Sciences* 107(43): 18278-18285.
17. Russ, G.R., Cheal, A.J., Dolman, A.M., Emslie, M.J., Evans, R.D., Miller, I.R., Sweatman, H. and Williamson, D.H. 2008, Rapid increase in fish numbers follows creation of world's largest marine reserve network, *Current Biology* 18(12): R514-R515.
18. Harrison, H.B., Williamson, D.H., Evans, R.D., Almany, G.R., Thorrold, S.R., Russ, G.R., Feldheim, K.A., van Herwerden, L., Planes, S., Srinivasan, M., Berumen, M.L. and Jones, G.P. 2012, Larval export from marine reserves and the recruitment benefit for fish and fisheries, *Current Biology* 22(11): 1023-1028.
19. Grech, A. and Marsh, H. 2008, Rapid assessment of risks to a mobile marine mammal in an ecosystem-scale marine protected area, *Conservation Biology* 22(3): 711-720.
20. Grech, A., Marsh, H. and Coles, R. 2008, A spatial assessment of the risk to a mobile marine mammal from bycatch, *Aquatic Conservation: Marine and Freshwater Ecosystems* 18(7): 1127-1139.
21. Dryden, J., Grech, A., Moloney, J. and Hamann, M. 2008, Rezoning the Great Barrier Reef World Heritage Area: does it afford greater protection for marine turtles, *Wildlife Research* 35: 477-485.
22. Sweatman, H. 2008, No-take reserves protect coral reefs from predatory starfish, *Current Biology* 18(14): R598-R599.
23. Osborne, K., Dolman, A.M., Burgess, S.C. and Johns, K.A. 2011, Disturbance and the dynamics of coral cover on the Great Barrier Reef (1995-2009), *PLoS ONE* 6(3): e17516.
24. Pitcher, C.R., Venables, B., Browne, M., Doherty, P. and De'ath, G. 2007, *Indicators of protection levels for seabed habitats, species and assemblages on the continental shelf of the Great Barrier Reef World Heritage Area*, Reef and Rainforest Research Centre, Cairns.
25. Great Barrier Reef Marine Park Authority 2013a, *Great Barrier Reef Biodiversity Conservation Strategy 2013*, GBRMPA, Townsville.
26. Great Barrier Reef Marine Park Authority 2012, *Vulnerability assessments*, GBRMPA, Townsville, <<http://www.gbrmpa.gov.au/about-the-reef/biodiversity/draft-biodiversity-conservation-strategy/vulnerability-assessments>>.
27. Great Barrier Reef Marine Park Authority 2013b, *Policies and position statements*, GBRMPA, Townsville, <<http://www.gbrmpa.gov.au/about-us/legislation-regulations-and-policies/policies-and-position-statements>>.
28. Great Barrier Reef Marine Park Authority, *Raine Island Climate change adaptation plan*, Unpublished draft.
29. Birtles, A., Curnock, M., Dobbs, K., Smyth, D., Arnold, P., Marsh, H., Valentine, P., Limpus, C., Hyams, W., Dunstan, A., Charles, D., Gatley, C., Mangott, A., Miller, D., Hodgson, A., Emerick, S. and Kendrick, A. 2005, 'Code of practice for the sustainable management of dugong and marine turtle tourism in Australia'.
30. Chaloupka, M.Y. 2003, *Phase 2 - Development of a Population Model for the Southern Great Barrier Reef Green Turtle Stock*, Research Publication No. 81 edn, Great Barrier Reef Marine Park Authority, Townsville.
31. Great Barrier Reef Marine Park Authority 2004, *Great Barrier Reef Marine Park Zoning Plan 2003*, GBRMPA, Townsville.
32. Australian Department of Environment, Sport and Territories 1995, *Review of funding arrangement for the Great Barrier Reef Marine Park Authority*, Great Barrier Reef Marine Park Authority, Townsville.
33. Management and Technology Consulting 1995, *Review of organisational arrangements in the Great Barrier Reef Marine Park Authority*, Townsville.
34. Whitehouse, J.F. 1993, *Managing multiple use in the coastal zone: a review of the Great Barrier Reef Marine Park Authority*, Australian Government Publishing Service, Canberra.
35. Australian Government 1998, *Commonwealth management of the Great Barrier Reef, The Auditor-General Audit Report No. 33*, Australian National Audit Office, Canberra.

36. Department of the Environment and Heritage 2006, *Review of the Great Barrier Reef Marine Park Act 1975, Review Panel Report*, Commonwealth of Australia, Canberra.
37. Hockings, M. and Gilligan, B. 2009, *Assessment of management effectiveness for the 2009 Great Barrier Reef Outlook Report*, Great Barrier Reef Marine Park Authority, Townsville.
38. Dalton, V. 2003, *Day-to-day management review*, Great Barrier Reef Marine Park Authority, Townsville.
39. Brown, R. 1997, *Independent review of the Great Barrier Reef Marine Park Authority: report submitted to the Hon. Sen. Robert Hill, Minister for the Environment*, R. Brown and Associates, Brisbane.
40. Whitehouse, J.F. 1992, *Review of the Magnetic Island Marina development*, Australian Government Publishing Service, Canberra.
41. Gilmour, A., Cunningham, J. and Lamond, D. 1991, *Day to day management of the Great Barrier Reef Marine Park: the role of Marine Park staff*, Macquarie University, Sydney.
42. Deloitte Touche Tohmatsu Ltd 1994, *Great Barrier Reef Aquarium management options*, Department of Environment, Sport and Heritage, Canberra.
43. Great Barrier Reef Marine Park Authority and Queensland Department of Environment 1998, *Day-to-day Management Coordination Unit evaluation and relocation: report and recommendations*, GBRMPA/QDOE, Townsville.
44. Great Barrier Reef Shipping Review Steering Committee 2001, *Review of ship safety and pollution prevention measures in the Great Barrier Reef*, Australian Maritime Safety Authority, Canberra.
45. Productivity Commission 2003, *Industries, land use and water quality in the Great Barrier Reef catchment: research report*, Productivity Commission, Canberra.
46. Futureye Pty Ltd, Teh-White, K., Houston, S., Baxter, C., Levine, J. and White, P. 2005, *Great Barrier Reef Marine Park Authority: Enhanced Community Partnerships*, Futureye Pty Ltd, Melbourne.
47. Australian Government 2003, *Commonwealth management of the Great Barrier Reef follow-up audit, The Auditor-General Audit Report No. 8*, Australian National Audit Office, Canberra.
48. Stanley, O. and Hansen, G. 1994, *Review of the Environmental Management Charge*, Great Barrier Reef Marine Park Authority, Townsville.
49. Tourism Review Steering Committee 1997, *Review of the marine tourism industry in the Great Barrier Reef World Heritage Area*, Great Barrier Reef Marine Park Authority, Townsville.
50. Sturgess, G.L. 1999, *The Great Barrier Reef Partnership: cooperation in the management of a World Heritage Area; a report into the review of the relationships of the Commonwealth and Queensland Governments in respect of the Great Barrier Reef*, Queensland Government, Brisbane.
51. Dobbs, K., Day, J., Skeat, H., Baldwin, J., Molloy, F., McCook, L.J., Johnson, M., Elliot, B., Skeat, A., Vohland, K., Wachenfeld, D. and Kenchington, R. 2011, *Developing a long-term outlook for the Great Barrier Reef, Australia: A framework for adaptive management reporting underpinning an ecosystem-based management approach*, *Marine Policy* 35(2): 233-240.



An aerial photograph of a coral reef system. The water is a vibrant turquoise color, with darker patches indicating deeper areas or different reef structures. Several small, sandy islands are visible, some with sparse vegetation. The reef extends from the foreground towards the horizon, where the water meets a clear blue sky.

## Chapter 4

### **Matters of national environmental significance in the Great Barrier Reef Region**



4.7.1	Marine turtles .....	4-33
4.7.2	Marine mammals.....	4-33
4.7.3	Sharks and rays .....	4-33
4.7.4	Seabirds .....	4-34
<b>4.8</b>	<b>Wetlands of international importance.....</b>	<b>4-34</b>
4.8.1	Shoalwater and Corio Bays Area .....	4-34
4.8.2	Bowling Green Bay Area .....	4-35
<b>4.9</b>	<b>Environmental processes .....</b>	<b>4-36</b>
<b>4.10</b>	<b>Realising connectivity in the landscape .....</b>	<b>4-41</b>
<b>4.11</b>	<b>Connections between matters of national environmental significance .....</b>	<b>4-42</b>
<b>4.12</b>	<b>Summary of conclusions .....</b>	<b>4-46</b>
	<b>References.....</b>	<b>4-47</b>

## Table of contents

<b>4.1</b>	<b>What are matters of national environmental significance?.....</b>	<b>4-3</b>
<b>4.2</b>	<b>World heritage properties.....</b>	<b>4-6</b>
4.2.1	Outstanding universal value .....	4-6
<b>4.3</b>	<b>The Great Barrier Reef Marine Park.....</b>	<b>4-8</b>
4.3.1	Biodiversity .....	4-9
4.3.2	Geomorphological features .....	4-14
4.3.3	Indigenous and historic heritage values .....	4-18
4.3.4	Community benefits of the environment.....	4-21
<b>4.4</b>	<b>National heritage places .....</b>	<b>4-23</b>
<b>4.5</b>	<b>Commonwealth marine areas .....</b>	<b>4-23</b>
<b>4.6</b>	<b>Listed migratory species .....</b>	<b>4-25</b>
4.6.1	Marine turtles .....	4-25
4.6.2	Estuarine crocodiles .....	4-28
4.6.3	Whales.....	4-28
4.6.4	Dolphins.....	4-28
4.6.5	Dugongs .....	4-29
4.6.6	Sharks.....	4-29
4.6.7	Seabirds.....	4-30
4.6.8	Shorebirds .....	4-32
<b>4.7</b>	<b>Listed threatened species and ecological communities.....</b>	<b>4-32</b>

Cover page image: Aerial view of Stanley Reef



**Extract from Great Barrier Reef Region Strategic Assessment terms of reference**

**2.1 Description of matters of national environmental significance**

2.1.1 Describe the extent to which the following relevant matters of national environmental significance, as defined in the Environment Protection and Biodiversity Conservation Act 1999 apply to the strategic assessment area:

- a) world heritage properties, including consideration of the outstanding universal value of the Great Barrier Reef World Heritage Area
- b) national heritage places
- c) wetlands of international importance
- d) listed threatened species and ecological communities
- e) listed migratory species
- f) Commonwealth marine areas
- g) the Great Barrier Reef Marine Park.

The description must:

- h) identify key terrestrial, coastal and marine environmental, biodiversity and heritage values and/or attributes which underpin the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area
- i) describe ecosystem processes considered critical to the functioning of the relevant matters of national environmental significance, including the Outstanding Universal Value of the Great Barrier Reef World Heritage Area
- j) provide sufficient information to allow an understanding of the connectivity between the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area.

## **4.1 What are matters of national environmental significance?**

The Great Barrier Reef is recognised both nationally and internationally as an important natural area, worthy of protection. Since the 1970s, the Australian Government has progressively recognised and protected the Great Barrier Reef as one of Australia's most precious ecosystems (Figure 4.1). This commenced with the introduction of specific legislation in 1975 and its nomination as a World Heritage Area in 1981. Throughout the 1980s, 1990s and early 2000s, progressive declaration of sections of the Great Barrier Reef Marine Park (the Marine Park) put in place arrangements for the long-term protection of the ecosystem.

Introduction of the *Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act) brought a new national focus to protecting and managing Australia's most prized environmental assets. These are referred to as matters of national environmental significance. In 2009, the Marine Park was included as a matter of national environmental significance.

Seven matters of national environmental significance are relevant to the Great Barrier Reef Region (the Region) and therefore considered in this assessment:

- world heritage properties
- the Great Barrier Reef Marine Park
- national heritage places
- Commonwealth marine areas
- listed migratory species
- listed threatened species and ecological communities
- wetlands of international importance.

All of the matters of national environmental significance are interconnected; each a part of the rich and complex Great Barrier Reef environment (Figure 4.2).

# OUR INTERNATIONAL AND NATIONAL OBLIGATIONS TO PROTECT THE GREAT BARRIER REEF

Over almost half a century, the Australian Government has worked to recognise and protect the remarkable attributes of the Great Barrier Reef. This diagram shows how the international concepts of world heritage and outstanding universal value, as well as the declaration of the Great Barrier Reef Marine Park and other matters of national environmental significance relate to the Great Barrier Reef.





Figure 4.1 Timeline for the matters of national environmental significance relevant to the Great Barrier Reef Region

# MATTERS OF NATIONAL ENVIRONMENTAL SIGNIFICANCE

Matters of national environmental significance are Australia's environmental assets that are recognised as being of importance to the nation as a whole. They are defined and protected under the Australian Government's national environmental law, the *Environment Protection and Biodiversity Conservation Act 1999*. Seven are relevant to the Great Barrier Reef.




**Figure 4.2 Matters of national environmental significance in the Great Barrier Reef Region**

In addition, many of these matters overlap spatially (Figure 4.3). For example, the Great Barrier Reef Marine Park makes up almost all (about 99 per cent) of the Great Barrier Reef World Heritage Area. Listed threatened and listed migratory species range throughout the Region, with some also present in the adjacent coastal zone or Commonwealth marine areas beyond the Region.

In this chapter, the extent to which each matter of national environmental significance applies to the Region is described in detail. A common set of key values and attributes relevant to the matters is presented. In addition, the environmental processes relevant to the matters are outlined.





**Figure 4.3 Spatial extent of the matters of national environmental significance for the Great Barrier Reef Region**

The matters of national environmental significance overlap across the Region and some extend well beyond its boundary.

## 4.2 World heritage properties

The Great Barrier Reef is a world heritage property encompassing both Commonwealth and Queensland areas seaward of low water mark (see Figure 1.1, Chapter 1). Another world heritage property, the Wet Tropics of Queensland, is adjacent to the Great Barrier Reef World Heritage Area. There are close natural and cultural connections between the two properties.

### 4.2.1 Outstanding universal value

The Great Barrier Reef World Heritage Area covers 348,000 square kilometres and includes both marine areas and islands. There are 1050 islands including some 600 continental islands, 300 coral cays and 150 mangrove islands. It is inscribed on the World Heritage List because of its natural outstanding universal value.

*Outstanding universal value is defined as cultural and/or natural significance which is so exceptional as to transcend national boundaries and to be of common importance for present and future generations of all humanity<sup>1</sup>*

Outstanding universal value is central to the way the World Heritage Convention is implemented. A property is considered 'to be of outstanding universal value' if it meets one or more of ten world heritage criteria and is inscribed on the World Heritage List. In addition, to be deemed to be of outstanding universal value 'a property must also meet the conditions of integrity and/or authenticity and must have an adequate protection and management system to ensure its safeguarding'.<sup>1</sup>

The United Nations Educational, Scientific and Cultural Organization's (UNESCO's) Operational Guidelines<sup>1</sup> recognise that 'no area is totally pristine and that all natural areas are in a dynamic state and to some extent involve contact with people' and that human activities 'may be consistent with the outstanding universal value of the area where they are ecologically sustainable'.

Recognition of the Great Barrier Reef's outstanding universal value was based on the natural world heritage criteria in place at the time — acknowledging the Reef's natural values, together with the strong ongoing links between Aboriginal and Torres Strait Islander peoples and their sea country. The criteria have been amended and renumbered since the Reef was inscribed. The criteria at the time of listing and the equivalent current criteria are presented in Table 4.1.

Only those values, or attributes, that are consistent with the four criteria for which the Great Barrier Reef was inscribed can be considered to be its world heritage values. However, given the broad scope of the criteria under which it was listed, almost all aspects of the Reef's environment contribute to its outstanding universal value. This includes the Region's biodiversity, geomorphological features, and aspects of Traditional Owner connections to the area, its environmental processes, its superlative natural phenomena and its aesthetic value. The notable exception is historic heritage values (for example, shipwrecks and light stations) which are not encompassed by the natural criteria.



**Table 4.1 World heritage criteria relevant to the Great Barrier Reef**

Short title	Criteria at time of listing	Current criteria
<b>Major stages of the Earth's evolutionary history</b>	(i) outstanding examples representing the major stages of the Earth's evolutionary history	(viii) be outstanding examples representing major stages of the Earth's history, including the record of life, significant ongoing geological processes in the development of landforms, or significant geomorphic or physiographic features
<b>Ecological and biological processes</b>	(ii) outstanding examples representing significant ongoing geological processes, biological evolution and man's interaction with his natural environment	(ix) be outstanding examples representing significant ongoing ecological and biological processes in the evolution and development of terrestrial, fresh water, coastal and marine ecosystems and communities of plants and animals
<b>Natural beauty and phenomena</b>	(iii) unique, rare or superlative natural phenomena, formations or features or areas of exceptional natural beauty, such as superlative examples of the most important ecosystems to man	(vii) contain superlative natural phenomena or areas of exceptional natural beauty and aesthetic importance
<b>Habitats for conservation of biodiversity</b>	(iv) habitats where populations of rare or endangered species of plants and animals still survive	(x) contain the most important and significant natural habitats for <i>in situ</i> conservation of biological diversity, including those containing threatened species of outstanding universal value from the point of view of science or conservation

The attributes that contribute to the property's outstanding universal value are interconnected and are distributed across the entire extent of the property.

A Statement of the Outstanding Universal Value of the Great Barrier Reef World Heritage Area<sup>2</sup> is the official statement adopted by the World Heritage Committee outlining how the property met the criteria for outstanding universal value, integrity and protection and management at the time of listing. The statement was prepared retrospectively. In line with International Union for Conservation of Nature (IUCN) guidelines<sup>3</sup>, the statement is based on information that was available in 1981 and addresses the criteria in place at the time of inscription. It does, however, adopt the contemporary numbering for the relevant criteria.

The statement provides guidance on the world heritage attributes of the property. It is reproduced in full at Appendix 3 and is summarised below.

#### **Criteria (i) now (viii) — major stages of the Earth's evolutionary history**

The Great Barrier Reef, extending 2300 kilometres along Queensland's coast, is a globally outstanding representation of the major stages of the Earth's evolutionary history. The area has been exposed and flooded by at least four glacial and interglacial cycles and, over the past 18,000 years, reefs have grown on the continental shelf. During glacial periods, sea levels dropped, extending the coastline further east and exposing the reefs as flat-topped hills of eroded limestone with large rivers meandering between them. During interglacial periods, rising sea levels caused the formation of continental islands and coral cays, and new phases of coral growth. Today's varied seascapes and landscapes have been moulded by changing climates and sea levels, and the erosive power of wind and water, over long time periods. A recent report *Geological and geomorphological features of outstanding universal value in the Great Barrier Reef World Heritage Area*<sup>4</sup> identifies the geological and geomorphological features of outstanding universal value, and provides an overview of the pressures affecting values. The outcomes are summarised in Section 4.3.2.

#### **Criteria (ii) now (ix) — ecological and biological processes**

The ecological and biological processes of the Region reflect the maturity of an ecosystem that has evolved over millennia. Globally significant marine fauna groups include more than 4000 species of molluscs, more than 1625 species of fish, as well as a great diversity of sponges, anemones, marine worms, crustaceans, and many other taxonomic groups. The establishment of vegetation on the cays and continental islands exemplifies the important role of birds in processes such as seed dispersal and plant colonisation. Biodiversity attributes and environmental processes which support the Great Barrier Reef are described in more detail in Section 4.9 and Section 4.10. Human interaction with the natural environment is illustrated by strong ongoing links between Aboriginal and Torres Strait Islander peoples and their land and sea country, and includes numerous shell deposits (middens), fish traps, story places and marine totems. A description of Indigenous heritage values is provided in Section 4.3.3.

#### **Criteria (iii) now (vii) — natural beauty and natural phenomena**

The Great Barrier Reef demonstrates superlative natural beauty above and below the water, providing spectacular scenery. It is one of a few living structures visible from space, appearing as a complex string of reefs along Australia's north-east coast. From the air, the vast mosaic patterns of reefs, islands and coral cays produce an unparalleled aerial panorama. The Whitsunday Islands provide a magnificent vista of green vegetated islands and spectacular sandy beaches. Hinchinbrook Island supports vast mangrove forests, rugged vegetated

mountains and lush rainforest gullies. The Reef's natural phenomena include annual coral spawning, migrating whales, nesting turtles, and significant spawning aggregations of many fish species.

#### **Criteria (iv) now (x) — habitats for conservation of biodiversity**

The enormous size and diversity of the Region means it is one of the richest and most complex natural ecosystems on Earth, and one of the most significant for biodiversity conservation. The diversity of habitats supports tens of thousands of marine and terrestrial species, many of which are of global conservation significance. For example, the waters of the Region provide major feeding grounds for one of the world's largest populations of the threatened dugong, and it is an important area for humpback whale calving. Furthermore, six of the world's seven species of marine turtle occur in the Region, with internationally important breeding grounds for green, loggerhead and hawksbill turtles. A more detailed description of Great Barrier Reef species and habitats is provided in Sections 4.3.1, 4.6, 4.7 and 4.8.

#### **Integrity**

At the time of inscription it was considered that to include virtually the entire Great Barrier Reef within the property was the only way to ensure the integrity of the coral reefs in all their diversity. The Statement of Outstanding Universal Value<sup>2</sup> also notes that some of the key ecological, physical and chemical processes that are essential for the long-term conservation of the marine and island ecosystems and their associated biodiversity occur outside the boundaries of the property. A description of environmental processes which support and connect the Great Barrier Reef environment is provided in Section 4.9.

#### **A better understanding of aesthetics**

In 2012, a World Heritage Centre/IUCN monitoring mission visited the Great Barrier Reef to assess the state of conservation of the World Heritage Area and to contribute to the strategic assessment process. The mission report noted the property's aesthetic values are less well understood than other aspects, and recommended that '*further work is needed in relation to identifying and documenting the attributes related to the aesthetic values of the property*'.<sup>5</sup> The committee's finding is consistent with the 1997 review of the outstanding universal value of the Great Barrier Reef.<sup>6</sup>

In response to the recommendations of the monitoring mission and the World Heritage Committee, the Australian Government commissioned a report to consider the aesthetic values of the World Heritage Area.

The report<sup>7</sup> broadly defines aesthetic values to include not only visual elements, but also experiential elements, such as responses of the community to a place. It is also important to note that the relevant world heritage criterion includes two different concepts: aesthetic beauty and superlative natural phenomena. The latter describes large and spectacular natural events such as large aggregations of single species. In the report, both concepts are analysed with respect to the attributes that define them, as well as their levels of sensitivity. The attributes are mapped conceptually, rather than geographically, conveying the interconnections between all elements within and adjacent to the property.

The report lists the environmental and experiential attributes against each of the elements in the Statement of Outstanding Universal Value for the property. The environmental attributes identified are: the Reef as an entity; coral reefs; continental islands; beaches; coral cays; water clarity (calmness, intensity of colour); marine animals (abundance, diversity, colour, size); blue holes; lagoon floors; mangroves; seagrass meadows; shoals; cliffs and rocky shores; bays; estuaries; rainforest; birds; and butterflies. The experiential attributes identified are: beauty; naturalness; tranquillity; solitude; remoteness; discovery; and inspiration.

This approach provides a different lens for considering impacts on aesthetic attributes, and thus provides the potential for recommended changes or additions to current management practices in order to ensure the elements that make up the relevant world heritage criterion are adequately protected.

### **4.3 The Great Barrier Reef Marine Park**

The Region includes the Great Barrier Reef Marine Park (the Marine Park) — one of the seven relevant matters of national environmental significance. The Region and the Marine Park cover the same area, with the exception of 13 coastal exclusion areas that are not within the Marine Park (see Figure 1.1). The Marine Park covers 344,400 square kilometres and includes the subsoil beneath the seabed extending to a depth of 1000 metres and the airspace above extending to a height of 915 metres. It does not include Queensland islands and internal waters.

In accordance with the *Great Barrier Reef Marine Park Act 1975*, the Marine Park is managed to protect and conserve the biodiversity, heritage values and environment of the Region. The term 'environment' is defined broadly and, as well as ecosystems, natural resources and heritage values, it encompasses physical resources — the area's geomorphological and geological aspects; and people and communities — social, economic and cultural aspects.

The Marine Park is a multiple-use area allowing ecologically sustainable use.

As a result, there is a very wide range of values relevant to this matter of national environmental significance. The values of the Marine Park are grouped into four categories:

- biodiversity, including habitats and species
- geomorphological features
- Indigenous and historic heritage
- community benefits of the environment, comprising the cultural, social and economic benefits derived from the Region's environment.

The following is a description of the values relevant to this matter of national environmental significance.

#### 4.3.1 Biodiversity

Biodiversity is the variety of life on Earth. It includes all living things and the way they interact with each other and their environment.

The Great Barrier Reef is one of the world's most diverse and remarkable ecosystems, with a wide range of habitats and many thousands of different species recorded.

##### Habitats

The range of habitats in the Marine Park can be grouped into eleven broad types, based on those defined in *Great Barrier Reef Outlook Report 2009*<sup>9</sup>:

- islands
- beaches and coastlines
- mangrove forests
- seagrass meadows
- coral reefs (less than 30 metres deep)
- deeper reefs (greater than 30 metres deep)
- lagoon floor
- shoals
- *Halimeda* banks
- continental slope
- open waters.

Within each of these there is great variation, with 70 different biological regions recognised — 30 within the coral reefs and 40 in surrounding areas (Figure 4.4 and Figure 4.5).

Each habitat is briefly described below.

**Islands** are an important component of the Great Barrier Reef ecosystem. Continental islands and mangrove islands are located relatively close to the coast, whereas cays are mostly further offshore and are associated with reefs. Several species of terrestrial plants and animals are endemic to Great Barrier Reef islands (such as *Pisonia* forests).<sup>10</sup> While there are more than 1050 islands within the outer boundary of the Marine Park, only 70 (those declared as Commonwealth islands) are part of the Marine Park. The remainder are under Queensland Government jurisdiction.



*Russell Island, a continental island in the Marine Park between Cairns and Innisfail*

The **beaches and coastlines** within the Marine Park stretch approximately 2300 kilometres along the mainland coast of Queensland and on the islands. Sandy shores typically occur on the exposed coastline, on islands and on reefs (as coral cays). They are the habitats that support a wide range of species including as nesting grounds for shorebirds, seabirds and marine turtles. Muddy shores are generally adjacent to river mouths and estuaries in areas sheltered from prevailing winds. They act as depositional areas for sediments and nutrients discharged from the catchment or transported along the coast. Rocky coasts are intermittently distributed, providing hard structure which resists the erosive forces of wind and waves and provides habitat for many sessile species such as oysters.

**Mangrove forests** are an intertidal habitat of trees and shrubs covering an estimated 2070 square kilometres along the coast of the Marine Park. They grow in sheltered areas where fine sediments accumulate and where there is inundation by seawater during the tidal cycle. The mangrove forests along the Great Barrier Reef coast are very diverse, with at least 39 mangrove species and hybrids recorded.<sup>11,12,13</sup> The highest diversity is in the far north of the Marine Park.<sup>13</sup> Mangroves provide essential structure and habitat for a range of terrestrial, marine and intertidal species and play a critical ecosystem role as sources of primary production; carbon sequestration; nursery and breeding sites for many species; filtering of suspended sediments from the water; and as physical barriers from storms and extreme weather events.<sup>14</sup> The mangrove forests along the Great Barrier Reef coast are an integral part of the Great Barrier Reef ecosystem.




*Mangrove forests along the Great Barrier Reef Marine Park coast*

**Seagrass meadows** are distributed widely throughout the Marine Park, especially in the protected sediment-covered areas.<sup>14,15</sup> They colonise a broad range of areas — rivers, inlets, reefs and intertidal, subtidal and deepwater areas.<sup>15,16</sup> Based on the limited surveys undertaken, shallow water seagrasses (less than 15 metres deep) are estimated to cover approximately 5700 square kilometres, and deepwater seagrasses (deeper than 15 metres) are estimated to cover up to 40,000 square kilometres, although at depths below 15 metres they generally become very sparse (less than five per cent cover).<sup>15,17,18,19,20,21</sup> Fifteen species of seagrass occur within the seagrass meadows of the Great Barrier Reef.<sup>20,22,23</sup> The meadows are habitat for a diverse range of species. They are primary producers, stabilise bottom sediments and are a store for carbon.<sup>24</sup>


**Coral reefs** are one of the most biologically diverse ecosystems on the planet. The Great Barrier Reef is the world's largest coral reef ecosystem, ranging over 14 degrees in latitude, comprising more than 2900 separate coral reefs and containing more than 400 species of hard coral. Coral reef habitats cover an area of 26,000 square kilometres — about seven per cent of the Marine Park. They can be broadly classified into inner, mid and outer shelf platform reefs and can be made up of different types of reef. Reef types, including fringing, inshore turbid, shelf, ribbon, deltaic, northern detached and submerged coral reefs, represent major stages in the Earth's evolutionary history and are examples of the Region's outstanding universal value.<sup>4</sup> They also provide vitally important habitat for a wide diversity of plants and animals. In addition to their biological value, coral reefs deliver a range of ecosystem services including shoreline protection, and provide community benefits from, for example, fisheries and tourism.





**Figure 4.4 Reef bioregions of the Great Barrier Reef**

Thirty reef bioregions have been identified in the Great Barrier Reef Marine Park. Each represents an area with distinct groups of plants and animals and physical features.



**Figure 4.5 Non-reef bioregions of the Great Barrier Reef**

Forty non-reef bioregions have been identified in the Great Barrier Reef Marine Park. These cover all the areas between and beyond reef areas. Each represents an area with distinct groups of plants and animals and physical features.

The **deeper reefs** that occur in the Region are poorly understood compared to their shallow water counterparts. Mesophotic coral reefs are characterised by the presence of light-dependent corals occurring in areas where there is limited light for photosynthesis (between 30 and 150 metres).<sup>25,26</sup> Modelling indicates mesophotic reefs are likely to be widespread along the Great Barrier Reef shelf edge and may constitute a high proportion of the total coral reef area in the Marine Park.<sup>27</sup> In fact, about 60 per cent (25,600 square kilometres) of the seabed where coral reefs are likely to grow is potential deep reef habitat compared to about 40 per cent (16,110 square kilometres) which is suitable for shallower corals.<sup>28</sup> In the northern part of the Marine Park, the steeply sloping drop-off at the edge of the continental shelf means there is limited area for mesophotic reefs, and they are generally shallower than 70 metres. Further south, where reefs are set back from the shelf edge, submerged reefs have developed on the shoulder of the continental shelf to depths of about 150 metres. Deeper, cold water coral reefs typically occur in depths where light does not penetrate and temperature is fairly constant at four degrees Celsius.<sup>29</sup> They can be hotspots of biodiversity on the deep seabed.<sup>30</sup>

The **lagoon floor** habitat makes up approximately 210,000 square kilometres of the Marine Park. It covers the area between reefs and the coast and has an average depth of between 20 and 40 metres. Scientists have catalogued 5300 species of marine organisms from non-reef habitats in the Region.<sup>31</sup> The lagoon habitat is variable and includes some outcrops of marine life forming clumps that rise above the seafloor with sponges, sea-whips, gorgonians (sea fans) and interreef gardens.<sup>32</sup> Crinoids (feather stars) are sometimes extremely abundant in offshore areas. The lagoon floor supports many species, such as nematodes and microbial communities, which are the basis of many food chains and are important elements of a healthy functioning ecosystem. Larger organisms use the lagoon floor for food and shelter and as a nursery habitat. These include shellfish, crabs, prawns, sea urchins, sea stars and sea cucumbers, sponges, worms and fish (including sharks and rays). The lagoon floor is an important habitat for commercial, traditional and recreational fishers.

**Shoals** cover approximately 278 square kilometres of the Marine Park. They are submerged features on the seafloor that attract and support many fish and other species in areas away from obvious emergent coral reefs.<sup>33,34,35</sup> They may be unconsolidated sediments of sand, rubble, rock or reef substrate, or structurally more complex and diverse communities of filter feeders and corals. Plants and animals that use the habitat include gorgonians, sponges, algae, macroalgae and seagrasses.

**Halimeda banks** comprise large areas of the far northern Great Barrier Reef, inshore of the Ribbon Reefs, and also in the central northern Great Barrier Reef.<sup>31</sup> They are dominated by a live veneer of the calcareous green algae *Halimeda* which when it dies forms banks, typically up to 20 metres thick, on the seafloor usually in waters 40 metres or more deep. This habitat is poorly studied.<sup>9</sup>

The **continental slope** is a complex area, composed of relic reefs, landslides, canyons and plateaux, that extends down to more than 2000 metres.<sup>36,37</sup> It comprises approximately 15 per cent of the Marine Park. There has been little investigation of this remote habitat or the deepwater seabed habitats beyond.<sup>9</sup>

**Open water** habitat is critical to the healthy functioning of the whole Great Barrier Reef ecosystem. In particular, it provides connectivity between other habitats, from the coast to beyond the continental slope. Open water is dominated by microorganisms (plankton) and supports a range of other plants and animals such as invertebrates, fish, reptiles and marine mammals.

In addition to these habitats within the Marine Park, some **terrestrial habitats** in the coastal zone adjacent to the Region directly support the biodiversity of the Marine Park (for example the trapping of nutrients and sediments in wetlands). These habitats are described below.<sup>38</sup>

- **Saltmarshes** are an important, highly productive, interface between the marine and terrestrial environments, providing feeding and breeding areas for many marine species.
- **Freshwater wetlands** slow the overland flow of water; cycle nutrients and sediments; are important dry season refugia for many species; and are used by some marine species for parts of their life cycle.
- **Forested floodplains** help slow, capture and recycle nutrients and sediments; protect the soil surface from the erosive forces of rain; and are important nursery areas for many species with connections to the Great Barrier Reef.
- **Heath and shrublands** help slow the overland flow of water; prevent erosion; recycle nutrients and sediments; and are important as buffers on steep coastal hill slopes.
- **Grass and sedgelands** are used for feeding and roosting by migratory birds; help slow the overland flow of water; and capture nutrients and sediments.
- **Woodlands** reduce flooding by slowing the overland flow of water, and regulate sediments and nutrient supply to the Great Barrier Reef.
- **Forests** contribute to the hydrological cycle through evapotranspiration, cloud formation and rainfall generation.
- **Rainforests** minimise soil loss from erosion, including binding and stabilising soils, and provide habitat for species that also use Great Barrier Reef islands.
- **Connecting waterbodies**, such as rivers, creeks, estuaries and floodplains, are the aquatic link between the marine and terrestrial environment, transporting water, nutrients and sediments, and providing a movement corridor, as well as feeding and breeding areas for some marine and coastal species. Aquatic connectivity is provided through surface waterbodies and groundwater.



## Species

Nearly all groups of marine plants and animals are abundantly represented in the Great Barrier Reef, with thousands of different species recorded. They range from microscopic plankton to massive whales and include benthic organisms (those that live on the seafloor) such as corals and seagrasses and pelagic organisms (those that live in the water column) such as larvae and some fishes. Some of the species in the Marine Park are listed nationally as migratory or threatened (Sections 4.6 and 4.7).

Table 4.2 is a summary of the key groups of Great Barrier Reef species. It is recognised that there are many new species yet to be discovered and named.

**Table 4.2 Plants and animals of the Great Barrier Reef**

Thousands of species make up the Great Barrier Reef ecosystem. Nowhere near all of them have been identified and described. For some, the number of species recorded is provided; for others the most up-to-date estimate is given.<sup>39,40</sup>

Plants and animals of the Great Barrier Reef	Number of species recorded
Whales and dolphins	more than 30
Dugongs	1
Sharks and rays	136
Bony fish	1625
Crocodiles	1
Marine turtles	6
Sea snakes	14 breeding species
Seabirds	20 nesting species
Echinoderms	630
Crustaceans	about 1300
Molluscs	as many as 3000
Worms	at least 500
Hard corals	411
Soft corals and sea pens	at least 150
Sponges	at least 2500
<b>Mangroves</b>	39
Seagrass	15
Marine macroalgae	630

### 4.3.2 Geomorphological features


The geomorphological features within the Region are closely linked to its biodiversity. Their diversity provides structure for the diversity of habitats and species, and some features, such as coral reefs and *Halimeda* banks, are biologically constructed. Geomorphological features can be broadly grouped into coral reefs, islands and shorelines, channels and canyons, river deltas, *Halimeda* banks and seagrass meadows. The following descriptions are summarised from Whiteway *et al.* 2013<sup>4</sup>.

**Coral reefs** are an important geomorphological feature of the Marine Park as well as foundational for its biodiversity values. They take on many forms (Figure 4.6) depending on past environmental conditions and stages of development. Their diversity of shape and size can be a function of substrate size, depth and, to a lesser extent, relative sea level history and carbonate productivity.<sup>41,42</sup>

Fringing reefs grow along the coasts of the mainland or continental high islands.<sup>43</sup> The Great Barrier Reef includes many fringing reefs (758 of the 2904 named reefs), but they are generally small with an average area of less than one square kilometre. Together they comprise just 350 square kilometres or 1.8 per cent of the total reef area.<sup>44</sup> Fringing reefs are more common around continental high islands than along the mainland coast, with 352 fringing reefs concentrated in the Whitsunday, Cumberland and Northumberland island groups.<sup>44</sup>

Inshore turbid zone reefs are typically located in shallow water (less than 10 metres) and usually within 10 kilometres of the coast. They occur in locations where reef development is affected by continual or repeated land-based sediment inputs, elevated turbidity and fluctuating salinities.





**Figure 4.6 Three-dimensional bathymetry of the Great Barrier Reef shelf**

Image courtesy of Dr Robin Beaman ([www.deeppreef.org](http://www.deeppreef.org))

Shelf reefs include a range of reef types in the mid-shelf area, including irregular reef patches, crescentic reefs, lagoonal reefs and planar reefs.

Ribbon reefs are barrier reefs on the steepest northern edge of the continental shelf. They are distributed over approximately 700 kilometres of the shelf edge between Cooktown and the Torres Strait, with individual ribbon reefs up to 28 kilometres long. Seaward of the ribbon reefs, the shelf quickly drops to considerable depth, reaching 1000 metres within a kilometre. Narrow passages, typically less than 800 metres wide, separate the reefs and allow water exchange between the Great Barrier Reef lagoon and the open ocean, including nutrient-rich waters upwelled from deeper waters off the shelf edge. The flows through these narrow channels are relatively strong. These geomorphological features, coupled with currents, provide conditions conducive to the formation of *Halimeda* banks.

Deltaic reefs are shelf-edge reefs with a triangular shape. Examples of deltaic reefs can be found in the northernmost 100 kilometres of the Great Barrier Reef shelf edge, where short deltaic reefs (less than four kilometres long) occur parallel to the shelf edge, separated by passages up to 200 metres wide and 35 metres deep. Strong currents are funnelled through the narrow passages between these reefs, transporting sediments into quieter back reef areas.<sup>45</sup> These deposits can then be colonised by reef communities.

The northern detached reefs are shelf-edge reefs that are separated from the main shelf edge by deep channels that may exceed 300 metres and be as much as six to seven kilometres wide.<sup>44,46</sup> They form on isolated pinnacles of continental crust, with different reef sizes and shapes depending on the substrate depth and geometry.<sup>44</sup>

Submerged reefs are 'drowned' reefs that formed during periods of lower sea level (in glacial periods), which currently have little or no upward coral growth.<sup>47,48</sup> They are most often found in waters from 20 to 120 metres deep (but can be deeper), with examples occurring both on the shelf and along the shelf edge. Submerged reefs include mesophotic and deeper water reefs and can take a variety of forms such as submerged barrier reefs, reef terraces, and isolated coral reef pinnacles and knolls.<sup>47</sup> Of the 2904 named reefs, almost one fifth (566 reefs) are classified as submerged.<sup>49</sup>

**Islands and shorelines** are distributed throughout the Great Barrier Reef. Carbonate reef islands (coral cays) are deposits of carbonate sediments mostly made up of the skeletal remains of reef organisms that have been deposited by waves and currents. There are about 300 coral cays throughout the Reef. They have formed on reef platforms after the mid-Holocene and are thus relatively young geomorphological features. They are geomorphologically diverse, varying in size, shape, sediment composition, location on the reef platform, elevation, age structure, the extent of consolidation or cementation, and the extent to which they have been colonised by vegetation. Vegetated coral cays tend to be more stable than those with no vegetation. Unvegetated cays are very dynamic and, in some cases, can be short-lived. Therefore, the number of reef islands can vary over time. It has been estimated that there are about 213 unvegetated coral cays and only 43 vegetated cays, with an additional 44 islands classified as low wooded islands.<sup>4</sup>



*Heron Island, a vegetated coral cay*

Mangroves occur in the intertidal environments predominantly along the continental coastline and have also colonised shorelines of many islands and reef platforms, including along the shorelines of high islands such as Hinchinbrook Island and Orpheus Island. As well as a feature of biodiversity, they play a role in the geomorphology of the Reef. The depositional environments associated with mangroves usually result in organic muds, often including the shells of various molluscs. Mangrove dominated shorelines on high islands occur throughout the Great Barrier Reef. They are best developed in sheltered locations such as embayments or island coasts away from the prevailing wind,<sup>50</sup> and behind emergent reef flats. Mangrove islands (fully covered by mangroves), in contrast, are rare in the Region, with Murdoch Island being the only example.


The area contains more than 600 continental high islands — outcrops isolated from the mainland coast during the last major sea level rise.<sup>44</sup> Some are within the Marine Park, most are Queensland islands. The continental high islands exhibit a range of shapes, sizes and rock types similar to that observed along the mainland coast. Examples of islands dominated by granite include Lizard Island, Hinchinbrook Island, Magnetic Island and the Palm Island and Whitsunday groups. Dunk Island comprises granite intrusions in basement metamorphic rocks. Islands underlain by sandstone are less common, the Flinders Island Group being an example.

Some islands, such as Lady Elliot Island, preserve sequences of gravel and shingle ridges that run parallel to the shore and grow seaward. These ridges are formed as coral fragments are transported onshore by waves and overwash during storm activity, with individual ridges formed during multiple events.<sup>51,52,53,54,55</sup> They can provide information about the frequency and severity of past storm events.

**Channels and canyons** are some of the lesser known geomorphological features of the Great Barrier Reef.

Palaeochannels are past river channels that have been inundated by sea level rise. They include the channel depression and the associated fluvial sedimentary features deposited by the former river, such as deposits and in-filled channels.


Karstic channels are shaped by the dissolution of layers of soluble bedrock, usually carbonate rock such as limestone composed of reef carbonates. In the Region, they are developed on reefs exposed during periods of low sea level (glacial periods). Landforms include steep slopes and gorges (for example the deep steep-sided passages between Hook and Hardy Reefs).<sup>41,56</sup> 'Blue holes' — deep circular depressions with steep sides that are thought to be submerged sinkholes — are also a karstic landform<sup>57</sup> and are rare both in the Region and globally. There are just three identified examples of blue holes within the Region, at Molar Reef, Cockatoo Reef and an unnamed reef, all located in the Pompey Complex east of Mackay. The blue hole at the unnamed reef is the deepest of the three, with an explored depth of 90 metres. It is one of the best examples in the world.<sup>56</sup>



*The Blue Hole at Cockatoo Reef in the Pompey Complex, east of Mackay*



Recent improvements in bathymetric mapping have revealed extensive and large submarine canyons, associated landslides and turbidite deposits within the Region (see Figure 4.6). Submarine canyons are important for sediment transport and hosting deepwater ecosystems. They can modify shelf-edge oceanography to produce upwellings.<sup>58</sup> They also preserve information about sea level change, as well as sediment and tectonic movements.<sup>59,60,61</sup> For example, recent research has identified areas of the northern shelf edge prone to collapse and possibly capable of generating a large tsunami.<sup>62</sup> Around the Ribbon Reefs, most submarine canyons are between five and 20 kilometres long, have average widths ranging between 900 and 8000 metres, and maximum incision depths of between 144 and 815 metres.<sup>58,63</sup> The largest submarine canyon is Bligh Canyon, extending beyond the continental shelf and into the Coral Sea Basin (Figure 4.7). It is more than 200 kilometres long, almost 10 kilometres wide, and has incised as much as 300 metres into the seabed.<sup>64</sup> It begins approximately 200 kilometres east of the Lockhart River mouth and extends out across the Region and into the Coral Sea Marine Reserve. Turbidite deposits form at the base of the continental shelf and are the result of sediment transport from the shelf, including underwater landslides, and are responsible for distributing vast amounts of sediment into the deep ocean. They also provide historical records about sedimentation in the adjacent shelf area, tectonic movements and responses to sea level.<sup>65</sup>



**Figure 4.7 Three-dimensional bathymetry of submarine canyons along the Great Barrier Reef shelf edge**  
Image courtesy of Dr Robin Beaman ([www.deeppreef.org](http://www.deeppreef.org))

Approximately 30 major rivers and several hundred smaller, often short-lived, streams drain into the Great Barrier Reef lagoon.<sup>66</sup> Many of these rivers and streams transport significant loads of terrestrial sediment, including bedload sediments (medium sands and coarser) that can accumulate to form **river deltas**. The combination of relatively high sediment yields and relatively low wave energy (except during cyclones), due to the protection of the outer barrier reefs, means sizeable deltas have formed at the mouths of many rivers, especially the Barron, Herbert, Burdekin and Fitzroy rivers.

**Halimeda banks** are formed by *Halimeda*, a genus of benthic green algae which produces calcified green deposits in a plant-like form.<sup>67</sup> It is believed *Halimeda* began growing on the Great Barrier Reef earlier than corals and, in some locations, Holocene *Halimeda* banks overlie older Pleistocene *Halimeda* deposits.<sup>68</sup>

**Seagrass meadows** are classed as a geomorphic feature because they stabilise the seabed by decreasing current velocity, thus allowing suspended sediment to settle to the seafloor.<sup>69</sup> Seagrass roots then stabilise the sediment. Seagrass meadows are also important as they support a range of carbonate secreting organisms that grow on their surface which contribute to the carbonate sediments within the Region. Seagrass meadows are present in water depths up to 60 metres, but particularly occur in areas with high sediment and nutrient availability, such as close to the mainland.<sup>69,70</sup>

### 4.3.3 Indigenous and historic heritage values

In 2008, the importance of the Great Barrier Reef's heritage values was formally recognised by an addition to the main object of the Great Barrier Reef Marine Park Act that required their long-term protection and conservation.

In line with the definition in the Great Barrier Reef Marine Park Act and the EPBC Act, the heritage values of the Marine Park include:

*'its natural and cultural environment having aesthetic, historic, scientific or social significance, or other significance, for current and future generations of Australians'.*

For the Great Barrier Reef Marine Park, the natural heritage referred to in the definition is encompassed by the biodiversity and geomorphological values discussed previously.

The Indigenous and historic components include both tangible heritage (such as places) and intangible heritage (such as experiences, traditions, historic events and stories). The values relevant to these two categories are discussed below.

#### Indigenous heritage


Indigenous heritage recognises the heritage of Aboriginal and Torres Strait Islander peoples who are the First Australians and Traditional Owners of the Great Barrier Reef. While Aboriginal and Torres Strait Islander peoples have some common heritage values, there are also many unique expressions of heritage between Aboriginal and Torres Strait Islander peoples. In addition, differences are also found between families, clans, groups and communities.

Indigenous heritage is a unique, dynamic and diverse living heritage. It continuously links generations of Aboriginal and Torres Strait Islander peoples through time and is an important part of their cultures.<sup>71</sup>

For Indigenous peoples, relationships with country, people, beliefs, knowledge, lore, language, totems, symbols, ways of living, sea, land, sky and objects all arise from a spiritual belief, and cultural practice and association.<sup>71</sup>

Heritage is a central element in Indigenous custom, and its conservation ensures continued respect for Indigenous ancestors and the ancestral beings who shaped the land and waterways.<sup>72</sup>

There are more than 70 Aboriginal and Torres Strait Islander Traditional Owner clan groups who hold heritage values for their land and sea country in and adjacent to the Region. These include cultural, spiritual, economic, social or physical values (or a mixture of these) and demonstrate continuing connections with the Great Barrier Reef and its natural resources. An example of the types of values Traditional Owners report is provided in Figure 4.8.



**Figure 4.8 Dominant values identified by Rockhampton Traditional Owner participants at a strategic assessment workshop**

Values raised most often are represented in larger text.

For Aboriginal and Torres Strait Islander peoples, the significance of individual features is derived from their interrelatedness within the cultural landscape.

*Cultural landscapes represent the combined works of nature and humankind, they express a long, intimate and diverse relationship between people and their natural environment.*<sup>73</sup>



Throughout this report, Indigenous heritage values are combined into four broad categories:

- cultural practices, observances, customs and lore
- sacred sites, sites of particular significance, places important for cultural tradition
- stories, songlines, totems and languages
- Indigenous structures, technology, tools and archaeology.

While this report presents information on heritage values under discrete headings, in reality Indigenous values cannot be separated. In addition, the natural environment described above is fundamental to Traditional Owners and their connection to land and sea country. All aspects are connected and interrelated, and information described for any one particular aspect should be viewed in this context.

**Cultural practices, observances, customs and lore** are comprised of intangible features such as skills, folklore, rituals, religious beliefs and intellectual traditions — passed down from generation to generation. Indigenous heritage is an intrinsic part of custom and continues to be observed by Indigenous people in sea country management. For example, controlling use of and access to sea country estates by Traditional Owners and non-Traditional Owners regulates resource use based on cultural practices and belief systems. It is consistent with the recognition of traditional customs, practices and lore. Seasonal and cultural use of marine resources and the opening or closing of harvesting seasons according to ecological events (for example, flowering of particular plants or the arrival of migratory bird species) continue to be practiced by Great Barrier Reef Traditional Owners.<sup>74</sup>

Important skills and traditional ecological knowledge are passed down from one generation to the next, enabling Aboriginal and Torres Strait Islander peoples to follow the same seasonal patterns as their ancestors. The passing on of these skills and knowledge means different resources are exploited at different times of the year. For hunting and fishing, this depends on which species are 'fat' at a given time.<sup>75</sup> Hunters and fishermen also use their skills and experience to determine if individuals have suitable body condition for use.

Traditional ecological knowledge is more than understanding availability of seasonal resources; the knowledge involved can fulfil multiple purposes. The Yuibera Traditional Owners, near Mackay, use the head of the cycad plant as a vegetable when ripe; however, the plant itself also signifies the presence of water which has implications for fire regimes.<sup>76</sup>

**Sacred sites, sites of particular significance and places important for cultural tradition** are tangible aspects of the Region's Indigenous heritage. Sacred sites are significant heritage places for Indigenous peoples and their enduring traditions. As an example, they may be creation or resting places for ancestral spirits, places that contain healing water and medicinal plants, burial grounds, traditional tracks of Aboriginal people's movements or sites associated with special events.<sup>77</sup>

Many are areas of great importance for the conservation of biodiversity across land and sea country, and many communities are unable to separate the reasons for protecting the spiritual connections between people and the Earth from the reasons for conserving this biodiversity. In most cases, natural and cultural heritage values of sites are inextricably entwined. They form a continuum rather than being separate entities.<sup>78</sup>

'Sea country' is not additional to clan estate on land — it is inseparable from it. As on land, sea country contains evidence of events that occurred during the Dreaming by which all geographic features, animals, plants and people were created. Sacred sites often relate to creation events, Dreaming tracks or songlines along which spiritual beings travelled during the creation period. A defining feature of sacred natural sites is that Aboriginal people have known about and cared for them since time immemorial. Aboriginal and Torres Strait Islander peoples have custodial responsibilities as part of their lore which tie them to country, thereby ensuring the maintenance of spiritual, cultural, biological and other values of such sites.<sup>78</sup>

Many groups have unique maritime and coastal cultures that have evolved over thousands of years. Worrungu Bay (near Cape Upstart) is a significant area for the Juru people. The bay is a women's meeting area. Juru women would collect shellfish from the swamp and walk to the sand dunes area and cook them on the fire.<sup>79</sup>

Cape Hillsborough National Park is known for containing burial grounds of the Yuibera clan and is a sacred place for Indigenous spirits. Mangrove areas along the coast are still used for men's ceremonies in the early wet season, and the eastern area of the beach contains fish traps that can only be seen during a very low tide.<sup>76</sup> There are also fish traps on Hinchinbrook Island, where they operate as a system of stone-walled pools that flood at high tide, trapping fish as the tide goes out.

Expressive social activities such as **stories, songlines, totems and languages**, as well as music and dance, are part of Indigenous heritage and everyday life and are an integral aspect of ceremonies.<sup>74</sup> For Aboriginal and Torres Strait Islander peoples all that is sacred is in the land, water, air and sea. Knowledge of the environment, places or objects may be passed down the generations through these mediums. Stories and songs, dance, dress, art and language connect people to a place or time. They provide history, learning and perspective.

Lizard Island is a sacred place, known as Jiigurru (or Dyiigurra), to the Dingaal (or Dingiil) Aboriginal people of North Queensland. During the Dingaal Dreamtime, the Lizard group of islands was formed. The island group is thought to be a stingray, with Jiigurru being the body and the other islands forming the tail.<sup>80</sup>

Some Aboriginal and Torres Strait Islander peoples can be identified by their totems, which can be any animal, plant, tree or object such as birds, marine turtles, sharks, crocodiles, fish, cordyline plant, pandanus tree, sand

dunes and wind. The diamond stingray (Yama) is the totem of the Wuthathi tribe (Shelbourne Bay, North Queensland) and many other sea creatures are totem emblems for Aboriginal and Torres Strait Islander peoples.<sup>80</sup> Totems are an important part of cultural identity, and they can be used in song, dance and music and on cultural implements. Some Aboriginal and Torres Strait Islander peoples' customs forbid eating the animal that is their totem, while others make exceptions for special occasions such as ceremonies. For some, their totems are their favoured form of sustenance.

**Indigenous structures, technology, tools and archaeology** are perhaps the more well-known and documented features in Indigenous heritage literature.

Archaeological sites can include:

- occupation (residential) sites — areas containing stone tools, food remains, ochre, charcoal, cooking stones and shells
- middens — deposits of food refuse, usually shellfish
- grinding grooves — evidence of tool making or food processing found on flat sections of rock
- rock art (which often tells Dreaming stories, and sometimes provides pictorial evidence of past rituals central to the lives of Traditional Owners)
- scarred trees as a result of bark being removed for food or to make canoes, water containers, shields or huts.

Tools and implements reflect the geographic location of each group and their trading interactions with other groups. There are distinct differences in the materials used in implements for daily activities (such as hunting, cooking and collecting) as people used resources that were available to them.

Archeological evidence documents the trade links between coastal and hinterland Aboriginal people. Nywaigi people have found many stone axes and grinding stones that people have been using for thousands of years. The stones were traded with neighbouring tribes from the mountains where the stones originated.<sup>79</sup> Nomadic Aboriginal people of the Whitsunday Islands, the Ngabo, built sturdy three-piece bark canoes that were capable of open sea journeys.<sup>81</sup>

Ancient rock art sites provide an insight into the chronicle heritage of Indigenous peoples, while oral histories transferred through time deliver traditional knowledge and understanding about tools or technology. Bindal Traditional Owners from the Townsville area have documented the presence of important rock art at Cape Ferguson. There are drawings of circles believed to represent shields and also bora rings and initiation.<sup>79</sup>

Mandingablbay Yidinji Traditional Owners, like other coastal Aboriginal groups in the Cairns area, developed a wide range of technologies from local material used for hunting, fighting, making substantial shelters, baskets, fish traps and tools. Single outrigger canoes were used extensively for fishing, hunting and travelling well into the twentieth century.<sup>81</sup>

### Historic heritage

For the purpose of this assessment, historic heritage relates to the occupation and use of an area since the arrival of European and other migrants. It illustrates the way in which the many cultures of Australian people have modified, shaped and created the cultural environment. By its very nature, it will continue to evolve to represent the flow of history and changing community perceptions.<sup>82</sup>

While some specific aspects of the Region's historic heritage have been well documented, knowledge of many historic places or events is limited. The following is a summary of the broad range of the Marine Park's historic heritage.

There are more than 470 **historic shipwrecks** (that is, shipwrecks greater than 75 years old) in the Marine Park.

The **World War II features and sites** of the Great Barrier Reef include shipwrecks, more than 200 aircraft wrecks, unexploded ordnances and structures on islands.

The **lightstations**, comprising the lighthouse, accommodation and other infrastructure, servicing the Marine Park are associated with the rich shipping and navigational history of the Great Barrier Reef. There is a range of lightstations along the Reef on Commonwealth and Queensland islands, including Commonwealth and state heritage-listed lightstations built in the 1800s, 'concrete tower' aids to navigation dating from between the 1920s and the 1960s, and 'steel frame' aids to navigation.

The continuing scientific importance of the Great Barrier Reef means **places of scientific significance** are of particular heritage value. They include research stations and expedition sites such as those of the 1928 Yonge expedition to Low Isles — the first detailed scientific study of a coral reef in the world.

**Places of historic significance** include Endeavour Reef where Captain Cook ran aground, and routes taken by Cook, Flinders and Bligh on their voyages through the Great Barrier Reef. Islands within the Region have also played a major role in the heritage of the Reef. Places of historic significance include those connected with Edmund Banfield on Dunk Island and Mrs Watson's cottage on Lizard Island. Places that illustrate changes in use of the Great Barrier Reef are also significant — from early guano mining on islands, through dugong and turtle factories, to limestone and granite mining, and oil exploratory leases.

**Places of social significance** are also part of the Marine Park's historic heritage, including iconic sites such as the Green Island underwater observatory — reputed to be one of the first underwater observatories in the world; the Cod Hole — an iconic dive site; John Brewer Reef — the site of the first floating hotel; and Whitehaven Beach — a spectacular and highly visited white sand beach in the Whitsundays.

#### 4.3.4 Community benefits of the environment

The strategic assessment explicitly considers the community benefits derived from the environment. This recognises that the Marine Park is a multiple-use marine protected area as well as the interconnectedness of people and their environment, as reflected in the definition of the environment in both the GBRMP Act and the EPBC Act:

*'Environment includes ecosystems and their constituent parts, including people and communities; natural and physical resources; the qualities and characteristics of locations, places and areas; heritage values of places; and the social, economic and cultural aspects of the above.'*

In 2005, the Millennium Ecosystem Assessment provided clear evidence of the links between healthy ecosystems and human wellbeing. It showed that at the most fundamental level, humans need functioning ecosystems for clean water, fresh air, food, shelter and climate regulation.<sup>83</sup> People also derive less tangible benefits from healthy ecosystems such as nature appreciation, opportunities for relaxation and enjoyment, and a better understanding of the complex world in which we live. Four types of ecosystem services were described by the Millennium Ecosystem Assessment: *supporting* (for example, food webs and habitats); *regulating* (for example, shoreline protection and climate regulation); *provisioning* (for example, seafood and pharmaceuticals) and *cultural* (for example, aesthetics and recreation).<sup>83</sup>

The Reef environment contributes much to the community's wellbeing, both locally and more indirectly throughout Australia and the world. The Reef's biodiversity, geomorphological features, heritage values and natural beauty supports people economically, provides them with food, protection and enriches their lives.

Benefits vary according to people's cultural connections, experiences, personal perspectives, and dependence upon and familiarity with the Region. Traditional Owners have long highlighted the benefits that their communities derive from the Region's environment, including access to the Reef's resources, employment and improved health outcomes.

*"Our concerns about the health of our people are directly connected to the ability of our people to access our traditional country. Being healthy means looking after our spiritual health and our physical health."* Girringun Aboriginal Corporation<sup>84</sup>

The Region's values considered important by the community were explored in a series of workshops and surveys with stakeholders and Traditional Owners in the latter half of 2012 (see Section 2.11), as well as through discussions with the Authority's advisory committees and through one-on-one talks. The range of information, views and experiences greatly assisted in determining the full set of values for the Region. In addition, the process strengthened a common understanding of the broad range of values that the Region supports and their current condition and trend. The values that consistently emerged from the Traditional Owner and stakeholder workshops (see Appendix 5) are presented in Table 4.3.

For the strategic assessment, concepts involved in human wellbeing have been adapted to group the cultural, social and economic benefits derived from the environment into the following broad elements: income and employment; access to Reef resources; understanding, appreciation and enjoyment; personal connection; health benefits; and aesthetics.

Reef-dependent activities such as commercial fishing, marine tourism and the Reef-based recreation industry generate **income and employment** for thousands of people both within and outside the Reef catchment, as the flow-on benefits reach far beyond the boundaries of the World Heritage Area. These industries play an important role in regional Queensland and rely on a healthy Reef ecosystem for long-term economic stability.<sup>85</sup> Tangible benefits are measured in terms of the financial contribution they make to the Queensland and Australian economies.

**Access to Great Barrier Reef resources** by Reef-dependent industries, Traditional Owners, recreational users and others enables users to visit the Reef and to derive benefit from it. An important consideration in accessing the Reef is the need to balance the requirements of the variety of uses.

**Understanding, appreciation and enjoyment** help people use the Reef's resources wisely. Understanding comes from learning — from reading, listening, sharing knowledge, observing or from direct experience. It may also develop through employment; place of residence; recreational experiences; family, personal or spiritual connections; or remotely through social media. Stewardship activities, together with scientific knowledge and cultural knowledge gained from stories passed from one generation to the next, provide a context for understanding the Reef and its values. It also allows reflection on what the Reef may have been like in the past; how it contributed to human wellbeing, and how it has responded to human activities. This understanding can provide direction for future management and use. Knowledge held by Traditional Owners, tourism operators, farmers, fishers, scientists, managers and catchment residents can promote widespread understanding of the Great Barrier Reef at local, national and international levels.

**Aesthetics** are also closely related to the way in which people value and enjoy the Reef. Attributes of aesthetic qualities emphasise psychological, social or cultural dimensions and may be determined by how people travel to a place, what they do when they get there, and what experiences they have. Places that are easy to access are less likely to provide opportunities for enjoying nature, solitude or tranquillity, but may enhance opportunities for socialising and personal comfort. Psychological, social or cultural dimensions of aesthetics also include a sense of history, a sense of place, inspiration, spiritual connections, opportunities for learning, relaxation, recreation and escapism.<sup>7</sup>

**Health benefits** of natural ecosystems are now well recognised in the literature and through initiatives like *Healthy Parks, Healthy People* which explore the many ways in which nature significantly contributes to human health and wellbeing.<sup>86</sup> Reef visitors and coastal residents benefit from relaxation and stress reduction through Reef-based recreational activities; healthy inputs to diets from freshly caught local seafood; and exercise from snorkelling, boating and island and beach walks.

**Personal connection** is derived from peoples' spiritual connections, cultural ties, employment, stewardship activities, places of residence and recreational activities. It links each individual stakeholder, visitor, local resident and Traditional Owner to the Great Barrier Reef environment.

**Table 4.3 Values consistently identified in Traditional Owner and stakeholder workshops, 2012**

Values	Aspects consistently identified
<b>Specific sites and habitats</b>	
Islands, beaches and coastlines; estuaries, deep water, bays, inlets and coral reefs	<p><b>Natural heritage:</b> obligation to have it there for our children</p> <p><b>Cultural:</b> significance for Traditional Owners (e.g. locations of fish traps, sites for traditional use of marine resources and ceremonies); other cultural heritage values (e.g. locations of shipwrecks, lighthouses, sites of Captain Cook's landings)</p> <p><b>Economic:</b> support commercial fishing; shipping (deep water); tourism and recreation</p> <p><b>Social:</b> support recreation (e.g. walking, camping, snorkelling, diving, fishing, reef-walking, wildlife watching, relaxation, spending time with family and friends); education; health; lifestyle; stewardship; traditional use</p> <p><b>Aesthetic:</b> reefs and islands can be seen from space; place of natural wonder; spectacular pristine beauty; awesome; spiritual; majestic and calming; looked upon with pride; unique habitats</p>
Wetlands	<p><b>Natural heritage:</b> important breeding and feeding grounds for a variety of fish species, migratory birds and other animals</p> <p><b>Cultural:</b> as part of Traditional Owner belief systems, wetlands contain physical materials such as medicines and food sources</p> <p><b>Economic:</b> support commercial fishing and tourism</p> <p><b>Social:</b> support recreation (e.g. fishing, wildlife watching, education)</p>
Seagrass meadows	<p><b>Natural heritage:</b> important breeding and feeding grounds for a variety of fish species, turtles, dugong and other animals</p> <p><b>Cultural:</b> support dugong and turtle populations and continued cultural expression for Traditional Owners</p> <p><b>Economic:</b> support commercial fishing and tourism</p> <p><b>Social:</b> support recreation (e.g. snorkelling, stewardship, education)</p>
<b>Species</b>	
Fishes, estuarine crocodiles, birds, whales, dolphins, dugongs, sharks, rays, sea snakes, marine turtles, corals	<p><b>Natural heritage:</b> having such a large protected area for species to move through without risk</p> <p><b>Cultural:</b> traditional hunting of some species and totemic and spiritual significance for Traditional Owners</p> <p><b>Economic:</b> support commercial fishing and tourism</p> <p><b>Social:</b> support recreation (e.g. snorkelling, diving, wildlife watching, education)</p> <p><b>Aesthetic:</b> 'wow' factor</p>
<b>Ecosystem processes</b>	
Connectivity	<p><b>Natural heritage:</b> linking biophysical processes, supporting biodiversity</p> <p><b>Cultural:</b> connection of people to land and sea through stories; also in terms of the way Indigenous people moved seasonally from inland areas to the sea in search of food. Connectivity occurs from west to east (i.e. catchment to sea) and from north to south (through travel and trade routes).</p>
Integrity	<p><b>Natural heritage:</b> linking biophysical processes; supporting biodiversity</p> <p><b>Cultural:</b> accessing and maintaining sites for traditional use of marine resources, ceremonies and stories</p>
Spawning (coral, fish)	<p><b>Natural heritage:</b> contributes to biodiversity</p> <p><b>Economic:</b> support commercial fishing and tourism</p>
Water quality	<p><b>Economic:</b> support commercial fishing and tourism</p> <p><b>Social:</b> support human health; recreation (e.g. swimming, snorkelling, diving)</p> <p><b>Aesthetic:</b> water colour and clarity</p>



## 4.4 National heritage places

The Great Barrier Reef World Heritage Area was one of 15 world heritage properties included as a national heritage place on the National Heritage List in 2007. The place has the same boundary as the World Heritage Area (see Figure 1.1, Chapter 1). While there are specific criteria that apply to the listing of national heritage places, the national heritage listing of the world heritage properties was done on the basis of those values identified by the World Heritage Committee. Therefore, for the purposes of this assessment, the values of the Great Barrier Reef national heritage place are taken to correspond to the world heritage criteria. As a result, the two matters of national environmental significance are addressed in a similar way in the report with the same set of values, or attributes, underpinning them.

The national heritage criteria identified as corresponding to those for which the property was world heritage listed are:

- the place has outstanding heritage value to the nation because of the place's importance in the course, or pattern, of Australia's natural or cultural history
- the place has outstanding heritage value to the nation because of the place's possession of uncommon, rare or endangered aspects of Australia's natural or cultural history
- the place has outstanding heritage value to the nation because of the place's potential to yield information that will contribute to an understanding of Australia's natural or cultural history
- the place has outstanding heritage value to the nation because of the place's importance in demonstrating the principal characteristics of:
  - a class of Australia's natural or cultural places; or
  - a class of Australia's natural or cultural environments
- the place has outstanding heritage value to the nation because of the place's importance in exhibiting particular aesthetic characteristics valued by a community or cultural group.

## 4.5 Commonwealth marine areas

All parts of the Great Barrier Reef Region beyond state waters (that is, greater than three nautical miles from the territorial sea baseline) are a matter of national environmental significance as a Commonwealth marine area (Figure 4.9).

The Commonwealth marine area also extends beyond the Region into the Torres Strait, Coral Sea and to the south of the Region. The environmental processes within the Region are connected with these areas, as are some of the Region's species. In addition, some users of the Region (for example, tourism operators and commercial, charter and recreational fishers) also use Commonwealth marine areas adjacent to the Region.

Within the Region, the Commonwealth marine area is a subset of the Marine Park. Protection of the environment is a common aspect shared by the Marine Park and the Commonwealth marine area. Management arrangements for the Commonwealth marine area within the Region are effectively the same as those for the Marine Park. Management of Commonwealth marine areas outside the Region is the responsibility of the Director of National Parks as part of the Commonwealth marine reserves network.

Because of these commonalities, the Great Barrier Reef Marine Park (Section 4.3) is generally used in this report as a surrogate for the Commonwealth marine area. Similar values are relevant to both areas.




Figure 4.9 Boundaries of the Commonwealth marine area within and outside the Region

## 4.6 Listed migratory species

The Great Barrier Reef Region supports 77 of the currently listed migratory species comprising six marine turtle species; 11 mammal species including the dugong and two inshore dolphins; five species of shark; 54 species of shorebirds and seabirds; and the estuarine crocodile (Table 4.4).

The fact that these species move during their life histories — sometimes very large distances — means they spend much of their time outside the Region and, hence, may be exposed to impacts well beyond the boundaries of the Region or even the nation.

**Table 4.4 Listed migratory species in the Great Barrier Reef Region**

The species marked with the symbol - <sup>T</sup> - are also listed threatened species (Table 4.7). The scientific name is presented in *italics*.

Species group	Migratory species
<b>Marine turtles</b>	Flatback turtle <sup>T</sup> ( <i>Natator depressus</i> ) Hawksbill turtle <sup>T</sup> ( <i>Eretmochelys imbricata</i> ) Leatherback turtle <sup>T</sup> ( <i>Dermochelys coriacea</i> ) Green turtle <sup>T</sup> ( <i>Chelonia mydas</i> ) Loggerhead turtle <sup>T</sup> ( <i>Caretta caretta</i> ) Olive ridley turtle <sup>T</sup> ( <i>Lepidochelys olivacea</i> )
<b>Crocodiles</b>	Estuarine crocodile ( <i>Crocodylus porosus</i> )
<b>Whales</b>	Antarctic minke whale ( <i>Balaenoptera bonaerensis</i> ) Bryde's whale ( <i>Balaenoptera edeni</i> ) Humpback whale <sup>T</sup> ( <i>Megaptera novaeangliae</i> ) Sei whale <sup>T</sup> ( <i>Balaenoptera borealis</i> ) Blue whale <sup>T</sup> ( <i>Balaenoptera musculus</i> ) Fin whale <sup>T</sup> ( <i>Balaenoptera physalus</i> ) Killer whale ( <i>Orcinus orca</i> ) Sperm whale ( <i>Physeter macrocephalus</i> )
<b>Dugong</b>	Dugong ( <i>Dugong dugon</i> )
<b>Dolphins</b>	Australian snubfin dolphin ( <i>Orcaella heinsohni</i> ) Indo-Pacific humpback dolphin ( <i>Sousa chinensis</i> )
<b>Sharks</b>	Longfin mako ( <i>Isurus paucus</i> ) Shortfin mako ( <i>Isurus oxyrinchus</i> ) White shark <sup>T</sup> ( <i>Carcharodon carcharias</i> ) Porbeagle ( <i>Lamna nasus</i> ) Whale shark <sup>T</sup> ( <i>Rhincodon typus</i> )
<b>Seabirds</b>	Black-naped tern ( <i>Sterna sumatrana</i> ) Brown booby ( <i>Sula leucogaster</i> ) Common noddy ( <i>Anous stolidus</i> ) Greater frigatebird ( <i>Fregata minor</i> ) Lesser crested tern ( <i>Sterna bengalensis</i> ) Little tern ( <i>Sterna albifrons</i> ) Northern giant petrel <sup>T</sup> ( <i>Macronectes halli</i> ) Red-footed booby ( <i>Sula sula</i> ) Sooty albatross <sup>T</sup> ( <i>Phoebastria fusca</i> ) Wandering albatross <sup>T</sup> ( <i>Diomedea exulans</i> ) White-bellied sea eagle ( <i>Haliaeetus leucogaster</i> ) Wilson's storm petrel ( <i>Oceanites oceanicus</i> ) Bridled tern ( <i>Onychoprion anaethetus</i> ) Caspian tern ( <i>Sterna caspia</i> ) Common tern ( <i>Sterna hirund</i> ) Grey-headed albatross <sup>T</sup> ( <i>Thalassarche chrysostoma</i> ) Lesser frigatebird ( <i>Fregata ariel</i> ) Masked booby ( <i>Sula dactylatra</i> ) Osprey ( <i>Pandion haliaetus</i> ) Roseate tern ( <i>Sterna dougalii</i> ) Southern giant petrel <sup>T</sup> ( <i>Macronectes giganteus</i> ) Wedge-tailed shearwater ( <i>Puffinus pacificus</i> ) White-winged black tern ( <i>Chlidonias leucopterus</i> )
<b>Shorebirds</b>	Bar-tailed godwit ( <i>Limosa lapponica</i> ) Caspian plover ( <i>Charadrius asiaticus</i> ) Common sandpiper ( <i>Actitis hypoleucos</i> ) Eastern curlew ( <i>Numenius madagascariensis</i> ) Eastern reef egret ( <i>Egretta sacra</i> ) Great knot ( <i>Calidris tenuirostris</i> ) Greenshank ( <i>Tringa nebularia</i> ) Grey-tailed tattler ( <i>Tringa brevipes</i> ) Lesser golden plover ( <i>Pluvialis dominica</i> ) Little curlew ( <i>Numenius minutus</i> ) Red knot ( <i>Calidris canutus</i> ) Ruddy turnstone ( <i>Arenaria interpres</i> ) Sharp-tailed sandpiper ( <i>Calidris acuminata</i> ) Terek sandpiper ( <i>Xenus cinereus</i> ) Whimbrel ( <i>Numenius phaeopus</i> ) Black-tailed godwit ( <i>Limosa limosa</i> ) Cattle egret ( <i>Ardea ibis</i> ) Curlew sandpiper ( <i>Calidris ferruginea</i> ) Eastern great egret ( <i>Ardea modesta</i> ) Glossy ibis ( <i>Plegadis falcinellus</i> ) Greater sand plover ( <i>Charadrius leschenaultii</i> ) Grey plover ( <i>Pluvialis squatarola</i> ) Latham's snipe ( <i>Gallinago hardwickii</i> ) Lesser sand plover ( <i>Charadrius mongolus</i> ) Marsh sandpiper ( <i>Tringa stagnatilis</i> ) Red-necked stint ( <i>Calidris ruficollis</i> ) Sanderling ( <i>Calidris alba</i> ) Swinhoe's snipe ( <i>Gallinago megala</i> ) Wandering tattler ( <i>Tringa incana</i> ) Wood sandpiper ( <i>Tringa glareola</i> )

### 4.6.1 Marine turtles

All six species of marine turtle that occur in the Region and adjacent areas (green turtle, hawksbill turtle, flatback turtle, leatherback turtle, loggerhead turtle and olive ridley turtle) are listed as both migratory and threatened species. Many migrate to nesting or foraging areas outside the Region. Important nesting, inter-nesting and foraging areas for four of the species have been identified in the Region.<sup>87</sup> Leatherback turtles and olive ridley turtles are rarely sighted within the Region — there are no known nesting sites, and very little is known about how they use the Great Barrier Reef. There is also very limited knowledge about foraging flatback turtles.



A brief description of each species is provided below.

The **green turtle** is found in tropical, subtropical and temperate waters around the world and is the most abundant of the six species of marine turtle found in the Region. Green turtles are slow growing, taking decades to mature to breeding adults. Age at maturity is estimated to be about 30 to 40 years. Two genetic stocks of green turtles breed within the Great Barrier Reef Marine Park — southern and northern stocks.<sup>88,89</sup> Within a feeding area, the proportion of adult females that prepare to breed in a year is variable and is a function of the Southern Oscillation Index two years before the breeding season.<sup>90,91</sup> Nesting for the southern stock is concentrated in the Capricorn Bunker group of islands; the total nesting population is expected to be approximately 8000 females in an average breeding season.<sup>92</sup> Nesting for the northern stock is concentrated on Raine Island and Moulter Cay, and there is significant fluctuation in the size of the annual nesting population. Size estimates of the nesting population in the waters around Raine Island have varied between 788 and 135,300 females.<sup>88</sup> These cays support one of the few remaining large breeding aggregations of green turtles in the world.<sup>89,92</sup> The southern stock is recovering at almost four per cent per annum. The northern stock is stable but showing signs of decline.<sup>92</sup> Green turtles are generalist feeders, eating predominantly seagrass but also algae, mangrove fruit and jellyfish. Southern stock turtles forage in waters mainly south of Cape Melville and eastwards to New Caledonia and sometimes Fiji.<sup>92</sup> Northern stock turtles forage in waters mainly north of Cape Melville, throughout Torres Strait and waters of eastern Indonesia and Papua New Guinea. Green turtles that nest in the Coral Sea have also been found foraging in the Region.

**Hawksbill turtles** that nest in the Region are part of the Torres Strait–northern Great Barrier Reef breeding stock. The number of females nesting annually at Milman Island, an important nesting site, has previously been estimated at between 100 and 500.<sup>93</sup> However, there is no published monitoring data since 2000. Hawksbill turtles eat primarily algae, but also feed on sponges, soft corals, seagrass and shellfish.<sup>94</sup> They grow very slowly and do not reach sexual maturity until after 31 years of age.<sup>95</sup> It is estimated that the breeding stock is declining by around three per cent per year.<sup>96</sup> Hawksbill turtles nesting in the Region have been found in Papua New Guinea and elsewhere in the Region.<sup>93</sup>

**Flatback turtles** are the only marine turtle that nests solely in Australia. The stock which nests in the Great Barrier Reef is known as the eastern Australian stock. There is little data on age at maturity for the species, but it is likely to be 20 or more years.<sup>97,98</sup> Three decades of research and monitoring of the eastern Australian stock at rookeries in the central Great Barrier Reef indicate that the population is stable.<sup>99</sup> All class sizes of flatback turtles are carnivorous, with post-hatchlings feeding on zooplankton, and adults feeding principally on soft-bodied invertebrates including soft corals, sea pens, holothurians and jellyfish.<sup>97</sup> The foraging distribution for the eastern Australian stock extends from Hervey Bay to Torres Strait and possibly into the Gulf of Papua.<sup>100</sup>



*Green turtle hatchling on Raine Island*



*Flatback turtle nesting*

**Loggerhead turtles** that inhabit the Great Barrier Reef are part of the eastern Australian stock which breeds in southern Queensland. Loggerhead turtles in the south-western Pacific Ocean are slow growing, taking about three decades to mature to breeding adults. In 1976–1977, the estimated population of nesting loggerhead turtles in the Great Barrier Reef was 3500; the population subsequently declined to fewer than 500 females in the 1999–2000 breeding season.<sup>101</sup> Annual monitoring shows the long-term decline has reversed at all monitored nesting beaches since 2001, when turtle excluder devices on otter trawlers became mandatory.<sup>102</sup> In eastern Australian coastal waters, loggerhead turtles feed principally on molluscs and crabs.<sup>103</sup> The foraging distribution for the eastern Australian stock extends from northern New South Wales to the Northern Territory, Papua New Guinea, south-west Pacific and Coral Sea.

There are no major rookeries for the **leatherback turtle** in Australia.<sup>104</sup> The progressive decline in breeding frequency recorded in eastern Australia — from low density, annual nesting recorded in the 1970s and 1980s to now, when no nesting is observed<sup>104</sup> — indicates the population that visits Australian waters is likely to be in decline. This correlates with reports of a significant decline in Pacific Ocean leatherback turtle populations,<sup>105</sup> and recorded declines in the numbers of animals interacting with shark control equipment set along the Queensland coast.<sup>106</sup> Leatherback turtles are one of the fastest growing marine turtles. Individuals from the eastern Pacific




population are estimated to reach maturity in 13 years.<sup>107</sup> Leatherback turtles are carnivorous and, in Australian waters, they feed extensively on colonial tunicates,<sup>108</sup> jellyfish and other soft-bodied invertebrates.<sup>109,110</sup>

There are two main breeding areas for **olive ridley turtles** in Australia, one in the Northern Territory with an estimated annual nesting density of a few thousand, and the other along the north-western coast of Cape York Peninsula between Weipa and Bamaga.<sup>111</sup> There are no records of olive ridley turtles nesting along the east coast of Australia, even though sexually mature females have been recorded in foraging areas off Townsville.<sup>111</sup> No Queensland rookeries are within national parks or similar protected habitat.<sup>111</sup> However, a substantial part of their east coast foraging range is contained within the Region, although there is no information upon which to conclude the population status of the species in the Region. Adult and large immature olive ridley turtles are carnivorous, feeding principally on molluscs and small crabs.<sup>111</sup>

The important feeding and breeding areas of marine turtles within the Region are presented in Figure 4.10.

The six marine turtle species are considered collectively in the report under the grouping of 'marine turtles'.



**Figure 4.10 Identified important feeding and breeding areas for four species of marine turtles within the Region**

The species mapped are the green, hawksbill, flatback, and loggerhead turtles.<sup>87</sup>

#### 4.6.2 Estuarine crocodiles

On the east coast of Queensland, estuarine crocodiles inhabit reef, coastal and inland waterways from north of Gladstone. Following extensive harvesting of wild populations in northern Australia between the 1940s and 1960s, management measures to protect the species were adopted by all states and the Australian Government. In Queensland, the species was fully protected in 1974.<sup>112</sup>

Estuarine crocodiles are considered temporary migrants in the Region. They occur widely but at low densities, principally along the mainland coast and on and around islands. Coastal river systems and their associated wetlands are their primary habitat and also where they nest. Crocodiles on islands are not likely to be a major component of the species population.<sup>113</sup>

#### 4.6.3 Whales

Eight listed migratory whale species are relevant to the Region, with each spending only part of their life history within its boundaries. Four of the species are also listed threatened species, namely blue whales, fin whales, humpback whales and sei whales. Australian populations of these species have previously declined, primarily as a legacy of past commercial exploitation.

**Blue, fin and sei whales** are wide-ranging oceanic species and, in general, spend summers in higher latitudes and winters in warmer tropical waters.<sup>114</sup> They are rarely sighted within the Region, and very little is known about their movements or use patterns there.

In contrast, **humpback whales** are the most commonly sighted listed marine mammal species in the Region. The east Australian population is recovering from severe depletion by commercial whaling in the 20<sup>th</sup> century.<sup>115</sup> Humpback whales migrate to the Region during the winter months to calve and mate.<sup>116</sup> Modelling of environmental suitability for humpback whales identified two core areas where there is a higher probability of humpback whales: offshore Proserpine extending south to Mackay within the inner reef lagoon (including the Whitsundays), and the Capricorn and Bunker groups of islands and reefs approximately 100 kilometres east of Gladstone.<sup>117</sup> Continued recovery of the population means humpback whales are being sighted more frequently in the Region, including in areas where they have not occurred for decades.



*Humpback whales are both a listed migratory and a listed threatened species*

The **Antarctic minke whale**, **Bryde's whale**, **killer whale** and **sperm whales** are believed to be seasonal migrants to the Region, occurring mostly during the winter months. However, there is very little information about these species. Killer whales are rarely sighted within the Region, although there have been anecdotal reports of the species near Moreton Bay, south of the Region, during humpback whale calving season. Sperm and Bryde's whales are known to move through the Region from records obtained through the joint Field Management Program's Marine Wildlife Stranding program and reports from the Authority's Sightings Network.<sup>118</sup>

Until recently, there was thought to be only one species of minke whale, referred to as *Balaenoptera acutorostrata*. There are now two generally accepted species, the common minke whale (*B. acutorostrata*) and the Antarctic minke whale (*B. bonaerensis*) — a listed migratory species. The dwarf minke whale, which is regularly sighted in the Region, is currently regarded as an undescribed subspecies of *B. acutorostrata* and is, therefore, not a listed migratory species.

In this report, all listed migratory and threatened whale species are generally considered as a group along with other whale species in the Region. The detailed understanding of the humpback whale relative to other whale species means it is used as an example in some assessments in this report.

#### 4.6.4 Dolphins

The **Australian snubfin dolphin** and **Indo-Pacific humpback dolphin** are listed migratory species that occur in the Region. They share similar life history traits and habitat preferences. They are long-lived, slow-growing and late-maturing; have low reproductive rates, low relative abundances, and high habitat and diet specificity; maintain small group sizes; and occupy small home ranges, with little overlap. Each localised population is estimated to

have fewer than 100 individuals.<sup>119,120</sup> These traits make them susceptible to a number of pressures in and adjacent to the Region. Along the east coast of Queensland, these dolphins are primarily found in shallow waters less than 20 metres deep, close to the coast, close to river and creek mouths and in the proximity of seagrass meadows.<sup>121</sup> The aggregation of snubfin dolphins around the mouth of the Fitzroy River is the southernmost identified resident population in Australian waters.<sup>120</sup>

There are currently no overall population estimates for the Australian snubfin or Indo-Pacific humpback dolphins in the Region;<sup>122</sup> however, there are local population estimates for Australian snubfin dolphin aggregations in Cleveland-Halifax Bays<sup>119</sup> and in Keppel Bay-Fitzroy River<sup>120</sup> of about 70 dolphins each. An aggregation has also been recorded at Princess Charlotte Bay-Bathurst Bay on Cape York Peninsula.<sup>123</sup> There have been population estimates for Indo-Pacific humpback dolphins in Cleveland Bay (50 or fewer)<sup>119</sup>; the Capricorn coast (about 64); Keppel Bay (about 107); and Port Curtis (about 85)<sup>124</sup>. Modelling suggests that current mortality in the populations of these species is greater than what they can sustain.<sup>119,120</sup>

All dolphin species are considered as a group in the report and, where relevant, the two listed species are discussed separately.

#### 4.6.5 Dugongs

The Great Barrier Reef Region supports globally significant populations of dugongs. They occur along the length of the Region, feeding mainly on seagrasses. Dugongs play an important ecological role in coastal marine ecosystems.

North of Cooktown, there is a healthy population, and the population trend over the last several decades of monitoring has been stable.<sup>125</sup> However, dugong populations south of Cooktown have been mostly declining for decades, with an estimated population of only 600 animals between the Daintree River and the southern limit of the Region in 2011,<sup>126,127</sup> compared with an estimate of 2059 when the previous survey was conducted in 2005. This is the lowest number recorded in this area since surveys began in 1986.

Dugongs are considered separately as a discrete key value throughout the report, and are the subject of a demonstration case study (see Chapter 9).

#### 4.6.6 Sharks

The **whale shark** is both a listed migratory and listed threatened species. It is the world's largest fish, with the largest recorded being 12 metres long.<sup>128</sup> Despite its size, it feeds primarily on plankton. The whale shark has a broad global distribution in tropical to warm-temperate waters. It is often seen far offshore, but also comes close to shore and sometimes enters lagoons of coral reefs. The species is widely distributed in Australian waters and is regularly sighted throughout the Region.



*Whale sharks are regularly sighted in the Region*

**White sharks** are also a listed migratory and listed threatened species. They are a large apex predator that grow to at least six metres and can weigh up to 2000 kilograms.<sup>129</sup> They are most frequently observed in coastal temperate and subtropical regions, but have been observed in tropical areas such as the Coral Sea, Papua New Guinea, the central Pacific, northern Brazil and the tropical south-west Indian Ocean.<sup>130</sup> They occur mainly in coastal waters but also spend significant time in the open ocean.<sup>130</sup> White sharks are rarely sighted in the Region, although there are a few reports in waters off central Queensland. Some animals tagged in New South Wales' waters have been tracked swimming north into the Great Barrier Reef, indicating connectivity over long distances.<sup>131</sup> The ranges of the **longfin mako**, **porbeagle** and **shortfin mako** extend well beyond the Region. The **porbeagle** is a wide-ranging, coastal and oceanic shark that is found from the surface down to 370 metres.<sup>129</sup> The **shortfin mako** is a wide-ranging oceanic and pelagic shark found in waters down to 650 metres.<sup>129</sup> The **longfin mako** is a widely-distributed, but rarely encountered, oceanic tropical shark.

The listed shark species are considered in the report as part of the larger group of sharks and rays.

#### 4.6.7 Seabirds

Within the Region, 23 seabird species are listed as migratory under the EPBC Act. In the description below, the species are divided into *inshore and coastal*, and *offshore and pelagic* foragers due to differences in feeding behaviour and associated implications for their ecology.<sup>132,133</sup> Seabirds and their habitat are also protected under three bilateral migratory bird agreements between Australia and Japan, China and the Republic of Korea.<sup>134</sup>

Seabirds are generally long-lived, characterised by late sexual maturity, small clutch sizes, slow chick growth rates and extended fledgling periods. They require habitats with suitable areas for both breeding and foraging. Specific requirements for breeding vary between species, but commonly include an area free from disturbance by terrestrial predators and storms and tide; the presence of others of their species, including potential mates; and suitable access to foraging grounds.<sup>135</sup>

The timing of seabird breeding activity in the Region is complex and poorly understood.<sup>132</sup> Most seabirds breed on relatively remote islands, with each species tending to nest in a specific habitat type. Coral cays are the preferred habitats of most seabirds, providing 73 per cent of the major nesting sites.<sup>135</sup> More than 75 cays have been identified as seabird breeding colonies, with 56 considered to be key sites and 20 minor sites (Figure 4.11).<sup>134,135</sup> The most significant nesting sites are Raine Island, Michaelmas Cay, the cays of the Swain Reefs and the islands of the Capricorn and Bunker Groups.<sup>134</sup> For the majority of species, peaks in breeding occur between October and April though, for some species, nesting can occur year round.<sup>132</sup> However, for some seabird species on the Great Barrier Reef, such as masked boobies in the Swain Reefs, nesting peaks during the winter months.<sup>136</sup>




Figure 4.11 Principal seabird breeding islands in the Region



### Inshore and coastal foraging seabirds

There are eight species of inshore and coastal foraging seabirds that nest within the Region, five of which are listed migratory species (Table 4.4). The Region supports more than 50 per cent of Australia's inshore foraging roseate terns, lesser crested terns and black-naped terns (Table 4.5).<sup>134,135</sup>

Generally, inshore and coastal foraging seabird species source food closer to their breeding colony compared with offshore and pelagic foragers. For example, in the Great Barrier Reef, crested terns mostly forage singularly or in pairs over shallow reef flats and coastal waters<sup>137</sup> within two to three kilometres of their colony<sup>138</sup>, although some venture up to 12 kilometres away<sup>133,138</sup>. Roseate terns and black-naped terns forage within three kilometres of the colony, and the lesser crested tern forages even closer.<sup>139</sup> Colonies of inshore and coastal foraging seabirds are also smaller, more numerous and more widely distributed.<sup>140</sup>

**Table 4.5 Estimated population of listed migratory inshore and coastal foraging seabirds known to breed in the Great Barrier Reef Region**


Source: Adapted from Congdon, 2008<sup>132</sup>

Species	Estimated population in the Region
Caspian tern ( <i>Hydroprogne caspia</i> )	70
Roseate tern ( <i>Sterna dougallii</i> )	6000
Black-naped tern ( <i>Sterna sumatrana</i> )	3900
Little tern ( <i>Sternula albigrons</i> )	1000
Lesser crested tern ( <i>Thalaseus bengalensis</i> )	6300

### Offshore and pelagic-foraging seabirds

There are 12 species of offshore and pelagic-foraging seabirds that nest within the Region, 10 of which are listed migratory species (Table 4.6).<sup>10</sup> The Region supports more than 50 per cent of offshore foraging black noddies and approximately 25 per cent of wedge-tailed shearwaters, and brown and masked boobies.<sup>134,135</sup>

Offshore and pelagic-foraging seabirds usually have single clutches and much slower growing chicks with longer fledging periods than seabirds with other methods of foraging.<sup>132</sup> Pelagic-foraging species such as sooty terns may live up to 32 years and larger species such as boobies and frigatebirds even longer.<sup>132</sup> Most offshore and pelagic-foraging seabird species do not become sexually mature or return to breed for between five to 12 years after fledging. The nesting period for some species, such as frigatebirds, can be up to six months.<sup>132</sup>



Red-footed boobies on Raine Island. From top of branch, two brown juveniles, a white adult and a light brown bird in an intermediate phase.

Characteristically, these species nest in large colonies, often spread over multiple, closely spaced islands that are close to abundant food and provide suitable nesting habitat. The most important example is the wedge-tailed shearwater, where approximately 500,000 individuals breed on the 13 islands of the Capricorn and Bunker Groups which comprise a staggering 66 per cent of the biomass of breeding seabirds in the Region.<sup>135,141</sup> They feed on pelagic fish whose abundance and distribution are determined by oceanographic upwellings.<sup>135</sup>

The listed migratory seabird species are considered in this report as part of the larger group of seabirds.

**Table 4.6 Estimated population of listed migratory offshore and pelagic-foraging seabirds known to breed in the Region**

Source: Adapted from Congdon, 2008<sup>132</sup> and King, 1993<sup>142</sup>

Species	Estimated population in the Region
Wedge-tailed shearwater ( <i>Ardenna pacifica</i> )	560,000
Red-footed booby ( <i>Sula sula</i> )	172
Brown booby ( <i>Sula leucogaster</i> )	18,500
Masked booby ( <i>Sula dactylatra</i> )	1100
Greater frigatebird ( <i>Fregata minor</i> )	20
Lesser frigatebird ( <i>Fregata ariel</i> )	2500
Sooty tern ( <i>Onychoprion fuscata</i> )	48,000
Bridled tern ( <i>Onychoprion anaethetus</i> )	13,900
Common noddy ( <i>Anous stolidus</i> )	46,000
Black noddy ( <i>Anous minutus</i> )	300,000

#### 4.6.8 Shorebirds

There are 41 species of shorebirds known to inhabit the Region, 30 of which are listed migratory species (Table 4.4). These species use a wide range of habitats within and adjacent to the Region, including beaches, rocky shores, estuaries, intertidal flats, coral cays and reefs, freshwater wetlands, grasslands, pasture land and sewage treatment plants.

Most are generalists, with several exceptions including the sooty oystercatcher, which are generally only found on rocky ocean shores.<sup>143</sup> The most crucial habitats for shorebirds are tidal flats. Loss or degradation of habitat in any of their feeding, breeding or staging areas can result in failure to breed or death from starvation or predation. Maintenance of these habitats in an undisturbed condition is particularly important for migratory shorebirds to enable them to build adequate body condition prior to their annual northward migration.

The majority of migratory shorebirds spend part of the year in Australia, departing for their breeding grounds in northern China, Mongolia, Siberia and Alaska between March and May. They arrive back in Australia between August and November after their return migration. Birds migrate along the East Asian-Australian Flyway, which includes numerous stopover sites for resting and feeding.<sup>144</sup>

There are no population estimates for the Region's shorebirds. Significant declines in some shorebird populations have been reported in Australia.<sup>145</sup>

Internationally significant numbers of shorebirds occur at a number of sites with the Region, particularly the islands off False Orford Ness in Cape York, Pelican Island and nearby islands, the Cairns foreshore, Cape Bowling Green, Burdekin River delta, Pioneer River to McEwan's Beach and Notch Point near Mackay, Shoalwater Bay and Broad Sound.<sup>143</sup>

The listed migratory shorebird species are considered in the report as part of the larger group of shorebirds.

### 4.7 Listed threatened species and ecological communities

Twenty-six species that are currently listed as threatened under the EPBC Act occur in the Great Barrier Reef Region. There are six marine turtle species, five marine mammal species, seven shark and ray species, and six seabird and shorebird species (Table 4.7).

The water mouse *Xeromys myoides*, a listed threatened species, may occur in coastal saltmarsh, mangroves and adjacent freshwater wetland habitats adjacent to the southern half of the Region. It is not considered in this assessment. None of the listed ecological communities are known from the Region.

**Table 4.7 Listed threatened species in the Great Barrier Reef Region**

The species marked with the symbol - <sup>M</sup> - are also listed migratory species (Table 4.4). The scientific name is presented in *italics*.

Status	Marine turtles	Marine mammals	Sharks and rays	Seabirds
<b>Vulnerable</b>	Flatback turtle <sup>M</sup> <i>Natator depressus</i> Green turtle <sup>M</sup> <i>Chelonia mydas</i> Hawksbill turtle <sup>M</sup> <i>Eretmochelys imbricata</i>	Fin whale <sup>M</sup> <i>Balaenoptera physalus</i> Humpback whale <sup>M</sup> <i>Megaptera novaeangliae</i> Sei whale <sup>M</sup> <i>Balaenoptera borealis</i> Subantarctic fur seal <i>Arctocephalus tropicalis</i>	Dwarf sawfish <i>Pristis clavata</i> Freshwater sawfish <i>Pristis pristis</i> Green sawfish <i>Pristis zijsron</i> Whale shark <sup>M</sup> <i>Rhincodon typus</i> White shark <sup>M</sup> <i>Carcharodon carcharias</i>	Northern giant petrel <sup>M</sup> <i>Macronectes halli</i> Sooty albatross <sup>M</sup> <i>Phoebastria fusca</i> Wandering albatross <sup>M</sup> <i>Diomedea exulans</i>
<b>Endangered</b>	Loggerhead turtle <sup>M</sup> <i>Caretta caretta</i> Leatherback turtle <sup>M</sup> <i>Dermochelys coriacea</i> Olive ridley turtle <sup>M</sup> <i>Lepidochelys olivacea</i>	Blue whale <sup>M</sup> <i>Balaenoptera musculus</i>	N/A	Grey-headed albatross <sup>M</sup> <i>Thalassarche chrysostoma</i> Southern giant petrel <sup>M</sup> <i>Macronectes giganteus</i>
<b>Critically endangered</b>	N/A	N/A	Grey nurse shark — east coast population <i>Carcharias taurus</i> Spear-tooth shark <i>Glypis glyphis</i>	Herald petrel <i>Pterodroma heraldica</i>

#### 4.7.1 Marine turtles

The six listed threatened marine turtle species which occur in the Region are also listed migratory species, and are discussed in Section 4.6.1.

#### 4.7.2 Marine mammals

The four listed threatened whale species which occur in the Region are also listed migratory species, and are discussed in Section 4.6.3.

The **subantarctic fur seal** is rarely seen within the Region, and those recorded are believed to have been transients. Dispersing juvenile and adult male sub-Antarctic fur seals are known to make long journeys of up to 3000 kilometres. Given its very rare occurrence in the Region, this species is not considered further in this report.

#### 4.7.3 Sharks and rays

The Region is within the range of the critically endangered east coast population of **grey nurse sharks**. The species is rarely sighted within the Region, with only some records in the central and southern parts.



*The grey nurse shark, a critically endangered species, is rarely sighted in the Region*

The **dwarf sawfish**, **freshwater sawfish** and **green sawfish** all occur in similar habitats within and adjacent to the Region.<sup>129</sup> Sawfish are commonly found in estuaries and freshwater rivers and are known to move easily between fresh and saltwater.<sup>129</sup> They are long-lived and mature late; have slow growth rates, low reproduction rates and low abundance; and can exhibit habitat and prey specificity. These life history traits make them susceptible to a number of impacts occurring in and adjacent to the Region.<sup>146</sup>

The **spear-tooth shark** belongs to the genus *Glyphis*, a small group of poorly known sharks. It is thought to have a fragmented distribution, confined to a few highly turbid, tidal rivers and estuaries in northern Australia.<sup>147</sup> The species has not been recorded in or near the Region since 1982, when it was recorded in the Bizant River flowing into Princess Charlotte Bay.<sup>147</sup> The spear-tooth shark is listed as critically endangered under the EPBC Act.

The **whale shark** and the **white shark** are listed as both threatened and migratory species, and are discussed in Section 4.6.6.

As with the listed migratory sharks (Section 4.6.6), all these threatened shark and ray species are considered in this report within the group 'sharks and rays'. Particular issues in relation to these threatened species are highlighted where relevant.

#### 4.7.4 Seabirds

Five of the six listed threatened seabird species which occur in the Region are also listed migratory species and are discussed in Section 4.6.7. There are very few sightings in the Region of the listed threatened seabirds. The **grey-headed albatross**, **northern giant petrel**, **sooty albatross**, **southern giant petrel** and **wandering albatross** are all primarily associated with offshore areas of the Region, near and beyond the continental shelf. Their most critical foraging habitat is considered to be waters south of 25 degrees latitude<sup>148</sup> (and, hence, outside the Region). There is no breeding habitat within the Region.<sup>148</sup>

There are few records of the **herald petrel** for the Great Barrier Reef. All have been at Raine Island where, from 1979 to 1987, 13 birds were banded. Of these, only three individuals have been recaptured at Raine Island (the last in 1985), and one has been recovered breeding on Round Island in the Indian Ocean in 2010.<sup>149</sup> Since 1987, there has been only one observed incidence of breeding by herald petrels on the island.<sup>150</sup>

As with the listed migratory seabirds, all these threatened seabird species are combined in this report in the grouping 'seabirds'.

### 4.8 Wetlands of international importance


#### 4.8.1 Shoalwater and Corio Bays Area

The Shoalwater and Corio Bays Area was listed under the Convention on Wetlands of International Importance (the Ramsar Convention) in 1996. The area extends over terrestrial and marine areas and part is within the Region (Figure 4.12).

It is the largest wilderness area within the Central Mackay Coast biogeographic area, and is representative of coastal, sub-coastal, aquatic landscapes and ecosystems which are relatively undisturbed habitat for significant plants and animals, including rare and threatened species.<sup>151</sup>


The Shoalwater and Corio Bays Area supports about 791 plant species and sub-species, 445 fish species, 22 frog species, 66 reptile species, 226 bird species, 24 mammal species and 23 bat species. Listed threatened marine species that occur at the site include the green turtle, hawksbill turtle, flatback turtle and loggerhead turtle. Dugongs, a listed migratory species, also occur in the area. Dozens of listed migratory bird species have been recorded at the site, and it is of international importance to the migratory eastern curlew, whimbrel and great knot. The area supports more than 20,000 shorebirds, most of them migratory. Six shorebird species occur in numbers greater than one per cent of their estimated flyway population size — grey-tailed tattler, bar-tailed godwit, eastern curlew, whimbrel, terek sandpiper, and Australian pied oystercatcher.<sup>152</sup>

The area is part of the traditional lands of the Darumbal people. The dune fields contain archaeological sites including shell middens, scatters of stone tools and dinner camp sites. The area is connected with the broader Great Barrier Reef ecosystem and beyond. For example, 69 adult green turtles previously recorded at distant breeding areas throughout the Coral Sea were recaptured between 2000 and 2004 while foraging in western Shoalwater Bay.<sup>153</sup>



Freshwater Beach, Shoalwater Bay





**Figure 4.12 Shoalwater and Corio Bays Area**


The Shoalwater and Corio Bays Area is a wetland of international importance, part of which is within the Region.

#### 4.8.2 Bowling Green Bay Area

The Bowling Green Bay Area is a wetland of international importance adjacent to the Region, listed under the Ramsar Convention in 1993. The site is located about 50 kilometres south-east of Townsville and covers more than 47,000 hectares, including Cape Bowling Green and parts of Cape Cleveland (Figure 4.13). The area is part of the traditional lands of the Juru and Bindal people.

Ninety-nine per cent of the Ramsar site is within a Queensland national park. At the northern extremities of Cape Bowling Green and Cape Cleveland, there are small lighthouse areas (approximately 80 hectares and three hectares, respectively). A further 208 hectares have been excised at the western end of Bowling Green Bay for the Australian Institute of Marine Science. Several small townships are encompassed by the area, but do not form part of it.

The area is identified as being regionally unique and internationally important for the diversity and extent of marine, estuarine and freshwater wetland types it supports. As one of the remaining, relatively intact wetlands adjacent to the Region, it supports the Great Barrier Reef ecosystem, including through nutrient assimilation and sediment stabilisation.



**Figure 4.13 Bowling Green Bay Area**

The Bowling Green Bay Area is a wetland of international importance directly adjacent to the Region.

A key feature of the Bowling Green Bay Area is Cape Bowling Green. During the last low sea level period, the Burdekin River flowed out to sea via Bowling Green Bay.<sup>44,154</sup> It diverted from its present course at Kelly Mount, flowing directly north into the Bay, past Cape Cleveland and out to open water through the exposed coral reefs. The Burdekin reverted back to its present course about 4000 years ago, resulting in what is now Australia's longest (20 kilometres) free-standing sand spit, Cape Bowling Green. The course of the Burdekin River is also important as it delivered enormous sediment loads which built up the 100 beach ridges connecting Cape Cleveland to the mainland, the second-most extensive beach ridge sequence on the Great Barrier Reef coast.

## 4.9 Environmental processes

Environmental processes are the forces, cycles, reactions and interactions that alter and shape a natural area, keeping it healthy and functioning. They support and connect the Great Barrier Reef environment, and interact with and influence each other. They can also be affected by activities within and adjacent to the Region — transferring effects onto the Region's values.

The following is a description of the principal environmental processes in the Region. It includes the ecosystem health processes identified in the *Outlook Report 2009*<sup>9</sup>, with minor updates and amendments, and the major geomorphological processes that are altering and shaping the Reef environment. In many cases, the ecological and geomorphological processes are one and the same; for example, reef building and erosion have both ecological and geomorphological aspects.

The major processes are illustrated in Figure 4.14. All these processes are critical to the Region, and are comprehensively considered throughout the report. They are individually assessed in Chapter 7.




Figure 4.14 Key environmental processes of the Great Barrier Reef Region

**Waves, currents and tides** all mix oceanic water and have an important effect on marine life. The Great Barrier Reef is part of a larger system of ocean circulation throughout the Pacific Ocean, which delivers nutrients and larvae from other regions as well as deep water into the Region. At the largest spatial scale (thousands of kilometres), major oceanic currents of the Coral Sea affect patterns of connectivity between reefs and the temperature of the Region's waters.<sup>155</sup> At small scales (centimetres to metres), turbulence can affect the settlement patterns of organisms such as corals. While surface currents are primarily driven by wind, deeper ocean currents are mainly driven by relative densities of seawater, affected by salinity and temperature.<sup>155</sup>

Upwelling of cold, nutrient-rich, sub-thermocline waters to the sea surface create 'hotspots' of marine primary production. In the Great Barrier Reef, upwelling intrusions include those on the central Great Barrier Reef which are enhanced during conditions of consistently low winds. During these conditions, the southward-flowing East Australian Current flows faster, lifting the thermocline closer to the surface, spilling cooler waters onto the shelf.<sup>155,156</sup>

**Cyclones** regularly affect coral reefs and other habitats at regional and local levels. In the past five years, there have been several cyclones of category three and above affecting the Great Barrier Reef. In addition to strong winds and rain, the powerful waves generated during cyclones can seriously damage habitats and geomorphological features, particularly coral reefs and shorelines.<sup>155</sup> **Wind** also plays a role; in particular, it can cause dramatic changes in the shape of islands and the coast and affect ocean currents.<sup>155</sup>



Damage to corals on Rib Reef by Cyclone Larry



**Sea level** is an important determinant of species and habitat distribution, as well as affecting individual foraging and reproduction activities. It varies naturally day-to-day with the tides and over longer time scales from the El Niño Southern Oscillation and the Pacific Decadal Oscillation. In addition, cyclonic winds can also cause storm surges, an onshore rise of water above the predicted astronomical tide. Taking a longer time period, over the past 100,000 years sea levels have risen and fallen many times, shifting the position of reef growth on the continental shelf.<sup>47</sup> Most modern reefs of the Great Barrier Reef formed as sea level crept over the shelf margin 9000 years ago, and coral recruits settled and re-colonised on ancient reefs.<sup>44</sup> However, variations in relative sea level have resulted in very different evolutionary histories for coral reefs over the period of *Holocene* growth. While *global* sea level reached its present position 6500 years ago, the loading on the continental margin caused subsidence and *relative* sea level reached its present position much later. In contrast, sea level on the inner shelf reached its present level 6500 years ago. Sea level is currently rising in Australian waters, especially in northern areas.<sup>157,158</sup>

**Sea temperature** is a key environmental factor controlling the distribution and diversity of marine life. It is critical to reef building and is one of the key variables that determine the north-south limits of coral reefs and coral reef diversity.<sup>159</sup> When temperature limits are exceeded, physiological processes may break down. For reef habitats, the most critical mechanism affected is the symbiotic association between animals (such as corals and clams) and the algae which live within their tissues and provide much of their nutrition. If sea temperatures exceed a certain threshold, these algae are expelled — an effect known as 'bleaching'.<sup>160</sup> Sea temperature also plays a role in ocean circulation, through deep ocean currents as cooler, denser water sinks to the bottom and warmer, less dense water rises.

**Freshwater inflow** from the creeks and rivers in the Great Barrier Reef catchment naturally form a thin layer of freshwater on the surface of the heavier seawater and, during flood events, may extend to mid-shelf and outer reefs. This can result in extensive fluctuations in ocean **salinity**, especially in intertidal and shallow habitats. Heavy rainfall directly on the ocean can also reduce surface salinity. At natural volumes and frequencies, freshwater inflow from catchments triggers essential migration, reproduction and recruitment of various fish and crustaceans.<sup>161</sup> Depending on geology and soil permeability, freshwater also moves as groundwater to estuaries or the sea. Some mangroves, saltmarsh plants and seagrasses depend on freshwater seepage. Some marine animals (for example, sea snakes) consume freshwater from submarine groundwater seepages.<sup>162</sup> Freshwater may also seep through the ocean floor from drowned river channels through 'wonky holes'.<sup>163</sup> Salinity is also a key driver of ocean circulation.<sup>155</sup>

**Sedimentation** — the inflow, dispersion and resuspension of sediments — has been a natural phenomenon in the Region since the current sea level was reached about 6500 years ago.<sup>44,164,165</sup> Inshore areas are most exposed to sedimentation, especially areas close to river mouths. Longshore drift, tides and currents redistribute sediment across the continental shelf and along the coast. The deposited sediments can be colonised by mangroves and seagrass meadows.

Levels of **light** control the depth range of marine plants (for example, seagrasses and algae), as well as all animals which have a symbiotic dependence on photosynthesis (for example, corals). Light attenuation is a measure of decrease in light availability per metre in the water column, and is influenced by depth and the amount of sediment, dissolved substances and plankton in the water.<sup>166</sup> Thus, it is a more limiting factor in inshore areas compared to offshore habitats less affected by catchment run-off. The availability of light is central to the productivity of seagrass meadows, as well as the symbiotic relationship between corals and algae and the reef building activities of corals, and calcareous and coralline algae.

**Nutrient cycling** plays a critical role in maintaining biodiversity. Most nutrient concentrations (for example, nitrogen and phosphorus) in the open ocean are very low — and, with the exception of upwelling areas, they are effectively nutrient deserts.<sup>167</sup> Low concentrations of nitrates, in particular, severely limit productivity. Open ocean coral reefs in non-upwelling areas accommodate nutrient deficiency by having a high level of nutrient cycling.<sup>167</sup> For reefs nearer land, additional nutrients are derived from terrestrial sources. Nitrates also boost algal production and can result in planktonic algal blooms that, in turn, trigger secondary blooms of planktonic consumers. Algal blooms can affect ecosystems when there is an overabundance of nutrients, causing deleterious effects known as eutrophication.<sup>168</sup>

The ocean plays an important role in global **carbon cycling**.<sup>169</sup> Carbon is exchanged in and out of the ocean daily, but it can also be stored there for millions of years. Overall, the ocean is a carbon 'sink' as it takes more carbon from the atmosphere than it gives out. Carbon dioxide from the atmosphere dissolves in the surface waters of the ocean where it stays in dissolved form, or gets taken up by organisms. Photosynthesis by phytoplankton (free-floating microscopic algae) in surface waters turns the carbon into organic matter. Other organisms use carbon to make calcium carbonate for their shells and skeletons. The use of carbon by biological and chemical processes allows more carbon dioxide to enter the water from the atmosphere. The living organic matter in the ocean eventually decomposes and is broken down into simpler forms of matter (detritus). Some sinks to the ocean floor, where it forms layers of carbon-rich sediments. This part of the carbon cycle can lock up carbon for millions of years.<sup>169</sup>

The total amount of carbon in the ocean is about 50 times greater than the amount in the atmosphere, and, at least half of the atmospheric oxygen comes from photosynthesis by marine plants. Currently, almost half the carbon emitted to the atmosphere by fossil fuel burning is absorbed and sequestered into the ocean.<sup>169,170</sup>



It is estimated that an increase in the amount of carbon dioxide absorbed by the ocean has already changed global **ocean acidity**, reducing the ocean's pH by 0.1 units compared to the long-term average. The acidity of the ocean is of vital importance to those marine animals that grow shells or skeletons — more acidic water reduces their ability to grow strong skeletons.<sup>171,172</sup>

**Redox (reduction-oxidation)** processes are chemical reactions where the oxidation state of atoms and elements are changed through the transfer of electrons.<sup>173</sup> They are vital chemical reactions fundamental to ecosystems. Processes such as photosynthesis, denitrification and respiration are all important marine redox reactions.

As part of the carbon cycle, most food webs are originally based on **primary production**: the production of food by photosynthesis using energy from the sun. In tropical marine ecosystems, contributions to primary production come not only from plants such as macroalgae, turf algae, seagrasses and mangroves, but also, in large part, from phytoplankton and corals which have microscopic algae within their tissues. Primary production is closely linked to concentrations of available inorganic nutrients.<sup>174</sup>

Marine sediments undergo vertical mixing through a process known as **bioturbation**. Bioturbation is the process by which sediment macro-biology mix surface sediments. Bioturbation can promote sediment oxidation and promote nutrient conversion. Bioturbation can help to both consolidate and bind unconsolidated sediments, and can partially unconsolidate bound surface sediments.

The role of **microbial processes** in trophic interactions is not well known, having received only very limited scientific study.<sup>175</sup> Marine snow, a continuous shower of mostly organic detritus falling from the upper layers of the water column, and the increasing frequency of diseases are visible outcomes of changes in bacterial levels in the water column and in benthic organisms.<sup>176</sup> However, most microbial processes remain invisible. Some microbial processes cause sub-surface water to become completely depleted of oxygen. Others, including the processing of detritus and the maintenance of biologically active substrates, are fundamental to the functioning of benthic ecosystems. Microbial processes are very responsive to organic and inorganic nutrient concentrations and environmental conditions.<sup>175</sup>

**Herbivory** is a key process for the health and resilience of coral reefs.<sup>177,178,179</sup> By feeding on the algae of a reef, herbivores (for example, some fish and green turtles) reduce the amount that competes with corals for space to establish and grow. Herbivores have a particularly important role in maintaining reef ecosystems because, without their constant presence, many reefs would be rapidly overtaken by algae.<sup>177,178</sup> Fish are important herbivores on the coral reefs of the Great Barrier Reef.<sup>179</sup> Dugongs are an important herbivore in seagrass meadow habitats.

**Particle feeding** is undertaken by a wide range of animals from the very large (whale sharks) to the microscopic (copepods). Most marine invertebrates, such as sea cucumbers, scallops, sponges, corals and many crustaceans (for example, prawns, crabs and lobsters) are particle feeders.<sup>167</sup> They are an important part of the energy and nutrient cycle, feeding on detritus, bacteria, plankton and particulate nutrients.<sup>167</sup>

**Predation** — animals consuming other animals — has a fundamental influence on marine ecosystems by controlling the abundance of many prey animals and causing a range of cascading effects through the food web.<sup>167</sup> Predators and top predators in coral reef ecosystems include most big bony fish and sharks, as well as a wide array of smaller finfish and invertebrates, seabirds, some marine turtles and marine mammals.

**Symbiosis**, the interdependence of different organisms for the benefit of one or both participants, is much more prevalent in the oceans than on land. Examples include the association between giant anemones and anemone fish and the *Chelonibia* barnacle found on marine turtles.<sup>180</sup> One of the most important symbioses is between hard corals and microscopic algae.<sup>181</sup> The algae photosynthesise like other green plants however, up to 95 per cent of the nutrients produced are used by the coral host organism.<sup>181,182</sup> In coral reefs, reef building is literally powered by the sun via algal photosynthesis. Symbiosis with algae also occurs in soft corals and anemones, as well as in an assortment of other animals like sponges, flatworms and molluscs.<sup>183</sup> This symbiosis is an example of **mutualism**, where both organisms benefit from the relationship.

In **parasitism**, another type of symbiosis, one organism (the parasite) benefits at the expense of another (the host). For example, parasitic isopod crustaceans live on the gills of many reef fish, gaining shelter and food. High numbers of parasites can be an indicator of poor environmental conditions.<sup>184</sup>

**Recruitment** occurs when juvenile organisms survive to be added to a population. It plays an important role in the replenishment of populations, reef building and connectivity.<sup>185</sup> For coral reefs, it is especially important after disturbances such as mass coral bleaching or cyclones. The intensity and timing of fish spawning events plays an important role in the large scale patterns of recruitment.<sup>186</sup>

**Competition** for all resources, including space, nutrients and food is always intense in tropical marine ecosystems. This is partly because they are diverse, so that individual species have many others to compete with, and also because the habitats are three-dimensional. Competition between corals and algae for space is a fundamental process on coral reefs.<sup>187</sup> For coral reefs to be maintained in the ecosystem, there must be continual settlement and growth of juvenile corals. This recruitment may be hampered if a reef becomes overgrown by algae.<sup>188</sup>

On long time scales, **evolution** is an important part of natural systems such as the Great Barrier Reef, operating in both time and space.<sup>189</sup> Speciation, mutation and adaptation are all important components of evolution.


**Extinction** of populations and species is another ecosystem process, caused by reductions in population size or genetic variation, and often associated with the removal or modification of habitats.<sup>190</sup>

**Reef building** is both a geomorphological and an ecological process including **calcification** by living coral, **erosion, deposition** and **accretion**.<sup>191</sup> Only a small proportion of a coral reef is living coral — the remainder is coral-based pavement, boulders, fragments, beach-rock accretions and sediment. Cyclone wave action can break coral and mobilise vast amounts of sediment so that reef shapes change and cays appear and disappear.<sup>44</sup> Coral-based sediment fills in depressions and can consolidate and accrete to form soft rock.<sup>44</sup>

**Erosion** can occur through physical processes such as waves, wind and currents; chemical processes such as ocean acidification; or biological processes such as bioerosion. Erosion during previous periods of lower sea levels has influenced the shape of the continental shelf and slope which, in turn, modifies behaviour of currents and upwellings.<sup>192</sup> Today, as coral reefs are growing they are also being eroded, usually through physical abrasion (such as by waves) and bioerosion (by molluscs, marine worms, sponges, crustaceans, echinoderms and fish). Fish are the most effective bioeroders, with one adult humphead parrotfish consuming more than five tonnes of structural reef carbonate per year.<sup>193</sup> This biological activity results in the breakdown of the reef substratum and nutrient cycling.<sup>194</sup> Erosion is also a major process in shaping the Marine Park's coastline.

On a geological timescale, **tectonic forces** such as continental drift, friction, subsidence and the vertical movement of the seafloor, have played a role in shaping the Australian continent, its mountain ranges and its continental shelf.

Ecological **connectivity** is the movement of biota and materials across and through landscapes and seascapes. It may be related to **migration** between breeding and foraging areas (for example, humpback whales and marine turtles), movement by ocean currents (for example, coral spawn, fish spawn) or dispersal movements by species (for example, dugongs). The diagram of the life cycle of the red emperor fish (Figure 4.15) illustrates the importance of a variety of coastal and marine ecosystems and their relationship to this commercially and recreationally important fish. Connectivity through migration forms an important part of the life cycle of a number of species that travel through and within the Region, including a number of species that are matters of national environmental significance (for example, marine turtles and whales). **Genetic connectivity**, the opportunity for a plant or animal to find and potentially produce viable offspring with other individuals of the same species, is a crucial process in the ecosystem. Currents can play a major role in the genetic connectivity of some plants and animals for example, transporting some larvae for thousands of kilometres. At the other extreme, site specific animals that bear live young are more restricted in their genetic connectivity.<sup>195</sup>



**Figure 4.15 Crossing the blue highway**

The journey of the red emperor — a fish popular with commercial and recreational fishers — is an example of the importance of connectivity to matters of national environmental significance. At different stages of its growth and development the species uses different habitats across the breadth of the Reef (© Russell Kelley [www.russellkelley.info](http://www.russellkelley.info) and the Australian Coral Reef Society [www.australiancoralreefsociety.org](http://www.australiancoralreefsociety.org)).

### The importance of connectivity

Connectivity between and within species and habitats is a key process in the Great Barrier Reef ecosystem. Connections may occur over short periods, from generation to generation, over seasons or in cycles. There are connections between estuarine and inshore habitats and those further offshore; north-south connections between habitats; connections between open water and seabed habitats; and larger scale connections to environments outside the Region such as the Hervey Bay area and further south, Torres Strait and the Coral Sea. Connectivity is important to every aspect of the Reef, including processes as different as nutrient flows, migration, larval dispersal and gene flow. Species or habitats with low natural connectivity are likely to be especially vulnerable to disturbance or local extinctions.

Migration is a key component of connectivity on a broad scale. A number of listed species live in the Reef for only part of the year or for part of their life — this includes the humpback whale, green, loggerhead and hawksbill turtles, and some seabirds. Some fish species, like marlin, are also highly mobile and travel well beyond the Region for parts of their life cycle.<sup>196</sup> Conservation threats to these migratory species often occur well beyond the Region. For example, some marine turtles that nest or forage in the Region may be injured or killed in areas hundreds, and even thousands, of kilometres away.<sup>92,102</sup> Other species, such as dugongs, can move along the coast, especially if they are in search of food, after impacts in their regular feeding areas.<sup>197,198,199</sup>

Larval corals and fishes disperse widely between reefs, usually carried by water movement, and often strongly influenced by larval behaviour.<sup>200</sup> There is now definitive evidence of the larvae of two coral reef fish species transferring from areas that have been closed to fishing to adjacent fished areas. This illustrates the benefits of no-take (green) zones for reef fish conservation and sustainable harvesting.<sup>201</sup>

The life cycles of many fish include movement between marine habitats and adjacent estuaries, pools and floodplain habitats. These species depend on these habitats remaining healthy, connected and accessible. Within the marine environment there are ecological connections between mangroves and seagrass meadows adjacent to coral reefs, including for some reef fish populations.<sup>202</sup>

Connectivity between habitats can also increase the resilience of the Reef ecosystem. For example, connectivity between mangroves and coral reefs provides benefits for herbivorous fish populations which contribute to coral reef resilience by grazing on algae.<sup>203</sup>

The loss and modification of coastal wetlands and the deterioration of connecting water bodies has reduced or severed the connectivity between marine and adjacent freshwater habitats. This makes the role of the protected wetland systems such as the Shoalwater and Corio Bays Area and the Cape Bowling Green Bay Area wetlands of international importance even more important.

Birds play a key role in connecting different ecosystems, both locally and across the oceans. On a local scale, birds distribute plant species. For example, about 63 per cent of the plant species on Milman Island in the northern Great Barrier Reef are believed to have been introduced by birds.<sup>204</sup> One key species is the pied imperial pigeon which roosts on islands within the Reef at night and feeds in mainland forests during the day. Migratory waders, some of which travel from as far afield as Siberia, provide connectivity to ecosystems across the globe. As with migratory marine species, populations of waders overwintering in Australia are affected by factors in other parts of their migratory journey.

Traditional Owners describe the importance of connectivity as part of their association with their land and sea country. It encompasses the biological connectivity vital to a healthy marine environment and the cultural connectivity that links them to the natural environment.

*“Cultural value is how it is all connected and makes us connected to country”*<sup>205</sup>

## 4.10 Realising connectivity in the landscape

The Great Barrier Reef and adjacent catchment encompasses a variety of ecosystem types that are connected both spatially and temporally. The Authority has built on the information presented in the *Informing the Outlook for Great Barrier Reef Coastal Ecosystems Report*<sup>38</sup> and the *Coastal Ecosystem Assessment Frameworks*<sup>206</sup> to map catchment coastal ecosystem areas most important for supporting the biodiversity and ecological processes of the World Heritage Area and Marine Park due to their proximity to, and connection with the ecosystems of the Great Barrier Reef (Figure 4.16).


The Authority has developed a method to categorise coastal ecosystems into distinct components based on the services and functions provided to the World Heritage Area and Marine Park<sup>206</sup>. The method enables an assessment of the Reef's ecological functions supported by coastal ecosystems, the risks to these functions, and the cumulative impacts at work across the catchment that are affecting the long-term health of the Great Barrier Reef over a range of scales.

The following steps were used in considering terrestrial ecosystems and their relationship to the World Heritage Area and Marine Park:

- identifying coastal ecosystems in the catchment using the *Coastal Ecosystems Assessment Framework*<sup>206</sup>
- describing the relationship between coastal ecosystems within an area to those adjacent to and downstream of that area



- identifying the environmental assets and characteristics, including biophysical, ecological and hydrological processes
- consideration of surface and groundwater recharge and discharge, and areas of high biodiversity value
- identifying and mapping land use constraints, natural hazards, important ecosystem types, waterways and stream orders
- identifying areas that support the Reef's biodiversity and ecological functions or contain native vegetation.



**Figure 4.16 A method for identifying and prioritising coastal ecosystem functional connections to the Great Barrier Reef World Heritage Area and the Great Barrier Reef Marine Park**

The maps, clockwise from top left, show (a) Mulgrave–Russell, (b) Haughton, (c) Fitzroy and (d) Baffle basins within the Great Barrier Reef catchment.

The analysis<sup>207</sup> takes into account values such as habitats, likelihood of erosion, highest astronomical tide, and land use for the Mulgrave-Russell<sup>208,209</sup>, Haughton<sup>210,211</sup>, Fitzroy<sup>212,213</sup>, and Baffle<sup>214,215</sup> basins. It represents the surface level only and does not include groundwater. The darker areas of the catchment shown here are of higher importance to the healthy functioning of the Great Barrier Reef ecosystem, supporting its biodiversity and ecological processes, because of their proximity to and connectivity with the Great Barrier Reef. The analysis takes into account wetlands and areas that are frequently inundated or flooded, as well as areas influenced by tidal processes including sea level changes and storm surges. It represents the surface level hydrology only and does not include groundwater.

## 4.11 Connections between matters of national environmental significance

As outlined above, the matters of national environmental significance, defined and protected under the EPBC Act, range from individual listed threatened and migratory species to the entire Great Barrier Reef Marine Park and World Heritage Area. The Marine Park and World Heritage Area provide critical habitat and ecosystem processes that support the life cycles of listed species. Conversely, listed species are important natural attributes of the Marine Park's environment and the outstanding universal value of the world heritage property. Accordingly, there is substantial overlap and connections between the matters of national environmental significance relevant to the Region. In addition, the matters of national environmental significance and the values that relate to them do not occur in isolation, but are embedded within complex and dynamic systems. They are distributed throughout the Region; there is no part that does not contain values of national significance.



Effective future protection of these matters relies on the integration of management measures and a focus on protection of the Great Barrier Reef environment as a whole.

Given the scale and complexity of the Region and the diversity of values that relate to the matters of national environmental significance, the report focuses on a suite of key values and attributes relevant to the Region's matters of national environmental significance.

For the strategic assessment, the key values and attributes of the seven matters of national environmental significance in the Region are grouped into four broad categories:

- biodiversity, including the Region's habitats and species — some of which are listed migratory or threatened species or are part of a wetland of international importance
- geomorphological features
- Indigenous and historic heritage
- community benefits of the environment, comprising the cultural, social and economic benefits derived from the Region's environment.

The key values and attributes presented in Table 4.8 are based on those identified in the description of each matter in Sections 4.2 to 4.8. The key environmental processes relevant to each matter are based on the description in Section 4.9, and are presented in Table 4.9.

For some of the matters of national environmental significance (for example, the listed threatened and listed migratory species), only a subset of the values and attributes are identified as relevant. For the Great Barrier Reef Marine Park and Commonwealth marine areas, all those listed are considered relevant. This is because of the broad definition of the 'environment' within the governing legislation of these two matters, including heritage, social and cultural aspects.

It is important to note that not all attributes of the outstanding universal value of the Great Barrier Reef world heritage property are identified individually in Table 4.8. Where relevant, certain attributes mentioned in the Statement of Outstanding Universal Value<sup>2</sup> have been grouped into an overarching value for the property. For example, the statement identifies attributes such as 'migrating whales' and 'humpback whale calving'. These and other relevant attributes have been grouped under the value 'whales'. In addition, although not specifically identified in Table 4.8, attributes such as those relating to the life cycle of particular species are considered throughout the assessment. Integrity, as a fundamental concept of outstanding universal value, is considered in Table 4.8 and, where relevant, throughout the report.

Inclusion of the terrestrial habitats that support the Great Barrier Reef as values reflects the important role these habitats have in maintaining ecosystem processes and supporting biodiversity within the Region.

**Table 4.8 Key values and attributes of matters of national environmental significance**

For the World Heritage Area, connections are based on the Statement of Outstanding Universal Value<sup>2</sup>. For the listed species, the key linkages (such as important habitats and components of their diets) are shown for each group of species. For wetlands of international importance, the connections are those discussed in the Ramsar Convention information sheet.<sup>151</sup>

Key values and attributes	World heritage properties					Listed migratory and threatened species											
	Criterion i (now viii)	Criterion ii (now ix)	Criterion iii (now vii)	Criterion iv (now x)	Integrity	Great Barrier Reef Marine Park	National heritage places	Commonwealth marine areas	Marine turtles	Estuarine crocodiles	Whales	Dolphins	Dugongs	Sharks and rays	Seabirds	Shorebirds	Wetlands of international importance
Biodiversity — Great Barrier Reef habitats																	
Islands		●	●	●	●	●	●	●	●	●					●	●	●
Beaches and coastlines			●		●	●	●	●	●	●					●	●	●
Mangrove forests			●	●	●	●	●	●	●	●				●	●	●	●
Seagrass meadows				●	●	●	●	●	●			●	●	●			●
Coral reefs (<30 m)		●	●	●	●	●	●	●	●					●			●

Key values and attributes	World heritage properties								Listed migratory and threatened species								Wetlands of international importance
	Criterion i (now viii)	Criterion ii (now ix)	Criterion iii (now vii)	Criterion iv (now x)	Integrity	Great Barrier Reef Marine Park	National heritage places	Commonwealth marine areas	Marine turtles	Estuarine crocodiles	Whales	Dolphins	Dugongs	Sharks and rays	Seabirds	Shorebirds	
Deeper reefs (>30 m)		●	●	●	●	●	●	●	●					●			
Lagoon floor				●	●	●	●	●				●	●	●			
Shoals				●	●	●	●	●				●		●	●		
<i>Halimeda</i> banks		●			●	●	●	●	●								
Continental slope					●	●	●	●									
Open waters			●	●	●	●	●	●	●	●	●	●	●	●	●		●
<b>Biodiversity — terrestrial habitats that support the Great Barrier Reef</b>																	
Saltmarshes						●		●		●					●	●	●
Freshwater wetlands						●		●		●					●	●	●
Forested floodplains						●		●									●
Heath and shrublands						●		●									
Grass and sedgeland						●		●								●	●
Woodlands						●		●									●
Forests						●		●									●
Rainforests			●			●		●									
Connecting water bodies				●	●	●	●	●	●	●		●	●	●	●	●	●
<b>Biodiversity — species</b>																	
Mangroves			●	●	●	●	●	●									●
Seagrasses				●	●	●	●	●	●				●				●
Macroalgae				●	●	●	●	●	●								
Benthic microalgae				●	●	●	●	●									
Corals			●	●	●	●	●	●									
Other invertebrates		●		●	●	●	●	●	●		●	●		●	●	●	●
Plankton and microbes				●	●	●	●	●			●						
Bony fish		●	●	●	●	●	●	●			●	●		●	●	●	●
Sharks and rays				●	●	●	●	●			●			●			
Sea snakes				●	●	●	●	●									
Marine turtles			●	●	●	●	●	●	●								●
Estuarine crocodiles				●	●	●	●	●		●							
Seabirds			●	●	●	●	●	●							●		
Shorebirds				●	●	●	●									●	●
Whales			●	●	●	●	●	●			●						
Dolphins				●	●	●	●	●				●					
Dugongs				●	●	●	●	●					●				●

Key values and attributes	World heritage properties								Listed migratory and threatened species									Wetlands of international importance
	Criterion i (now viii)	Criterion ii (now ix)	Criterion iii (now vii)	Criterion iv (now x)	Integrity	Great Barrier Reef Marine Park	National heritage places	Commonwealth marine areas	Marine turtles	Estuarine crocodiles	Whales	Dolphins	Dugongs	Sharks and rays	Seabirds	Shorebirds		
Geomorphological features																		
Coral reefs	●		●		●	●	●	●										
Islands and shorelines	●		●		●	●	●	●	●							●	●	
Channels and canyons	●				●	●	●	●										
River deltas	●				●	●	●	●										
Halimeda banks	●				●	●	●	●										
Seagrass meadows	●				●	●	●	●	●								●	
Indigenous heritage																		
Cultural practices, observances, customs and lore		●			●	●	●	●	●	●	●	●	●	●	●	●	●	
Sacred sites, sites of particular significance, places important for cultural tradition		●			●	●	●	●									●	
Stories, songlines, totems and languages		●			●	●	●	●	●	●	●	●	●	●	●	●	●	
Indigenous structures, technology, tools and archaeology		●			●	●	●	●									●	
Historic heritage																		
Places of historic significance — historic shipwrecks						●		●										
Places of historic significance — World War II features and sites						●		●										
Places of historic significance — lightstations						●		●										
Places of historic significance — other						●		●									●	
Places of scientific significance (research stations, expedition sites)						●		●										
Places of social significance — iconic sites						●		●										
Community benefits of the environment																		
Income						●		●									●	
Employment						●		●									●	
Understanding						●		●										
Appreciation			●			●	●	●									●	
Enjoyment			●			●	●	●										
Access to Reef resources						●		●										
Personal connection						●		●										
Health benefits						●		●										
Aesthetics			●		●	●	●	●	●	●	●	●	●	●	●	●		

**Table 4.9 Key environmental processes relevant to matters of national environmental significance**

For the World Heritage Area, connections are based on the Statement of Outstanding Universal Value<sup>2</sup>. For listed species, processes that have a major supporting role in maintaining the species are shown (for example, the role that beaches play in the nesting of listed marine turtles). For wetlands of international importance, the connections shown are those discussed in the Ramsar Convention information sheet.<sup>151</sup>

	World heritage properties					Listed migratory and threatened species											
Key environmental processes	Criterion i (now viii)	Criterion ii (now ix)	Criterion iii (now vii)	Criterion iv (now x)	Integrity	Great Barrier Reef Marine Park	National heritage places	Commonwealth marine areas	Marine turtles	Estuarine crocodiles	Whales	Dolphins	Dugongs	Sharks and rays	Seabirds	Shorebirds	Wetlands of international importance
Waves, currents and tides	●	●			●	●	●	●	●						●	●	
Cyclones	●	●			●	●	●	●	●	●		●	●	●	●	●	
Wind	●	●			●	●	●	●									
Sedimentation	●	●			●	●	●	●	●				●			●	●
Sea level	●	●			●	●	●	●	●	●						●	●
Sea temperature		●			●	●	●	●	●	●							
Light		●			●	●	●	●	●				●				
Nutrient cycling		●			●	●	●	●									●
Ocean acidity		●			●	●	●	●									
Freshwater inflow and salinity		●			●	●	●	●									●
Microbial processes		●			●	●	●	●									
Particle feeding		●			●	●	●	●									
Primary production		●			●	●	●	●	●				●				
Herbivory		●			●	●	●	●	●				●				
Predation		●			●	●	●	●	●	●	●	●		●	●	●	
Symbiosis		●			●	●	●	●									
Competition		●			●	●	●	●	●	●	●	●	●	●	●	●	
Connectivity	●	●			●	●	●	●	●	●	●	●	●	●	●	●	●
Recruitment		●			●	●	●	●	●	●	●	●	●	●	●	●	●
Reef building	●	●	●		●	●	●	●									
Tectonic forces	●	●	●		●	●	●	●									

## 4.12 Summary of conclusions

- The seven matters of national environmental significance relevant to the Great Barrier Reef Region are interconnected and overlapping. They range from individual listed threatened and migratory species to the entire Great Barrier Reef Marine Park and world heritage property.
- The matters of national environmental significance are distributed throughout the Region, and there is no part that does not contain values relevant to them.
- A total of 62 key values relevant to the matters of national environmental significance in the Region are identified. Twenty key environmental processes are also identified.
- The key values and attributes are combined into one comprehensive set which forms the basis of the assessments throughout the report. They are grouped into four broad categories: biodiversity, including the Region's habitats and species; geomorphological features; Indigenous and historic heritage values; and community benefits derived from the environment.



## References

1. UNESCO Intergovernmental Committee for the Protection of the World Cultural and Natural Heritage 2012, *Operational guidelines for the implementation of the World Heritage Convention*, UNESCO, <<http://whc.unesco.org/archive/opguide12-en.pdf>>.
2. Department of Sustainability, Environment, Water, Population and Communities 2012, *Statement of Outstanding Universal Value: Great Barrier Reef World Heritage Area*, DSEWPac, Canberra.
3. International Union for the Conservation of Nature 2010, *Guidance on the preparation of retrospective statements of outstanding universal value for world heritage properties*, IUCN, Gland, Switzerland.
4. Whiteway, T., Smithers, S., Potter, A. and Brooke, B. 2013a, *Geological and geomorphological features of outstanding universal value in the Great Barrier Reef World Heritage Area: Technical report prepared for the Department of Sustainability, Environment, Water, Population and Communities*, Geoscience Australia and James Cook University, Townsville.
5. UNESCO 2012, *Mission report Great Barrier Reef*, UNESCO, Paris.
6. Lucas, P.H.C., Webb, T., Valentine, P.S. and Marsh, H. 1997, *The outstanding universal value of the Great Barrier Reef World Heritage Area*, Great Barrier Reef Marine Park Authority, Townsville.
7. Context Pty Ltd 2013, *Defining the aesthetic values of the Great Barrier Reef*, Department of Sustainability, Environment, Water, Population and Communities, Canberra.
8. UNESCO 2013a, *Provisional agenda, State of conservation of World Heritage properties inscribed on the World Heritage List*, UNESCO, Paris, France.
9. Great Barrier Reef Marine Park Authority 2009, *Great Barrier Reef Outlook Report 2009*, GBRMPA, Townsville.
10. Turner, M. and Batianoff, G.N. 2007, Vulnerability of island flora and fauna of the Great Barrier Reef to climate change, in *Climate change and the Great Barrier Reef: a vulnerability assessment*, eds J.E. Johnson and P.A. Marshall, Great Barrier Reef Marine Park Authority and Australian Greenhouse Office, Townsville, pp. 621-666.
11. Lovelock, C. and Clarke, S. 1993, *Field guide to the mangroves of Queensland*, Australian Institute of Marine Science, Townsville.
12. Wolanski, E., Mazda, Y., Furukawa, K., Ridd, P., Kithaka, J., Spagnol, S. and Stieglitz, T. 2001, Water circulation in mangroves and its implications for biodiversity, in *Oceanographic processes of coral reefs: physical and biological links in the Great Barrier Reef*, ed. E. Wolanski, CRC Press, Boca Raton, Florida, USA, pp. 53-76.
13. Duke, N.C. 2006, *Australia's mangroves: the authoritative guide to Australia's mangrove plants*, University of Queensland, Brisbane.
14. Duke, N.C. and Larkum, A.W.D. 2008, Mangroves and seagrasses, in *The Great Barrier Reef: biology, environment and management*, eds P. Hutchings, M. Kingsford and O. Hoegh-Guldberg, CSIRO Publishing, Collingwood, pp. 156-170.
15. Waycott, M., Collier, C., McMahon, K., Ralph, P., McKenzie, L.J., Udy, J. and Grech, A. 2007, Vulnerability of seagrasses in the Great Barrier Reef to climate change, in *Climate Change and the Great Barrier Reef: a vulnerability assessment*, eds J.E. Johnson and P.A. Marshall, Great Barrier Reef Marine Park Authority and Australian Greenhouse Office, Townsville, pp. 193-235.
16. McLeod, J.J. 1997, Large scale seagrass dieback, in *Report to governments on actions necessary for dugong conservation in the Great Barrier Reef and Hervey Bay/Great Sandy Strait*, Great Barrier Reef Marine Park Authority, Townsville.
17. Coles, R.G., Lee Long, W.J., McKenzie, L.J., Short, M., Rasheed, M.A. and Vidler, K. 1995, *Distribution of deep-water seagrass habitats between Cape Weymouth and Cape Tribulation, northeastern Queensland*, Department of Primary Industries (Queensland), Cairns.
18. Coles, R., Lee Long, W., McKenzie, L.J., Roelofs, A. and De'ath, G. 2000, Stratification of seagrasses in the Great Barrier Reef World Heritage Area, northeastern Australia, and the implications for management, *Biologia Marina Mediterranea* 7(2): 345-348.
19. Coles, R., Beumer, J., McKenzie, L.J. and Rasheed, M. 2005, Seagrass fisheries habitat in Queensland coastal waters and issues for their protection and management, in *Proceedings of the Rainforest Meets Reef: Joint Conference of CRC Reef and Rainforest CRC, 22-24 November 2005, Townsville*, eds. L. Goggin and T. Harvey, CRC Reef Research Centre, Townsville, pp.29.
20. Collier, C. and Waycott, M. 2009, *Drivers of change to seagrass distributions and communities on the Great Barrier Reef, literature review and gaps analysis. Report to the Marine and Tropical Sciences Research Facility*, Reef and Rainforest Research Centre, Cairns.
21. Veron, J.E.N., DeVantier, L.M., Turak, E., Green, A.L., Kininmonth, S., Stafford-Smith, M. and Peterson, N. 2009, Delineating the Coral Triangle, *Galaxea* 11(2): 91-100.
22. Waycott, M., McMahon, K., Mellors, J., Calladine, A. and Kleine, D. 2004, *A guide to tropical seagrasses of the Indo-West Pacific*, James Cook University, Townsville.
23. Lanyon, J. 1986, *Guide to the identification of seagrasses in the Great Barrier Reef Region*, Great Barrier Reef Marine Park Authority, Townsville.
24. Great Barrier Reef Marine Park Authority 2012a, *A vulnerability assessment for the Great Barrier Reef: Seagrass*, GBRMPA, Townsville.
25. Hinderstein, L.M., Marr, J.C.A., Martinez, F.A., Dowgiallo, M.J., Puglise, K.A., Pyle, R.L., Zawada, D.G. and Appeldoorn, R. 2010, Theme section on "Mesophotic coral ecosystems: Characterization, ecology and management", *Coral Reefs* 29(2): 247-251.
26. Kahng, S.E., Garcia-Sais, J.R., Spalding, H.L., Brokovich, E., Wagner, D., Weil, E., Hinderstein, L. and Toonen, R.J. 2010, Community ecology of mesophotic coral reef ecosystems, *Coral Reefs* 29(2): 255-275.
27. Bridge, T.C.L. 2012, *Habitats and sessile benthic megafaunal communities in the mesophotic zone of the Great Barrier Reef World Heritage Area*, PhD Thesis, James Cook University, Townsville.
28. Harris, P.T., Bridge, T.C.L., Beaman, R.J., Webster, J.M., Nichol, S.L. and Brooke, B.P. 2013, Submerged banks in the Great Barrier Reef, Australia, greatly increase available coral reef habitat, *ICES Journal of Marine Science* 70(2): 284-293.
29. Pickrell, J. 2009, *The drowned reef: World's largest fossil*, *Cosmos* Apr 1, <<http://www.cosmosmagazine.com/features/drowned-reef/>>.
30. Beaman, R. 2009, *Understanding coldwater coral ecosystems on the Great Barrier Reef margin: Progress Report for AINGRA09006*, Australian Institute of Nuclear Science and Engineering, New South Wales.
31. Pitcher, C.R., Doherty, P., Arnold, P., Hooper, J., Gribble, N.A., Bartlett, C., Browne, M., Campbell, N., Cannard, T., Cappel, M., Carini, G., Chalmeres, S., Cheers, S., Chetwynd, D., Colegax, A., Coles, R., Cook, S., Davie, P., De'ath, G., Devereux, D., Done, B., Donovan, T., Ehrke, B., Ellis, N., Ericson, G., Fellegara, I., Forcey, K., Furey, M., Gledhill, D., Good, N., Gordon, S., Haywood, M., Hendricks, M., Jacobsen, I., Johnson, J., Jones, M., Kininmonth, S., Kistle, S., Last, P., Leite, A., Amrks, S., McLeod, I., Oczkowicz, S., Robinson, M., Rose, C., Seabright, D., Sheils, J., Sherlock, M., Skelton, P., Smith, D., Smith, G., Speare, P., Stowar, M., Strickland, C., Van der Geest, C., Venables, W., Walsh, C., Wassenberg, T.J., Welna, A. and Yearsley, G. 2007, *Seabed biodiversity on the continental shelf of*

the Great Barrier Reef World Heritage Area.

AIMS/CSIRO/QM/QDPI CRC Reef Research Task final report, CSIRO Marine and Atmospheric Research, Hobart.

32. Cappel, M. and Kelley, R. 2001, Connectivity in the Great Barrier Reef World Heritage Area: An overview of pathways and processes, in *Oceanographic processes of coral reefs: Physical and biological links in the Great Barrier Reef*, ed. E. Wolanski, CRC Press, Boca Raton, Florida, pp. 161-187.

33. Speare, P. and Stowar, M. 2007, *Preliminary findings from the first baseline survey of the Magnetic Shoals*, Reef and Rainforest Research Centre Ltd, Cairns.

34. Cappel, M., De'ath, G., Stowar, M.J., Johannson, C. and Doherty, P.J. 2009a, *The influence of zoning (closure to fishing) on fish communities of the deep shoals and reef bases of the southern Great Barrier Reef Marine Park. Part 2: development of protocols to improve accuracy in baited video techniques used to detect effects of zoning*, Reef and Rainforest Research Centre, Cairns.

35. Cappel, M., MacNeil, M.A., Stowar, M.J. and Doherty, P.J. 2009b, *The influence of zoning (closure to fishing) on fish communities of the deep shoals and reef bases of the southern Great Barrier Reef. Part 1: Baited video surveys of the Pompey, Swain and Capricorn-Bunker groups of reefs off Mackay and Gladstone. Report to the Marine and Tropical Sciences Research Facility*, Reef and Rainforest Research Centre, Cairns and Australian Institute of Marine Science, Townsville.

36. Beaman, R.J., Webster, J.M. and Wust, R.A.J. 2008, New evidence for drowned shelf edge reefs in the Great Barrier Reef, Australia, *Marine Geology* 247(1-2): 17-34.

37. Webster, J.M., Beaman, R.J., Bridge, T., Davies, P.J., Byrne, M., Williams, S., Manning, P., Pizarro, O., Thornborough, K., Woolsey, E., Thomas, A. and Tudhope, S. 2008, From corals to canyons: The Great Barrier Reef margin, *EOS, Transactions, American Geophysical Union* 89(24): 217-218.

38. Great Barrier Reef Marine Park Authority 2012b, *Informing the outlook for Great Barrier Reef coastal ecosystems*, GBRMPA, Townsville.

39. Pitcher, C.R., Austin, M., Burrige, C.Y., Bustamante, R.H., Cheers, S.J., Ellis, N., Jones, P.N., Koutsoukos, A.G., Moeseneder, C.H., Smith, G.P., Venables, W. and Wassenberg, T.J. 2008, *Recovery of seabed habitat from the impact of trawling in the far northern section of the Great Barrier Reef Marine Park: CSIRO final report to GBRMPA*, CSIRO Marine and Atmospheric Research, Cleveland, Queensland.

40. Hutchings, P., Kingsford, M. and Hoegh-Guldberg, O. 2008, *The Great Barrier Reef: biology, environment and management*, CSIRO Publishing, Collingwood.

41. Hopley, D. 1982, *Geomorphology of the Great Barrier Reef: Quarternary development of coral reefs*, John Wiley-Interscience, New York.

42. Hopley, D. 1983, Morphological classification of shelf reefs: A critique with special reference to the Great Barrier Reef, in *Perspectives on Coral Reefs*, ed. D.J. Barnes, Brian Clouston, Canberra, pp. 180-199.

43. Smithers, S.G. 2011, Fringing reefs, in *Encyclopedia of Modern Coral Reefs*, ed. D. Hopley, Springer, Berlin, pp. 430-446.

44. Hopley, D., Smithers, S.G. and Parnell, K.E. 2007, *The geomorphology of the Great Barrier Reef: Development, diversity and change*, Cambridge University Press, Cambridge, UK.

45. Maxwell, W.G.H. 1970, Deltaic patterns in reefs, *Deep Sea Research* 17: 1005-1018.

46. Veron, J.E.N. 1978, Deltaic and dissected reefs of the Far Northern Region, *Philosophical Transactions of the Royal Society of London, Series B* 284(999): 23-37.

47. Abbey, E., Webster, J.M. and Beaman, R.J. 2011, Geomorphology of submerged reefs on the shelf edge of the Great Barrier Reef: The influence of oscillating Pleistocene sea-levels, *Marine Geology* 288(1): 61-78.

48. Bridge, T. and Guinotte, J. 2013, *Mesophotic coral reef ecosystems in the Great Barrier Reef World Heritage Area: Their potential disturbance and possible role as refugia from disturbance. Research Publication No. 109*, Great Barrier Reef Marine Park Authority, Townsville.

49. Hopley, D., Parnell, K.E. and Isdale, P.J. 1989, The Great Barrier Reef Marine Park: dimensions and regional patterns, *Australian Geographical Studies* 27: 47-66.

50. Spenceley, A.P. 1982, *The geomorphological and zonal development of mangrove swamps in the Townsville Area, North Queensland*, James Cook University, Townsville.

51. Nott, J. 2006, Tropical Cyclones and the Evolution of the Sedimentary Coast of Northern Australia, *Journal of Coastal Research* 22(1): 49-62.

52. Chappell, J., Chivas, A., Wallensky, E., Polach, H. and Aharon, P. 1983, Holocene palaeo-environmental changes, central to north Great Barrier Reef inner zone, *BMR journal of Australian geology and geophysics* 8: 223-235.

53. Nott, J. and Hayne, M. 2001, High frequency of 'super cyclones' along the Great Barrier Reef over the past 5000 years, *Nature* 413(6885): 508-511.

54. Chivas, A., Chappell, J., Polach, H., Pillans, B. and Flood, P. 1986, Radiocarbon evidence for the timing and rate of island development, beach-rock formation and phosphatization at Lady Elliot Island, Queensland, *Marine Geology* 69: 273-287.

55. Hayne, M. and Chappell, J. 2001, Cyclone frequency during the last 5000 years at Curacao Island, north Queensland, Australia, *Palaeogeography, Palaeoclimatology, Palaeoecology* 168: 207-219.

56. Hopley, D. 1997, The age of the outer ribbon reef surface, Great Barrier Reef, Australia: implications for hydro-isostatic models, in *The Proceedings of the 3rd International Coral Reef Symposium*, ed. D. L. Taylor, Miami, Florida, USA, pp.23-28.

57. Backshall, D., Barnett, J., Davies, P., Duncan, D., Harvey, N., Hopley, D., Isdale, P., Jennings, J. and Moss, R. 1979, Drowned dolines: the blue holes of the Pompey Reefs, Great Barrier Reef, *Journal of Australian Geology and Geophysics* 4: 99-109.

58. Puga-Bernabeu, A., Webster, J.M., Beaman, R.J. and Guilbaud, V. 2013, Variation in canyon morphology on the Great Barrier Reef margin, north-eastern Australia: Influence of slope and barrier reefs, *Geomorphology* 191: 35-50.

59. Henrich, R., Cherubini, Y. and Meggers, H. 2010, Climate and sea level induced turbidite activity in a canyon system offshore the hyperarid Western Sahara (Mauritania): The Timiris Canyon, *Marine Geology* 275(1-4): 178-198.

60. Pierau, R., Hanebuth, T.J.J., Krastel, S. and Henrich, R. 2010, Late Quarternary climatic events and sea-level changes recorded by turbidite activity, Dakar Canyon, NW Africa, *Quarternary Research* 73(2): 385-392.

61. Pierau, R., Henrich, R., Preiu-Daimler, I., Krastel, S. and Geersen, J. 2011, Sediment transport and turbidite architecture in the submarine Dakar Canyon off Senegal, NW-Africa, *Journal of African Earth Sciences* 60(3): 196-208.

62. Puga-Bernabeu, A., Webster, J.M. and Beaman, R.J. 2012, Potential collapse of the upper slope and tsunami generation on the Great Barrier Reef margin, north-eastern Australia, *Natural Hazards* 66(2): 557-575.

63. Puga-Bernab  ,  ., Webster, J.M., Beaman, R.J. and Guilbaud, V. 2011, Morphology and controls on the evolution of a mixed carbonate-siliciclastic submarine canyon system, Great Barrier Reef margin, north-eastern Australia, *Marine Geology* 289(1-4): 100-116.

64. Whiteway, T., Smithers, S., Potter, A. and Brooke, B. 2013b, Beaman R, personal communication, in *Geological and geomorphological features of Outstanding Universal Value in the Great Barrier Reef World Heritage Area* Geoscience Australia, Canberra, and James Cook University, Townsville, pp. 113.

65. Webster, J.M., Beaman, R.J., Puga-Bernab  ,  ., Ludman, D., Renema, W., Wust, R.A.J., George, N.P.J., Reimer, P.J., Jacobsen, G.E. and Moss, P. 2012, Late Pleistocene history of turbidite sedimentation in a submarine canyon off the northern Great Barrier Reef, Australia, *Palaeogeography, Palaeoclimatology, Palaeoecology* 331-332: 75-89.

66. Furnas, M. 2003, *Catchments and corals: terrestrial runoff to the Great Barrier Reef*, Australian Institute of Marine Science, Townsville.

67. Drew, E.A. 1983, *Halimeda* biomass, growth rates and sediment generation on reefs in central Great Barrier Reef province, *Coral Reefs* 2: 101-110.
68. Marshall, J.F. and Davies, P.J. 1988, *Halimeda* bioherms of the Northern Great Barrier Reef, *Coral Reefs* 6: 139-148.
69. Coles, R., McKenzie, L.J. and Campbell, S. 2003, The seagrasses of eastern Australia, in *World Atlas of Seagrasses*, eds E.P. Green and T.S. Frederick, United Nations Environment Programme World Conservation Monitoring Centre and University of California Press, Berkeley, California.
70. Lee Long, W.J., Mellors, J.E. and Coles, R.G. 1993, Seagrasses between Cape York and Hervey Bay, Queensland, Australia, *Australian Journal of Marine and Freshwater Research* 44(1): 19-31.
71. New South Wales National Parks and Wildlife Service 2003, *Guidelines for Aboriginal Heritage Impact Assessment*, NSWNPWS, Sydney.
72. Indigenous Heritage Section of the Australian Heritage Commission 2002, *Ask first: a guide to respecting Indigenous heritage places and values*, Australian Heritage Commission, Canberra.
73. UNESCO 2013b, *Cultural landscape*, UNESCO, Paris, France, <<http://whc.unesco.org/en/culturallandscape/#1>>.
74. Smyth, D. 2001, Management of Sea Country: Indigenous people's use and management of marine environments, in *Working on Country: Contemporary Indigenous Management of Australia's Lands and Coastal Regions*, eds R. Baker, J. Davies and E. Young, Oxford University Press, Melbourne, pp. 60-73.
75. Gray, F. and Zann, L. 1988, Traditional knowledge of the marine environment in northern Australia, in *Workshop Series*, eds Anonymous, Great Barrier Reef Marine Park Authority, Townsville, pp.196.
76. Reef Catchments Mackay Whitsunday 2010, *Working together on country*, Reef Catchments Mackay Whitsunday, Queensland, <<http://reefcatchments.com.au/files/2013/02/WorkingTogetherOnCountry.pdf>>.
77. Wild, R. and McLeod, C. (eds) 2008, *Sacred natural sites: Guidelines for protected area managers, Series No 16*, International Union for the Conservation of Nature, Gland, Switzerland.
78. Aplin, G. 2002, *Heritage: Identification, conservation and management*, Oxford University Press, Melbourne.
79. NQ Dry Tropics 2008, *Healthy Land Yarn: A landholder guide to Aboriginal cultural heritage in the Burdekin Dry Tropics NRM region*, NQ Dry Tropics, Townsville.
80. Pring, A. 2002, *Astronomy and Australian Indigenous Peoples (draft)*, Astronomic Society of South Australia, Adelaide, <<http://www.assa.org.au/media/2912/aaaip.pdf>>.
81. Great Barrier Reef Marine Park Authority 2014, *Traditional Owners of the Great Barrier Reef*, GBRMPA, Townsville, <<http://www.gbrmpa.gov.au/our-partners/traditional-owners/traditional-owners-of-the-great-barrier-reef>>.
82. State of the Environment 2011 Committee Australia *state of the environment 2011: Independent report to the Australian Government Minister for Sustainability, Environment, Water, Population and Communities*, Department of Sustainability, Environment, Water, Population and Communities, Canberra.
83. Millenium Ecosystem Assessment 2005, *Ecosystems and human well-being: Synthesis*, Island Press, Washington DC.
84. Girringun Aboriginal Corporation 2005, *Girringun Region Sea Country statement; TUMRA accreditation manuscript for the Great Barrier Reef Marine Park Authority*, Girringun Aboriginal Corporation, Cardwell.
85. Day, J., Wren, L. and Volhand, J. 2012, Chapter 2: Section1 - Community engagement in safeguarding the world's largest reef: the Great Barrier Reef, Australia, in *World Heritage: Benefits beyond borders*, ed. A. Galla, UNESCO Publishing, Cambridge University Press, pp. 18-29.
86. Marshall, N., Bohensky, E., Goldberg, J., Gooch, M., Lankester, A., Pert, P., Scherl, L. and Tobin, R. 'SELTMP 2011: Social and economic conditions on the Great Barrier Reef', Commonwealth Scientific and Industrial Research Organisation, Townsville, Unpublished report.
87. Dobbs, K., Fernandes, L., Slegers, S., Jago, B., Thompson, L., Hall, J., Day, J., Cameron, D., Tanzer, J., Macdonald, F. and Limpus, C. 2007, Incorporating marine turtle habitats into the marine protected area design for the Great Barrier Reef Marine Park, Queensland, Australia, *Pacific Conservation Biology* 13(3): 293-302.
88. Limpus, C.J., Miller, J.D., Parmenter, C.J. and Limpus, D.J. 2003, The green turtle, *Chelonia mydas*, population of Raine Island and the northern Great Barrier Reef: 1843-2001, *Memoirs of the Queensland Museum* 49(1): 349-440.
89. Chaloupka, M.Y., Bjørndal, K.A., Balazs, G.H., Bolten, A.B., Ehrhart, L.M., Limpus, C.J., Suganuma, H., Tröng, S. and Yamaguchi, M. 2008, Encouraging outlook for recovery of a once severely exploited marine megaherbivore, *Global Ecology and Biogeography* 17(2): 297-304.
90. Limpus, C. and Nicholls, N. 2000, ENSO regulation of Indo-Pacific green turtle populations, in *Applications of Seasonal Climate Forecasting in Agricultural and Natural Ecosystems*, eds G. Hammer, N. Nicholls and C. Mitchell, Kluwer Academic Publishers, Dordrecht, pp. 399-408.
91. Limpus, C.J. and Nicholls, N. 1988, The southern oscillation regulates the annual numbers of green turtles (*Chelonia mydas*) breeding around northern Australia, *Australian Journal of Wildlife Research* 15(2): 157-161.
92. Limpus, C.J. 2008a, *A biological review of Australian marine turtle species, 2 Green turtle, Chelonia mydas (Linnaeus)*, Environmental Protection Agency, Brisbane.
93. Limpus, C.J. 2009a, *A biological review of Australian marine turtle species, 3 Hawksbill turtle, Eretmochelys imbricata*, Environmental Protection Agency, Brisbane.
94. Bell, I.P. 2012, Algivory in hawksbill turtles: *Eretmochelys imbricata* food selection within a foraging area on the Northern Great Barrier Reef. *Marine Ecology* 34(1): 43-55.
95. Limpus, C.J. 1992, The hawksbill turtle (*Eretmochelys imbricata*) in Queensland: population structure within a southern Great Barrier Reef feeding ground, *Wildlife Research* 19: 189-506.
96. Miller, J.D., Limpus, C.J. and Bell, I.P. 2008, Nesting biology of *Eretmochelys imbricata* in the northern Great Barrier Reef, in *Australian hawksbill turtle population dynamics project*, eds C.J. Limpus and J.D. Miller, Environmental Protection Agency, Brisbane, pp. 41-84.
97. Limpus, C.J. 2007, *A biological review of Australian marine turtle species, 5 Flatback turtle, Natator depressus (Garman)*, Environmental Protection Agency, Brisbane.
98. Limpus, C.J., Fleay, A. and Baker, V. 1984, The flatback turtle, *Chelonia depressa*, in Queensland: reproductive periodicity, philopatry and recruitment, *Australian Wildlife Research* 11: 579-587.
99. Limpus, C.J., Parmenter, C.J. and Limpus, D.J. 2002, The status of the flatback turtle, *Natator depressus*, in eastern Australia, in *Proceedings of the twentieth annual symposium on sea turtle biology and conservation*, eds A. Mosier, A. Foley and B. Brost, National Oceanic and Atmospheric Administration Technical Memorandum NMFS-SEFSC 477, Miami, Florida, USA, pp.140-142.
100. Limpus, C.J., Parmenter, C.J., Baker, V. and Fleay, A. 1983, The flatback turtle, *Chelonia depressa*, in Queensland: post-nesting migration and feeding ground distribution, *Australian Wildlife Research* 10: 557-561.
101. Limpus, C.J. and Limpus, D.J. 2003, Loggerhead turtles in the Equatorial and Southern Pacific Ocean: a species in decline, in *Loggerhead sea turtles*, eds A.B. Bolten and B.E. Witherington, Smithsonian Institution, Washington DC, pp. 199-209.
102. Limpus, C.J. 2008b, *A biological review of Australian marine turtle species, 1 Loggerhead turtle, Caretta caretta (Linnaeus)*, Environmental Protection Agency, Brisbane.
103. Limpus, C.J., deVilliers, D.L., deVilliers, M.A., Limpus, D.J. and Read, M.A. 2001, The loggerhead turtle, *Caretta caretta*, in Queensland: observations on feeding ecology in warm



temperate waters, *Memoirs of the Queensland Museum* 46(2): 631-645.

104. Limpus, C.J. 2009b, *A biological review of Australian marine turtle species, 6 Leatherback turtle, Dermochelys coriacea* (Vandelli), Environmental Protection Agency, Brisbane.

105. Spotila, J.R., Reina, R.D., Steyermark, A.C., Plotkin, P.T. and Paladino, F.V. 2000, Pacific leatherback turtles face extinction, *Nature* 405(6786): 529-530.

106. Hamann, M., Limpus, C., Hughes, G., Mortimer, J. and Pilcher, N. 2006, *Assessment of the conservation status of the leatherback turtle in the Indian Ocean and South-East Asia*, IOSEA Marine Turtle MoU Secretariat, Bangkok.

107. Zug, G.R. and Parham, J.F. 1996, Age and growth in leatherback turtles *Dermochelys coriacea* (Testudines: Dermochelyidae): a skeletochronological analysis, *Chelonian Conservation and Biology* 2(2): 244-249.

108. Prince, R.I.T. 2004, Stranding of small juvenile leatherback turtle in Western Australia, *Marine Turtle Newsletter* (104): 3-5.

109. Limpus, C.J. and McLachlan, N.C. 1979, Observations on the leatherback turtle, *Dermochelys coriacea* (L.), in Australia, *Australian Journal of Wildlife Research* 6(1): 105-116.

110. Limpus, C.J. 1984, A benthic feeding record from neritic waters for the leathery turtle (*Dermochelys coriacea*), *Copeia* (2): 552-553.

111. Limpus, C.J. 2008c, *A biological review of Australian marine turtle species, 4 Olive ridley turtle Lepidochelys olivacea* (Eschscholtz), Environmental Protection Agency, Brisbane.

112. Letts, G.A. 1987, The management of crocodilians in Australia, introductory comments, in *Wildlife management: Crocodile and alligators*, eds G.J.W. Webb, S.C. Manolis and P.J. Whitehead, Surrey Beatty and Sons, New South Wales, pp. 103-105.

113. Miller, J. and Bell, I.P. 1997, Crocodiles in the Great Barrier Reef World Heritage Area, in *State of the Great Barrier Reef World Heritage Area workshop, 27-29 November, 1995*, Townsville, eds D. Wachenfeld, J. Oliver and K. Davis, Great Barrier Reef Marine Park Authority, Townsville, pp.248-255.

114. Department of Environment and Heritage 2005, *Blue, fin and sei whale recovery plan: 2005 - 2010*, DEH, Canberra.

115. Department of the Environment and Heritage 2005, *Humpback Whale Recovery Plan 2005 - 2010*, DEH, Canberra.

116. Noad, M.J., Dunlop, R.A., Cato, D. and Paton, D. 2008, *Abundance estimates of the east Australian humpback whale population: Final report for the Australian Department of the Environment, Water, Heritage and the Arts*, University of Queensland, Brisbane.

117. Smith, J.N., Grantham, H.S., Gales, N., Double, M.C., Noad, M.J. and Paton, D. 2012, Identification of humpback whale breeding and calving habitat in the Great Barrier Reef, *Marine Ecology Progress Series* 447: 259-272.

118. Department of Environment and Heritage Protection 2012, *Marine wildlife strandings*, DEHP, Brisbane, <[http://www.ehp.qld.gov.au/wildlife/caring-for-wildlife/marine\\_strandings.html](http://www.ehp.qld.gov.au/wildlife/caring-for-wildlife/marine_strandings.html)>.

119. Parra, G.J., Corkeron, P.J. and Marsh, H. 2006, Population sizes, site fidelity and residence patterns of Australian snubfin and Indo-Pacific humpback dolphins: Implications for conservation, *Biological Conservation* 129(2): 167-180.

120. Cagnazzi, D., Parra, G.J., Westley, S. and Harrison, P.L. 2013, At the heart of the industrial boom: Australian snubfin dolphins in the Capricorn Coast, Queensland, need urgent conservation action, *PLoS ONE* 8(2): e56729.

121. Great Barrier Reef Marine Park Authority 2012c, *A vulnerability assessment for the Great Barrier Reef: Inshore dolphins*, GBRMPA, Townsville.

122. Parra, G.J. 2005, *Behavioural ecology of Irrawaddy, Orcaella brevirostris* (Owen in Gray, 1866), and Indo-Pacific humpback dolphins, *Sousa chinensis* (Osbeck, 1765), in northeast Queensland, Australia: A comparative study, PhD thesis, James Cook University, Cairns.

123. Parra, G.J., Schick, R. and Corkeron, P.J. 2006, Spatial distribution and environmental correlates of Australian snubfin and Indo-Pacific humpback dolphins, *Ecography* 29(3): 396-406.

124. Cagnazzi, D. 2010, *Conservation status of Australian snubfin dolphin, Orcaella heinsohni, and Indo-Pacific humpback dolphin, Sousa chinensis, in the Capricorn Coast, Central Queensland, Australia*, PhD, Southern Cross University, NSW.

125. Marsh, H., Hodgson, A., Lawler, I.R., Grech, A. and Delean, S. 2007, *Condition, status and trends and projected futures of the dugong in the northern Great Barrier Reef and Torres Strait: Including identification and evaluation of the key threats and evaluation of available management options to improve its status*, Report to the Marine and Tropical Sciences Research Facility, Reef and Rainforest Research Centre, Cairns.

126. Sobtzyk, S., Hagihara, R., Grech, A. and Marsh, H. 2012, *Aerial survey of the urban coast of Queensland to evaluate the response of the dugong population to the widespread effects of the extreme weather events of the summer of 2010-11, Final report to the Australian Marine Mammal Centre and the National Environmental Research Program*, James Cook University, Townsville.

127. Marsh, H., De'ath, G., Gribble, N.A. and Lane, B. 2005, Historical marine population estimates: triggers or targets for conservation? The dugong case study, *Ecological Applications* 15(2): 481-492.

128. Karbhari, J.P. and Josekutty, C.J. 1986, On the largest whale shark *Rhincodon typus* Smith landed alive at Cuffe Parade, Bombay, *Marine Fisheries Information Service, Technical and Extension Series* 66: 31-35.

129. Last, P.R. and Stevens, J.D. 2009, *Sharks and rays of Australia*, 2nd edn, CSIRO, Collingwood.

130. Bruce, B.D. 2008, The biology and ecology of the white shark (*Carcharodon carcharias*), in *Sharks of the Open Ocean: Biology, fisheries and conservation*, eds M.D. Camhi, E. Pikitch and E.A. Babcock, 13th edn, Blackwell Publishing, New Jersey, USA, pp. 69.

131. CSIRO Marine and Atmospheric Research 2008, *Summary of CMAR white shark tagging in Australian waters*, CMAR, Aspendale, <<http://www.cmar.csiro.au/tagging/whitesharks/taggedsharks.html>>.

132. Congdon, B.C. 2008, Seabirds, in *The Great Barrier Reef: biology, environment and management*, eds P.A. Hutchings, M.J. Kingsford and O. Hoegh-Guldberg, CSIRO, Collingwood, pp. 359-368.

133. Hulsman, K. 1988, The structure of seabird communities: an example from Australian waters, in *Seabirds and other marine vertebrates, competition, predation and other interactions*, ed. J. Burger, Columbia University Press, New York, USA, pp. 59-91.

134. Turner, M., Green, R. and Chin, A. 2006, Birds, in *The state of the Great Barrier Reef*, ed. A. Chin, Great Barrier Reef Marine Park Authority, Townsville.

135. Hulsman, K., O'Neill, P. and Stokes, T. 1997, Current status and trends of seabirds on the Great Barrier Reef, in *Proceedings of the State of the Great Barrier Reef World Heritage Area Workshop, 27-29 November, 1995*, Townsville, Queensland, eds D. Wachenfeld, J. Oliver and K. Davis, Great Barrier Reef Marine Park Authority, Townsville, pp. 259-282.

136. O'Neill, P., Heatwole, H., Preker, M. and Jones, M. 1996, *Populations, movements and site fidelity of brown and masked boobies on the Swain Reefs, Great Barrier Reef, as shown by banding recoveries*, CRC Reef Research Centre, Townsville.

137. Surman, C.A. and Wooller, R.D. 2003, Comparative foraging ecology of five sympatric terns at a sub-tropical island in the eastern Indian Ocean, *Journal of Zoology* 259: 219-230.

138. Hulsman, K. 1983, *Survey of the seabird colony of the Capricornia section of the Great Barrier Reef Marine Park. II Population parameters and some management options. Research report for the Great Barrier Reef Marine Park Authority*, Griffith University School of Australian Environmental Studies, Nathan.

139. Hulsman, K. 1984, *Survey of the seabird colony of the Capricornia section of the Great Barrier Reef Marine Park. III*



Population parameters and management strategies. Research report for the Great Barrier Reef Marine Park Authority, Griffith University School of Australian Environmental Studies, Nathan.

140. Higgins, P.J. and Davies, S.J.J.F. (eds) 1996, *Handbook of Australian, New Zealand and Antarctic birds. Volume 3: Snipe to Pigeons*, Oxford University Press, Melbourne.
141. Dyer, P.K., O'Neill, P. and Hulsman, K. 2005, Breeding numbers and population trends of wedge-tailed shearwater (*Puffinus pacificus*) and black noddy (*Anous minutus*) in the Capricorn Cays, southern Great Barrier Reef, *Emu* 105: 249-257.
142. King, B.R. 1993, The status of Queensland seabirds, *Corella* 17(3): 65-92.
143. Geering, A., Agnew, L. and Harding, S. 2007, *Shorebirds of Australia*, CSIRO Publishing, Collingwood.
144. Great Barrier Reef Marine Park Authority 2012d, *A vulnerability assessment for the Great Barrier Reef: Shorebirds*, GBRMPA, Townsville.
145. Wilson, H.B., Kendall, B.E., Fuller, R.A., Milton, D.A. and Possingham, H.P. 2011, Analyzing variability and the rate of decline of migratory shorebirds in Moreton Bay, Australia, *Conservation Biology* 25(4): 758-766.
146. Tobin, A.J., Simpfendorfer, C.A., Mapleston, A., Currey, L., Harry, A.V., Welch, D.J., Ballagh, A.C., Chin, A., Szczecinski, N., Schlaff, A., White, J. and Moore, B. 2010, *A quantitative ecological risk assessment of sharks and finfish of the Great Barrier Reef World Heritage Area inshore waters: a tool for fisheries and marine park managers. Identifying species at risk and potential mitigation strategies*, Marine and Tropical Sciences Research Facility, Cairns.
147. Stevens, J.D., Pillans, R.D. and Salini, J.P. 2005, *Conservation assessment of Glyptis sp A (speartooth shark), Glyptis sp. C (northern river shark), Pristis microdon (freshwater sawfish) and Pristis zijsron (green sawfish): final report to the Department of the Environment and Heritage*, CSIRO Marine Research, Hobart.
148. Department of Sustainability, Environment, Water, Population and Communities 2011, *National recovery plan for threatened albatrosses and giant petrels 2011-2016*, DSEWPC, Hobart.
149. Dobbs, K.A. 2005, Recoveries of seabirds banded between 1978 and 1987 at Raine Island, MacLennan and Moulter Cays and Sandbanks No. 7 and 8, northern Great Barrier Reef, Australia, *Corella* 29: 65-72.
150. Batianoff, G.N. and Cornelius, N.J. 2005, Birds of Raine Island: population trends, breeding behaviour and nesting habitats, *Proceedings of the Royal Society of Queensland* 112: 1-29.
151. Department of Defence, Environmental Protection Agency, Great Barrier Reef Marine Park Authority and Environment Australia 1999, *Information sheet on Ramsar Wetlands (RIS)*, Environment and Heritage Section, Department of Defence, Canberra.
152. O'Neill, P. 2009, *State of the Environment Report for Shoalwater Bay Training Area 2008*, Department of Defence, Canberra.
153. Limpus, C.J., Limpus, D.J., Arthur, K.E. and Parmenter, C.J. 2005, *Monitoring green turtle population dynamics in Shoalwater Bay, 2000-2004*, Great Barrier Reef Marine Park Authority, Townsville.
154. Hopley, D. 1970, *The geomorphology of the Burdekin Delta, North Queensland*, James Cook University of North Queensland, Department of Geography Monograph Series No:1, Townsville.
155. Kingsford, M.J. and Wolanski, E. 2008, Oceanography, in *The Great Barrier Reef: Biology, environment and management*, eds P. Hutchings, M. Kingsford and O. Hoegh-Guldberg, CSIRO Publishing, Collingwood, pp. 28-39.
156. Berkelmans, R., Weeks, S.J. and Steinberg, C.R. 2010, Upwelling linked to warm summers and bleaching on the Great Barrier Reef, *Limnology and Oceanography* 55(6): 2634-2644.
157. Hobday, A.J. and Lough, J.M. 2011, Projected climate change in Australian marine and freshwater environments, *Marine and Freshwater Research* 62(9): 1000-1014.
158. Church, J.A., White, N.J., Hunter, J.R. and McInnes, K.L. 2012, Sea level, in *A marine climate change impacts and adaptation report card for Australia 2012*, eds E.S. Poloczanska, A.J. Hobday and A.J. Richardson, CSIRO, Canberra, pp. 27-46.
159. Bellwood, D.R., Hughes, T.P., Connolly, S.R. and Tanner, J. 2005, Environmental and geometric constraints on Indo-Pacific coral reef biodiversity, *Ecology Letters* 8(6): 643-651.
160. van Oppen, M.J.H. and Lough, J.M. (eds) 2009, *Coral bleaching: patterns, processes, causes and consequences*, Springer, Berlin.
161. Robins, J.B., Halliday, I., Staunton-Smith, J., Mayer, D.G. and Sellin, M.J. 2005, Freshwater-flow requirements of estuarine fisheries in tropical Australia: a review of the state of knowledge and application of a suggested approach, *Marine and Freshwater Research* 56(3): 343-360.
162. Lillywhite, H.B., Babonis, L.S., Sheehy III, C.M. and Tu, M. 2008, Sea snakes (*Laticauda* spp.) require fresh drinking water: implication for the distribution and persistence of populations, *Physiological and Biochemical Zoology* 81(6): 785-796.
163. Stieglitz, T. 2005, Submarine groundwater discharge into the near-shore zone of the Great Barrier Reef, Australia, *Marine Pollution Bulletin* 51(1-4): 51-59.
164. Maughan, M., Brodie, J. and Waterhouse, J. 2008, *Reef exposure model for the Great Barrier Reef lagoon*, Australian Centre for Tropical Freshwater Research, James Cook University, Townsville.
165. Kroon, F.J., Kuhnert, P.M., Henderson, B.L., Wilkinson, S.N., Kinsey-Henderson, A., Abbott, B., Brodie, J.E. and Turner, R.D.R. 2012, River loads of suspended solids, nitrogen, phosphorus and herbicides delivered to the Great Barrier Reef lagoon, *Marine Pollution Bulletin* 65(4-9): 167-181.
166. Fabricius, K.E. 2005, Effects of terrestrial runoff on the ecology of corals and coral reefs: Review and synthesis, *Marine Pollution Bulletin* 50(2): 125-146.
167. Hoegh-Gulberg, O. and Dove, S. 2008, Primary production, nutrient recycling and energy flow through coral reef ecosystems, in *The Great Barrier Reef: Biology, environment and management*, eds P. Hutchings, M. Kingsford and O. Hoegh-Guldberg, CSIRO Publishing, Collingwood, pp. 59-73.
168. Brodie, J.E., Devlin, M., Haynes, D. and Waterhouse, J. 2011, Assessment of the eutrophication status of the Great Barrier Reef lagoon (Australia), *Biogeochemistry* 106(2): 281-302.
169. National Aeronautics and Space Administration 2013, *Carbon Cycle*, NASA, <<http://science.nasa.gov/earth-science/oceanography/ocean-earth-system/ocean-carbon-cycle/>>.
170. Sarmiento, J.L., Hughes, T.M.C., Stouffer, R.J. and Manabe, S. 1998, Simulated response of the ocean carbon cycle to anthropogenic climate warming, *Nature* 393(6682): 245-249.
171. De'ath, G., Lough, J.M. and Fabricius, K.E. 2009, Declining coral calcification on the Great Barrier Reef, *Science* 323(5910): 116-119.
172. Orr, J.C., Fabry, V.J., Aumont, O., Bopp, L., Doney, S.C., Feely, R.A., Gnanadesikan, A., Gruber, N., Ishida, A., Joos, F., Key, R.M., Lindsay, K., Maier-reimer, E., Matear, R., Monfray, P., Mouchet, A., Najjar, R.G., Plattner, G.K., Rodgers, K.B., Sabine, C.L., Sarmiento, J.L., Schlitzer, R., Slater, R.D., Totterdell, I.J., Weirig, M.F., Yamanaka, Y. and Yool, A. 2005, Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms, *Nature* 437(7059): 681-686.
173. Boyer, R. *Interactive concepts in biochemistry*, John Wiley and Sons, <<http://www.wiley.com/college/boyer/0470003790/index.htm>>.
174. Brodie, J., De'ath, G., Devlin, M., Furnas, M. and Wright, M. 2007, Spatial and temporal patterns of near-surface chlorophyllain the Great Barrier Reef lagoon, *Marine and Freshwater Research* 58(4): 342-353.
175. Webster, N.S. and Bourne, D.G. 2012, Microbes, in *A marine climate change impacts and adaptation report card for Australia 2012*, eds E.S. Poloczanska, A.J. Hobday and A.J. Richardson, CSIRO, Canberra.


176. Fabricius, K.E. and Wolanski, E. 2000, Rapid smothering of coral reef organisms by muddy marine snow, *Estuarine, Coastal and Shelf Science* 50(1): 115-120.
177. Hughes, T.P., Rodrigues, M.J., Bellwood, D.R., Ceccarelli, D., Hoegh-Guldberg, O., McCook, L.J., Molschanivskyj, N.A., Pratchett, M.S., Steneck, R.S. and Willis, B. 2007, Phase shifts, herbivory, and the resilience of coral reefs to climate change, *Current Biology* 17(4): 360-365.
178. McCook, L.J., Folke, C., Hughes, T., Nyström, M., Obura, D. and Salm, R. 2007, Ecological resilience, climate change and the Great Barrier Reef, in *Climate Change and the Great Barrier Reef: A vulnerability assessment*, eds J.E. Johnson and P.A. Marshall, Great Barrier Reef Marine Park Authority and Australian Greenhouse Office, Townsville, pp. 75-96.
179. McCook, L.J. 1999, Macroalgae, nutrients and phase shifts on coral reefs: scientific issues and management consequences for the Great Barrier Reef, *Coral Reefs* 18: 357-367.
180. Epibiont Research Cooperative 2007, *A synopsis of the literature on the turtle barnacles (Cirripedia: Balanomorpha: Coronuloidea) 1758-2007*, ERC Special Publication Number 1.
181. Stanley, G.D. 2006, Photosymbiosis and the evolution of modern coral reefs, *Science* 312: 857-858.
182. Muscatine, L. 1973, Nutrition of corals, in *Biology and geology of coral reefs*, eds O.A. Jones and R. Endean, Academic Press, New York, USA, pp. 77-115.
183. LaJeunesse, T.C. 2002, Diversity and community structure of symbiotic dinoflagellates from Caribbean coral reefs, *Marine Biology* 141: 387-400.
184. Lewis, J., Hoole, D. and Chappell, L.H. 2003, Parasitism and environmental pollution: parasites and hosts as indicators of water quality, *Parasitology* 126(7): S1-S3.
185. Hughes, T.P., Baird, A.H., Dinsdale, E.A., Molschanivskyj, N.A., Pratchett, M.S., Tanner, J.E. and Willis, B.L. 2000, Supply-side ecology works both ways: The link between benthic adults, fecundity, and larval recruits, *Ecology* 81(8): 2241-2249.
186. Beverton, R.J.H. 1991, Fish resources: threats and protection, *Netherlands Journal of Zoology* 42(2-3): 139-175.
187. McCook, L.J. 2001, Competition between corals and algal turfs along a gradient of terrestrial influence in the nearshore central Great Barrier Reef, *Coral Reefs* 19: 419-425.
188. Birrell, C.L., McCook, L.J., Willis, B.L. and Harrington, L. 2008, Chemical effects of macroalgae in larval settlement of the broadcast spawning coral *Acropora millepora*, *Marine Ecology Progress Series* 362: 129-137.
189. Veron, J.E.N. 2008, *A reef in time: the Great Barrier Reef from beginning to end*, Harvard University Press, Cambridge, USA.
190. Frankham, R. and Ralls, K. 1998, Conservation biology - Inbreeding leads to extinction, *Nature* 392(6675): 441-442.
191. Birkeland, C.E. (ed) 1997, *Life and death of coral reefs*, Chapman and Hall, New York.
192. Weeks, S.J., Bakun, A., Steinberg, C.R., Brinkman, R. and Hoegh-Guldberg, O. 2010, The Capricorn Eddy: a prominent driver of the ecology and future of the southern Great Barrier Reef, *Coral Reefs* 29: 975-985.
193. Bellwood, D.R., Hoey, A.S. and Choat, J.H. 2003, Limited functional redundancy in high diversity systems: resilience and ecosystem function on coral reefs, *Ecology Letters* 6(4): 281-285.
194. Edmonds, T.H. 1990, Comment: The Queensland rainforest and wetlands conflict: Australia's external affairs power: domestic control and international conservation, *Environmental Law* 20(22840): 387-413.
195. Ponder, W., Hutchings, P. and Chapman, R. 2002, *Overview of the conservation of Australian marine invertebrates: a report for Environment Australia*, Australian Museum, Sydney.
196. Domeier, M.L. and Speare, P. 2012, Dispersal of adult black marlin (*Istiompax indica*) from a Great Barrier Reef spawning aggregation, *PLoS ONE* 7(2): e31629.
197. Meager, J.J., Limpus, C.J. and Sumpton, W.D. 2013, *A review of the population dynamics of dugongs in southern Queensland*, Department of Environment and Heritage Protection, Brisbane.
198. Marsh, H. 1995, The life history, pattern of breeding and population dynamics of the dugong, in *Population and Biology of the Florida Manatee*, eds T. O'Shea, B.B. Ackerman and H. Percival, US Fish and Wildlife Service, Washington, DC, pp. 75-83.
199. Preen, A.R. and Marsh, H. 1995, Response of dugongs to large-scale loss of seagrass from Hervey Bay, Queensland, Australia, *Wildlife Research* 22: 507-519.
200. Howells, E., van Oppen, M.J.H. and Willis, B.L. 2009, High genetic differentiation and cross-shelf patterns of genetic diversity among Great Barrier Reef populations of *Symbiodinium*, *Coral Reefs* 28(1): 215-225.
201. Harrison, H.B., Williamson, D.H., Evans, R.D., Almany, G.R., Thorrold, S.R., Russ, G.R., Feldheim, K.A., van Herwerden, L., Planes, S., Srinivasan, M., Berumen, M.L. and Jones, G.P. 2012, Larval export from marine reserves and the recruitment benefit for fish and fisheries, *Current Biology* 22(11): 1023-1028.
202. Olds, A.D., Connolly, R.M., Pitt, K.A. and Maxwell, P.S. 2012a, Primacy of seascape connectivity effects in structuring coral reef fish assemblages, *Marine Ecology Progress Series* 462: 191-203.
203. Olds, A.D., Pitt, K.A., Maxwell, P.S. and Connolly, R.M. 2012b, Synergistic effects of reserves and connectivity on ecological resilience, *Journal of Applied Ecology* 49: 1195-1203.
204. Card, M.A. 2001, *Vegetation dynamics of Milman Island, northern Great Barrier Reef, Australia*, School of Tropical Biology, James Cook University, Townsville.
205. Barry, G. 2012, *Transcript of an interview with a Traditional Owner from Mamu Country*.
206. Great Barrier Reef Marine Park Authority 2012e, *Great Barrier Reef Coastal Ecosystems Assessment Framework*, GBRMPA, Townsville.
207. Great Barrier Reef Marine Park Authority 2013a, *Scoping a methodology to identify priority areas that support the Great Barrier Reef World Heritage Area*, GBRMPA, Townsville.
208. Great Barrier Reef Marine Park Authority 2013b, *Mulgrave-Russell Basin Assessment. Wet Tropics NRM region: Assessment of ecosystem services within the Mulgrave-Russell basin focusing on understanding and improving the health and resilience of the Great Barrier Reef*, GBRMPA, Townsville.
209. Great Barrier Reef Marine Park Authority 2013c, *Coastal Ecosystem Management. Mount Peter, Cairns: review of coastal ecosystem management to improve the health and resilience of the Great Barrier Reef World Heritage Area*, GBRMPA, Townsville.
210. Great Barrier Reef Marine Park Authority 2013d, *Haughton Basin Assessment: Burdekin Dry Tropics Natural Resource Management Region*, GBRMPA, Townsville.
211. Great Barrier Reef Marine Park Authority 2013e, *Coastal ecosystems management. Lower Burdekin Floodplain: Review of coastal ecosystem management to improve the health and resilience of the Great Barrier Reef World Heritage Area*, GBRMPA, Townsville.
212. Great Barrier Reef Marine Park Authority 2013f, *Fitzroy Basin Assessment, Fitzroy Basin Association Natural Resource Management region: Assessment of ecosystem services within the Fitzroy Basin focusing on understanding and improving the health and resilience of the Great Barrier Reef*, GBRMPA, Townsville.
213. Great Barrier Reef Marine Park Authority 2013g, *Fish habitat connectivity case study: Lower Fitzroy basin*, GBRMPA, Townsville.
214. Great Barrier Reef Marine Park Authority 2013h, *Baffle Basin Assessment. Burnett-Mary Regional Management Group NRM region: Assessment of ecosystem services within the Baffle Basin focusing on understanding and improving the health and resilience of the Great Barrier Reef*, GBRMPA, Townsville.
215. Great Barrier Reef Marine Park Authority 2013i, *Development case study: Baffle Basin*, GBRMPA, Townsville.



## Chapter 5

---

### Drivers and activities





## Table of contents

<b>5.1</b>	<b>Background .....</b>	<b>5–3</b>
<b>5.2</b>	<b>Drivers of change .....</b>	<b>5–3</b>
5.2.1	Drivers relevant to the Region .....	5–3
5.2.2	Climate change .....	5–4
5.2.3	Economic growth .....	5–8
5.2.4	Population growth .....	5–11
5.2.5	Technological development .....	5–13
5.2.6	Societal attitudes .....	5–15
<b>5.3</b>	<b>Activities adjacent to the Region .....</b>	<b>5–16</b>
5.3.1	Agriculture .....	5–16
5.3.2	Aquaculture .....	5–19
5.3.3	Urban development .....	5–20
5.3.4	Industrial development and resource extraction .....	5–21
5.3.5	Port activities .....	5–22
<b>5.4</b>	<b>Activities within the Region .....</b>	<b>5–25</b>
5.4.1	Traditional use of marine resources .....	5–25
5.4.2	Tourism .....	5–26
5.4.3	Fishing — commercial .....	5–30
5.4.4	Fishing — recreational .....	5–35
5.4.5	Recreation .....	5–37
5.4.6	Shipping .....	5–41
5.4.7	Defence activities .....	5–46
5.4.8	Research activities .....	5–49
<b>5.5</b>	<b>Summary of conclusions .....</b>	<b>5–49</b>
	<b>References .....</b>	<b>5–51</b>

Cover page image: North Reef lighthouse on North Reef Island



**Extract from Great Barrier Reef Region Strategic Assessment terms of reference**

**3. Assessment of impacts on matters of national environmental significance**


**3.1 Actual and potential impacts**

- a) describe the environmental, social, cultural and economic drivers affecting the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area.
- b) describe and analyse the actual and potential impacts on the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area, in the strategic assessment area, including:
  - i. impacts from past, present and future activities
  - .....

## 5.1 Background

As outlined in Chapter 4, the Great Barrier Reef Region (the Region) comprises a rich mosaic of biodiversity, geomorphology and heritage values. The condition of those values determines the quality of the cultural, social and economic benefits the community derives from the environment (such as aesthetics, income, appreciation and enjoyment). As a result of external drivers and activities both within the Region and beyond its boundaries, a number of impacts are diminishing the condition of the Region's values and attributes and the quality of the benefits they provide. This chain of cause and effect is illustrated in Figure 5.1.

Understanding these causal relationships helps forecast the future condition of the Region's values and attributes, including likely changes in the community benefits derived from the Region.



**Figure 5.1 The influence of drivers and activities**

## 5.2 Drivers of change

Drivers are overarching causes of change in the environment. They can affect the environment indirectly by changing the way people undertake activities that affect the environment (indirect drivers), or by directly changing conditions in the environment itself (direct drivers). They can also work in concert with one another and operate across a range of scales. An understanding of their positive and negative effects is fundamental to understanding the past, present and future condition of the values and attributes underpinning matters of national environmental significance.

### 5.2.1 Drivers relevant to the Region

The drivers most relevant to the Great Barrier Reef Region were identified by examining the *Australia State of the Environment 2011*<sup>1</sup> and relevant peer-reviewed literature, and through input from key Great Barrier Reef natural and social scientists. The five drivers analysed are:

- climate change
- economic growth
- population growth
- technological developments
- societal attitudes.

Climate change is a 'direct driver' that has direct and ongoing effects on the environment as well as indirect effects mediated by other processes or activities. The other four drivers are 'indirect drivers', influencing people's activities that, in turn, affect the environment. There are links between all these drivers, with each one influencing the others. For example, technological developments can play a role in climate change, economic growth,

population growth and societal attitudes. Similarly, population growth affects climate change, economic growth and societal attitudes.

This description of drivers relevant to the Region has been informed by the CSIRO's recent analysis of the environmental,<sup>2</sup> social and economic conditions that are predicted to substantially change the way people live (megatrends).<sup>2</sup>

### 5.2.2 Climate change

The Earth's climate has always been changing. Ice ages ending, glaciers melting and sea levels changing are natural phenomena. However, changes in the Earth's climate have typically occurred over millennia. In fact, the stable climate over the past 1000 years has, in part, enabled the rapid expansion of human populations and development.

It is now almost certain that the climate is changing at a rate unprecedented in the geological record.<sup>3</sup> A rapidly changing climate poses substantial risks and challenges for individuals, societies and nations and the ecosystems that support them.

Increased concentrations of greenhouse gases (particularly carbon dioxide) in the atmosphere result in more heat being trapped, increasing the Earth's temperature. Atmospheric carbon dioxide is at a concentration unprecedented within the past 650,000 years.<sup>4,5</sup> The rapid increase in emissions of carbon dioxide and other greenhouse gases, since the Industrial Revolution, has amplified their natural role in retaining heat within the Earth's atmosphere.<sup>3</sup>

#### Trends

Increases in the concentration of greenhouse gases in the Earth's atmosphere are driving changes in a number of variables that can affect the Region's values. Variables of particular significance are:

- sea temperature
- ocean acidity
- sea level
- ocean currents
- tropical storm intensity
- weather variability — rainfall, wind, droughts and flooding.

The indicative trends in these variables and the uncertainties around these trends are summarised in Figure 5.2 and described in detail in the relevant parts of Chapter 6.

The climate change variables listed do not work in isolation from one another. For example, as more carbon dioxide is released into the atmosphere, air and sea temperatures rise. This warming causes ice to melt and thermal expansion in the oceans which, in turn, causes the sea level to rise.

#### Implications for the Region's values

Climate change is a direct and indirect driver for coral reef ecosystems such as the Great Barrier Reef, and there have already been some serious effects on the Region's biodiversity values for example, coral bleaching in 1998 and 2002.<sup>6</sup> The future implications for biodiversity values depend on the rate and extent of increases in greenhouse gas concentrations, because this is the factor driving the change. Although change in ecosystem state is likely to be a gradual process, there is evidence to suggest that the reduced resilience of ecosystems increases their vulnerability to catastrophic events. For example, a severe cyclone may push a coral reef towards a tipping point, beyond which it is unable to rebound.

Potential climate change effects for species groups and habitats have been considered in *Climate change and the Great Barrier Reef: a vulnerability assessment*<sup>7</sup> and in many recent scientific studies.<sup>8,9,10</sup> The effects, both individually and combined, are likely to have far-reaching consequences for the Region's environment. The 2013 Water Quality Scientific Consensus Statement concluded that 'key Great Barrier Reef ecosystems are showing declining trends in condition due to continuing poor water quality, cumulative impacts of climate change and increasing intensity of extreme events'.<sup>11</sup>

The projected vulnerability of a number of the Region's habitats and species presented in Figure 5.2 shows not all components of the Great Barrier Reef are affected equally. Corals and seabirds are considered to be some of the most vulnerable species to the predicted changes. Many other species, however, will also be negatively affected. Molluscs, for example, will have a reduced capacity to develop hard shells due to ocean acidification.

Reef-building corals are highly vulnerable to several environmental factors driven by increasing greenhouse gas concentrations — increasing temperature, ocean acidification, and increased frequency of severe storms. As the climate changes, the capacity of hard corals to grow and reproduce will be increasingly compromised, with flow-on effects on other species dependent on coral reefs.

The frequency and severity of coral bleaching is predicted to increase under climate change, with potentially serious consequences for the Reef.<sup>12,13,14,15</sup> Coral bleaching can affect large areas (known as mass bleaching events). Bleaching is not always fatal for corals, but has been one of the main causes of coral death around the world in the past 20 years.<sup>8,14,16</sup>

Recent assessments predict that reefs could experience temperature-induced bleaching events twice per decade by about 2020, and annual bleaching events could occur by mid-century.<sup>17</sup> Patterns of bleaching will vary by location. In favourable conditions, recovery of corals following bleaching is likely, and reef habitats may recover if the frequency remains at fewer than about two events per decade. However, severe degradation of Great Barrier Reef corals and coral reef habitats is likely to occur before, rather than after, the time when bleaching events occur annually, since there are a large number of additional pressures and impacts on coral reefs.<sup>8,17,18</sup>

Implications for the Region's values at different concentrations of atmospheric carbon dioxide can be summarised as:

- **350 parts per million:** Optimum limits for coral reef ecosystems are at or below this concentration. This would require a lowering of global carbon dioxide concentrations.
- **400 parts per million** (close to the current concentration): The frequency of severe bleaching is likely to increase, with rising summer temperatures leading to the dominance of thermally tolerant species. While coral reef ecosystems are likely to be affected by a number of impacts related to climate change, they are expected to remain coral dominated in many areas. This concentration of atmospheric carbon dioxide is only slightly above the average level reached globally in March 2013,<sup>19</sup> and there is already evidence of effects on the Reef, such as declining calcification rates, that are suggested to be caused by temperature stress and ocean acidification.<sup>20</sup>
- **450 parts per million:** It is predicted that the diversity of corals on reefs will decline under the combined effects of elevated temperatures and ocean acidity.<sup>1</sup> Ocean acidification is likely to further affect the growth of most calcifying organisms.<sup>1,21</sup> This level of atmospheric carbon dioxide poses an extreme risk for coral reef ecosystems and tropical coastal habitats.

The major impacts of climate change are described in further detail in Chapter 6.

#### Implications for activities and industries

Reef-dependent activities, including tourism, fishing, recreation and traditional use, are vulnerable to the negative effects that ocean acidification, sea level rise, more frequent extreme weather and warming sea temperatures may have on Reef condition.<sup>7,22</sup>

The Reef-based tourism industry is very concerned about the impacts of climate change on its businesses and livelihoods, including degradation of reef sites, poor recovery of bleached sites as a result of other stresses, and a loss of marketing appeal as a high-quality reef destination.<sup>23</sup>

It is likely fishing activities will be highly sensitive to climate change, including projected changes in fish abundance, survivorship<sup>24,25,26</sup>, size and distribution, disruptions to shallow water nurseries and loss of coral reef habitats, as well as changes in cyclone and storm activity<sup>7,27,28</sup>.


Traditional Owners are concerned about rising temperatures altering the seasonality and availability of marine resources, as well as the potential loss of totemic species and the possible displacement of their coastal communities due to rising sea levels.<sup>27</sup>

Climate change science is a rapidly expanding field, and there is improved understanding of the implications for the Great Barrier Reef and Reef-dependent industries.<sup>29</sup> Tools are being developed and applied to help communities and industries recognise their vulnerabilities and adaptation needs. For example, a climate change vulnerability assessment of the East Coast Otter Trawl Fishery<sup>30</sup> has been published, and key stakeholders have been engaged in a climate change adaptation planning process.<sup>31</sup>

An examination of the extreme weather events in the summer of 2010–2011 highlights the potential impacts on communities and industries from the predicted increased frequency of these events. As a result of cyclone Yasi and the central Queensland floods, the summer's extreme weather damaged 27 per cent of Queensland's road network and 4750 kilometres of the state's railway lines.<sup>32</sup> It was also responsible for 11 port closures.<sup>33</sup> Many resorts, jetties and marinas were damaged, along with more than 4000 houses.<sup>33</sup>

Two Reef-dependent industries were particularly affected by that season's cyclones and floods.<sup>34</sup> A survey of 145 commercial fishers and 62 marine tourism operators showed the most significant impacts across each industry were the three to four months of lost operations after the extreme weather. Many fishers were unable to fish because of the large amounts of debris and sediment remaining in the water, and the reduced catchability<sup>28</sup> of some Reef-associated target species. Tourism operators were far less able to attract tourists to their destination due to the perception that tourism experiences had been affected Reef-wide.<sup>34</sup>

Flooding rains, intense cyclones and rising sea levels may have serious impacts on regional industries regardless of whether they are directly dependent on the Reef. However, the people who depend on the Reef for their activities may face additional vulnerabilities linked to deteriorating Reef condition. Patterns of use may also change in the wake of extreme weather and climate change, with the potential to create new 'hotspots' of pressure on the Great Barrier Reef ecosystem.<sup>35</sup>



**Figure 5.2 Projected trends in climate change variables and the Region's biodiversity values**

The average monthly global carbon dioxide concentration reached 396 parts per million (ppm) in March 2013.<sup>36</sup> A concentration of 550 ppm carbon dioxide is predicted to be reached by about 2100.<sup>3,19</sup> Concentrations could potentially exceed this value, resulting in even more serious effects on the Region's ecosystem. In certain scenarios, tipping points for ecosystems can occur suddenly in response to a severe disturbance, hence causing a rapid change.



## The science of carbon dioxide concentrations and coral reefs


The pre-industrial concentration of atmospheric carbon dioxide was approximately 280 parts per million.<sup>37</sup> Since then, the concentration of atmospheric carbon dioxide has been increasing, and reached 396 parts per million in March 2013.<sup>19</sup>

Figure 5.3 shows atmospheric concentrations since 1958, and the increasing annual mean growth rate in carbon dioxide from one of the global observatories. In addition to changing the climate, an increasing concentration of carbon dioxide in the atmosphere leads to ocean acidification.

Coral reefs are sensitive to climate change and ocean acidification. The consequences of rising concentrations of atmospheric carbon dioxide put reefs at increasing risk of serious decline. When the global atmospheric carbon dioxide concentration exceeded approximately 320 parts per million, widespread temperature-induced mass coral bleaching (leading to mortality) began to be observed by the scientific community.<sup>38</sup> Bleaching began on most reefs worldwide when carbon dioxide concentrations reached approximately 340 parts per million.<sup>38</sup>

If carbon dioxide concentrations reach 450 parts per million, scientists predict reefs will be in rapid and terminal decline worldwide as the result of multiple synergies arising from mass bleaching, ocean acidification and other environmental impacts.<sup>4</sup> It is predicted that damage to shallow reef communities will become extensive, with consequent reductions of biodiversity followed by extinctions.<sup>1</sup> Reefs will cease to be large-scale nursery grounds for fish, threatening food security for millions of people around the world, and no longer provide the same community benefits. There will be knock-on effects to ecosystems associated with reefs, to other pelagic and benthic ecosystems, to coastal protection and reef-dependent industries and communities.<sup>4,38</sup>

Based on a current predicted trajectory, a concentration of 550 parts per million of carbon dioxide could be reached by about 2100.<sup>3</sup> Emissions are continuing to grow and, on the current trend, 450 parts per million carbon dioxide could be reached by about 2040.



**Figure 5.3 Mean atmospheric concentrations and the annual mean increase of carbon dioxide, 1958–2013**

The data was collected from Mauna Loa, Hawaii. This is the longest record of direct measurements of carbon dioxide concentrations in the atmosphere. In the second graph, mean annual increase is the difference between the start of January and the end of December of that year. Decadal averages of growth in carbon dioxide concentrations are represented by the horizontal black bars. (Source: NOAA and Scripps Institution of Oceanography<sup>39</sup>)

### Tourism industry addressing climate change

Recognising the threat that climate change poses to the health of the Great Barrier Reef, industries that depend on it are proactively taking action to tackle climate change. Innovators within the marine tourism industry, tourism industry associations and key tourism agencies worked with the Authority to form the Great Barrier Reef Tourism Climate Change Action Group in 2006, to encourage industry action on climate change. Implementation of the Group's *Great Barrier Reef Tourism Climate Change Action Strategy 2009–2012*<sup>40</sup> has delivered a range of products specific to the Great Barrier Reef and its tourism industry, including a carbon emissions calculator, climate change case studies, and climate change operator workshops.

The emissions calculator assists Great Barrier Reef tourism operators to reduce their emissions and adapt to climate change. The online calculator allows operators to calculate their carbon footprint, access information on how to reduce their emissions, and track the results of any changes they make.

Lady Elliot Island Eco Resort, on the Region's southernmost island, is showcased in the climate change case study on sustainable island resorts. The resort's operators were inspired to make operational changes in response to climate change. After undertaking an energy audit, they replaced diesel generators with a large hybrid solar power system in 2008. A follow-up audit in 2009 showed they had reduced the resort's non-renewable energy use by 32 per cent. By mid-2012, they had reduced the resort's diesel consumption by almost 70 per cent.

In 2007, the Authority's Reef HQ Aquarium set out to achieve a 50 per cent reduction in energy usage, motivated by concerns about climate change. By raising the air-conditioning temperature, undertaking minor building works and improving lighting arrangements, the aquarium has substantially reduced energy consumption. By June 2012, power usage had been cut by 27 per cent.

A rooftop photovoltaic system has also been progressively installed, and now totals 1230 square metres of solar panels. Reef HQ has become a registered solar power station, and its 205 kilowatt peak system should offset about 20 per cent of the aquarium's current power use. This initiative, combined with the installation of a new stored chilled water air conditioning system, means the target of a 50 per cent reduction by the end of 2013 is expected to be met.



Solar panels on the roof of Reef HQ Aquarium

### 5.2.3 Economic growth

Queensland's economy is currently worth \$260 billion per annum, and is principally based on mining, construction, tourism and agriculture.<sup>41,42</sup>

Queensland has had the highest long-term average economic growth rate of any Australian state or territory for over 20 years.<sup>41</sup> The state has had an average annual growth rate of 4.2 per cent over the last decade, and has outpaced the economic growth rate of both the rest of Australia and the Organisation for Economic Co-operation and Development group of nations for the past 20 years.<sup>41</sup>

Much of Queensland's economic activity takes place in the Great Barrier Reef catchment, although this activity is mostly not dependent on the Reef environment. The state's strong export trade is dominated by mining and agriculture. Three-quarters of Queensland's exports go to Asia, the most populous region in the world, with new trading opportunities opening up in Latin America and the Caribbean region. Japan is the state's largest trading partner, followed (in order) by China, India, South Korea and Taiwan.<sup>41</sup> About 80 per cent of the world's seaborne metallurgical coal exports are from Queensland<sup>41</sup>, shipped through the Great Barrier Reef.

Reef-dependent industries in the catchment and the World Heritage Area — tourism, commercial fishing, recreation and research activities — make a significant contribution to the state and national economy, especially in terms of employment. In 2012, these uses combined contributed \$5.7 billion to the Australian economy and employed almost 69,000 people (see Table 7.9 in Chapter 7).<sup>43</sup> These Reef-dependent industries provide benefits to associated industry sectors through demand for services such as vessel manufacture and maintenance, catering and retail.

In 2011, most people living in the catchment were employed in retail trade, health care and social assistance, or construction. Over the past six years, there have been fewer catchment residents employed in manufacturing, agriculture, forestry and fishing; and more employed in the mining and minerals sector, particularly in the Gladstone and Isaac local government areas.<sup>41,43</sup>

### Future trends

According to the 2013–2014 Queensland State Budget,<sup>44</sup> growth in the gross state product in 2011–2012 was mainly driven by increased business investment. Since 2010–2011, business investment has been dominated by the start of construction of three coal seam gas-to-liquefied natural gas projects, with a combined capital expenditure of more than \$60 billion. The staged completion of the three projects is projected to mean that business investment will fall in each year from 2013–2014 to 2015–2016. The first liquefied natural gas exports are scheduled for 2014–2015 and export volumes are expected to increase substantially in the following year, driving double digit growth in exports in 2015–2016. Although business investment is forecast to decrease in 2013–2014 and 2015–16, investment conditions are forecast to improve outside the resources sector, based on projected improvements in global economic conditions and projected depreciation of the Australian dollar.

The state budget forecast is that coal exports will continue to grow, supported by strong demand from China and the completion of the Eagle Downs and Grosvenor (Phase 1) coal mines. It also notes that major risk surrounds the timing and rate of liquefied natural gas investment. Growth in other sectors of the Queensland economy may not be enough to prevent Queensland's economic growth slowing more rapidly than currently expected, especially in 2014–2015.

The Queensland Government has indicated its aim is to continue growth in the state's economy by doubling the value of agricultural production by 2040,<sup>45</sup> expanding the resources industry, growing the construction sector through expedited planning processes, and enhancing tourism through marketing campaigns and the development of new tourism destinations and products.<sup>41</sup>

For agriculture in the catchment, it has been forecast that the pastoral sector is likely to grow and further intensify.<sup>42</sup> On the other hand, further large-scale changes from low-intensity agriculture to sugar production are forecast to be relatively unlikely in the medium to long term. In recent years, growth in intensive production of fruit and vegetables (horticulture) has generally exceeded demand.<sup>42</sup> While strategies are being developed to export product, their perishable nature and the expense of processing and transport will likely constrain large-scale growth.<sup>42</sup> As with other sectors of the Queensland economy, there is a degree of uncertainty around future trends in agriculture, especially in relation to global economic trends and the value of the Australian dollar.


There are a number of resource development projects proposed or under assessment. Changing economic circumstances mean it is difficult to predict the number that will reach construction and production. Growth in the mining and resources sector, in and adjacent to the Region's catchment, would more than likely require expanded port capacity and associated infrastructure along the coast adjacent to the southern half of the Region (Figure 5.4).<sup>41</sup>

In particular, Queensland has 98 per cent of Australia's proven coal seam gas reserves, and economic activity associated with the development of coal seam gas projects has grown substantially over the last decade (Figure 5.5). A Bureau of Resources and Energy Economics report anticipates that Australia will become the world's largest liquid natural gas exporter by 2020.<sup>46</sup> Curtis Island, within the World Heritage Area, is the site for the world's first project converting coal seam gas to liquefied natural gas.<sup>41</sup>

The continued and increasing global demand for coal<sup>47</sup> has resulted in mine expansions, new mines and additional mine proposals. The projected export volumes for coal in 2025 range from 267 to 383 million tonnes for thermal coal, and 260 to 306 million tonnes for metallurgical coal. Over this period, production of both thermal and metallurgical coal is projected to increase significantly in the Bowen, Surat and Galilee basins in Queensland, which export coal through the Great Barrier Reef.<sup>46</sup>

It is recognised that the recent slowing of the resources sector and the uncertainties around the realisation of potential projects, mean that growth in the sector may not be as substantial as has been forecast.

Changing global market trends also influence commercial fishing patterns in the Great Barrier Reef, especially the nature and level of demand for wild-caught product and the worldwide expansion of aquaculture fisheries.<sup>22</sup> Market conditions, price trends and competition have not been favourable enough to trigger widespread expansion of aquaculture in the Region's catchment.<sup>42</sup>



**Figure 5.4 Major resource projects planned or underway in Queensland<sup>41</sup>**

There are a number of major projects planned or underway that would greatly expand the volume of cargo handled by ports adjacent to the Region.

Tourism was identified in the early 2000s as having the greatest capacity for growth, and that may occur as the global financial situation improves.<sup>48</sup> The Queensland Government recognises tourism as a pillar of the state's economy and has identified a number of actions as catalysts and drivers for tourism investment, infrastructure and access. Actions of particular relevance to the Region and its tourism industry are:

- developing a Commercial Aviation Plan and a Queensland Drive Tourism Strategy
- ensuring a strategic approach to cruise shipping and superyacht priorities, infrastructure and supply chains
- including tourism in the State Planning Policy
- encouraging local governments to provide for appropriate tourism opportunities in their planning schemes (for example, a regional plan for Cape York)
- working with agencies to ensure tourism is considered in developing government plans and programs of economic and community infrastructure priorities
- developing a Queensland Ecotourism Plan.<sup>49</sup>

The value of tourism and commercial fishing in the Great Barrier Reef has remained relatively stable over the past six years.<sup>43</sup> It is difficult to predict future trends given uncertainty about the value of the Australian dollar, long-term implications of the global financial uncertainty and trends in Reef condition. Some onshore tourism sectors are expected to benefit from growth in the resources sector, particularly business-related travel, however, these benefits may not flow on to the marine sector.



## Implications for the Region's values

The scale and scope of the resources boom has led to changes in land use with the Great Barrier Reef catchment, particularly in the Fitzroy, Burdekin and Mackay–Whitsunday areas. It has also created unprecedented demands for water, power and new infrastructure including roads, railways and large-scale ports.<sup>50</sup> Any expansions to ports and associated infrastructure would result in the need for increased dredging and shipping in the Region (Sections 5.3.5 and 5.4.6). If not properly managed, increased industrial development can result in impacts to coastal and marine systems, adversely affecting the Region's productivity, biodiversity, connectivity and aesthetics at regional scales.<sup>50</sup>




Figure 5.5 Wells drilled in Queensland, 1954–2011<sup>51</sup>

## 5.2.4 Population growth

Indigenous Australians have lived in the Great Barrier Reef catchment for thousands of years, using its marine and coastal resources for food, shelter and cultural activities. European settlement commenced more than 150 years ago, resulting in an increased number of people living in the catchment. The estimated total population of the Great Barrier Reef catchment was 1,165,115 people as at 30 June 2012, which is approximately 25 per cent of Queensland's total population.<sup>52</sup>

The far northern part of the catchment (from north of Port Douglas and Mossman to Cape York) is sparsely populated. The two largest communities are Cooktown and Hope Vale, with 2339 and 858 residents respectively.<sup>52</sup> The southern part of the catchment from Port Douglas to Bundaberg is more heavily populated, including six major urban centres with populations of between 50,000 and 190,000 residents. The largest of these is Townsville, with an estimated population of 189,931, growing at an average of about 2.2 per cent every year.<sup>52</sup> This compares to a Queensland average growth rate of 1.7 per cent and a national average of 1.4 per cent.<sup>53</sup>

A profile of each of the catchment's natural resource management areas is presented in Table 5.1, including total population, average age and income of residents, and number of dwellings. The highest income earners live in the Mackay–Whitsundays natural resource management area, corresponding with higher resource-based activity. This coastal area also supports large numbers of fly-in fly-out miners.

Table 5.1 Population, housing and income within natural resource management areas of the Great Barrier Reef catchment, 2011<sup>54</sup>

Natural resource management area	Population	Average age	Number of dwellings	Average individual income
Cape York	8,396	34	3,730	\$37,794
Wet Tropics	186,514	37	85,352	\$39,311
Northern Gulf	43,729	41	20,374	\$33,628
Burdekin (also known as Burdekin Dry Tropics)	237,758	36	103,031	\$44,222
Mackay Whitsunday	112,796	37	45,665	\$47,831
Fitzroy	211,341	36	92,829	\$45,344
Burnett Mary	273,260	42	128,317	\$31,626

### Future trends

Much of the Great Barrier Reef catchment is expected to experience annual population growth rates of 1.6 per cent or higher in coming years.<sup>52</sup> In contrast, the national rate of population growth is expected to slow from 1.4 per cent to 1.2 per cent per annum over the next 40 years. Growth is forecast to be focused in the major regional centres (Figure 5.6) and many of the state's fastest growing local government areas are within the catchment, mostly in southern areas<sup>55</sup> (Figure 5.7). The Gladstone and Isaac local government areas are expected to expand particularly rapidly, due to increased activity in the resources sector.

As well as employment opportunities, lifestyle and amenity factors are likely to be attracting people to live in coastal areas.<sup>56</sup>

### Implications for the Region's values

Population growth in the catchment is likely to drive changes in a number of activities relevant to the Region and its values. These will range from more direct use of the Region to widespread impacts on coastal habitats that support the Great Barrier Reef.

With an increasing population comes an expanded urban footprint required to accommodate the increasing number of residents. Although urban development occupies only a small proportion of the catchment, its footprint has more than doubled in the last decade, especially in coastal areas.<sup>50</sup> This development — including its supporting infrastructure and services, such as roads, water, sewerage and power — affects catchment habitats that support the Great Barrier Reef, and has implications for the Region's values at a local level.

With regard to direct use of the Region, it is likely the trend of increasing recreational vessel ownership<sup>57</sup> will continue in the catchment. This is predicted to include faster growth in vessel numbers in the more southern parts of the catchment (Section 5.4.5). The trend towards people owning larger, better equipped vessels increases the potential for recreational users to access the outer reefs of the Region.

With increasing use of the Region comes increasing demand for coastal infrastructure (for example, roads, marinas and boat ramps) including in new sections of the coast. High demand and long wait times at popular access points can result in use being spread to adjacent, less popular areas as people choose to spend more time travelling and less time queuing.<sup>58</sup>

The movement of people into areas adjacent to the Region results in a higher proportion of new residents who may have less knowledge of its management arrangements than longer-term residents. However, there is evidence that some of the people moving to the catchment are 'tree-changers' and 'sea-changers', and that they have deliberately chosen to live adjacent to the Great Barrier Reef, often for amenity or lifestyle reasons rather than income.<sup>56</sup> These new residents are likely to become engaged with the Reef and its management.

Global population increases and demographic shifts are also likely to drive changes relevant to the Great Barrier Reef, ranging from increased emissions, pollution and marine debris to increased demand for seafood and increased shipping of export cargo through the Reef.





Figure 5.6 Predicted population growth in regional centres of the catchment, 2006–2031<sup>52</sup>



**Figure 5.7 Forecast average annual population growth by local government area, 2006–2031<sup>55</sup>**


### 5.2.5 Technological development

Technological development is the application of scientific knowledge to create tools to solve specific social, economic or environmental problems. Technological advances have brought major changes to the way people communicate, work, learn, travel and spend leisure time.

Technology has changed the way we learn about, manage and use the Region and its resources. Examples of its influence include:

- Reef managers, visitors, tourism operators, commercial and recreational fishers, researchers, students and teachers all benefit from the latest maps and satellite imagery of the Reef.
- Information about the Reef can be communicated instantaneously through social media such as Facebook, Twitter, blogs and online news reporting.
- The public can become involved in campaigns about the Reef through online advocacy organisations.
- Reef HQ Aquarium's videoconferencing facility enables students around the world to learn about the Great Barrier Reef.
- Global positioning systems allow safer navigation of the Reef, and the ability to more reliably locate sites and share locations with others. This technology also provides opportunities for sharing spatial information about the Reef and how it is used.
- Satellite-based vessel monitoring systems are used to track large ships and some commercial fishing vessels moving through the Great Barrier Reef.

- Satellite telemetry and acoustic arrays enable greater understanding of the movements and habitat use of species of conservation concern and those of economic importance.<sup>59,60</sup>
- Researchers use state-of-the-art satellite imagery, oceanographic instruments, laboratory equipment and portable weather stations to better understand, explain and predict changes in Reef condition, significantly improving understanding of the Region and contributing to its management.
- Light detection and ranging (lidar) mapping from low-flying aircraft delivers fast, high resolution bathymetry mapping in areas that are too shallow for vessel-based multibeam mapping.<sup>61</sup>
- Advanced diving technologies and remotely operated vehicles enable scientists and others to explore the length, breadth and depth of the Great Barrier Reef like never before.
- Development of antifouling alternatives that do not contain tributyltin (TBT) and have lower concentrations of copper, reduce shipping impacts.
- Turtle excluding devices on the suction openings of dredges reduce the effects of dredging on large mobile marine animals.<sup>62</sup>
- Improvements in fishing gear have reduced bycatch in trawl nets (for example, through turtle excluder devices and other bycatch reduction devices) and improved seafood product quality.<sup>62</sup>
- Technological developments have increased the efficiency of fishing operations to catch target species.<sup>63</sup>
- Newer engines, mobile refrigeration, depth sounders, side scanning sonar and positioning systems have increased the range, endurance and efficiency of boat and shore-based fishing.
- The combination of depth sounders and global positioning systems have improved fishers' ability to find fish and accurately relocate previous fishing sites and target deep shoals and fish aggregation areas.
- Reef-based tourism relies on technological advances, such as the latest dive gear, digital cameras and global positioning systems, to provide high quality visitor experiences.
- Low emission engines, which are more fuel efficient and less polluting, are increasingly being used by Reef users. Combined with more efficient hull designs, this is increasing vessel range.
- Web-based emissions calculators, such as those developed by the Authority, guide Reef-dependent industries in ways to reduce carbon emissions.
- Island resorts use new technologies, such as solar power and sewage treatment, to reduce their ecological footprint.
- In the catchment, advances in farming technology are reducing the use of fertilisers and pesticides, helping to slow and reverse negative trends in Reef water quality.<sup>64</sup>
- In the resources sector, new technologies are advancing mineral exploration and mining across the catchment, resulting in new mine sites and the reopening of mines that were not profitable using older technology.
- To improve catchment water quality and ecosystem health outcomes, the design of urban stormwater management devices are being improved and applied by local government to new developments.



### Future trends

Relationships between industry, government and community form and change over time, responding to and driving technological change, and altering land use and infrastructure development. New technology emerges from a combination of innovation, political imperatives, public acceptability and societal uptake. Future changes in technology and consumer demand will undoubtedly drive changes in the way people use the Region, the areas they want to use and the types of infrastructure installed; however, the nature of these changes is unclear. As an



example, greater interest in renewable energy may prompt applications for wind turbines over the ocean, or infrastructure using wave or tidal power. Changes in consumer demand and engineering advances may alter plans for tourism infrastructure.

### **Implications for the Region's values**

The development and spread of scientific knowledge and technologies can have significant implications for ecological systems, human use and wellbeing. New technologies can drive both positive and negative changes relevant to the Region's environment.

Technologies which better guide and monitor shipping traffic, enhance visitor experiences, reduce carbon emissions, monitor Reef use and contribute to our collective understanding of the Reef, enable the Region's values to be better protected and managed. Rapid improvements in spatial technology are providing an increasing number of spatial datasets for management. Changing vessel and navigational technology is likely to change the spatial patterns of fishing, tourism and recreational use, including allowing vessels to travel further offshore and better focus their use on preferred locations.

## **5.2.6 Societal attitudes**

Societal attitudes operate at international, national and local scales, and are shaped by cultural and social norms, institutional arrangements, economic imperatives and politics. They may be strongly influenced by external sources, particularly the mass media.<sup>65</sup> Societal attitudes significantly influence the potential for an individual, group or community to take action to help conserve natural assets such as the Great Barrier Reef.

Societal attitudes about the Reef have changed dramatically in the past and will continue to do so into the future. For thousands of years, societal attitudes about the Reef were those held by the Traditional Owner clan groups whose customary estates include sea country within the Region. Their culture and lore was reflected in ongoing stewardship and custodianship of the Reef environment. Traditional Owners continue to maintain a close and dynamic connection to their sea country, which integrates nature, heritage and culture.

The attitudes of early European explorers were principally shaped by their anxiety about being shipwrecked, due to the sheer size and complexity of the coral reef system. By the late 1800s, non-Indigenous Australians saw the Reef as a bountiful resource to exploit through whaling, dugong and turtle harvesting, pearling and commercial fishing. It was not until the early part of the twentieth century that they also began to explore its natural wonders in earnest, through science, recreation and tourism. This appreciation of the Reef flourished during the 1940s, 1950s and 1960s, and continues today.<sup>66</sup>

By the mid-1960s, Australians were beginning to express concerns about the future of the Great Barrier Reef, particularly with respect to outbreaks of crown-of-thorns starfish and the possibility of drilling for oil.<sup>66</sup> This growing public affinity for the Reef, and sense of responsibility for its future, led to the establishment of the *Great Barrier Reef Marine Park Act* (the Act) in 1975 and subsequent progressive protection of the Great Barrier Reef as a marine park.

Legislation such as the Act, non-regulatory incentives for behaviour change and international agreements and conventions (for example, the World Heritage Convention, the Kyoto Protocol and trade agreements) reflect societal attitudes, and play a major role in shaping the condition of the Great Barrier Reef.

### **Future trends**

Future trends regarding dominant societal attitudes are difficult to predict. International, national and local leaders who understand the issues and what is at stake can exert a strong influence on people's attitudes. Societal attitudes can also gradually change with persistent targeted messages reaching all sectors of society.<sup>65</sup>

Awareness of threats to the Reef, and willingness to accept responsibility for maintaining and restoring Reef health, are important pre-conditions for behavioural change towards the Reef.<sup>67</sup> Connections with the environment, feelings of trust, notions of fairness and respect also have a great influence on the way people respond to management initiatives and policies. Examples of social variables that may drive different outcomes for the Region include:

- national and state legislation
- international, national and state agreements, policies or guidelines
- mass media messages
- perceptions of the need for and fairness of environmental regulations
- perceptions of Reef condition
- feelings of empowerment to affect change
- individual and collective participation in environmental initiatives
- personal willingness to change behaviour towards the environment
- personal sense of place and attachment to local environments
- societal attitudes towards illegal or harmful activities on or adjacent to the Reef.

Before a person is willing to change activities or behaviours which have adverse impacts on the environment, they first have to recognise there is a problem. A study of the public's perception of the Reef found there is a high level

of acceptance that the community has a responsibility to look after the Reef.<sup>68</sup> Thirty-nine per cent of residents in Queensland coastal communities believe their activities at home impact on the Reef, while 23 per cent believe their activities at work impact on the Reef.<sup>69</sup>

### Implications for the Region's values

Changes in societal attitudes on a global scale through to attitudinal changes in a coastal community can all drive changes in activities relevant to the Region's environment.

Stewardship actions driven by community and industry are critically important for changing prevailing attitudes and supporting management initiatives to maintain and enhance the Region's values. Stewardship is a central tenet of many of the management programs for the Region, and the growing interest in these programs reflects shifts in thinking towards ecologically sustainable development, human wellbeing and a healthy, vibrant Great Barrier Reef.

## 5.3 Activities adjacent to the Region

This section examines activities within Great Barrier Reef catchment that are likely to be causing impacts or pressures on the values of the Region (Figure 5.8). The activities described are:

- agriculture
- aquaculture
- urban development
- industrial development and resource extraction
- port activities (undertaken in both the catchment and the Region)

The summary includes an examination of trends and indirect impacts on the Region's values. A more comprehensive analysis of these activities is presented in the Queensland Government's Coastal Zone Strategic Assessment.





Figure 5.8 Activities within and adjacent to the Region

### 5.3.1 Agriculture

Agriculture within the Great Barrier Reef catchment is an important producer of food and, to a lesser extent, fibre. In 2010–11, Queensland's agriculture had a total gross value of \$9.5 billion<sup>70</sup>, much of which would have been derived from the catchment. The Great Barrier Reef catchment was first developed for agriculture when cattle and sheep grazing were introduced at the time of European settlement in the 1850s.<sup>71</sup> Intensive agriculture (sugar cane) started about 10 years later.<sup>71</sup> However, it was probably not until the early to mid-1900s and the introduction of heavy machinery that land clearing accelerated, and intensification of the use of the coastal zone began in earnest.<sup>71</sup> This change in the use of the catchment has been recorded in sediments and coral cores taken in the Great Barrier Reef as changes in the sediment signatures<sup>72</sup>, in nutrients<sup>73</sup> and in coral communities.<sup>74</sup> Today, more than 80 per cent of the catchment supports some form of agriculture (Figure 5.9).

Grazing is the most extensive land use, occurring in more than 74 per cent of the catchment. It is particularly extensive in the larger dry tropical catchments of the Fitzroy and Burdekin rivers but is a significant portion of most catchments, even in the Wet Tropics. It is the dominant land use in Cape York, covering about 50 per cent of the Great Barrier Reef catchment in that area (Figure 5.9).



**Figure 5.9 Land use in the Great Barrier Reef catchment, 2009 and 1999<sup>50</sup>**

Grazing is the predominant land use in the catchment. Intensive agriculture is confined to a relatively small area, mainly close to the coast. For much of the catchment, comprehensive land use mapping was undertaken in 2009. For the Cape York natural resource management area, the most recent information is from 1999.



Intensive agriculture occurs in about five per cent of the catchment. It is largely confined to the lower coastal floodplain of the Great Barrier Reef catchment, south of Port Douglas. The coastal plains with the most intensive agricultural production are the Mackay Whitsunday, Wet Tropics and Burnett Mary regions. Extensive grain and cotton areas in the upper Fitzroy catchment and, to a lesser extent, mixed cropping on the Atherton Tableland in the Barron catchment and upland parts of the Mary catchment are the exceptions, where intensive agriculture is inland of the coastal plain (Figure 5.9).

### Trends

Over the 160 years of European settlement in the catchment, agricultural use has become more widespread and more intensive. Historically, broadscale land clearing and intensive cropping on the coastal floodplain have affected coastal habitats and the processes that support the Great Barrier Reef environment.

Between 1990 and 1999, the area under sugar cultivation grew by 30 per cent from 323,000 to 424,000 hectares. In the same period, there was also rapid growth in banana cultivation, mainly in the Tully and Johnstone River catchments, south of Cairns, with the area under cultivation more than doubling so that the area now produces most of Australia's bananas.<sup>71</sup> This expansion in intensive agriculture correlates strongly with the peak of broadscale tree clearing. The rates of broadscale clearing in Queensland increased until the late 1990s (around 750,000 hectares per year in 1999–2000) when the *Vegetation Management Act 1999* (Qld) was introduced. By 2007–08, implementation of this legislation had significantly reduced the rate of clearing to about 123,000 hectares per year and, with policies for wetland protection, led to greatly reduced rates of loss of coastal ecosystems. The overall loss of wetlands, however, has been significant in the coastal floodplain, with more than 80 per cent lost in many Great Barrier Reef basins<sup>50</sup> — noting that the rate of wetland loss has slowed in recent years. The full ramifications of these losses on the health of Great Barrier Reef ecosystems remain to be quantified. Work over many years has identified that prior to the implementation of the *Vegetation Management Act* broadscale clearing was having a significant impact on water quality flowing to the Great Barrier Reef. A 2008 scientific consensus report on water quality in the Great Barrier Reef<sup>75</sup> identified that this legislation was responsible for reducing the extensive land clearing of previous decades, and had been a critical element in governments beginning to address the impacts of land use on water quality in the Great Barrier Reef as reported from the 1970s.

Recently, the Queensland Government proposed a suite of changes to the *Vegetation Management Act* which includes repealing regulations that apply to clearing high value regrowth on freehold and Indigenous lands; allowing broadscale clearing for 'high value' intensive agricultural production; and promoting self-assessment of areas that contain remnant or high value regrowth.<sup>76</sup>

There is also a strong relationship between agricultural chemical inputs (for example, fertilisers) and increasing intensification of agricultural activity (for example, moving from grazing to cropping). The use of fertilisers throughout the southern and central sections of the Great Barrier Reef catchment increased exponentially from the 1940s to about 2000 (Figure 5.10). After this period, it began to decline.<sup>75</sup> There are a number of factors contributing to a decline in fertiliser use. The issue of water quality decline in the Great Barrier Reef came to the fore in the early 2000s with the release of the Authority's Water Quality Action Plan.<sup>77</sup> This plan highlighted the need for significant reductions in nutrient inputs to the Great Barrier Reef and led to the joint Australian and Queensland government's *Reef Water Quality Protection Plan* (Reef Plan) in 2003. The early 2000s also saw two major multi-million dollar Australian Government funded programs in the Great Barrier Reef catchment: the Natural Heritage Trust; and the National Action Plan for Salinity and Water Quality. Both these programs supported Reef Plan actions and provided assistance to farmers and other land managers through, for example, landcare groups or local government, to improve land management activities and water quality. This focus on improving land management and identifying issues around excess fertiliser application led to an improved attention by industry on best practices, and a general and sustained reduction in application rates.<sup>78</sup> Additionally, as noted above, the Queensland *Vegetation Management Act*, which came into effect in early 2000, significantly reduced the rate of broadscale tree clearing in the state.<sup>75</sup> More recently, Australian Government initiatives such as *Caring for Our Country* and the Queensland Government's regulatory measures and extension programs have been aimed at improving land management practices by farmers and graziers.

Land use data shows the area under intensive agriculture (that is, where fertiliser applications are made) remained stable with little or no increase between 1999 and 2009.<sup>50</sup> There were a number of market forces that put downward pressure on fertiliser use and potentially led to a drop in fertiliser application rates, including worldwide declines in sugar prices through the early 2000s, and increasing costs of fertiliser.<sup>78,79</sup>

As reported in Section 5.2.3, it is forecast the pastoral sector is likely to grow and further intensify in the catchment in the medium to long term, but that large-scale changes to sugar production are relatively unlikely. Large-scale growth in intensive production of fruit and vegetables (horticulture) is predicted to be unlikely.<sup>42</sup>

### Impacts


Over the years, agricultural development in the catchment has resulted in significant loss, modification and fragmentation of terrestrial habitats that support the Great Barrier Reef, which has affected the health of the Reef's inshore ecosystems.<sup>50</sup> Extensive broadscale clearing of forests, woodlands and forested floodplains has increased erosion and, therefore, sediment loads in rivers, streams, freshwater wetlands and, ultimately, the Great Barrier Reef.<sup>50</sup> A significant area of forested floodplain and wetland habitats has been lost or modified for intensive agriculture.<sup>50</sup> In addition to flow-on effects to the Region from clearing these habitats, activities such as infilling and building channels, levees and bunds have affected many ecological processes that support the Great



Barrier Reef, such as water flow, groundwater recharge and discharge, nutrient and sediment cycling, and connectivity.<sup>80</sup>

Diffuse source pollution from agriculture is the main source of excess nutrients, fine sediments and pesticides, including herbicides, entering the Great Barrier Reef Region.<sup>11</sup> The trends in loads of these pollutants entering the Great Barrier Reef steadily increased with agricultural development of the catchment. In 2012, nutrient and sediment loads were estimated to be about five or six times natural levels.<sup>81</sup> Together, increased sediments, nutrients and pesticides have had serious impacts on marine ecosystems, particularly coral reefs and seagrass beds. For example, recent research has shown the timing of shifts or degradation of coral communities on inshore reefs is strongly correlated with increases in terrestrial run-off.<sup>74</sup> Furthermore, there is increasing evidence of a link between increased nutrients and crown-of-thorns starfish outbreaks<sup>82</sup> and degradation of seagrass meadows<sup>83</sup>. In some cases, agricultural activities have also resulted in the exposure of acid sulphate soils.<sup>84</sup>

Recent improvements in land management and agricultural practices are recognised in the 2013 Water Quality Scientific Consensus Statement<sup>11</sup> which concluded that *'the use of improved land and agricultural management practices is proven to reduce run-off of suspended sediment, nutrients and pesticides at the paddock scale'*. These improvements will potentially mean reduced future impacts on the Region's ecosystems.



**Figure 5.10 Fertiliser use in the Great Barrier Reef catchment**

After decades of increasing fertiliser use in the Great Barrier Reef catchment, calculations indicate the amounts used are now declining. The 1910 to 1990 data was derived from Pulsford.<sup>85</sup> The 1999, 2009 and 2010 data points were estimated using Queensland Land Use Mapping Program data<sup>86</sup>, the Fertilizer Industry Association of Australia application rates for different land uses<sup>87</sup> and Incitec Pivot published figures<sup>88</sup>.

### 5.3.2 Aquaculture

Aquaculture refers to an operation for the propagation, rearing, keeping or breeding of an aquatic organism. It is Australia's fastest growing primary industry.<sup>89</sup> Although there has been limited marine-based aquaculture within the Region, and no facilities are in operation at present, land-based aquaculture occurs in the catchment, principally for prawns, barramundi, redclaw and freshwater fish.<sup>90,91</sup>

Aquaculture operations are located close to the coast in a number of areas in the southern half of the catchment, typically where there is access to good water supplies. There are approximately 700 hectares of prawn ponds currently in production adjacent to the Great Barrier Reef.<sup>90</sup>

#### Trends

While over the last decade there has been little spatial expansion of land-based aquaculture adjacent to the Region, overall production has increased.<sup>90</sup> Before 2010–2011, prawn aquaculture experienced strong annual production increases; however, in 2010–11 production fell by 25 per cent, mainly due to cyclones.<sup>90</sup> Several marine-based aquaculture operations have been proposed or have begun over the past two decades (for example fish, pearls, sponges) but none are in operation at present, primarily due to economic or environmental sustainability issues.

#### Impacts

The potential impacts on the Great Barrier Reef environment associated with land-based aquaculture facilities include: increased loads of sediment and nutrients (nitrogen and phosphorus) in discharged wastewater; clearing or modifying coastal habitats; modifying hydrologic processes; disturbing acid sulphate soils; introducing marine species; genetic pollution; and introducing disease.<sup>92</sup>

While the loads of sediment and nutrients discharged to waterways leading to the Great Barrier Reef are relatively small in comparison to those derived from other agricultural sources (for example grazing, cane farming, horticulture), their impact locally can be significant.

Significant work is currently underway investigating waste treatment methods that can reduce the load of sediment and nutrients discharged by these facilities. Settlement ponds are already being utilised by new farming ventures to treat this waste, whilst further work investigating the commercial feasibility of algal bioremediation is also underway.

The major sustainability issue is the same as for other industrial or agricultural activity that discharges sediment and nutrients to Great Barrier Reef waters. Given that the majority of the inshore waters in the southern two-thirds of the Great Barrier Reef Marine Park already exceed their assimilative capacity, the ability for these waters to assimilate additional loads of sediment and nutrients discharged by these facilities is limited.

Actions to offset environmental impacts and deliver net benefits have been proposed as mechanisms to allow the expansion of this industry, while meeting the long-term goal that by 2020 the water quality entering the reef from broadscale land use has no detrimental impact on the health and resilience of the Great Barrier Reef.

The Great Barrier Reef Marine Park (Aquaculture) Regulations were introduced in 2000 to ensure the discharge of waste from land-based aquaculture facilities did not significantly impact the plants and animals of the Marine Park. In 2005, the then Minister for the Environment and Heritage accredited Queensland Law as providing the requisite degree of protection for the plants and animals of the Great Barrier Reef Marine Park. Based on this accreditation, Queensland is now solely responsible for the assessment and approval of all land-based aquaculture facilities that discharge waste to waterways that lead to the Marine Park.

### 5.3.3 Urban development

Urban development refers to the construction or expansion of a town or city, including buildings, houses, roads, water and electricity supply. The six major population centres within the Great Barrier Reef catchment — Cairns, Townsville, Mackay, Rockhampton, Gladstone and Bundaberg — account for about 42 per cent of the population. They are the areas of most intensive urban development.

#### Trends

Population growth in the Great Barrier Reef catchment continues to occur at rates faster than the Australian average, especially along the coast (Figure 5.7). The catchment's current population is expected to grow by approximately 44 per cent to 1.74 million by 2031.<sup>93</sup> The majority of this population is expected to live on or near the coast. Population growth in the regional centres of the catchment (Figure 5.6) is driving expansion of its urban footprint. For example, Townsville's population is projected to grow from approximately 191,000 to 300,000 by 2031. As a result, the number of dwellings in Townsville is predicted to increase from 74,205 in 2011 to 121,914 in 2031.<sup>93</sup>

#### Impacts

Although urban development occupies only a small proportion of Queensland's total land area, it can be a source of impacts in relation to the Region's values at a local scale.

By its very nature, urban development in coastal areas modifies the local habitats that support the Great Barrier Reef ecosystem — this includes clearing and infilling, modifying water flow and changing aquatic connectivity between coastal and marine habitats.<sup>50</sup>

Local government is primarily responsible for managing large sewage treatment plants in urban areas along the Great Barrier Reef coast, and has the responsibility of permitting smaller sewage treatment facilities in peri-urban areas. Local councils continue to work to minimise the amount of pollutants entering the Region's waters from urban areas; however, these areas remain a localised source of nutrients, pesticides and litter entering the marine system.<sup>94</sup> Such pollution can enter the Region via sewage, stormwater and wastewater discharge. While the contribution of sewage discharge to the Region's overall nutrient loads is relatively small, it often occurs in more disturbed areas already subject to multiple other impacts. In 2002, the Queensland State Coastal Management Plan<sup>95</sup> identified the need for improving sewage management for environmental and human health reasons. It required that all sewage treatment plants discharging into coastal waters that enter the Marine Park be upgraded to remove excess nutrients by 2010. This involved significant investment by the Queensland Government and local councils, with assistance from the Australian Government, to upgrade facilities in major population centres. It has resulted in significant reductions in the nutrient load from urban areas.

There remains a number of smaller communities along the coast that discharge secondary treated sewage to waterways that lead to the Great Barrier Reef, or that are serviced by septic systems. In these communities, it is currently not considered to be economically viable to upgrade to tertiary treatment.<sup>96</sup> As Queensland's population continues to grow along the coast, this is likely to be an increasing issue for water quality.

All island resorts discharging sewage from marine outfalls into the Marine Park meet tertiary standards.<sup>22</sup>

Other local-scale impacts from urban development include exposure of potential acid sulphate soils<sup>51</sup>, increased light<sup>97</sup>, and atmospheric pollution, increased noise and visual disturbance.

### Land-based sewage management in the Great Barrier Reef and its catchment

In 1991, the Authority introduced a policy requiring tourism resort operators to have a permit to discharge sewage into the Marine Park and to have upgraded their sewage treatment plants to tertiary treatment standard by 2002. This decision was based on the best science at the time, which had begun to identify issues with changes in the nutrient and sediment concentrations in coastal waters of the Region. It was also in response to a specific case of seagrass communities expanding because of the retention of diluted effluent on the reef flat around a resort, and localised effects on coral reefs adjacent to other resorts.<sup>98,99,100</sup> An assessment of sewage discharges into the Region undertaken in 2000, identified 16 other sewage treatment plants that discharged into coastal waters in or adjacent to the Marine Park that needed upgrading to achieve the Authority's tertiary treatment policy.<sup>101</sup>

In parallel, the Queensland Government implemented a process for improving coastal management and identified critical issues, such as the need to improve management of sewage discharges, with the release of its position paper on managing the coast.<sup>102</sup> This led to the first Queensland State Coastal Management Plan which came into effect in August 2002, and required all sewage treatment plants discharging into coastal waters that enter the Marine Park to upgrade facilities to a tertiary treatment standard by 2010.<sup>95</sup>



*Northern wastewater treatment plant, Cairns*

Over the subsequent decade, about \$620 million was invested in upgrading sewage treatment plants to a tertiary treatment standard in the three largest coastal cities adjacent to the Great Barrier Reef — Cairns, Townsville and Mackay. All levels of government and the coastal communities have contributed to these initiatives. Most population centres that discharge sewage via waterways that lead to the Reef now treat their sewage to tertiary standard.<sup>22</sup> Until 2011, the Queensland Government had provided about 50 per cent of the capital expenditure to fund these upgrades. Some upgrades are still underway in smaller population centres (for example, Sarina is undergoing a \$40 million upgrade). The Australian Government has supported local governments in some of these upgrades; for example, contributing \$40 million to the Townsville City Council as part of the \$170 million redevelopment of the Mt St John sewage treatment plant. Most of the remaining funds for upgrading or building new sewage treatment plants have come from the coastal communities themselves, through local government rates.

While inputs of nutrients from sewage treatment plants accounted for only a small percentage of the overall nitrogen load entering the Great Barrier Reef from its catchment (less than four per cent of the total nitrogen load and less than one per cent of the total phosphorus load<sup>22</sup>), the reduction of loads at localities adjacent to high population centres is substantial. For example, in the Townsville local government area, introduction of tertiary treatment reduced the total nitrogen and total phosphorus loads by 81 per cent<sup>103</sup> which equates to an approximate 42 per cent load reduction from the Black and Ross rivers catchment. A reduction in the nutrient load from these facilities is likely to improve the overall ecosystem resilience of these areas to cumulative impacts. In total, the upgrades have removed an estimated 834 tonnes of nutrients per annum (approximately 80 per cent of the original total nutrient load from this source) that would otherwise have entered the Region.

### 5.3.4 Industrial development and resource extraction

For the purpose of this report, industrial development refers to the construction, operation or expansion of commercial industries, excluding agriculture, ports, tourism, fishing and aquaculture. Historically, there have been extensive small-scale mining operations through much of the catchment, including gold, tin, nickel and uranium mines.<sup>104</sup> At present, very limited mining occurs within the Great Barrier Reef coastal zone, and no mining occurs in the Great Barrier Reef World Heritage Area. A more detailed discussion of ongoing mining activities in the coastal zone of the World Heritage Area can be found in Section 5.2.5 of the Coastal Strategic Assessment

Report. Much of the supporting infrastructure for mining and industry is located in coastal areas and, in the past two decades, major State Development Areas have been declared in Gladstone (1993), Townsville (2003), Abbot Point (2008) and Gladstone–Curtis Island (2008).<sup>105</sup> State Development Areas are defined areas established to promote economic development in Queensland.

### Trends

Trends in industrial development in the catchment are described in Section 5.2.3.

### Impacts

Mining and industrial development can lead to increased pressure for expanded urban and service infrastructure to cater for more workers.<sup>106</sup> With regard to resource extraction, although the footprint of a mine or well may be relatively small, installation of the supporting infrastructure can lead to fragmentation, modification or loss of local coastal ecosystems.<sup>50</sup> If not properly managed, industrial development adjacent to the Great Barrier Reef can also result in localised impacts such as the exposure of potential acid sulphate soils; light pollution<sup>97</sup>; atmospheric pollution; artificial barriers to estuarine flow; discharge of pollutants; coastal reclamation; and marine debris.<sup>50,107</sup> Coastal industrial development can also diminish aesthetic values for users of the Region.<sup>108</sup>

### 5.3.5 Port activities

As an island nation, Australia is dependent on maritime trade. Consequently, ports and their associated infrastructure are of significant economic and social importance to Australia. A number of Queensland's ports, including those in the Region, are considered nationally significant for cargo throughputs and contributions to the national economy.<sup>109</sup>

There are 12 trading ports in the World Heritage Area, managed by four port authorities — all Queensland Government-owned corporations. Of these, eight are located at least partly in the Region, and only the minor ports of Cooktown and Quintell Beach in Cape York are located within the Marine Park (Table 5.2). In 2011–12, ports within the Great Barrier Reef Region accounted for 76 per cent of the total throughput for all Queensland ports combined. This amounted to 199.8 million tonnes of imports and exports through the Region (Figure 5.11).<sup>109</sup>

**Table 5.2 Trading ports located within, or partly within, the Great Barrier Reef World Heritage Area, Region and Marine Park**

A trading port refers to a port that is associated with a pilotage area managed by Maritime Safety Queensland.

Port	Great Barrier Reef World Heritage Area	Great Barrier Reef Region	Great Barrier Reef Marine Park
Quintell Beach	●	●	●
Cape Flattery	●	●	
Cooktown	●	●	●
Cairns	●		
Mourilyan	●		
Lucinda	●	●	
Townsville	●	●	
Abbot Point	●	●	
Mackay	●	●	
Hay Point	●	●	
Port Alma	●		
Gladstone	●		

In terms of infrastructure, operational capacity and size, the largest ports on the Region's coast are Abbot Point, Gladstone, Hay Point and Townsville. The Gladstone, Abbot Point and Hay Point ports are major hubs for the export of coal.<sup>46</sup> Hay Point is one of the largest coal export terminals in the world,<sup>110</sup> handling more than 80 million tonnes of coal in 2011–12.<sup>109</sup> During 2010–11, it had the highest value of coal exports among all Queensland ports at \$18 billion.<sup>47</sup>



The land and ocean-based activities associated with ports include:

- rail and road networks
- terminals, loading and unloading facilities
- land reclamation
- trestle structures
- dredging and disposal of dredge material
- storage and waste facilities, cargo holding facilities, stockpiles
- safety and navigational aids and lighting
- monitoring buoys
- tug boat and shipping berths
- ship departure channels and anchorages.

Activities associated with ports cut across many of the jurisdictional boundaries relevant to the Region, occurring on land, as well as in the World Heritage Area, the Region and the Marine Park.

Figure 5.12 illustrates the jurisdictional boundaries relevant to the operation of ports around Mackay.





Figure 5.11 Queensland trading ports in the Great Barrier Reef World Heritage Area, Region and Marine Park and their throughput, 2011–12<sup>109</sup>



**Figure 5.12 Port infrastructure at Mackay and the Port of Hay Point**

Using Mackay and Hay Point as examples, the facilities and management arrangements relating to the operation of Great Barrier Reef ports often cut across the boundaries for the World Heritage Area, the Region and the Marine Park.

## Trends

There has been major expansion of ports on the Region's coast over the past two decades. Between 1988–89 and 2010, total export tonnage from these ports grew by 297 per cent.<sup>111</sup> In the same period, the value of the exports increased by 550 per cent. Increases in bulk commodity exports from these ports are driving increases in shipping.

As at June 2013, five of the 12 commercial ports (Cairns, Townsville, Abbot Point, Hay Point and Gladstone) had active proposals for port expansions, driven mainly by growth in the resources sector. There are also proposals for two new ports (Wongai in Cape York and the Fitzroy terminal project) adjacent to the Region. A proposal to develop a port on Balacclava Island has been withdrawn, and it is noted that not all the current proposals may proceed.

A Bureau of Resources and Energy Economics report found the total capacity of planned infrastructure projects progressing through the approval process exceeds the projected volumes of commodity exports out to 2025.<sup>46</sup> However, a lower than expected ability to make use of this capacity was identified as a key risk, with predictions that further capacity expansion may be required to compensate for the lack of consistent throughput. This finding is similar to a recent study by the Queensland Department of State Development, Infrastructure and Planning that found Queensland's three major coal ports (Hay Point, Gladstone and Abbot Point) operated at only 52 per cent of their combined capacity in 2011–12, and that Abbot Point only operated at one quarter of its total capacity.<sup>112</sup> However, it should be noted that the industry was heavily impacted by severe weather and flooding between 2011 and 2012, hence these numbers may under represent the capacity. The exception is liquid natural gas, where the Bureau of Resources and Energy Economics report predicts that under a high market share scenario, there is insufficient time to build adequate capacity by 2020, despite planned infrastructure capacity exceeding projected export volumes.<sup>46</sup>

The draft of ships is a significant factor limiting navigable waterways, and the worldwide trend towards longer, deeper draft ships<sup>113</sup> affects port access requirements. In order to accommodate deeper draft ships, some ports may require more capital and ongoing maintenance dredging into the future.

## Impacts

Ports and associated access channels have been dredged since their establishment, well before the Marine Park was declared. Today, both capital and maintenance dredging are undertaken within and adjacent to the Marine Park. Sites for the disposal of dredge material (sediments from the seafloor) may be at sea or on land. Existing ocean disposal sites are located within and adjacent to the Marine Park. Most large-scale dredging and dredge material disposal are associated with the larger and busier ports such as Cairns, Townsville, Abbot Point, Hay Point and Gladstone.<sup>47</sup>

The dredging itself, plus the disposal of dredge material and its later resuspension, can have direct effects on ecosystems, such as coral reefs<sup>114,115</sup> and other habitats. Examples include: removal of existing habitats such as seagrasses;<sup>116</sup> disturbance to the seabed; increased underwater noise;<sup>117</sup> reduced water quality; transport and resuspension of contaminants and nutrients; burial and smothering of life on the seafloor, both at the disposal site and in surrounding areas after resuspension;<sup>118</sup> and increased turbidity.<sup>118,119</sup>

Impacts to the marine environment from the installation and maintenance of port infrastructure and general port activities may include: clearing, modifying and fragmenting coastal habitats; reclamation of marine areas; exposure of potential acid sulphate soils; creation of artificial habitats; alteration of natural coastal processes; the risk of large and small chemical and oil spills; marine debris; injury or death of marine wildlife; altered light regimes; displacement of other Marine Park users; and diminished aesthetic values for users and nearby communities. The impacts of ports are also directly linked to impacts associated with shipping and ship anchorages.

## 5.4 Activities within the Region

This section examines activities within the Great Barrier Reef Region that are likely to be causing impacts or pressures on the values of the Region (Figure 5.8). The activities described are:

- traditional use of marine resources
- tourism
- fishing — commercial
- fishing — recreational
- recreation
- shipping
- defence activities
- research activities.

Some of the activities are directly dependent on the Region's natural resources (Reef-dependent) and others are carried out regardless of the natural environment (not Reef-dependent). Examples of Reef-dependent activities are traditional use of marine resources, tourism, fishing (commercial and recreational), recreation and research activities. Reef-dependent activities are likely to be more sensitive to changes in the condition of the Region's values.<sup>120</sup>

### 5.4.1 Traditional use of marine resources

Traditional Owner connection to sea country within the Region continues to be practised and maintained according to traditional customs and spiritual lore, reflecting the ongoing stewardship and custodianship.

Traditional use of marine resources is the undertaking of activities as part of Aboriginal and Torres Strait Islander peoples' customs or traditions, for the purpose of satisfying personal, domestic or communal needs. It includes fishing, collecting (for example, shellfish) and hunting, as well as looking after cultural and heritage sites.

Marine resources have distinct cultural uses. Traditional Owners use these resources to practise their living maritime culture, provide traditional food for their families and educate the younger generation about traditional and cultural rules, protocols and activities in sea country.

For more than 40,000 years, the traditional use of marine resources has been conducted sustainably. Impacts such as coastal development, habitat degradation, boat strikes, pollution, netting and sedimentation as well as legacy impacts have affected Traditional Owners' use of the marine environment. Traditional Owners are now working in partnership with the Authority to conserve and protect species and ecosystems critical to the health of people, culture and country, including through development and implementation of Traditional Use of Marine Resources Agreements.

For example, since European settlement in Australia, the dramatic reduction in species such as marine turtles and dugongs has reduced the capacity of these populations to withstand the multitude of impacts that affect them, such as habitat degradation, boat strikes, pollution, netting, sedimentation and the effects of coastal development. In recognition of the pressures on these species, some Traditional Owner groups have taken steps to reduce, temporarily suspend or cease traditional hunting of them. In their Traditional Use of Marine Resources Agreement, the Woppaburra Traditional Owners of the Keppel Island region have specified that green turtle harvest will be limited and there will be no take of dugong at all:

*"We are fulfilling our customary obligations by taking an active role in managing the marine resources in our sea country for the future. Under our Dharumbal–Woppaburra Traditional Use of Marine Resources Agreement, we have agreed to limit green turtle harvest in specific areas under a traditional authority system, and to not harvest dugongs. We recognise that marine turtles and dugongs are under threat from many pressures and we are committed to the conservation and protection of these and all marine species. We are proud of our sea country and welcome you. We work in partnership with the Australian and Queensland governments, and we ask you to help keep our shared sea country healthy by following zoning rules and doing all you can to minimise the impacts of your visit."<sup>121</sup>*

Other Traditional Owner groups — such as the Gooreng Gooreng, Gurang, Taribelang Bunda and Bailai peoples who have an agreement covering the Port Curtis Coral Coast — have voluntarily agreed not to hunt dugongs for five years from 2011, in response to that year's extreme weather and the number of dugongs that were stranded dead in eastern Queensland.

### **Trends**

Traditional Use of Marine Resources Agreements, and one marine Indigenous Land Use Agreement, apply to approximately 13 per cent of the Great Barrier Reef Marine Park. Due to the success of the Authority's Reef Rescue Land and Sea Country Indigenous Partnership Program, other Traditional Owner groups are currently working to develop their own agreements, which will result in a significant increase in the area covered. With increased development in remote areas and changes to infrastructure, there is potential for the level and type of traditional use along the coast to change.

### **Benefits**

The continuing sea country management and custodianship of the Great Barrier Reef by Aboriginal and Torres Strait Islander Traditional Owners provides immeasurable benefit.

Many Traditional Owners use marine resources to practise their sustainable 'living maritime culture'; provide traditional food for families; and educate younger generations about traditional and cultural rules, protocols and activities in sea country.

Traditions are of high cultural importance, while social sharing during special events that require traditional resources is also critical to maintaining culture. Traditional Owners hold many cultural, economic and spiritual connections to the Region; establishing effective partnerships with them helps protect cultural and heritage values, conserve biodiversity and enhance the resilience of the Great Barrier Reef.

### **Impacts**

Any impacts attributable to traditional use of marine resources undertaken according to customs and traditions, are considered to have only minor or localised effects on values. This is distinct from any illegal poaching of species of conservation concern undertaken without the customary approval of the relevant Traditional Owners.

There have been some recent examples of disputes in multiple-use areas, where the activities of tourism operators and visitors conflicts with Traditional Owner cultural use of marine resources in the sea country areas where they express their native title rights.<sup>122</sup>

## **5.4.2 Tourism**

In this report, tourism is defined as the commercial activities that provide transport, accommodation or services to people who are visiting the Region principally for enjoyment. Non-commercial recreational activities are discussed in Section 5.4.5.

Australia's rich natural resources and high number of natural world heritage sites, based on a comparatively pristine natural environment, are recognised as some of the nation's tourism strengths.<sup>123</sup> For the Region, its long-term attractiveness as a tourism destination is largely based on the Great Barrier Reef's reputation as the world's largest and best known coral reef ecosystem — one that has spectacular and diverse species — combined with a high standard of tourism operations and protected area management. The Region's tourism industry is almost exclusively nature-based, with coral reefs and islands as the focus. There are opportunities to see iconic wildlife such as whales, turtles, sharks and seabirds — many of which are matters of national environmental significance — and to go boating, diving, snorkelling, fishing, sailing, hiking, camping or enjoy various water sports.

The industry offers a wide range of tourism experiences, from cruise ships and live-aboard vessels to day trips on high speed catamarans, fishing charters and kayaking tours. While tourism visitation occurs across most of the Great Barrier Reef, activity is consistently focused on a small portion of the Marine Park, with more than 85 per cent of all tourism activity management occurring in about seven per cent of the Region (Figure 5.13).<sup>124</sup>






## Trends

Commercial tourism began on the Great Barrier Reef in the 1890s. Since that time, vessels and technologies available for viewing the Reef environment and enjoying the Region have changed substantially. The number of visitors has increased, along with the size and geographic distribution of the industry.


Total tourism visitation to the Great Barrier Reef Marine Park in 2011 was 1.842 million.<sup>124</sup> Tourism visitation rose gradually over the 10 years prior to 2005, and has subsequently declined by about eight per cent per year (Figure 5.14). The declines have been attributable to a range of factors, including the high exchange rate of the Australian dollar, increased competition from international destinations, extreme weather events and the global financial crisis. In 2012, there were signs of a recovery in tourism visitation, however it was patchy in its extent.

The forecast outlook is for a slow recovery bolstered by the emerging markets of China and India (Figure 5.15), recent improved availability of airline capacity on some Asian airline routes, and a strengthening domestic market.



**Figure 5.14 Number of tourism visitor days to the Great Barrier Reef Marine Park, 1994–2012**<sup>124</sup>

A 'visitor day' is a visit by one tourist for one day. For overnight visits, each day is counted separately (e.g. a three-day visit by a tourist represents three visitor days). A half-day visit refers to visitors who undertake a trip of less than three hours, and exempt from the environmental management charge includes young children and trade familiarisations. Ongoing improvements in the way the environmental management charge information is recorded has progressively allowed more accurate differentiation of visitation. This figure does not include stand-alone coral viewing activities and scenic flights, estimated at more than 0.2 million. It also does not include the estimated 2.3 million passenger transfers conducted each year through the Region to and from islands.




**Figure 5.15 International visitation to the Great Barrier Reef catchment, 2002–2011**<sup>126</sup>

There has been a marked decline in Japanese visitors to the Reef catchment over the last decade. Visitation by Chinese tourists is increasing.

The Authority works in partnership with the tourism industry to improve standards for the protection and presentation of the Great Barrier Reef. Operators who are independently certified with Ecotourism Australia's ECO Certification Program at Ecotourism and Advanced Ecotourism levels, and who meet best practice standards when undertaking their tourism operations, are invited by the Authority to be recognised as high standard operators. The High Standard Tourism program began in 2004 with 19 tourism operators. The number of

operators involved has steadily increased to 60 in 2012. During that year, these operators carried 66 per cent of tourists visiting the Reef (Figure 5.16).

Increased economic activity associated with the resources industries throughout Australia has resulted in competition in sourcing labour for the tourism industry.<sup>125</sup> This has the potential to affect the level of expertise of tourism crew and their knowledge and understanding about the Great Barrier Reef, its values and its management arrangements. It may also affect the industry's ability to source adequate capital.<sup>125</sup>



**Figure 5.16 Percentage of visitors to the Marine Park carried on high standard tourism operations, 2004–2012**

Since the Authority's High Standard Tourism program began in 2004, the increasing number of recognised high standard tourism operators has resulted in a higher proportion of tourists using these operators to visit the Reef. (Visitor numbers shown includes coral viewing and scenic flights.)

### Benefits

Tourism is the most significant direct use of the Region — both in terms of economic value and employment (see Table 7.9 in Chapter 7). In 2012, tourism in the Reef catchment and World Heritage Area generated approximately \$6.4 billion in direct expenditure, \$5.2 billion value-added (that is, profit generated) and an equivalent of more than 64,000 full time jobs.<sup>43</sup>

The tourism industry is a key partner in the Reef's protection and management. On behalf of the Authority, the industry collects about \$7 million each year from tourists through an environmental management charge (see Chapter 3).<sup>124</sup> These funds directly contribute to management of the Great Barrier Reef Marine Park.

Tourism operators are actively involved as stewards of the Reef. Through the High Standard Tourism program, operators are increasingly working to voluntary best practice standards in their activities. In addition, tourism operators are leading the way in responding to climate change by reducing and mitigating their carbon emissions. The Association of Marine Park Tourism Operators, in partnership with the Authority, has undertaken targeted control of crown-of-thorns starfish. All these actions improve the sustainability of the industry and the health and resilience of the Reef.

Many operators also participate in research and monitoring programs, such as the Eye on the Reef environmental monitoring program, the Sightings Network and visitor surveys. Despite the financial pressures experienced through the downturn in tourism, 23 tourism operators continued to actively gather reef health data in early 2012 through different components of the Eye on the Reef monitoring program, including the water quality data presented in Figure 6.15 in Chapter 6.

Importantly, the tourism industry makes much of the vast area of the Great Barrier Reef accessible to visitors; without it, many visitors simply would not be able to enjoy or experience the Region's values. The industry, therefore, plays a key role in fulfilling Australia's world heritage obligation to 'present' the Great Barrier Reef World Heritage Area. Most tourism programs involve education and interpretation activities, aimed at increasing appreciation and understanding of the natural environment and sustainable practices that support the Reef.

### Impacts

Concern about tourism impacts in the Great Barrier Reef arose during the 1980s and 1990s because of rapidly increasing tourism visitor numbers, forecasts of exponential future growth and widely reported damage to popular reefs and adjacent islands. With the introduction of more intensive management of popular areas, limits on the number of vessels and group sizes, an increase in supporting infrastructure and the adoption of best practices by operators, many of these concerns have been avoided or mitigated. As a result, the impacts associated with tourism activities today are generally regarded as low risk and are concentrated in a few intensively managed areas.<sup>22,127</sup>

Tourism can cause localised impacts through anchor damage to coral reefs and seagrass meadows; poorly supervised activities, such as diving and snorkelling; and disturbance to wildlife including whales, marine turtles and seabirds, some of which are matters of national environmental significance. These impacts have been largely reduced by regulation (for example, whale approach distances), site management arrangements (such as group size limits at locations, use of moorings, seasonal seabird closure areas); permit arrangements (for example, fish feeding guidelines); education; and the adoption of best practices for activities (such as diving and snorkelling).

In recent years, reduced profitability across the industry has increased the risks associated with tourism-related structures in the Region (such as pontoons, jetties, underwater observatories and moorings). As structures age, they require more investment in maintenance to ensure they are not a threat to the surrounding environment and do not impact on amenity and presentation values of a location. In addition, coastal development, marinas and ancillary services associated with marine tourism can result in flow-on impacts on the environment within the Region, including dredging and disposal of dredge material, clearing or modifying coastal habitats and decreased water quality.

Tourism use of the Region also has the potential to impact on or displace other users, such as commercial fishers, Traditional Owners and recreational users, particularly in high use areas. There have been some recent examples of conflict between the activities of tourism operations and those of Traditional Owners exercising their traditional hunting rights.<sup>122</sup> There are also potential conflicts between sectors within the industry, such as charter fishers and site-based tourism. In areas experiencing growth or peaks in visitation, some users also express concern about popular sites being crowded because of intensive tourism activity.<sup>128</sup>

To minimise impacts on biodiversity, sewage discharge standards for all users, including tourism operations, have been improved. Currently, discharge at sea (more than one nautical mile from any reef or island and the mainland) remains necessary for many tourism operations, as there are insufficient land-based facilities to service the fleet's pump-out requirements.

A number of compliance incidents involving the tourism industry are reported annually, particularly from the more intensively used Cairns and Whitsunday areas. Reports are typically about breaches of marine parks permits, unpermitted activity, plan of management offences (such as undertaking activities not in accordance with group and vessel size limits), issues around payment of the environmental management charge, groundings and moorings offences. Tourism-related compliance incidents have been relatively stable over recent years, with about 140 reported each year since 2009–10.<sup>129</sup>

### 5.4.3 Fishing — commercial

The Great Barrier Reef supports a diversity of fisheries including commercial, recreational, charter and Indigenous fisheries, and is a major part of Queensland's fishing activity. Fishing is the major extractive use of the Region. Recreational fishing targets many of the same species as commercial fishing and is considered separately in Section 5.4.4. The activities of charter and Indigenous fishers in the Region account for a relatively small component of the overall fisheries take.

Commercial fishing is widespread across the Great Barrier Reef (Figure 5.17). Of the major commercial fisheries in the Region, trawl, net, line and pot are the largest (Table 5.3).

The Authority has a regulatory role in the management of commercial fishing through its Zoning Plan, Regulations and permits for a limited number of fisheries. Zoning provides the framework for extractive use of the Marine Park, delineating areas available for fishing and the types of activities that can occur. Within this framework, direct fisheries management is primarily the responsibility of Fisheries Queensland within the Queensland Department of Agriculture, Fisheries and Forestry. Gear and size restrictions apply, as well as other rules such as controls on fishing effort or total allowable catch to manage catch in some commercial fisheries. The Authority has an advisory role to other agencies in relation to the management of commercial fisheries in the Region.

The East Coast Trawl Fishery is Queensland's largest and most valuable commercial fishery, with about 400 licensed vessels producing more than 7000 tonnes of seafood product from the state's east coast. About 60 per cent of the fishery is undertaken in the Region.<sup>130</sup> The trawl fleet operates in coastal, lagoon floor and upper continental slope habitats of the Great Barrier Reef. Prawns make up most of the trawl fishery's retained catch (85 per cent by weight of the targeted catch in 2010).<sup>130</sup> Like most prawn trawl fisheries, the total weight of bycatch caught in the Queensland fishery is not measured routinely. Bycatch can comprise hundreds of species, a majority of which are caught very infrequently and in low numbers.<sup>131</sup> Since the early 2000's bycatch reduction devices are used in all trawl nets. In 2002 the estimated annual weight of bycatch was 10,000 tonnes<sup>132</sup> caught in just under 68,000 fishing days. Given the number of fishing days in the fishery has fallen to under 35,000 fishing days in 2013 the estimated weight of bycatch is now estimated to be just over 5,000 tonnes. Queensland fisheries provisions are used to manage the trawl fishery (as for other fisheries in the Region), and trawling is restricted to the General Use Zone (light blue) in the Marine Park.

The commercial component of the East Coast Inshore Fin Fish Fishery is Queensland's third most valuable commercial fishery and is conducted in the tidal waters of the east coast between the northern tip of Cape York and the Queensland–New South Wales border. Large mesh netting is the predominant fishing method of this fishery, targeting a range of inshore bony fish and tropical shark species. Some species are also taken commercially by hook and line. In 2009–10, the Queensland east coast catch included shark species (453 tonnes); mullet species (1754 tonnes); whiting (339 tonnes); barramundi (234 tonnes); bream (165 tonnes); blue



threadfin (152 tonnes); school mackerel (136 tonnes) and king threadfin (135 tonnes).<sup>133</sup> Bycatch has previously been estimated at less than 20 per cent for netting operations that target mullet, whiting, small mackerels, barramundi and mixed estuary species.<sup>134</sup> However, the need to continue to reduce the bycatch of species of conservation concern such as sawfish, dugongs, marine turtles and inshore dolphins (most being matters of national environmental significance) is recognised, and is a high priority for industry and management. Changes since 2009 in management of the fishery have included new net fishing arrangements, such as improved attendance requirements, changes to managing the take of sharks, and new and amended possession and size limits.

More than 125 bony fish species are caught by line from reef habitats in the Great Barrier Reef, but only a few of them are specifically targeted by the quota-managed commercial sector of the Coral Reef Fin Fish Fishery.<sup>135</sup> These include coral trout species for the international live food fish trade (commercial catch of 797 tonnes in 2010–11), red throat emperor (commercial catch of 256 tonnes in 2010–11), and a range of other reef fish such as red emperor and several cod species (commercial catch of all other reef fish species of 426 tonnes in 2010–11).<sup>135</sup> Other reef fish species not targeted commercially can be caught incidentally, and are either retained (by-product) or discarded.

Reef sharks may be caught incidentally while fishing for reef fish and are generally discarded.<sup>136</sup> A number of species of sharks may also be targeted for their fins; however, in the Region there are landing requirements that the carcasses must be retained with their fins to ensure full use of the shark. Seven fish species are prohibited from commercial take for conservation reasons, iconic values or ciguatera (poisoning) concerns, including humphead Maori wrasse and Queensland groper. In accordance with the Great Barrier Reef Marine Park Regulations 1983, all groupers over one metre in length are protected from fishing in the Marine Park.

The commercial fishery for Spanish mackerel uses line gear and is a highly selective fishery for this, the largest mackerel in Queensland. Spanish mackerel are known to aggregate in large numbers to spawn at particular reefs in the Region. The main commercial catch is from these reefs, between Ingham and Bowen. A quota system caps the commercial catch from Queensland's east coast at 544 tonnes per year.

Other commercial fisheries that use line gear and partly occur in the Region are the Rocky Reef Fin Fish Fishery (this predominantly targets snapper) and the Deepwater Fin Fish Fishery.

### Trends


Global fisheries trends influence those for the Great Barrier Reef. As wild-caught fisheries throughout the world continue to be fully exploited or over exploited,<sup>137</sup> the economic value of the Region's fisheries resources and the pressure to exploit them (legally and illegally) may increase.<sup>22</sup> International demand for wild-caught Queensland seafood may increase the targeting of additional species or increase demand for intensive aquaculture within the Region and its catchment.<sup>22</sup> Expected growth in aquaculture around the world (for example, aquaculture-raised coral trout are expected to be commercially viable in the near future<sup>138</sup>) may also lead to diversification in the reef line fishery and increased targeting of other species.

The commercial Coral Reef Fin Fish Fishery has reported significantly reduced catch rates of coral trout in areas affected by cyclones such as Hamish in March 2009 and Yasi in early 2011, which can have flow-on effects to other areas. For example, after cyclone Hamish in 2009, some active commercial live coral trout fishers moved northward to escape cyclone impacted areas where lower catch rates were being experienced.<sup>28</sup>

Fishing effort in the Region's otter trawl fishery has been relatively constant for the past five years, at about 20,000 boat days.<sup>130</sup> The peak trawl fishing effort was in 1997 (more than 73,400 boat days) (Figure 5.18). Total fishing effort is nominally limited by a cap which is distributed among licensed operators in tradeable effort units. There is, however, a significant amount of latent effort in the fishery, with approximately 40 per cent of the available total effort units not used in 2010.<sup>130</sup> This means trawl fishing effort could rise substantially under current management arrangements in response to more favourable economic conditions, such as lower foreign exchange rates.

Some fisheries (such as trawl) already catch a substantial amount of their product in the far north of the Region (Figure 5.17), and have done so for many years. Fishing effort in other fisheries may spread northwards to take advantage of catch availability and improving market access (due to improved infrastructure), or to offset other factors such as impacts of extreme weather and recreational uses displacing some commercial fishers close to urban areas.

Economic factors influence patterns of commercial fishing operations. For example, recent increases in fuel prices<sup>140</sup> have affected profitability and, in some cases, methods and areas of operation of Great Barrier Reef fisheries. The strength of the Australian dollar has also put pressure on commercial fisheries. In this economic environment, exported product is less profitable and there is increasing competition in the local market from cheap imports. Fishing business owners also have to compete for crew with the resources sector.



**Figure 5.17 Spatial distribution of catch by commercial line, net, pot and trawl fisheries, Great Barrier Reef Region, 2011<sup>139</sup>**


The amount of fisheries product taken from different areas in the Great Barrier Reef varies for each of the major fisheries. Net and pot fisheries are undertaken close to the coast, whereas line fishing and trawling extend further offshore. The tonnages shown are for each fisheries 'grid' (a 30 nautical mile square), excluding the zones closed to fishing. Note: commercial bait netting data includes Conservation Park Zones (other netting is restricted in these areas).

Some fisheries (such as trawl) already catch a substantial amount of their product in the far north of the Region (Figure 5.17), and have done so for many years. Fishing effort in other fisheries may spread northwards to take advantage of catch availability and improving market access (due to improved infrastructure), or to offset other factors such as impacts of extreme weather and recreational uses displacing some commercial fishers close to urban areas.

Economic factors influence patterns of commercial fishing operations. For example, recent increases in fuel prices<sup>140</sup> have affected profitability and in some cases methods and areas of operation of Great Barrier Reef fisheries. The strength of the Australian dollar has also put pressure on commercial fisheries. In this economic environment, exported product is less profitable and there is increasing competition in the local market from cheap imports. Fishing business owners also have to compete for crew with the resources sector.

**Table 5.3 Commercial fisheries of the Great Barrier Reef Region, 2011**

Type of fishery	Fishery	No. of active primary licences (Great Barrier Reef component)	Estimated proportion of Qld fishery within Great Barrier Reef	Retained commercial catch in the Great Barrier Reef, 2011	Main target species
<b>Otter trawl on or near seabed</b>	Trawl (otter)	196	~60%	3602 tonnes	prawns, scallops, Moreton Bay bugs, squids
<b>Net (mainly large mesh net)</b>	East Coast Inshore Fin Fish (net component)	224	~40%	1787 tonnes	barramundi, sharks, grey mackerel, threadfin salmon, bait species
<b>Line</b>	Mainly the Coral Reef Fin Fish, Spanish Mackerel, and East Coast Inshore Fin Fish (line component)	270	Reef~95%, Inshore~50%	1741 tonnes	Reef: coral trouts, cods, emperors and snappers Spanish mackerel inshore: barramundi, sharks, mackerels, cobia, pelagics
<b>Pot</b>	Mud crab, blue swimmer crab and spanner crab	212	Mud and blue swimmer~65%; Spanner <20%	1012 tonnes	mud crab, blue swimmer crab, spanner crab and three-spot crab
<b>Collection</b>	Coral	25	99%	~68 tonnes	'live rock' and potentially hundreds of species including hard and soft corals
	Marine aquarium fish	27	~60%	~78,000 fish, mostly juveniles	potentially hundreds of species but mostly damselfish and anemone fish, wrasses and angelfish
	Trochus	-	100%	no reported catch	large herbivorous gastropod
	Tropical rock lobster	7	100%	141 tonnes	ornate rock lobster
	Sea cucumber (beche-de-mer)	6	100%	387 tonnes	sea cucumbers (mainly white teatfish and burrowing blackfish)



**Figure 5.18 Annual trawl fishing effort in the Great Barrier Reef Region, 1990–2011<sup>130</sup>**

Data is based on fisher logbook records.

### Benefits

Fisheries product from the Region is important to local communities, as well as domestic and international markets. It is a major component of the total seafood catch from the Queensland east coast, representing about 95 per cent of the reef line fishery, 60 per cent of the trawl fishery and 40 per cent of the net fishery.<sup>22</sup> In a 2008 survey of catchment residents, 92 per cent had eaten seafood in the previous 12 months. The vast majority of those who bought seafood preferred to buy fresh seafood caught in Queensland.<sup>141</sup>

Commercial fishing is a multi-million dollar industry in the Region. In 2011–12, the economic contribution of the commercial fishing industry and aquaculture to the national economy was estimated to be \$160.3 million.<sup>43</sup> This value-added figure is the gross value of production minus tax on production and other factors such as operating costs and labour. Separate recent valuations of the major fishery sectors (calculated on the price paid to fishers at the first point of sale), reported the value of the East Coast Inshore Fin Fish Fishery to be about \$19.6 million a year<sup>133</sup>; the reef line fishery at about \$31 million a year<sup>142</sup>, and the trawl fishery at about \$110 million each year.<sup>143</sup> In 2012, commercial fishing in the Region was estimated to have generated the equivalent of 975 full-time jobs<sup>43</sup>, although data on employment is limited and estimates for secondary employment are lacking.

Fishers have a high attachment to their industry, and most have been involved in the industry for a long time (more than 20 years).<sup>34,144</sup> They are also generally very reliant on the industry, with most fishers receiving more than 75 per cent of their household income from fishing.<sup>34,145</sup> Fishing is more than just a job for most fishers. In a 2008 survey of inshore fishers in the Region, 63 per cent preferred to be at sea than on the land; 98 per cent stated they liked being a fisher; 93 per cent stated fishing is a lifestyle, not just their job; and 79 per cent felt proud to tell others they were a commercial fisher. Seventy per cent also stated they would still be fishing in three years.<sup>144</sup>

Although commercial fishers pay annual fees to the Queensland Government for their licence, commercial fisheries in Queensland are not managed on a cost recovery basis.

### Impacts

Commercial fishing is the largest extractive use of the Region. It also has other direct impacts such as the death of discarded catch and physical impacts on habitats. The total commercial catch retained by line, trawl, net and pot fisheries from the Great Barrier Reef Region was about 8100 tonnes in 2011, using more than 60,800 days of commercial fishing effort.<sup>139</sup>

Commercial fishing effort is not uniformly distributed in the Region (Figure 5.17). The areas fished depend on many factors including zoning, licence conditions, distance from shore, weather, access to distribution networks, distribution of the target species and fishers' preferences. This can result in a concentration of fishing effort and associated impacts in popular areas open to fishing and close to the coast.

While all commercial fisheries have made significant progress towards achieving better sustainability outcomes as a result of improved management arrangements, illegal fishing activities continue to undermine these outcomes. Based on knowledge gained through compliance and enforcement activities and on intelligence received, illegal commercial fishing within the Great Barrier Reef Marine Park remains a serious concern. Priority areas for enforcement include non-compliance with zoning requirements within the Coral Reef Fin Fish Fishery, and with net attendance requirements in the East Coast Inshore Fin Fish Fishery. Reliable information from individuals who work within the commercial fishing industry indicates illegal activity in the reef line fishery continues to be significant despite the recent low number of offences detected. A number of skippers and dory operators in the commercial Coral Reef Fin Fish Fishery continue to be non-compliant, even after receiving court-imposed fines — many of which remain unpaid. Those fishers who are intent on illegal activity continually modify their behaviour and employ surveillance avoidance tactics.<sup>129</sup>



There are particular concerns about non-compliance in the commercial netting special management areas which are critical to species of conservation concern and matters of national environmental significance, such as dugongs and the Australian snubfin dolphin. The remoteness of some of these locations makes enforcement particularly challenging; for example, in the Princess Charlotte Bay Special Management Area which is an area with significant populations of dugong and inshore dolphins — both species of conservation concern and matters of national environmental significance.<sup>146,147</sup> Management issues in relation to Princess Charlotte Bay are discussed in the relevant demonstration case study (see Section 9.6).

The stock status of a number of key targeted species, including grey mackerel and king threadfin, are classified as uncertain or undefined, while snapper are classified as overfished in Queensland.<sup>148</sup> There are also community and/or scientific concerns about the population status of some targeted species, including some mackerels<sup>149,150</sup>, snapper<sup>151</sup> and coral trout<sup>152</sup>. Between 1989 and 2003, some 300 to 620 tonnes of coral trout were estimated to have been discarded annually by the commercial Coral Reef Fin Fish Fishery in the Region.<sup>153</sup> Though the ecological significance of this level of discard is unknown, declines in some Reef predatory species, including coral trout, suggests predation in reef-related food webs may have altered significantly.<sup>22</sup>

The paucity of information about the stock status of sharks in the East Coast Inshore Fin Fish Fishery is of particular concern as the fishery takes a wide range of shark species, each with a different productivity and susceptibility to fishing.<sup>154</sup> Collection of basic information about the take of different shark species in this fishery has improved over the past five years, and many of the species taken are not considered at high risk from fishing, though some still are, such as the scalloped hammerhead. Information on interactions with rarer species (such as bignose and mako sharks) is difficult to obtain, and the population status of these species is largely unknown.

Concerns about overcapacity in some Reef-related fisheries<sup>155,156</sup>, and the assessment that some species are overfished or that their stock status is uncertain, indicates that there remains some ecological risks and unsustainable impacts on some retained species.

Management changes have led to a significant reduction in the ecological impacts of the trawl fishery (see Section 9.10); however, some concerns remain, particularly for 11 species of rays and skates, two species of Balmain bugs, two species of sea snakes and a poorly known upper continental slope habitat in the south-east of the Region.<sup>132</sup> The amount of effort is the main determinant of impacts on the Great Barrier Reef ecosystem from the trawl fishery, and risks could increase if effort levels rise.<sup>132</sup>

The main impacts on the Region's values from commercial net fishing are entanglement of species of conservation concern and bycatch.<sup>157</sup> The fishery is known to interact with dugongs, dolphins (particularly inshore dolphins), whales, marine turtles, crocodiles and green and freshwater sawfish — all of which are matters of national environmental significance.

Any death of a species with depleted populations, such as dugong, Australian snubfin dolphin and sawfish, are of particular concern. Commercial fishers are required to report interactions with such species — records of marine wildlife strandings in Queensland confirm fishing interactions continue to be a cause of death of marine wildlife in the Region.<sup>158</sup> Under-reporting by some commercial fishers of interactions is a recognised issue.<sup>156</sup> Discards of fishery species and most bycatch are not required to be reported. While there are a number of research projects providing important fisheries information<sup>154,159</sup>, there is currently no independent fishery observer program in the Region to complement the research.

Information to develop harvest strategies for the sustainable use of many target species is limited. The *Privacy Act 1988* limits the disclosure and use of some commercial fisher logbook data on catch and effort. This makes it difficult to develop an accurate representation of all fishing in the Region, particularly fine-scale patterns of use, and to assess cumulative impacts of all activities in a local area.

Several major research programs and studies have improved our understanding of the environmental effects of fishing, including on the broader ecosystem.<sup>160,161</sup> This information has informed reviews of management arrangements for commercial fishing activities.

#### 5.4.4 Fishing — recreational

Recreational fishing is one of the most popular activities on the Great Barrier Reef. Statewide, an estimated 703,000 residents went fishing in the 12 months prior to June 2010.<sup>162</sup> Residents living adjacent to the Reef fished for a total of approximately 2.6 million days in the 12 months between October 2010 and September 2011. Approximately 100,000 of these fisher days were concentrated in Mackay coastal waters.<sup>162</sup> Recreational fishing is also popular with visitors in areas adjacent to the Region; however, little is known about their activities.

In Queensland, the most common recreational fishing method (80 per cent) is line fishing (including the use of hooks and lures), followed by fishing with pots (13 per cent).<sup>162</sup> Together, fishing with cast nets, hand collection, pumps and spades, diving using spears, and hand collection comprised only seven per cent of all fishing effort. In the Great Barrier Reef Region, coral trout, red throat emperor, tropical snapper, morwong and sweetlip are commonly targeted by recreational fishers.<sup>162</sup> Hundreds of other species of bony fish, sharks and other animals are also caught.<sup>157,162</sup>


## Trends

Queensland-wide, the number of people fishing fell between 2000 and 2010. However, numbers remained about the same for those areas where residents are likely to be accessing the Region; that is, the Fitzroy, Mackay, Northern and Far North<sup>162</sup> (Figure 5.19).

The popularity of fishing in the Region is reflected in participation rates. Information collected during a 2010 statewide recreational fishing survey<sup>162</sup>, showed 28 per cent of residents who have lived for five years or more in the Mackay region (including Bowen, Mackay, Proserpine and the Whitsunday islands) fished in the 12 months prior to the survey. This was the highest participation rate for any region in Queensland, and much higher than the statewide average of 17 per cent. Other regions adjacent to the Great Barrier Reef also had relatively high participation rates of 20 to 23 per cent. Most of the fishing was in local marine waters from boats.

The 2010 survey indicated that people are catching fewer fish for a similar level of effort compared to a decade ago.<sup>162</sup> This may reflect lower abundances of some target species after many decades of fishing pressure.

A typical fishing trip is less than 20 kilometres, and often involves boating to a location and staying there to fish. However, some fishing trips cover hundreds of kilometres and last for several days.<sup>163</sup> A trend towards larger, more powerful, more fuel-efficient vessels,<sup>163</sup> combined with improvements in safety, mean recreational fishers may be likely to fish further from the mainland. Recreational fishing in the remote far north of the Region is increasing, due to factors such as improved road access, catch availability and infrastructure improvements. In addition, improved navigation and positioning systems make it more likely that fishers will be able to relocate specific fishing sites and increase their effectiveness at catching fish.



**Figure 5.19 Number of recreational fishers by Queensland residential region in 2000 and 2010<sup>162</sup>**

A recreational fisher is defined as anyone who fished once or more in the 12 months prior to April 2000 or June 2010.

## Benefits

Recreational fishing provides economic and social benefits to Queensland's communities.<sup>164</sup> For generations, fishing on the Great Barrier Reef has been an important recreational activity for coastal residents and visitors. Most people indicate that relaxation, socialising, catching fish and excitement are their reasons for fishing recreationally.<sup>164</sup> In 2008, more than half of all people visiting the Great Barrier Reef for recreational purposes went fishing.<sup>22</sup>

Economic contributions from recreational fishing are difficult to estimate. In 2012, all recreational activities (including fishing) were estimated to generate \$330 million, and the employment equivalent of 2724 full-time jobs (see Table 7.9 in Chapter 7).<sup>43</sup> It is likely that a significant portion of this economic activity is related to recreational fishing.

Many recreational fishers have contributed to research, monitoring and management programs (for example, the CapReef program in the Central Queensland area).<sup>165</sup> Recreational fishers also contribute to caring for Australia's waterways and fish stocks, including those in the Great Barrier Reef Region.<sup>166</sup> There can be considerable health and wellbeing benefits for individuals who fish for recreation.<sup>167</sup> There is also some contribution to management costs, with \$17.75 from each vessel registration in Queensland going to Fisheries Queensland for enhancing recreational fishing.<sup>168</sup>


## Impacts

Recreational fishing is a significant extractive use of the Region, removing marine animals from various parts of the food chain. In 2010–11, an estimated 13.3 million fish were caught by recreational fishers who reside in Queensland, plus 8.3 million animals of other species including crustaceans (such as crabs, prawns and lobster) and cephalopods (for example, squid and octopus).<sup>162</sup> A substantial, but as yet unquantified, portion of these total catches occurred in the Region and, for some species, most of the catch was taken there. The main target

species are coral trout and cod, emperor, tropical snapper, barramundi, bream, mackerel, whiting, crab, lobster and bait fish.<sup>22</sup> While the impacts resulting from each recreational fishing visit are relatively minor, the large number of participants is likely to contribute to a cumulative effect on Great Barrier Reef values. Technological improvements (for example, widespread use of positioning systems and sounders) have also allowed more efficient and targeted fishing by recreational fishers. Further, the species targeted by recreational fishers are mostly the same as those targeted by commercial fishers, increasing cumulative impacts on those species. Recreational fishery harvest levels have contributed to concerns about the population status of some species. Recreational fishing activities contribute to other impacts on values such as the death of discarded species, including some species of conservation concern. This includes drowning of marine turtles in poorly designed crab pots, bycatch of sharks and rays and incidental catch of seabirds. The recreational catch of sharks is highest in the northern part of the Region.<sup>162</sup> A large proportion of the recreational catch is subsequently released (for example, 60 per cent for inshore fishing), either through the increasingly popular practice of 'catch and release' or because they are undersized or unwanted species.<sup>58</sup> High release rates have been reported for snapper, barramundi, stripey snapper, Australian bass, sharks, and cod and grouper; the highest rates are for sharks with 94 per cent released.<sup>162</sup> Survival rates of the released species vary.<sup>169</sup> The ecosystem effects and cumulative impacts of recreational fishing are poorly understood, but are likely to be most concentrated in inshore areas close to major population centres and a relatively small number of offshore reefs. Local depletion, particularly of some inshore species, is of concern in some areas.<sup>22</sup> Increasing numbers of seasonal recreational fishers may increase the pressure on specific species and locations, but these impacts are also largely unquantified.

With regard to physical damage to the environment, the most likely impact resulting from recreational fishing is localised anchor damage at popular fishing grounds. This damage is likely to be more significant on fragile fringing reefs or nearshore islands. Fishing gear may also damage fragile species such as corals. Discarded fishing gear (from all fishery sectors) contributes to marine debris in the Region, and can entangle or kill marine animals<sup>170,171</sup>, including species of conservation concern.

Illegal recreational fishing continues to be of concern — activities in no-take zones are a priority for enforcement. Illegal fishing has the potential to impact on the Region's values, including eroding the resilience benefits of the network of no-take marine reserves.<sup>172</sup> Recreational fishing in protected zones continues to be the most frequent type of offence, particularly in those areas close to large population centres. Detected recreational fishing offences across all zones are increasing (Figure 5.20). Incident information shows most recreational fishers detected illegally fishing are resident in communities adjacent to the Marine Park, and that there are some fishers targeting no take (green) zones at night.<sup>129</sup>




**Figure 5.20 Recreational fishing compliance incidents by marine park zone, 2008–2012**

Each incident may include multiple offenders and/or multiple offences.

### 5.4.5 Recreation

Recreation is defined as an independent visit to the Region for enjoyment that is not part of a commercial operation. It is distinct from tourism, where a visitor pays to use a commercial operation (Section 5.4.2).


People living adjacent to the Region, as well as domestic and international visitors, use the Region for a wide range of recreational activities, including swimming, motorised boating and fishing, snorkelling, sailing, diving and jetskiing (Figure 5.21). The fisheries aspects of recreation are addressed in Section 5.4.4.



**Figure 5.21 Main activities of visitors to the Great Barrier Reef<sup>58</sup>**

A study in 2008 estimated there were more than 14 million recreational visits to the Marine Park by catchment residents every year.<sup>173</sup> This does not include visits by independent travellers from outside the catchment (such as retired, long-term holiday makers ('grey nomads') and cruising yachters).<sup>58</sup>


Because most recreational vessels are less than five metres in length, recreational use is concentrated in inshore areas close to urban centres. The most popular destinations are islands, followed by reefs, shoals, cays and wrecks (Figure 5.22). In 2008, almost half the vessel-based trips departed from the area between and including Cairns and Townsville.<sup>173</sup> Cairns, Townsville and Mackay have the highest numbers of boat registrations in Queensland (Figure 5.23).



**Figure 5.22 Main destinations of recreational vessel-based trips, 2003 and 2008<sup>173</sup>**

Most people who visit the Region for recreation go fishing or visit islands, reefs, shoals, rocks or wrecks.





**Figure 5.23 Vessel registrations in the Great Barrier Reef catchment, 2011**

The number of vessel registrations is shown by local government area.

## Trends

The number of recreational visits from residents in the catchment appears to have risen substantially in recent years, most likely as a result of three factors: population growth; an increase in the proportion of the population visiting the Region; and a rise in the average number of visits each person makes.<sup>173</sup>

The average number of visits by residents from the Great Barrier Reef catchment increased from 14 visits per person per year in 2003 to 15.5 in 2008. Half of those who visit from the catchment do so more than seven times in a year, while 17 per cent visit more than once a week.<sup>173</sup> Increasing recreational vessel ownership in the catchment (Figure 5.24) is likely to be contributing to the growing proportion of residents visiting the Region.

The relative popularity of different recreational activities has changed in recent years. Motorised boating has become more popular, and the proportional use of sailing boats has fallen (Figure 5.21).<sup>173</sup> Activities like shell collecting and reef walking are now far less popular than in the early twentieth century.<sup>174</sup> Between 2003 and 2008, the number of visits to destinations such as reefs, shoals and wrecks increased, while visits to islands declined (Figure 5.22).<sup>173</sup> Visiting creeks, inlets and rivers also became more popular. These trends are likely to continue.




Figure 5.24 Number of recreational vessels registered in the Great Barrier Reef catchment, 2001–2011<sup>57</sup>

## Benefits

The opportunity to enjoy the Region's environment is of enormous social and cultural value to Queensland residents, other Australians and international visitors.<sup>58</sup> The Great Barrier Reef is viewed as beautiful, pristine and unique by those who have already visited the Reef and those who are yet to visit.<sup>22</sup> In 2012, recreational activities (including fishing) was the second largest direct use of the World Heritage Area, generating \$330 million and the employment equivalent of 2724 full-time jobs (see Table 7.9 in Chapter 7).<sup>43</sup> Most of the contribution associated with recreational activity was derived from expenditure on equipment, largely reflecting the purchase of boats and maintenance or repair of recreational equipment.

Recreational users contribute to protection and management of the Marine Park through programs such as Clean up Australia Day, Order of Underwater Coral Heroes Volunteers (OUCH, a non-profit organisation based in the Whitsundays), the Strandings Hotline and the Sightings Network.

## Impacts

The Great Barrier Reef Outlook Report 2009 lists the likely impacts of recreational use as *'localised but frequent anchor damage to corals and seagrass meadows.....along with littering, boat strikes on marine mammals and turtles, and damage to corals when snorkelling and diving. There is some risk of introducing species through fouling on recreational vessels, especially those from overseas'*.<sup>22</sup> There are also potential impacts from the grounding of small vessels, oil and chemical spills, vessel sewage discharge and disturbance of wildlife.<sup>58</sup> Recreational use on islands can result in disturbance of seabirds and other wildlife, trampling of coastal vegetation and the introduction of weeds and feral animals.<sup>58</sup> Given the concentration of recreational use near the coast and regional centres, it follows that the main impacts are also likely to be in those areas.

There is the potential for Indigenous and historic heritage values to be affected by recreational use.<sup>58</sup> For example, Traditional Owner connections to their sea country, including access to resources, are being interrupted at least partly because of increasing recreational use.<sup>58</sup> Previously, some historic shipwrecks have been disturbed and artefacts removed, particularly during the 1950s to the 1970s.

While management arrangements have significantly reduced the likelihood of conflicts of use between recreational and other users in popular areas, there continues to be some localised negative interactions.

With increasing use of the Marine Park comes an increasing demand for coastal infrastructure to access the area (for example, roads, marinas and boat ramps). Construction and operation of these facilities can affect the Great

Barrier Reef ecosystem through damage to coastal habitats, dredging, disposal and resuspension of dredge material, reclamation and adverse effects on water quality. These facilities can also result in changes to patterns and intensity of use within the Region.


Queensland's Shark Control Program, established in 1962, aims to reduce the number of potentially dangerous sharks at popular swimming beaches and, thereby, lower the likelihood of shark bites. A combination of 338 drumlines and 37 mesh nets is currently deployed across 84 beaches in Queensland.<sup>175,176</sup> Thirty of these beaches are within the World Heritage Area, located at: Cairns (seven beaches); Townsville and Magnetic Island (eight beaches); Mackay (five beaches); Capricorn Coast (nine beaches); and Gladstone (one beach). In addition to catching targeted shark species, nets and drumlines catch non-target species, including other shark and rays species, marine turtles, dugongs, whales and dolphins — some of which are matters of national environmental significance. Drumlines tend to catch more loggerhead turtles. Nets tend to catch more marine mammals and green turtles. The effects of these nets on dugongs are described in the dugong demonstration case study (see Section 9.3) and on the snubfin dolphin in Section 7.1.5.

Compliance incidents associated with recreational activities, other than fishing, include vessel groundings and sinkings, pollution discharges, entering restricted areas or zones, approaching whales too closely and island national park offences such as camping without a permit, lighting campfires, taking domestic animals ashore and using generators.<sup>129</sup> The annual number of reported offences (excluding recreational fishers) has steadily declined over recent years from 130 in 2009–10 to 82 in 2011–12.<sup>177</sup>

#### 5.4.6 Shipping

Thousands of domestic and international ships transit the Region every year, carrying export goods, servicing coastal and inland communities and transporting passengers. Shipping, as described in this report, includes vessels greater than 50 metres in overall length, or carrying specialised product regardless of length (for example, oil tankers, chemical or liquefied gas carriers). It includes cruise ships and superyachts.

Shipping cargo includes coal, sugar, iron ore, timber, oil, chemicals, live cattle and general cargo. The four busiest ports in relation to commercial vessel visits in 2011–12 were the ports of Gladstone (1453 visits), Hay Point (809 visits), Townsville (747 visits) and Cairns (720 visits) (Figure 5.25). Although Cairns is one of the busiest ports on the Great Barrier Reef coast, it is not considered a major port due to the relatively small throughput (Section 5.3.5). Gladstone, Hay Point, Townsville and Abbot Point are considered the major ports due to their relatively high annual throughput.




**Figure 5.25 Total commercial vessel calls to ports adjacent to the Region, 2011–12<sup>109</sup>**

The graph represents total commercial vessel visits, excluding those made by cruise ships. Note: A vessel or ship 'call' relates to an arrival at port; whereas 'movements' include all reportable movements within the Great Barrier Reef, including movements on the way to and within the port.

The Region is one of the world's most regulated shipping areas. The Authority, together with the Australian Maritime Safety Authority and Maritime Safety Queensland, works closely to protect the marine environment from the potential adverse consequences of shipping operations. The Reef was designated a Particularly Sensitive Sea Area by the International Maritime Organization in 1990, and shipping traffic is confined to a Designated Shipping Area (Figure 5.26) unless otherwise permitted by the Authority.

Within the Designated Shipping Area, anchoring a ship does not require a permit from the Authority. Maritime Safety Queensland has designated 154 ship anchorages adjacent to some of the ports along the Region's coast (Figure 5.26). All but 12 are within the Marine Park. Including swing room, the anchorages cover about 1200 square kilometres. They confine the impacts arising from anchoring for these ports, such as disturbance to wildlife, physical damage to benthos, light pollution, waste discharge from a vessel, exotic species and diseases, noise pollution, user conflict and aesthetic impacts on the seascape.



**Figure 5.26 Designated Shipping Area and designated ship anchorages, Great Barrier Reef Region**

Over 150 ship anchorages have been designated by Maritime Safety Queensland adjacent to some of the ports along the Great Barrier Reef coast.



Measures to increase navigational safety and reduce the risk of ship groundings and collisions have included compulsory pilotage, recommended pilotage, an automatic identification system and mandatory vessel reporting and monitoring. These management arrangements mean there have been few shipping incidents relative to the large number of ship movements in the Region.<sup>22</sup>

In response to the *Shen Neng I* incident in 2010, when a large coal carrier grounded on a reef, the capabilities of the Great Barrier Reef and Torres Strait vessel traffic service were extended to the southern boundary of the Region.

### Trends

The number of ship voyages undertaken through the Region has increased substantially over the past 10 years (Figure 5.27 and Figure 5.28), driven mainly by industrial and mining activity. In 2011–12, Ports Australia estimated that more than 4300 ships called at Great Barrier Reef ports.<sup>109</sup>

In recent years, there have been various forecasts of shipping activity in and adjacent to the Region.<sup>46,178,179</sup> Based on projected export capacities, information from existing development proposals and predictions for the Region's four major ports, shipping in the Region is forecast to significantly increase over the next 10 to 15 years (Figure 5.29). This is likely to be driven by growth in the mining and liquefied natural gas industry, port expansions and increases in trade.<sup>47</sup>

The dynamic nature of shipping and the industries it services means there is likely to be continual changes in the types of bulk and packaged cargo carried through the Region. This is likely to include a greater range of substances that are noxious or hazardous to the environment. There is also a global trend towards longer, deeper draft ships.

The draft North-East Shipping Management Plan<sup>180</sup>, released in August 2013, examines shipping-related risks in the Great Barrier Reef, the Torres Strait and the Coral Sea (within Australia's Exclusive Economic Zone) and identifies protective measures to address them. It is a collaborative plan developed by the Australian Maritime Safety Authority in cooperation with other responsible government agencies, including the Authority.

Cruise ships comprise a minor component of the commercial vessels transiting the Region. In 2011–12, cruise ships represented only 1.5 per cent of the commercial vessel calls to Great Barrier Reef ports.<sup>109</sup> The cruise shipping industry is predicted to grow over coming decades.<sup>181</sup>

### Benefits

Ships that transit the waters of the Great Barrier Reef provide a service to communities adjacent to the Region, transporting export and import cargo as well as cruise ship passengers. An estimated \$38 billion of Australia's export trade is carried annually through the Region.<sup>182</sup> The economic activity generated by this shipping traffic provides a range of social and economic benefits to catchment communities and beyond.

### Impacts

There are many factors that increase the likelihood of impacts as a result of shipping activity in the Region. These include:

- limited (but improving) navigational facilities
- extensive areas of shallow water
- strong trade winds, cyclones and complex tidal streams
- reduced visibility during storm conditions
- some narrow shipping lanes.<sup>111</sup>

To date, the impacts of shipping have mainly related to: grounding of large vessels with the associated physical damage and pollution from toxic antifoulant paint; small chemical spills; large and small oil spills; increased noise; vessel strikes on wildlife; vessel-based waste discharge; the introduction of exotic marine species; and marine debris.<sup>47</sup> The increasing prevalence of ships anchoring off ports is likely to increase disturbance to the seabed, visual disturbance, potential for pest introduction, light pollution, interference with species' behaviour, and the displacement of others such as recreational users and commercial fishers.<sup>179</sup>

There is emerging evidence of additional impacts from ship operations worldwide, for example, the leaching of biocides from ships' antifouling coatings, vessel-animal interactions including collisions, and loss of 'communication space' for marine animals as a result of vessel noise.<sup>183</sup>

As a result of the comprehensive management arrangements in place, incidents involving ships are relatively infrequent, with annual totals of 10 or fewer in recent years.<sup>22</sup> However, given the projected increases in shipping activity it is critical that management arrangements keep pace with growth in the industry. Of major concern is the potential for ship groundings, major oil or chemical spills and the introduction of invasive marine species.<sup>184</sup> While the likely frequency of serious incidents is very low, the location and nature of an incident could mean that its environmental effects are very high, causing long-term damage to the Region's environment on a local and regional scale, and the social and economic benefits derived from it.




Figure 5.27 Polled ship positions, Great Barrier Reef Region, 2000<sup>185</sup>

### Proactive and collaborative planning for increased shipping

Driven primarily by Queensland's industrial and mining activity, shipping in the Region is projected to increase by about 150 per cent by 2032, resulting in approximately 10,097 commercial vessels calling at Great Barrier Reef ports<sup>178</sup> (Figure 5.27). Australia is currently the world's largest exporter of black coal, and Queensland supplies more than 75 per cent of Australia's metallurgical coal resources.<sup>46</sup> During the first decade of this century, Australian coal exports increased from 195 million tonnes in 2000–01 to 284 million tonnes in 2010–11.<sup>46</sup> Continued growth in shipping activity through the Region presents an increased risk of incidents, including ship-sourced pollution and damage from groundings.<sup>180</sup>

In anticipation of the increased risk, government agencies with jurisdiction over shipping activities in the north-eastern waters of Australia have developed the draft North–East Shipping Management Plan.<sup>180</sup> The draft plan applies to the Great Barrier Reef, the Torres Strait and the Coral Sea (within Australia's Exclusive Economic Zone). It examines known, potential and cumulative shipping-related impacts such as collisions, groundings, release of air emissions and other pollutants, marine pest introduction, wildlife disturbance, altered aesthetic value, and wildlife collisions, as well as impacts on Indigenous, cultural and social values.

The draft plan demonstrates a proactive approach to addressing the projected increased risks from shipping, outlining the safety and environmental standards that need maintaining or improving. It identifies a number of measures to reduce risks from shipping, including:

- additional areas for consideration of **pilotage requirements**





Figure 5.28 Polled ship positions, Great Barrier Reef Region, 2010<sup>186</sup>

- increased resources for **port state control inspections** to ensure all ships meet the applicable standards
- increased focus on areas related to **navigational risk** (such as fatigue, passage planning and navigational equipment)
- **additional protection for the Coral Sea** through seeking categorisation as a Particularly Sensitive Sea Area and implementing associated protective measures such as ship reporting and routing requirements
- using emerging **ship tracking technology** to improve awareness of shipping movements in the Coral Sea and to provide early alerting of ship breakdowns
- more active management of traffic such as a '**traffic organisation service**' to minimise collision risk, particularly in confined waters such as Hydrographers Passage
- working with industry to introduce the need for trading ships to be equipped with **Electronic Chart Display and Information Systems**, an alternative to paper nautical charts which assist with navigation through ship position monitoring and alarms alerting navigation officers when a ship exceeds pre-defined safety boundaries
- working with industry to introduce the need for trading ships travelling through protective locations to be equipped with **bunker oil tanks** which reduce the risk of an oil spill.

The current Great Barrier Reef and Torres Strait vessel traffic service is considered effective at preventing shipping incidents — almost 11,000 ship movements were monitored in 2011–12, without significant incident.<sup>180</sup>



**Figure 5.29 A projection of shipping increases at major ports adjacent to the Great Barrier Reef Region, 2012–2032<sup>178</sup>**

The category 'all commercial vessels' includes coal carriers, bulk carriers, container carriers, vehicle carriers, general cargo ships, tankers and cruise ships. Fishing, other tourism and recreational vessels are not included. All commercial vessel projections are based on 4.8 per cent annual growth. Coal vessel projections are based on 7.2 per cent annual growth. It is recognised that these projections may be higher than the eventual shipping traffic, depending on variables such as economic conditions.

#### 5.4.7 Defence activities

The Australian Defence Force has operated and trained in the Great Barrier Reef Region for more than 100 years, and the Region is predicted to increase in importance as a defence training area over the next 25 years.<sup>187</sup> The Authority is notified of all defence activities in the Region and can issue directions to the Australian Defence Force to minimise risks to the Great Barrier Reef.

Operational activities in the Region include ocean surveillance, maritime search and rescue missions, and hydrographic survey and charting. Defence forces also provide critical support for border protection activities such as environment and fisheries protection, immigration controls, and biosecurity. Australian Navy, Army and Air Force bases at Cairns and Townsville serve as the key platforms for defence operational activities in the Region.

Training activities are regularly undertaken in designated areas of the Region, covering less than four per cent of the area (Figure 5.30). While most of the designated defence training areas within or adjacent to the Region are small, the Shoalwater Bay Defence Training Area near Rockhampton is one of Australia's largest, and is regularly used. Every other year, Shoalwater Bay, and other defence training sites in the Region, host the combined Australia and United States of America training exercise called Talisman Sabre, which is Australia's largest defence training exercise.

The Australian Defence Force has a moratorium on the use of high explosives in the World Heritage Area, except in the Shoalwater Bay training area, where they continue to be used intermittently and in major training exercises. Stringent environmental procedures are followed when high explosives are used, including a requirement to maintain a safety zone for marine mammals and reptiles. There are dedicated shipboard and aerial observers who communicate any marine wildlife sightings to other participants.

#### Trends

The Australian Defence Force is currently enhancing its capabilities in amphibious landings and other ship-to-shore or coastal manoeuvres, including the acquisition of new landing vessels.<sup>188</sup> The key amphibious training areas in or adjacent to the Region are Shoalwater Bay and Cowley Beach, and the Army's major deployable force is based in Townsville. As a result, more frequent and intensive amphibious training exercises are expected in the Region in the coming years. Additional training areas may also be required to develop and maintain this new amphibious capability. An increase in near-shore vessel movements increases the risk of a vessel grounding and subsequent oil spill, as well as creating more sediment disturbance which can affect nearby seagrasses or corals. It also has the potential to result in increased disturbance to wildlife.

The United States of America has recently shifted its global military focus to enhance its capabilities in the Asia-Pacific region.<sup>188</sup> As a result, combined training exercises between Australian and U.S. forces are expected to increase in frequency and intensity. Shoalwater Bay and other sites in the Region will, therefore, increase in importance for major exercises such as Talisman Sabre. The Authority provided environmental impact advice on the 2013 tour and, to date, the exercises have shown no substantial environmental impacts. While U.S. forces operate under Australian Defence Force instructions during combined training exercises, any visiting foreign force increases the complexity of communications and, therefore, the risk of impacts.






Figure 5.30 Defence training areas relevant to the Region

The Australian Defence Force is increasingly aware of its contribution to greenhouse gas emissions, with the Royal Australian Navy trialling renewable fuels for ships. It is expected to introduce alternative, low-carbon fuels as suitable replacements become commercially available, and where consistent with operational needs.

While defence training activities are well-managed and have negligible impacts on the Great Barrier Reef, the predicted intensification of defence activities in the Region coincides with a decline in the Region's ecosystem health caused by a range of other pressures. The Australian Defence Force is working with the Authority to review the risks posed by defence activities in light of new information about the Region's declining ecosystem resilience and cumulative impacts.

### Benefits

Defence activities in the Region directly contribute to the training and operation of Australia's defence services. In addition, the acquisition of Shoalwater Bay in 1965 has provided environmental benefits. The land component remains largely undisturbed and is able to maintain natural environmental processes. It also supports high biodiversity, including internationally significant migratory species and wetlands, and has stunning landscape features.<sup>189</sup> Defence has contributed to research efforts by conducting population surveys of key species in both Shoalwater Bay and the Coral Sea. Operational activities can also help, directly and indirectly, to achieve management objectives for the Region including hydrographic surveys, charting, ocean surveillance, maritime search and rescue, defence aid to the civil community, and fisheries and border protection patrols.

Defence bases at Cairns and Townsville strongly support these regional economies. Little is known of the economic benefits of small-scale defence training activities to the coastal communities adjacent to the Region, although Talisman Sabre 2013 was predicted to contribute \$4 million to the Rockhampton economy, and \$200,000 to the Townsville economy.<sup>190</sup> Periodic visits from the United States of America, New Zealand and Singapore naval ships to ports at Townsville and Cairns also generate short-term economic benefits, related to increased visitation and tourism.<sup>191</sup>

### Impacts

Defence activities are well planned and well resourced, so incidents causing environmental harm to the Marine Park are rare. Standard operating procedures and contingency plans cover all defence activities, and any incidents are promptly reported and closely investigated.

However, by their nature, defence activities do pose risks which must be continually monitored and managed. In September 2005, a workshop – *Assessment of the risks of Defence activities in the Great Barrier Reef World Heritage Area* – was held. Using the Australian Standard risk assessment protocol, it was found that the potential impact of greatest concern was the introduction of marine pests through ballast water or hull fouling. The Australian Defence Force employs stringent quarantine measures to reduce this risk.<sup>192</sup>

Other impacts that may be of concern at a local or regional scale include:

- debris and residue from expendable stores, such as bullet casings or sonar buoys
- death, injury or disruption to marine life from explosives
- disturbance to marine animals from loud or low-flying aircraft
- discharge of sewage, food scraps and other wastes from ships
- oil spill from a ship grounding or collision
- loss of civilian life or property, particularly if civilians stray into defence training areas when exercises are taking place.

There is a range of legacy impacts associated with past defence activities. Most significant is the presence of large amounts of unexploded ordnance (such as shells, missiles and bombs) and chemical warfare agents which were deliberately dumped at sea at the end of World War II.<sup>193,194</sup> The largest post-war dumpsites were offshore from Cairns and Townsville, however, dumps occurred throughout the Region.<sup>187</sup> Chemical warfare agents were also dumped off Bowen and Proserpine in the late 1940s.<sup>187</sup> Some records were kept about the locations of these dumpsites, but the details are not reliable in terms of precise locations, quantities and types of materials dumped. In addition to dumped materials, Australia laid thousands of sea mines in the Great Barrier Reef during World War II. Mine sweeping activities after the war resulted in the Great Barrier Reef being declared safe for shipping by 1948<sup>195</sup>, but navigational charts still note former mined areas which may be unsafe for bottom activities such as trawling or anchoring. Encounters with World War II sea mines are extremely rare, and it is likely that they now pose a very low risk to humans or the environment. However, mines are occasionally discovered and require the Department of Defence and the Authority resources to neutralise; most recently, in 2013 at Night Island near Lockhart River, and in 2002 at Sudbury Reef near Cairns<sup>196</sup>.

These legacy issues are increasingly important to the Australian Defence Force. A pilot study is currently underway at John Brewer Reef (offshore Townsville) to evaluate the risks posed by unexploded ordnance. The Australian Defence Force is also working closely with researchers worldwide to evaluate the risks posed by World War II era sea mines in tropical environments. The Authority will continue to work with the Australian Defence Force to address the risks posed by both historic and contemporary defence activities.

### 5.4.8 Research activities

The Great Barrier Reef is an area of high scientific interest because of its biological and ecological diversity, geomorphology and cultural heritage. Scientific research has made a substantial contribution to the way the Reef is understood, managed and used. Monitoring also plays a key role in tracking trends in the Region's values and effects on those values.

Research programs range from small, independent student projects to large, multi-agency collaborative programs with teams of world-leading scientists.<sup>43</sup>

A network of six island research stations located at Lizard Island, Low Isles, Green Island, Orpheus Island, Heron Island and One Tree Island has been integral to research activities on the Reef. Eighty per cent of scientific research has been conducted around Lizard, Heron and Orpheus islands.<sup>197</sup>

#### Trends

The amount of research conducted in the Region, and the focus of that research, is determined by a number of variables including: the priorities of funding bodies such as the National Environmental Research Program and the Australian Research Council; the priorities of research end-users such as government agencies; and the research interests and capacities of scientists in universities and research institutions. There is a strong history of research programs which inform and help improve management of the Region.<sup>198</sup> Although past research focused primarily on studying the biophysical environment, there is also a history of social and economic studies. This recognises that effective natural resource management requires an understanding of social and economic systems as well as natural systems.

Technological advances continue to change the way research is conducted. For example, DNA mapping allows greater use of non-lethal sampling methods<sup>199</sup>, and remotely-sensed data reduces the need for field science.<sup>200,201</sup>

#### Benefits

Research and monitoring of the Great Barrier Reef environment continues to contribute to global knowledge about individual species, coral reef systems and tropical marine ecology. An improved understanding of the Region's environment and how its components interact and respond to changing conditions has contributed substantially to its protection and management. In addition, the results of targeted and applied research are providing managers with information to better measure the outcomes of management initiatives.

A range of academic institutions and government agencies undertake research about the Great Barrier Reef. In 2012, more than \$98 million of revenue was generated through the conduct of scientific research into the World Heritage Area, supporting more than 881 full-time equivalent jobs<sup>43</sup> (see Table 7.9 in Chapter 7). This is likely to be an underestimate of the total economic contribution of research activities as it is derived from only four major Australian research institutions involved in Great Barrier Reef research, namely: the Australian Institute of Marine Science; James Cook University's ARC Centre of Excellence in Coral Reef Studies; the Heron Island Research Station operated by the University of Queensland; and the Lizard Island Research Station operated by the Australian Museum.

The activities of other research institutions and government agencies make significant additional economic contributions. In 2010, research funding for Reef-related projects from Australian Research Council grants alone totalled more than \$8 million.<sup>202</sup>

Because the Great Barrier Reef is accessible and diverse, it provides a valuable training ground for researchers — supporting domestic and overseas students.

#### Impacts

The concentration of research activities, such as sampling, around research stations, has the potential to contribute to local depletion of some species and some minor, localised impacts on habitats. Little is known about the cumulative impacts of research activities at any particular location, however, given the scale of activities, overall impacts are likely to have only localised effects.

## 5.5 Summary of conclusions

Key drivers directly and indirectly influencing the Great Barrier Reef Region are:

- **Climate change.** Reef-dependent values and activities are vulnerable to the negative effects on Reef condition of ocean acidification, sea level rise, more frequent extreme weather and warming sea temperatures. The frequency and severity of associated impacts such as coral bleaching are predicted to increase, with serious consequences for the Reef likely before mid-century.
- **Economic growth.** Queensland's economy continues to grow rapidly and much of its economic activity takes place in the Great Barrier Reef catchment. Predicted growth in Queensland's economy relies on expansions in port capacity, shipping, coastal infrastructure and urban utilities adjacent to the Region.
- **Annual population growth** in the Reef catchment is 1.6 per cent compared with Australia's growth rate of less than 1.4 per cent. Combined with economic growth, population growth will increase pressure on the Region's environment, with potential effects on biodiversity, connectivity and aesthetics. An increasing population will likely mean increased recreational use of the Region.

- **Technological developments** contribute to our understanding of the Reef and enable the Region's values to be better protected and managed. They also improve access to previously remote areas for fishing and recreation, placing additional pressure on vulnerable habitats, and targeted fish and other species.
- Prevailing **societal attitudes** play a key role in people's use of and attitude towards the Reef. Stewardship actions implemented by community and industry are critically important for influencing attitudes and supporting management initiatives to maintain and enhance the Region's values.

Activities that affect the Region's values may be undertaken within the Region (direct uses) or beyond its boundaries.

- **Agriculture** within the catchment is an important producer of food and, to a lesser extent, fibre. Over the 160 years of European settlement in the catchment, agricultural use has become more widespread and more intensive. Land clearing and altering water courses have affected many ecological processes that support the Reef. There have been significant improvements in agricultural land use practices in recent years, such as those supported by the implementation of the *Reef Water Quality Protection Plan*.
- Although there is no marine-based **aquaculture** within the Region, there are land-based operations in the catchment. Within a local area, these may increase sediment and nutrient loads in discharged wastewater, modify hydrologic processes, introduce marine species or diseases and cause genetic pollution.
- **Urban development.** The six major population centres within the Great Barrier Reef catchment — Cairns, Townsville, Mackay, Rockhampton, Gladstone and Bundaberg — account for about 42 per cent of the population. While urban areas occupy only a small proportion of the catchment, development of housing and associated utilities increases land clearing, modifies water flows and affects connectivity between coastal and marine habitats.
- **Industrial development.** Much of the supporting infrastructure for mining and industry is located in coastal areas adjacent to the Region and, in the past two decades, four major State Development Areas have been declared in the catchment. If not properly managed, industrial development adjacent to the Great Barrier Reef could result in the discharge of pollutants, potential acid sulphate soils, artificial barriers to estuarine flow and coastal reclamation.
- There are 12 trading **ports** in or adjacent to the Region. Many are of economic importance, especially to export industries such as mining and agriculture. Over the past two decades, total export tonnage has grown by about 300 per cent. Five ports currently have active proposals for port expansions, and there are proposals for two new ports. Port operations and infrastructure can lead to impacts such as clearing, modifying and fragmenting of coastal habitats, reclamation of marine areas, alteration of natural coastal processes, as well as requiring capital and maintenance dredging.
- **Traditional use of marine resources.** Traditional Owners use coastal and marine resources to practice their living maritime culture, provide traditional food for their families and educate the younger generation about traditions, rules, protocols and activities. Impacts such as coastal development, habitat degradation, vessel strikes on wildlife, pollution, netting and sedimentation affect Traditional Owners' use of the Reef.
- **Tourism** is the most economically significant direct use of the Region, with about 2 million tourists visiting the Region each year. It offers a wide range of tourism experiences, focused on a small portion of the Region, and plays an important role in presenting the Region's world heritage values. Many tourism operators are active stewards of the Reef and more than 60 per cent of visitors travel to the Reef with a certified high standard tourism operator. The impacts associated with tourism activities, such as localised anchor damage and wildlife disturbance, are generally regarded as low risk, concentrated in a few intensively managed areas.
- The Great Barrier Reef supports a diversity of **fishing** activities including commercial, recreational, charter and Indigenous fisheries. Fisheries product from the Region is important to local communities, as well as domestic and international markets. Reef-based commercial fishing is mainly comprised of trawl, net, line and pot fisheries and is widely distributed across the Region. Net fishing impacts include bycatch and entanglement of species of conservation concern. Trawl bycatch can comprise hundreds of species, however, many are caught infrequently. Many species are targeted by both recreational and commercial fishers, increasing cumulative impacts on those species. Illegal fishing continues to be of concern as it erodes the benefits of no-take marine reserves.
- **Recreation other than fishing** includes swimming, motorised boating, snorkelling, sailing, diving and jetskiing. The most popular destinations are islands, reefs, shoals, cays and wrecks. Potential recreational impacts include localised anchor damage, littering, boat strikes on wildlife, and damage to corals when snorkelling and diving.
- **Ships** transiting the Reef carry export goods, service coastal and inland communities and transport passengers. The Region is one of the world's most regulated shipping areas. While the number of ship voyages increased substantially over the past 10 years — driven mainly by industrial activity — improvements in management arrangements have meant there have been relatively few shipping incidents.



- The Great Barrier Reef is a critical part of Australia's **defence** training programs. Training activities are regularly undertaken in a few designated areas of the Region, covering less than four per cent of its area. Recognising the potential for impacts such as noise, anchor damage and disturbance to wildlife, defence activities have undergone rigorous environmental assessments resulting in environmental management plans and protocols.
- **Research activities** contribute to global knowledge about Reef species and tropical marine ecology. An improved understanding of the Reef and how its components interact and respond to changing conditions contributes substantially to the Region's protection and management. Any impacts are minor and localised, mainly around research stations.

## References

1. Garnaut, R. 2008, *The Garnaut Climate Change Review*, Cambridge University Press, Melbourne.
2. Hajkowicz, S.A., Cook, H. and Littleboy, A. 2012, *Our future world: Global megatrends that will change the way we live, The 2012 revision*, CSIRO, Australia.
3. Intergovernmental Panel on Climate Change 2007a, *Climate change 2007 synthesis report, Summary for policymakers*, IPCC Secretariat, Gland, Switzerland.
4. Hoegh-Guldberg, O., Mumby, P.J., Hooten, A.J., Steneck, R.S., Greenfield, P., Gomez, E., Harvell, C.D., Sale, P.F., Edwards, A.J., Caldeira, K., Knowlton, N., Eakin, C.M., Iglesias-Prieto, R., Muthiga, N., Bradbury, R.H., Dubi, A. and Hatzioles, M.E. 2007, Coral reefs under rapid climate change and ocean acidification, *Science* 318: 1737-1742.
5. Intergovernmental Panel on Climate Change 2007b, *Climate Change 2007: Human and natural drivers of climate change*, IPCC, Gland, Switzerland, <[http://www.ipcc.ch/publications\\_and\\_data/ar4/wg1/en/sprsspm-human-and.html](http://www.ipcc.ch/publications_and_data/ar4/wg1/en/sprsspm-human-and.html)>.
6. Berkelmans, R., De'ath, G., Kininmonth, S. and Skirving, W.J. 2004, A comparison of the 1998 and 2002 coral bleaching events on the Great Barrier Reef: Spatial correlation, patterns and predictions, *Coral Reefs* 23(1): 74-83.
7. Johnson, J.E. and Marshall, P.A. (eds) 2007, *Climate change and the Great Barrier Reef: a vulnerability assessment*, Great Barrier Reef Marine Park Authority and Australian Greenhouse Office, Townsville.
8. De'ath, G., Fabricius, K.E., Sweatman, H. and Puotinen, M. 2012, The 27-year decline of coral cover on the Great Barrier Reef and its causes, *Proceedings of the National Academy of Sciences* 109(44): 17995-17999.
9. Koch, M., Bowes, G., Ross, C. and Zhang, X. 2013, Climate change and ocean acidification effects on seagrasses and marine macroalgae, *Global Change Biology* 19(1): 103-132.
10. Doropoulos, C., Ward, S., Diaz-Pulido, G., Hoegh-Guldberg, O. and Mumby, P.J. 2012, Ocean acidification reduces coral recruitment by disrupting intimate larval-algal settlement interactions, *Ecology Letters* 15(4): 338-346.
11. Brodie, J., Waterhouse, J., Schaffelke, B., Kroon, F., Thorburn, P., Rolfe, J., Johnson, J., Fabricius, K., Lewis, S., Devlin, M., Warne, M. and McKenzie, L.J. 2013, *2013 Scientific Consensus Statement: Land use impacts on Great Barrier Reef water quality and ecosystem conditions*, Reef Water Quality Protection Plan Secretariat, Brisbane.
12. Carilli, J.E., Norris, R.D., Black, B., Walsh, S.M. and McField, M. 2010, Century-scale records of coral growth rates indicate that local stressors reduce coral thermal tolerance threshold, *Global Change Biology* 16: 1247-1257.
13. Hoegh-Guldberg, O. and Bruno, J.F. 2010, The impact of climate change on the world's marine ecosystems, *Science* 328: 1523-1528.
14. Baker, A.C., Glynn, P.W. and Riegl, B. 2008, Climate change and coral reef bleaching: an ecological assessment of long-term impacts, recovery trends and future outlook, *Estuarine, Coastal and Shelf Science* 80(4): 435-471.
15. Hoegh-Guldberg, O. 1999, Climate change, coral bleaching and the future of the world's coral reefs, *Marine and Freshwater Research* 50(8): 839-866.
16. Wilkinson, C. (ed) 2008, *Status of coral reefs of the world: 2008*, Global Coral Reef Monitoring Network and Reef and Rainforest Research Centre, Townsville.
17. van Hooidonk, R., Maynard, J.A. and Planes, S. 2013, Temporary refugia for coral reefs in a warming world, *Nature Climate Change* 3(4): 508-511.
18. Baskett, M.I., Nisbet, R.M., Kappel, C.V., Mumby, P.J. and Gaines, S.D. 2010, Conservation management approaches to protecting the capacity for corals to respond to climate change: a theoretical comparison, *Global Change Biology* 16: 1229-1246.
19. Dlugokencky, E. & Tans, P. 2014, *Trends in atmospheric carbon dioxide: Recent Global CO<sub>2</sub>*, Earth System Research Laboratory, National Oceanic and Atmospheric Administration, <<http://www.esrl.noaa.gov/gmd/ccgg/trends/global.html>>.
20. De'ath, G., Lough, J.M. and Fabricius, K.E. 2009, Declining coral calcification on the Great Barrier Reef, *Science* 323(5910): 116-119.
21. Sheehan, P., Jones, R.N., Jolley, A., Preston, B., Clarke, M., Durack, P., Islam, S. and Whetton, P.H. 2008, Climate change and the new world economy: Implications for the nature and timing of policy responses, *Global Environmental Change* 18(3): 380-396.
22. Great Barrier Reef Marine Park Authority 2009, *Great Barrier Reef Outlook Report 2009*, GBRMPA, Townsville.
23. Tourism Industry Council 2008, *Climate change response manual for Great Barrier Reef Marine Park operators*, TIC, Brisbane.
24. Kikkawa, T., Kita, J. and Ishimatsu, A. 2004, Comparison of the lethal effect of carbon dioxide and acidification on red sea bream (*Pagrus major*) during the early developmental stages, *Marine Pollution Bulletin* 48(1-2): 108-110.
25. Ishimatsu, A., Kikkawa, T., Hayashi, M., Lee, K. and Kita, J. 2004, Effects of carbon dioxide on marine fish: larvae and adults, *Journal of Oceanography* 60(4): 731-741.
26. Guinotte, J.M. and Fabry, V.J. 2008, Ocean acidification and its potential effects on marine ecosystems, *Annals of the New York Academy of Sciences* 1134(1): 320-342.
27. Fenton, M., Kelly, G., Vella, K. and Innes, J. 2007, Climate change and the Great Barrier Reef: industries and communities, in *Climate change and the Great Barrier Reef: a vulnerability assessment*, eds J.E. Johnson and P.A. Marshall, Great Barrier Reef Marine Park Authority and Australian Greenhouse Office, Townsville, pp. 745-771.
28. Tobin, A.J., Schlaff, R., Tobin, R., Penny, A., Ayling, T., Ayling, A., Krause, B., Welch, D., Sutton, S., Sawynok, B., Marshall, N.A., Marshall, P.A. and Maynard, J.A. 2010a, *Adapting to change: minimising uncertainty about the effects of rapidly-changing environmental conditions on the Queensland Coral Reef Fin Fish Fishery, Final Report to the Fisheries Research and Development Corporation, Project 2008/103*, James Cook University, Townsville.
29. Poloczanska, E.S., Hobday, A.J. and Richardson, A.J. 2012, *Marine climate change in Australia, impacts and adaptation responses. 2012 report card*, CSIRO, Canberra.
30. Morison, A.K. and Pears, R.J. 2012, *Assessment of the ecological vulnerability of the East Coast Otter Trawl Fishery to climate change: a brief synthesis of information and results of an*

expert workshop, Great Barrier Reef Marine Park Authority, Townsville.

31. Great Barrier Reef Marine Park Authority 2012a, *Climate change adaptation: Outcomes from the Great Barrier Reef Climate Change Action Plan 2007-2012*, GBRMPA, Townsville.

32. Department of Transport and Main Roads 2011, *Annual report 2010-11*, Department of Transport and Main Roads, Brisbane.

33. Gooch, M., Vella, K., Marshall, N.A., Tobin, R. and Pears, R. 2012, A rapid assessment of the effects of extreme weather on two Great Barrier Reef industries, *Australian Planner* 50(3): 198-215.

34. Marshall, N.A. and Tobin, R. 2012, "More than meets the eye": The social and economic impacts of recent natural disasters on marine resource dependent industries of the Great Barrier Reef Region, Great Barrier Reef Marine Park Authority, Townsville.

35. Great Barrier Reef Marine Park Authority 2011a, *Extreme weather and the Great Barrier Reef*, GBRMPA, Townsville.

36. National Oceanic and Atmospheric Administration 2013, *Trends in Atmospheric Carbon Dioxide*, NOAA, Washington, DC, USA, <<http://www.esrl.noaa.gov/gmd/ccgg/trends/weekly.html>>.

37. Etheridge, D.M., Steele, L.P., Langenfelds, R.L., Francey, R.J., Barnola, J.M. and Morgan V I. 1996, Natural and anthropogenic changes in atmospheric CO<sub>2</sub> over the last 1000 years from air in Antarctic ice and firn, *Journal of Geophysical Research* 101: 4115-4128.

38. Veron, J.E.N., Hoegh-Guldberg, O., Lenton, T.M., Lough, J.M., Obura, D.O., Pearce-Kelly, P., Sheppard, C.R., Spalding, M., Stafford-Smith, M.G. and Rogers, A.D. 2009, The coral reef crisis: The critical importance of <350 ppm CO<sub>2</sub>>, *Marine Pollution Bulletin* 58(10): 1428-1436.

39. Tans, P. & Keeling, R. 2013, *Trends in atmospheric CO<sub>2</sub>: Mauna Loa, Hawaii*, National Oceanic and Atmospheric Administration: Earth System Research Laboratory and Scripps Institute of Oceanography, <[www.esrl.noaa.gov/gmd/ccgg/trends/](http://www.esrl.noaa.gov/gmd/ccgg/trends/)>.

40. Great Barrier Tourism Climate Change Action Group 2009, *Great Barrier Reef tourism climate change action strategy 2009-2012*, GBRMPA, Townsville.

41. Queensland Department of Employment, Economic Development and Innovation 2012, *Queensland in the 21st Century*, DEEDI, Brisbane, <<http://www.business.qld.gov.au/invest/benefits-business-queensland/queenslands-pro-business-economy/>>.

42. Binney, J. 2012, *The economic and social impacts of protecting environmental values in the Great Barrier Reef catchment waterways and the reef lagoon*, Report prepared for Queensland Department of Environment and Heritage Protection, Marsden Jacob Associates, Brisbane.

43. Deloitte Access Economics 2013, *Economic contribution of the Great Barrier Reef*, Great Barrier Reef Marine Park Authority, Townsville.

44. Queensland Government 2013a, *State Budget 2013-2014: Budget Strategy and Outlook*, Queensland Government, Brisbane.

45. Department of Agriculture, Fisheries and Forestry 2012a, *Queensland's agriculture strategy: A 2040 vision to double the value of production*, Discussion paper for consultation, DAFF, Brisbane.

46. Bureau of Resources and Energy Economics 2012, *Australian bulk commodity exports and infrastructure: outlook to 2025*, BREE, Canberra.

47. Great Barrier Reef Marine Park Authority 2012b, *Ports and shipping information sheet*, GBRMPA, Townsville.

48. Productivity Commission 2003, *Industries, land use and water quality in the Great Barrier Reef catchment: research report*, Productivity Commission, Canberra.

49. Queensland Government 2012a, *Delivering on DestinationQ*, Queensland Government, Brisbane.

50. Great Barrier Reef Marine Park Authority 2012c, *Informing the outlook for Great Barrier Reef coastal ecosystems*, GBRMPA, Townsville.

51. Department of Natural Resources and Mines 2013, *Wells drilled in Queensland 1954-2011*, DNRM, Brisbane, <<http://mines.industry.qld.gov.au/mining/production.htm>>.

52. Government Statistician 2013, *Queensland Regional Profile: GBR LGAs Region*, Queensland Treasury and Trade, Brisbane.

53. Office of Economic and Statistical Research 2012a, *Population growth highlights and trends, Queensland 2012*, Queensland Treasury and Trade, Brisbane.

54. Australian Bureau of Statistics 2013, *Census of Population and Housing 2011: Data and analysis*, ABS, Canberra, <<http://www.abs.gov.au/websitedbs/censushome.nsf/home/quickstats?opendocument&navpos=220>>.

55. Queensland Government Statistician's Office 2014, *Queensland Regional Database: population estimates 1991 to 2012 and population projections 2011 to 2036: local government areas (2014) and Statistical Area Level 2*, Queensland Treasury and Trade, Brisbane, <<http://www.qgso.qld.gov.au/products/qld-regional-database/qld-regional-database/index.php>>.

56. Bohnet, I., Pert, P. and Schultz, P. 2010, *Sea- and tree-change phenomena in the Wet Tropics: Impacts of urbanisation and mitigation potential*, Final Project Report to the Marine and Tropical Sciences Research Facility, Reef and Rainforest Research Centre, Cairns.

57. Queensland Government (Unpublished data), 'Vessel Registration Data by LGA for December 2000, 2006 and 2011. Vessel Registration Data by Postcode for 2007, 2009, 2011. Data retrieved from Department of Transport and Main Roads (DTMR) Transport Registration and Integrated Licensing System (TRAILS)'.

58. Great Barrier Reef Marine Park Authority 2012d, *Recreation management strategy for the Great Barrier Reef Marine Park*, GBRMPA, Townsville.

59. Heupel, M.R., Simpfendorfer, C.A. and Fitzpatrick, R. 2010, Large-Scale Movement and Reef Fidelity of Grey Reef Sharks, *PLoS ONE* 5(3): 9650.

60. Heupel, M.R., Simpfendorfer, C.A. and Hueter, R.E. 2003, Running before the storm: blacktip sharks respond to falling barometric pressure associated with Tropical Storm Gabrielle. *Journal of fish biology* 63: 1357.

61. Stieglitz, T.C., Kelley, R., Beaudoin, J., Hughes-Clarke, J.E. and Kininmonth, S. 2006, Multibeam bathymetry mapping of the Great Barrier Reef seascape: A pilot study, in *Ocean Sciences Meeting Program and Abstracts, American Geophysical Union, 20-24 Feb, Honolulu, Hawaii, USA*, eds. Anonymous, American Geophysical Union, Washington, DC, USA, .

62. Morison, A.K. 2011, *Ecological risk assessment of the East Coast Otter Trawl Fishery in the Great Barrier Reef Marine Park: report of the Phase II Expert Workshop on Risk to Ecosystem Processes, held on 15 November 2010, Brisbane*, Unpublished report to the Great Barrier Reef Marine Park Authority.

63. O'Neill, M.F. and Leigh, G.M. 2007, Fishing power increases continue in Queensland's East Coast Trawl fishery, Australia, *Fisheries Research* 85(1-2): 84-92.

64. Department of the Premier and Cabinet 2013, *Great Barrier Reef Report Card 2011, Reef Water Quality Protection Plan*, Reef Water Quality Protection Plan Secretariat, Brisbane.

65. McCombs, M. 2004, *Setting the agenda: The mass media and public opinion*, Polity Press, Cambridge, UK.

66. Trigger, D., Schubert, M. and Meurk, C. Unpublish, *Shifting baselines in the non-scientific literature*, Great Barrier Reef Marine Park Authority, Townsville.

67. Emtage, N. and Reghenzani, J. 2008, *Wet Tropics Sustainable Agriculture Survey Interim Report: A survey of rural landholders in the Wet Tropics Natural Resource Management region. Report to the Marine and Tropical Sciences Research Facility*, Reef and Rainforest Research Centre, Cairns.

68. Young, J. and Temperton, J. 2008, *Measuring community attitudes and awareness towards the Great Barrier Reef 2007*, Great Barrier Reef Marine Park Authority, Townsville.
69. Young, J. and Mar, R. 2010, *Community perceptions of climate change and the effects on the Great Barrier Reef. Report prepared for the Great Barrier Reef Marine Park Authority*, Colmar Brunton, Brisbane.
70. Office of Economic and Statistical Research 2012b, *Agriculture (a): Gross value of production by commodity, Queensland, 2000-01 to 2010-11*, Queensland Treasury and Trade, Brisbane, <<http://www.oesr.qld.gov.au/products/tables/agriculture-gross-value-production/index.php>>.
71. Gilbert, M. and Brodie, J. 2001, *Population and major land use in the Great Barrier Reef catchment area: spatial and temporal trends*, Great Barrier Reef Marine Park Authority, Townsville.
72. McCulloch, M., Fallon, S., Wyndham, T., Hendy, E., Lough, J.M. and Barnes, D. 2003, Coral record of increased sediment flux to the inner Great Barrier Reef since European settlement, *Nature* 421(6924): 727-730.
73. Marion, G. 2007, *The nitrogen isotopic composition of the organic matrices of coral skeleton: A proxy for historical nitrogen provenance in tropical coastal oceans*, PhD thesis, University of Queensland, Brisbane.
74. Roff, G., Clark, T.R., Reymond, C.E., Zhao, J., Feng, Y., McCook, L.J., Done, T.J. and Pandolfi, J.M. 2013, Palaeoecological evidence of a historical collapse of corals at Pelorus Island, inshore Great Barrier Reef, following European settlement, *Proceedings of the Royal Society B: Biological Sciences* 280(1750): 2012-2100.
75. Brodie, J.E., Binney, J., Fabricius, K., Gordon, I., Hoegh-Guldberg, O., Hunter, H., O'Reagain, P., Pearson, R., Quirk, M., Thorburn, P., Waterhouse, J., Webster, I. and Wilkinson, S. 2008, *Synthesis of evidence to support the Scientific Consensus Statement on water quality in the Great Barrier Reef*, Reef Water Quality Protection Plan Secretariat, Brisbane.
76. Queensland Government 2013b, *Vegetation Management Framework Amendment Bill 2013*, Queensland Government, Brisbane, <<http://www.legislation.qld.gov.au/Bills/54PDF/2013/VegetMgmtFrAmwAB13.pdf>>.
77. Great Barrier Reef Marine Park Authority 2001a, *Great Barrier Reef Catchment Water Quality Action Plan. A Report To Ministerial Council on targets for pollutant loads*, GBRMPA, Townsville.
78. Brodie, J. 2007, *Nutrient management zones in the Great Barrier Reef catchment: a decision system for zone selection: report to the Department of Environment and Heritage*, James Cook University, Townsville.
79. Hildebrand, C. 2002, *Report: Independent assessment of the sugar industry*, Assessment Secretariat, Canberra.
80. Great Barrier Reef Marine Park Authority 2013, *Haughton Basin Assessment: Burdekin Dry Tropics Natural Resource Management Region*, GBRMPA, Townsville.
81. Kroon, F.J., Kuhnert, P.M., Henderson, B.L., Wilkinson, S.N., Kinsey-Henderson, A., Abbott, B., Brodie, J.E. and Turner, R.D.R. 2012, River loads of suspended solids, nitrogen, phosphorus and herbicides delivered to the Great Barrier Reef lagoon, *Marine Pollution Bulletin* 65(4-9): 167-181.
82. Fabricius, K.E., Okaji, K. and De'ath, G. 2010, Three lines of evidence to link outbreaks of the crown-of-thorns seastar *Acanthaster planci* to the release of larval food limitation, *Coral Reefs* 29: 593-605.
83. Great Barrier Reef Marine Park Authority 2012e, *A vulnerability assessment for the Great Barrier Reef: Seagrass*, GBRMPA, Townsville.
84. Powell, B. and Martens, M. 2005, A review of acid sulfate soil impacts, actions and policies that impact on water quality in Great Barrier Reef catchments, including a case study on remediation at East Trinity, *Marine Pollution Bulletin* 51: 149-164.
85. Pulsford, J.S. 1996, *Historical nutrient usage in coastal Queensland river catchments adjacent to the Great Barrier Reef Marine Park*, Great Barrier Reef Marine Park Authority, Townsville.
86. Department of Environment and Resource Management 2012, *Queensland land use mapping program*, DERM, Brisbane, <<http://www.derm.qld.gov.au/science/lump/>>.
87. Fertilizer Industry Federation of Australia 2000, *Application rates*, FIFA, Canberra, <<http://www.fertilizer.org.au/>>.
88. Incitec Pivot Fertilisers 2013, *Incitec Pivot Fertilisers*, IPF, Melbourne, <<http://incitecpivotfertilisers.com.au>>.
89. Food and Agriculture Organization of United Nations 2005, *National Aquaculture Sector Overview, Australia. Fact Sheets*, FAO, Rome.
90. Wingfield, M. 2012, *Ross Lobegeiger Report to Farmers: Aquaculture productivity survey Queensland 2010-2011*, Department of Agriculture, Fisheries and Forestry, Brisbane.
91. Department of Agriculture, Fisheries and Forestry 2012b, *Aquaculture industry profile*, DAFF, Brisbane, <[http://www.daff.qld.gov.au/28\\_15105.htm](http://www.daff.qld.gov.au/28_15105.htm)>.
92. Lucas, J.S. and Southgate, P.C. 2005, *Aquaculture Farming Aquatic Animals and Plants*, Wiley-Blackwell, West Sussex, UK.
93. Office of Economic and Statistical Research 2013, *Data tables: Project dwellings (medium series) by local government area, Queensland, 30 June, 2006 to 2031*, Queensland Treasury and Trade, Brisbane, <<http://www.oesr.qld.gov.au/products/tables/proj-dwellings-medium-series-lga-qld/index.php>>.
94. Townsville City Council 2010, *Black Ross (Townsville) Water Quality Improvement Plan: Improving Water Quality from Creek to Coral*, Townsville City Council: Creek to Coral, Townsville 2010, TCC, Townsville.
95. Environment Protection Agency 2002, *State Coastal Management Plan: Queensland's Coastal Management Policy*, EPA, Brisbane.
96. Department of Energy and Water Supply 2013, *Queenslands water sector: a 30-year strategy. Discussion paper - shaping our water future*, DEWS, <[http://www.dews.qld.gov.au/\\_data/assets/pdf\\_file/0013/31144/30yr-water-strategy-discussion-paper.pdf](http://www.dews.qld.gov.au/_data/assets/pdf_file/0013/31144/30yr-water-strategy-discussion-paper.pdf)>.
97. Kamrowski, R.L., Limpus, C.J., Moloney, J. and Hamann, M. 2012, Coastal light pollution and marine turtles: Assessing the magnitude of the problem, *Endangered Species Research* 19: 85-98.
98. Steven, A.D. and van Woesik, R. 1990, *A multi-disciplinary examination of the Hayman Island fringing reef: influence of a secondary sewage discharge*, Great Barrier Reef Marine Park Authority, Townsville.
99. van Woesik, R. 1989, *A preliminary examination of the sedimentology, reef growth and hydrography of Green Island*, Great Barrier Reef Marine Park Authority, Townsville.
100. Great Barrier Reef Marine Park Authority 2001b, *Great Barrier Reef Water Quality: Current Issues*, GBRMPA, Townsville.
101. Waterhouse, J. and Johnson, J. 2002, Sewage discharges in the Great Barrier Reef Region, *Water* 29(5): 43-49.
102. Environmental Protection Agency 1999, *Queensland's coast: managing its future: a position paper on coastal management in Queensland*, Environmental Protection Agency, Brisbane.
103. Vis, M. 2013, *Personal Communication*, Townsville City Council.
104. Brodie, J., Pearson, R., Lewis, S.E., Bainbridge, Z.T., Waterhouse, J. and Prange, J. 2009, *Water quality research: baseline synthesis and year 1 summary, Report to the Marine and Tropical Sciences Research Facility*, Reef and Rainforest Research Centre, Cairns.
105. Department of State Development, Infrastructure and Planning 2012, *Current State Development Areas*, Department of Local Government, Community Recovery and Resilience, Brisbane, <<http://www.dlg.qld.gov.au/land-for-industry/current-state-development-areas.html>>.
106. Franks, D.M., Brereton, D. and Moran, C.J. 2010, Managing the cumulative impacts of coal mining on regional



communities and environments in Australia, *Impact Assessment and Project Appraisal* 28(4): 299-312.

107. Eberhard, R. 2012, *Fitzroy River estuary development proposals: A review of issues*, Fitzroy Basin Association, Rockhampton.

108. Context Pty Ltd 2013, *Defining the aesthetic values of the Great Barrier Reef*, Department of Sustainability, Environment, Water, Population and Communities, Canberra.

109. Ports Australia 2012, *Trade statistics for 2011/12*, Ports Australia, Sydney, <<http://www.portsaustralia.com.au/tradestats/>>.

110. Brunner, R.W.J. 2011, *Sustainability in the design of bulk coal ports: an Australian case study*, North Queensland Bulk Ports Corporation, Mackay, Queensland.

111. Department of Sustainability, Environment, Water, Population and Communities 2013, *State Party Report on the state of conservation of the Great Barrier Reef World Heritage Area (Australia) property ID N154 in response to World Heritage Committee decision WHC 35 COM 7B.10*, DSEWPac, Canberra.

112. Queensland Government 2012b, *Great Barrier Reef ports strategy economic analysis*, Department of State Development, Infrastructure and Planning, Brisbane.

113. Rodrigue, J.P., Comtois, C. & Slack, B. 2013, *The Geography of Transport Systems*, Hofstra University, Department of Global Studies and Geography, <<http://people.hofstra.edu/geotrans/>>.

114. Erftemeijer, P.L., Riegl, B., Hoeksema, B.W. and Todd, P.A. 2012, Environmental impacts of dredging and other sediment disturbances on corals: a review, *Marine Pollution Bulletin* 64(9): 1737-1765.

115. Foster, T., Smith, A., Jury, M. and Driscoll, A. 2011, Overview of PIANC Report 108 - dredging and port construction around coral reefs, in *Coasts and ports 2011 : Diverse and developing, Proceedings of the 20th Australasian Coastal and Ocean Engineering Conference and the 13th Australasian Port and Harbour Conference*. eds. Anonymous , Engineers Australia, Barton, ACT, pp.370-375.

116. Erftemeijer, P.L. and Lewis, R.R. 2006, Environmental impacts of dredging on seagrasses: A review, *Marine Pollution Bulletin* 52(12): 1553-1572.

117. Slooten, L., Benjamins, S. and Turek, J. 2011, *Potential impacts of Project Next Generation on Hector's dolphins and other marine mammals*, Otago University, Otago, NZ.

118. Bak, R. 1978, Lethal and sublethal effects of dredging on reef corals, *Marine Pollution Bulletin* 9: 14-16.

119. Rogers, C.S. 1990, Responses of coral reefs and reef organisms to sedimentation, *Marine Ecology Progress Series* 62: 185-202.

120. Marshall, N.A. and Marshall, P.A. 2007, Conceptualizing and operationalizing social resilience within commercial fisheries in northern Australia, *Ecology and Society* 12(1): Art. 1.

121. Barry, G. 2012, *Transcript of an interview with Woppaburra Traditional Owners*.

122. Packham, L. 2013, *Calls for action on traditional turtle hunting after tourists witness slaughter on Green Island*, April 23, The Cairns Post, Cairns.

123. World Economic Forum. 2013, *The travel and tourism competitiveness report 2013: Reducing barriers to economic growth and job creation*, WEF, Geneva, Switzerland.

124. Great Barrier Reef Marine Park Authority 2012f, *Environmental management charge*, GBRMPA, Townsville, <<http://www.gbrmpa.gov.au/zoning-permits-and-plans/environmental-management-charge>>.

125. Tourism Research Australia 2011, *State of the industry 2011*, TRA, Canberra.

126. Tourism Research Australia Unpublished report, 'International visitation in catchment 2002-2011 IVS, TRA, Canberra'.

127. Harriott, V.J. 2002, *Marine tourism impacts and their management on the Great Barrier Reef*, CRC Reef, Townsville.

128. Ormsby, J., Moscardo, G., Pearce, P. and Foxlee, J. 2004, *A review of research into tourist and recreational uses of*

*protected natural areas GBRMPA research on-line publication No. 79*, Great Barrier Reef Marine Park Authority, Brisbane.

129. Great Barrier Reef Marine Park Authority 2012g, *Field management quarterly report (IN-CONFIDENCE)*, GBRMPA, Townsville.

130. Queensland Department of Employment, Economic Development and Innovation 2011a, *Annual status report 2011: East Coast Otter Trawl Fishery*, DEEDI, Brisbane.

131. Robins, J.B. and Campbell, M.J., McGilvray, J.G. 1999, Reducing Prawn-trawl Bycatch in Australia: An overview and an Example from Queensland, *Marine Fisheries Review* .

132. Pears, R.J., Morison, A.K., Jebreen, E.J., Dunning, M.C., Pitcher, C.R., Courtney, A.J., Houlden, B. and Jacobsen, I.P. 2012a, *Ecological risk assessment of the East Coast Otter Trawl Fishery in the Great Barrier Reef Marine Park: summary report*, Great Barrier Reef Marine Park Authority, Townsville.

133. Department of Sustainability, Environment, Water, Populations and Communities 2012, *Assessment of the East Coast Inshore Fin Fish Fishery*, Commonwealth of Australia, Canberra.

134. Halliday, I.A., Ley, J.A., Tobin, A.J., Garrett, R.N., Gribble, N.A. and Mayer, D.A. 2001, *The effects of net fishing: addressing biodiversity and bycatch issues in Queensland inshore waters*, Fisheries Research and Development Cooperation Project 97/206, Department of Primary Industries, Deception Bay, Queensland.

135. Mapstone, B.D., McKinlay, J.P. and Davies, C.R. 1996, *A description of commercial reef line fishery logbook data held by the Queensland Fisheries Management Authority*, CRC Reef Research Centre and Department of Tropical Environment Studies and Geography, James Cook University, Townsville.

136. Heupel, M.R., Williams, A.J., Welch, D., Mapstone, B.D., Ballagh, A. and Simpfendorfer, C.A. 2008, *Compilation of information on the interaction of reef sharks with the reef line fishery in the Great Barrier Reef Marine Park: final report to the Great Barrier Reef Marine Park Authority*, Great Barrier Reef Marine Park Authority, Townsville.

137. Fisheries and Aquaculture Department 2007, *The state of the world's fisheries and aquaculture 2006*, Food and Agriculture Organization, Rome, Italy.

138. Sim, S.Y. 2004, *Regional developments and update*, April-June edn, Asia-Pacific Marine Finfish Aquaculture Network, Bangkok, Thailand, <<http://library.enaca.org/AquacultureAsia/Articles/April-June-2004/10mfan-ezine-april04.pdf>>.

139. Department of Agriculture, Fisheries and Forestry 2012c, 'Unpublished data'.

140. Australian Institute of Petroleum Ltd N, *Facts about diesel prices*, AIP, Canberra, <[http://www.aip.com.au/pricing/facts/Facts\\_about\\_Diesel\\_Prices.htm](http://www.aip.com.au/pricing/facts/Facts_about_Diesel_Prices.htm)>.

141. Tobin, R.C., Beggs, K., Sutton, S.G. and Penny, A. 2010b, *Baseline socio-economic data for Queensland east-coast inshore and rocky reef fishery stakeholders. Part D: Seafood consumers*, Technical Report No. 8, Fishing and Fisheries Research Centre, James Cook University, Townsville.

142. Department of Agriculture, Fisheries and Forestry 2012d, *Commercial line fishing in Queensland*, DAFF, Brisbane, <<http://www.daff.qld.gov.au/fisheries/commercial-fisheries/queenslands-commercial-fisheries/line-fisheries>>.

143. Department of Agriculture, Fisheries and Forestry 2012e, *Commercial trawling in Queensland*, DAFF, Brisbane, <<http://www.daff.qld.gov.au/fisheries/commercial-fisheries/queenslands-commercial-fisheries/rawl-fishery>>.

144. Tobin, R.C., Beggs, K., Sutton, S., Penny, A., Maroske, J. and Williams, L. 2010c, *Baseline socioeconomic data for Queensland east coast inshore and rocky reef fishery stakeholders, Part B: Charter fishers*, Fishing and Fisheries Research Centre, James Cook University, Townsville.

145. Tobin, R., Sutton, S., Penny, A., Williams, L., Maroske, J. and Nilsson, J. 2010d, *Baseline socio-economic data for Queensland east-coast inshore and rocky reef fishery stakeholders. Part A: commercial fishers*, FRDC Project No. 2007/048, Technical Report No. 5, Fishing and Fisheries Research Centre, James Cook University, Townsville.



146. Great Barrier Reef Marine Park Authority 2012h, *A vulnerability assessment for the Great Barrier Reef: Inshore dolphins*, GBRMPA, Townsville.
147. Great Barrier Reef Marine Park Authority Unpublished report, 'A vulnerability assessment for the Great Barrier Reef: Dugongs'.
148. Queensland Department of Employment, Economic Development and Innovation 2011b, *Stock status of Queensland's fisheries resources 2011*, DEEDI, Brisbane.
149. Great Barrier Reef Marine Park Authority 2011b, *A vulnerability assessment for the Great Barrier Reef: grey mackerel*, GBRMPA, Townsville.
150. Campbell, A.B., O'Neill, M.F., Staunton-Smith, J., Atfield, J.C. and Kirkwood, J. 2012, *Stock assessment of the Australian East Coast Spanish mackerel (Scomberomorus commerson) fishery*, Department of Agriculture, Fisheries and Forestry, Brisbane.
151. Campbell, A.B., O'Neill, M.F., Sumpton, W.D., Kirkwood, J. and Wesche, S. 2009, *Stock assessment summary of the Queensland snapper fishery (Australia) and management strategies for improving sustainability*, Department of Employment, Economic Development and Innovation, Brisbane.
152. Andersen, J. 2012, *Reef fishery in crisis as stocks drop*, 16 January 2012, Townsville Bulletin, Townsville.
153. Brown, I.W., Sumpton, W.D., McLennan, M., Welch, D.J., Kirkwood, J., Butche, A., Mapleston, A., Mayer, D., Begg, G., Campbell, M., Haliday, I. and Sawynok, W. 2008, *National strategy for the survival of released line-caught fish: tropical fish species, Fisheries Research and Development Corporation Project No. 2003/019*, Department of Primary Industries and Fisheries, Brisbane.
154. Tobin, A.J., Simpfendorfer, C.A., Mapleston, A., Currey, L., Harry, A.V., Welch, D.J., Ballagh, A.C., Chin, A., Szczecinski, N., Schlaff, A., White, J. and Moore, B. 2010e, *A quantitative ecological risk assessment of sharks and finfish of the Great Barrier Reef World Heritage Area inshore waters: a tool for fisheries and marine park managers. Identifying species at risk and potential mitigation strategies*, Marine and Tropical Sciences Research Facility, Cairns.
155. Pascoe, S., Dichmont, C.M., Brooks, K., Pears, R.J. and Jebreen, E. 2012, Management objectives of Queensland fisheries: putting the horse before the cart, *Marine Policy* 37: 115.
156. Pears, R.J., Morison, A.K., Jebreen, E.J., Dunning, M.C., Pitcher, C.R., Courtney, A.J., Houlden, B. and Jacobsen, I.P. 2012b, *Ecological risk assessment of the East Coast Otter Trawl Fishery in the Great Barrier Reef Marine Park: Technical report*, Great Barrier Reef Marine Park Authority, Townsville.
157. Queensland Department of Employment, Economic Development and Innovation 2011c, *Annual status report 2010: East Coast Inshore Fin Fish Fishery*, DEEDI, Brisbane.
158. Department of Environment and Heritage Protection 2012, *Marine wildlife strandings*, DEHP, Brisbane, <[http://www.ehp.qld.gov.au/wildlife/caring-for-wildlife/marine\\_strandings.html](http://www.ehp.qld.gov.au/wildlife/caring-for-wildlife/marine_strandings.html)>.
159. Courtney, A.J., Haddy, J.A., Campbell, v., Roy, D.P., Tonks, M.L., Gaddes, S.W., Chilcott, K.E., O'Neill, M.F., Brown, I.W., McLennan, M., Jebreen, E.J., van der Geest, C., Rose, C., Kistler, S., Turnbull, C.T., Kyne, P.M., Bennett, M.B. and Taylor, J. 2007, *Bycatch weight, composition and preliminary estimates of the impact of bycatch reduction devices in Queensland's trawl fishery*, Department of Primary Industries and Fisheries, Brisbane.
160. Mapstone, B.D., Davies, C.R., Little, L.R., Punt, A.E., Smith, A.D.M., Pantus, F., Lou, D.C., Williams, A.J., Jones, A., Ayling, A.M., Russ, G.R. and McDonald, A.D. 2004, *The effects of line fishing on the Great Barrier Reef and evaluations of alternative potential management strategies.*, CRC Reef Research Centre, Townsville.
161. Pitcher, C.R., Doherty, P., Arnold, P., Hooper, J., Gribble, N.A., Bartlett, C., Browne, M., Campbell, N., Cannard, T., Cappel, M., Carini, G., Chalmeres, S., Cheers, S., Chetwynd, D., Colegax, A., Coles, R., Cook, S., Davie, P., De'ath, G., Devereux, D., Done, B., Donovan, T., Ehrke, B., Ellis, N., Ericson, G., Fellegara, I., Forcey, K., Furey, M., Gledhill, D., Good, N., Gordon, S., Haywood, M., Hendricks, M., Jacobsen, I., Johnson, J., Jones, M., Kininmonth, S., Kistler, S., Last, P., Leite, A., Amrks, S., McLeod, I., Oczkowicz, S., Robinson, M., Rose, C., Seabright, D., Sheils, J., Sherlock, M., Skelton, P., Smith, D., Smith, G., Speare, P., Stowar, M., Strickland, C., Van der Geest, C., Venables, W., Walsh, C., Wassenberg, T.J., Welna, A. and Yearsley, G. 2007, *Seabed biodiversity on the continental shelf of the Great Barrier Reef World Heritage Area. AIMS/CSIRO/QM/QDPI CRC Reef Research Task final report*, CSIRO Marine and Atmospheric Research, Hobart.
162. Taylor, S., Webley, J. and McInnes, K. 2012, 2010 *State wide recreational fishing survey*, Department of Agriculture, Fisheries and Forestry, Brisbane.
163. Rolfe, J., Gregg, D., and Tucker, G., 2011, *Valuing local recreation in the Great Barrier Reef, Australia, Research Report No. 102 August 2011*, Crawford School of Economics and Government, Australian National University, Canberra.
164. Sutton, S.G. 2006, *An assessment of the social characteristics of Queensland's recreational fishers*, CRC Reef Research Centre, Townsville.
165. Infofish Australia 2012, *About CapReef*, Infofish Australia, Fitzroy, Queensland, <<http://info-fish.net/capreef/about-capreef/>>.
166. Barwick, M., Sawynok, B., Sawynok, S., Ansell, D. and Prokop, F. 2009, *Angling for conservation: Review and extension of conservation and sustainability-focussed initiatives which have been funded, supported or undertaken by Australia's recreational fishing sector*, Fisheries Research and Development Corporation, Canberra.
167. McManus, A., Prof., Hunt, W., Dr., Storey, J. and White, J. 2011, *Identifying the health and well-being benefits of recreational fishing*, FRDC Project Number: 2011/217, Centre of Excellence for Science, Seafood and Health; Curtin Health Innovation Research Institute; Curtin University, Perth.
168. Maritime Safety Queensland 2012, *Registration for recreational boats*, MSQ, Brisbane, <<http://www.msq.qld.gov.au/Registration/Recreational-ships.aspx>>.
169. Sumpton, W.D., Brown, I.W., Mayer, D.G., McLennan, M.F., Mapleston, A., Butcher, A.R., Welch, D.J., Kirkwood, J.M., Sawynok, B. and Begg, G.A. 2010, Assessing the effects of line capture and barotrauma relief procedures on post-release survival of key tropical reef species in Australia using recreational tagging clubs, *Fisheries Management and Ecology* 17: 77-88.
170. Meager, J.J. and Limpus, C.J. 2012a, Marine wildlife stranding and mortality database annual report 2011, III Marine turtles, *Conservation and Technical Data Report* 3: 1-46.
171. Meager, J.J. and Limpus, C.J. 2012b, Marine wildlife stranding and mortality database annual report 2011, I Dugong, *Conservation and Technical Data Report* 2011(1): 1-30.
172. Ayling, A.M. and Choat, J.H. 2008, *Abundance patterns of reef sharks and predatory fishes on differently zoned reefs in the offshore Townsville region*, Great Barrier Reef Marine Park Authority, Townsville.
173. Lawrence, K., van Putten, I. and Fernbach, M. 2010, *Profiles of recreational use of the Great Barrier Reef Marine Park*, Great Barrier Reef Marine Park Authority, Townsville.
174. Fernbach, M. and Nairn, K. 2007, *Reef recollections: An oral history of the Great Barrier Reef*, Great Barrier Reef Marine Park Authority, Townsville.
175. Department of Sustainability, Environment, Water, Population and Communities 2009, *Death or injury to marine species following capture in beach meshing (nets) and drum lines used in Shark Control Programs*, DSEWPA, Canberra, <<http://www.environment.gov.au/biodiversity/threatened/ktp/shark-control-programs.html#attachment>>.
176. Department of Agriculture, Fisheries and Forestry 2012f, *Shark control equipment and locations*, DAFF, Brisbane, <<http://www.daff.qld.gov.au/fisheries/services/shark-control-program/shark-control-equipment-and-locations>>.
177. Great Barrier Reef Marine Park Authority 2011c, *Annual report*, GBRMPA, Townsville, <<http://www.gbrmpa.gov.au/resources-and-publications/publications/annual-reports/>>.
178. Polglaze, Griffin, Miller and Associates 2012, *Great Barrier Reef shipping: review of environmental implications*, PGM Environment, Safety Bay, Western Australia.

179. GHD Pty Ltd 2013, *Ship anchorage management in the Great Barrier Reef World Heritage Area*, Great Barrier Reef Marine Park Authority, Townsville.
180. North-East Shipping Management Group 2013, *North-east shipping management plan (draft for consultation)*, Australian Maritime Safety Authority, Canberra.
181. Deloitte Access Economics 2012, *The economic contribution of the cruise sector to Australia*, Deloitte Access Economics, Canberra.
182. Office of Economic and Statistical Research 2012c, *Overseas export of goods by port value, Queensland Ports, 2001-02 to 2011-12*, Queensland Treasury and Trade, Brisbane, <<http://www.oesr.qld.gov.au/regions/mackay/tables/os-export-goods-port-qld-ports/index.php>>.
183. Hatch, L.T., Clark, C.W., Van Parijs, S.M., Frankel, A.S. and Ponirakis, D.W. 2012, Quantifying loss of acoustic communication space for right whales in and around a U.S. National Marine Sanctuary. *Conservation Biology* 26(6): 983-994.
184. Great Barrier Reef Marine Park Authority 2011d, *Grounding of the Shen Neng 1 on Douglas Shoal, April 2010: impact assessment report*, GBRMPA, Townsville.
185. Australian Maritime Safety Authority a, *Australian Ship Reporting records (AUSREP) for the calendar year 2000*, AMSA, Canberra.
186. Australian Maritime Safety Authority b, *Australian Ship Reporting records (AUSREP) for the calendar year 2010*, AMSA, Canberra.
187. URS Australia Pty Ltd 2006, *Strategic environmental assessment of defence activities in the Great Barrier Reef World Heritage Area*, Directorate of Environmental Stewardship, Department of Defence, Canberra.
188. Department of Defence 2013, *Defence White Paper*, Commonwealth of Australia, Canberra.
189. O'Neill, P. 2009, *State of the Environment Report for Shoalwater Bay Training Area 2008*, Department of Defence, Canberra.
190. Aurecon Australia 2012, *Talisman Sabre 2013: Public environment report (draft for consultation)*, Aurecon Australia, Canberra.
191. Townsville Enterprise 2012, *Townsville North Queensland Region: destination tourism strategy 2012-2016*, Townsville Enterprise, <<http://townsvilleenterprise.com.au/Files/232480%20%20DTS%20-%20Townsville%20North%20Queensland%20FINAL.PDF>>.
192. Australian Defence Force 2009, *Maritime activities environment management plan and supporting planning guides: Ship planning guides and procedure cards*, ADF, Canberra.
193. Fairfax Media 1947, *Dangerous tasks for men of the Navy*, 18 June, Sydney Morning Herald, Sydney.
194. Royal Australian Navy 2013, *HMAS Bungaree*, Royal Australian Navy, Canberra, <<http://www.navy.gov.au/hmas-bungaree>>.
195. News Ltd 1948, *Mine sweeping ends*, 7 August, The Cairns Post, Cairns.
196. Department of Defence 2002, *Royal Australian Navy detonates World War II mine on Great Barrier Reef*, DoD, Brisbane.
197. Great Barrier Reef Marine Park Authority 2012i, *Scientific research: information sheet*, GBRMPA, Townsville.
198. Woodley, S.J., Williams, D.M., Harvey, T. and Jones, A. 2006, *World Heritage research: making a difference. CRC Reef research, education and capacity building 1999-2006*, CRC Reef Research Centre, Townsville.
199. Lundgren, P. 2011, *Genetics and genetic tools in coral reef management: A synthesis of current research and its application in the management of coral reefs*, Great Barrier Reef Marine Park Authority, Townsville.
200. Brando, V.E., Schroeder, T., Blondeau-Patissier, D., Clementson, L. and Dekker, A.G. 2011, *Reef Rescue Marine Monitoring Program: using remote sensing for GBR wide water quality. Final report for 2010/11 activities*, CSIRO, Canberra.
201. Phinn, S., Dekker, A., Brando, V., Roelfsema, C. and Scarth, P. 2004, *MR2: Remote Sensing for Moreton Bay*, CRC for Coastal Zone, Estuary and Waterways Management, Brisbane.
202. Australian Research Council 2012, *National Competitive Grants Program Dataset*, Australian Research Council, Canberra, <[http://www.arc.gov.au/general/searchable\\_data.htm](http://www.arc.gov.au/general/searchable_data.htm)>.



## Chapter 6

## Impacts on the values





6.8.3	Assessment of outcomes arising from cumulative impacts .....	6–66
<b>6.9</b>	<b>Effects on matters of national environmental significance.....</b>	<b>6–67</b>
6.9.1	World heritage properties .....	6–67
6.9.2	Great Barrier Reef Marine Park .....	6–68
6.9.3	National heritage places.....	6–68
6.9.4	Commonwealth marine area .....	6–69
6.9.5	Listed migratory and listed threatened species.....	6–69
6.9.6	Wetlands of international importance .....	6–69
<b>6.10</b>	<b>Key information gaps .....</b>	<b>6–69</b>
6.10.1	Biodiversity .....	6–70
6.10.2	Geomorphological features .....	6–71
6.10.3	Indigenous and historic heritage values .....	6–71
6.10.4	Community benefits of the environment .....	6–71
6.10.5	Cumulative impact assessment .....	6–71
6.10.6	Processes to address information gaps .....	6–72
<b>6.11</b>	<b>Summary of conclusions .....</b>	<b>6–73</b>
<b>References</b> .....		<b>6–76</b>

## Table of contents

<b>6.1</b>	<b>Background .....</b>	<b>6–3</b>
6.1.1	Context .....	6–3
6.1.2	Challenges in assessing impacts.....	6–3
6.1.3	Terminology .....	6–4
<b>6.2</b>	<b>Impacts considered.....</b>	<b>6–4</b>
6.2.1	Identifying likely past and present impacts .....	6–4
<b>6.3</b>	<b>Legacy impacts .....</b>	<b>6–9</b>
<b>6.4</b>	<b>Impacts on biodiversity values .....</b>	<b>6–10</b>
6.4.1	Climate change impacts .....	6–10
6.4.2	Catchment run-off .....	6–15
6.4.3	Degradation of coastal ecosystems..	6–26
6.4.4	Direct use.....	6–29
6.4.5	Assessment of impacts on biodiversity.....	6–46
<b>6.5</b>	<b>Impacts on geomorphological features .....</b>	<b>6–49</b>
<b>6.6</b>	<b>Impacts on Indigenous and historic heritage values .....</b>	<b>6–51</b>
6.6.1	Indigenous heritage values.....	6–51
6.6.2	Historic heritage values .....	6–54
<b>6.7</b>	<b>Impacts on community benefits of the environment .....</b>	<b>6–54</b>
<b>6.8</b>	<b>Cumulative impacts .....</b>	<b>6–55</b>
6.8.1	Qualitative models .....	6–55
6.8.2	Spatial approaches.....	6–58

Cover page image: Aerial view of Townsville



**Extract from Great Barrier Reef Region Strategic Assessment terms of reference**

**3. Assessment of impacts on matters of national environmental significance**

**3.1 Actual and potential impacts**

- .....
- b) describe and analyse the actual and potential impacts on the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area, in the strategic assessment area, including:
    - i. impacts from past, present and future activities
    - ii. direct, indirect, consequential and cumulative impacts
    - iii. the likely impacts of climate change
  - c) describe the spatial and temporal scale at which impacts and their effects on the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area, are occurring or are likely to occur
  - d) identify key information gaps and processes to address critical information needs.

## **6.1 Background**

### **6.1.1 Context**

The Great Barrier Reef Region is a large and complex natural system with multiple uses that is subject to many impacts. The state of the Region's biodiversity, geomorphological features and Indigenous and historic heritage values, and the benefits derived from the environment, are constantly changing in response to a complex suite of interactions between drivers, activities and impacts.

To develop a full understanding of the impacts and their effects on the Region's values the individual and cumulative effects of all impacts are considered in this chapter. The focus is on the impacts on the ecological system (that is, its biodiversity values) as this is fundamental to the matters of national environmental significance in the Region. An assessment of the impacts on key biodiversity values informs the assessments of impacts on geomorphological features and Indigenous and historic heritage values. Flow-on effects on the community benefits derived from the environment are also described.

To date, understanding of cumulative impacts in the Region has been limited. The assessment of management effectiveness for the *Great Barrier Reef Outlook Report 2009* identified the extent to which cumulative impacts are being addressed as '*the weakest indicator across the entire management effectiveness assessment*'<sup>1</sup>. The independent review of the Authority's current management arrangements, carried out as part of the strategic assessment process (see Chapter 8), arrived at the same conclusion and further stated that processes to address cumulative impacts were '*problematic for most issues*'. Both qualitative and spatial modelling are employed in this chapter to consider cumulative impacts (Section 6.8).

### **6.1.2 Challenges in assessing impacts**

Specific challenges in assessing impacts, and cumulative impacts in particular, in the Great Barrier Reef Region include:

- the diversity of values and attributes underpinning matters of national environmental significance relevant to the Region
- the availability of information on the condition and trend of a number of values and attributes to allow development of a 'baseline' for the assessment
- the complex interactions, processes and relationships of the hugely diverse species, habitats and bioregions that make up the ecosystem
- the number of drivers, activities and impacts acting on the values and attributes
- limited understanding of the relationships between the ecological and human systems
- the multiple scales at which processes and impacts occur and interact with ecological and human systems
- limited understanding of the cause-and-effect relationships of multiple or 'compounding' impacts on the Region's values
- the dynamic nature of systems and the rapid pace of change
- the delayed response time of species, habitats and systems to change


- a limited understanding of the thresholds for species, habitats and ecosystem health.

### 6.1.3 Terminology

In describing and assessing impacts, the following types are considered, adapted from *A guide to undertaking strategic assessments*<sup>2</sup>:

- Direct impacts — where the loss or modification of values is a direct result of an action within the strategic assessment area (for example, dredging and disturbing wildlife).
- Indirect impacts — can be either:
  - from actions outside the strategic assessment area with ‘downstream’ effects in the area (for example, modifying supporting terrestrial habitats, urban and industrial discharge)
  - as a result of another direct impact (for example, an oil spill resulting from the grounding of a ship).
- Consequential impacts — where the impact arises from an action made possible by an initial direct impact (for example, anchor damage from ships now able to visit an area after dredging).
- Cumulative impacts — the successive and combined effects of impacts on the environment, taking into account direct, indirect and consequential impacts and the incremental and compounding effects of these impacts over time.

Direct, indirect and consequential impacts are diagrammatically represented in Figure 6.1.



**Figure 6.1 Direct, indirect and consequential impacts**

The strategic assessment considers all types of impacts on the Region's values. Using ports and shipping as an example: dredging to deepen a port channel is a direct impact, a plume of sediment from that dredging is an indirect impact, and any anchor damage to the seafloor as a result of a ship visiting the port following the dredging is a consequential impact.

## 6.2 Impacts considered

### 6.2.1 Identifying likely past and present impacts

The set of past, present and future impacts affecting or likely to affect the Region's values was developed by consolidating the outcomes of:

- the *Outlook Report 2009*<sup>1</sup>
- the analysis of drivers and activities within and adjacent to the Region presented in Chapter 5
- workshops involving managers and technical experts in relation to future monitoring needs for the Region.

The descriptors used are based on those used in the *Outlook Report 2009*, with some amendments and updates. The final 40 impacts considered are listed in Table 6.1 and matched to their likely causes, either direct drivers or activities, in Table 6.2.

Each of the impacts is individually assessed in relation to its effects on biodiversity (Section 6.4), geomorphological features (Section 6.5) and Indigenous and historic heritage values (Section 6.6). The flow-on effects on community benefits, including aesthetics, is discussed in Section 6.7. The focus is on the impacts on

the ecological system (that is, its biodiversity values) as this is fundamental to the matters of national environmental significance in the Region.

It is important to recognise that there is great variation in the spatial extent of the impacts — some operate in only a few areas within the Region, while others are, or have the potential to be, widespread. These variations are indicated in Table 6.1 and discussed throughout the assessment.

As demonstrated in Table 6.2, the past and present impacts can be grouped into four broad categories:

- climate change
- catchment run-off
- degradation of coastal ecosystems
- direct use.

Although some of the impacts fall into more than one category, they have been grouped this way to help structure the assessment.

The most significant legacy impacts on the Region from historic activities, both within the Region and the catchment, are described in Section 6.3. The continuing influence these impacts have on the condition of the Region's values is taken into account in the assessments later in the chapter.

**Table 6.1 Impacts considered in the assessment**

Note — Scale refers to Reef-wide: likely to cause effects through much of the Region; Regional: likely to cause effects in extended areas; and Local: likely to cause effects in small areas (for example a reef or bay).

Abbreviated title	Impact	Scale
Acid sulphate soils	Exposure and subsequent oxidation of potential acid sulphate soils	Local
Altered ocean currents	Altered ocean currents due to climate change or anomalies related to the El Niño-Southern Oscillation, and altered coastal water movement at a local scale	Reef-wide
Artificial barriers to flow	Artificial barriers to riverine and estuarine flow including breakwalls, weirs, dams, gates, ponded pastures, and weeds causing changes to hydrology, groundwater and ecological connectivity	Regional
Atmospheric pollution	Atmospheric pollution, including coal dust	Local
Coastal reclamation	Coastal land reclamation, including for ports and groynes	Local
Cyclone activity	Cyclone activity	Regional
Dredging	Dredging of the seafloor	Local
Disposal and resuspension of dredge material	Sea dumping of dredge material including smothering, loss and modification of seabed habitats and resuspension	Local
Exotic species and diseases	Introduction of exotic species and diseases from aquaculture operations, hull fouling, ballast release, imported bait and release of aquarium specimens to the Region, plus the introduction of weeds and feral animals to islands	Regional
Extraction — death of discarded species	Death of non-retained species from fishing, collecting, hunting, scientific sampling and Queensland's Shark Control Program	Reef-wide
Extraction — fishing in spawning aggregations	Fishing in unprotected fish spawning aggregations	Regional
Extraction — herbivores	Retained take (extraction) of herbivores through commercial, recreational and traditional use	Reef-wide
Extraction — lower order predators	Retained take (extraction) of lower order predators (e.g. coral trout and snapper) through commercial, recreational and traditional fishing	Reef-wide
Extraction — lower trophic orders	Retained take (extraction) of lower trophic orders (e.g. scallops, sea cucumbers and prawns) through commercial, recreational and traditional fishing	Reef-wide
Extraction — top order predators	Retained take (extraction) of top order predators (e.g. sharks) through commercial, recreational and traditional fishing and the Queensland Shark Control Program	Reef-wide
Illegal fishing and poaching	Illegal fishing, collecting and poaching (foreign or domestic) including of species of conservation concern	Reef-wide
Increased freshwater inflow	Increased freshwater inflow from prolonged or heavy rainfall including flood events, and from changes to catchment ecosystems; resulting in reduced salinity	Regional
Increased sea temperature	Increased sea temperature due to climate change	Reef-wide
Light impacts (artificial)	An increased amount of artificial light	Local

Abbreviated title	Impact	Scale
Marine debris	Manufactured material discarded, disposed of or abandoned in the marine and coastal environment (including discarded fishing gear and plastics)	Reef-wide
Mining	Whilst mining of the Great Barrier Reef was halted 50 years ago and subsequent coral growth has obliterated most impacts, mining on high and low islands for guano based phosphate rock has a continuing impact. In addition, many of the significantly important beach ridge sequences are seen as a source of building materials by developers	Local
Modifying supporting terrestrial habitats	Clearing or modifying supporting terrestrial habitats such as wetlands, saltmarshes, mangroves and sand dunes — this also includes trampling and damage from recreational vehicle use	Regional
Noise pollution	Noise from human activities, both below and above water	Local
Nutrients from catchment run-off	Nutrients entering the Region in run-off from the catchment	Reef-wide
Ocean acidification	Increasing acidity of the Region's waters	Reef-wide
Outbreak of crown-of-thorns starfish	Outbreak of crown-of-thorns starfish (i.e. when the density exceeds about 30 starfish per hectare)	Regional
Outbreak of disease	Outbreak of naturally occurring diseases	Local
Outbreak or bloom of other species	Outbreak of naturally occurring or native species, excluding crown-of-thorns starfish	Local
Pesticides from catchment run-off	Pesticides (including herbicides, insecticides, fungicides) entering the Region in run-off from the catchment	Regional
Physical damage — fishing	Physical damage to marine habitats from fishing, such as the effects of trawling on or near the seabed	Reef-wide
Physical damage — other	Physical damage to coral reefs and seafloor habitats including from anchoring of vessels of any size, grounding of small vessels, diving and snorkelling. Does not include physical damage for fishing or ship grounding	Local
Physical damage — ship grounding	Grounding of ships including physical damage and the dislodging of antifoulants	Local
Rising sea level	Rising sea level	Reef-wide
Sediments from catchment run-off	Sediments entering the Region in run-off from the catchment	Reef-wide
Spill — large chemical	Chemical spills that trigger a national or regional response or are more than 10 tonnes	Regional
Spill — large oil	Oil spills that trigger a national or regional response or are more than 10 tonnes	Regional
Spill — small chemical and oil	Chemical and oil spills that do not trigger a national or regional response and are less than 10 tonnes	Local
Urban and industrial discharge	Point and diffuse-source land-based discharge of pollutants from urban and industrial land use and mining, including polluted water, sewage, wastewater and stormwater	Local
Vessel strike on wildlife	Death or injury to wildlife as a result of being struck by a vessel of any type or size	Local
Waste discharge from vessels	Waste discharged from a vessel into the marine environment	Local
Wildlife disturbance	Disturbance to wildlife including from snorkelling, diving, fish feeding, walking on islands and beaches, and the presence of boats	Local



### Impacts identified by Traditional Owners and stakeholders

During development of the strategic assessment, the Authority sought advice from a range of stakeholders and Traditional Owners (see Section 2.11 and Appendix 5) regarding impacts on the Region's values.

Participants at stakeholder and Traditional Owner workshops identified the impacts considered to be affecting the Region's values. Those mentioned consistently across all workshops were:

- climate change
- extreme weather events
- declining water quality
- coastal development, including urban development and increased industrialisation of the coast
- port development and associated increases in shipping
- mining
- overfishing.

Key impacts identified by Traditional Owners in particular included:

- dispossession from country and loss of access to traditional grounds
- unauthorised entry to sites of cultural significance
- coastal development
- vandalism, including damage and graffiti
- marine debris and litter
- ships and boats cutting across storylines
- over fishing
- declining water quality
- extreme weather events.

In a follow-up survey completed by 118 participants, climate change was rated as the most serious threat by 34 per cent of respondents, water quality by 23 per cent, while 12 per cent considered extreme weather the most serious.



*In 2011 cyclone Yasi, an extreme weather event, produced damaging winds and waves that affected much of the coastline between Cairns and Townsville.*

**Table 6.2 Impacts and their sources**

Broad categories used in the assessment are shown (on left); however impacts may fall into more than one category.

\* indicates impact is likely to be the result of a variety of activities.

	Impacts	Direct driver	Activities in the Region													
			Activities in the catchment													
			Climate change	Agriculture	Aquaculture	Urban development	Industrial development	Port activities	Traditional use	Tourism	Fishing — commercial	Fishing — recreational	Recreation	Shipping	Defence activities	Research activities
Climate change	Altered ocean currents	●			●		●									
	Cyclone activity	●														
	Increased sea temperature	●														
	Ocean acidification	●														
	Rising sea level	●														
Catchment run-off	Increased freshwater inflow	●	●		●	●										
	Nutrients from catchment run-off		●	●	●	●										
	Outbreak of crown-of-thorns starfish		●		●	●				●	●					
	Pesticides from catchment run-off		●		●	●										
	Sediments from catchment run-off		●	●	●	●	●									
	Urban and industrial discharge			●	●	●	●									
Degradation of coastal ecosystems	Acid sulphate soils		●	●	●	●	●									
	Artificial barriers to flow		●	●	●	●	●									
	Atmospheric pollution		●		●	●	●						●			
	Coastal reclamation				●	●	●		●	●	●	●				
	Light impacts (artificial)				●	●	●						●			
	Modifying supporting terrestrial habitats		●	●	●	●	●		●	●	●	●				
Direct use	Dredging				●	●	●							●		
	Dumping and resuspension of dredge material						●							●		
	Exotic species and diseases*			●					●	●	●	●	●			
	Extraction — death of discarded species									●	●					●
	Extraction — fishing in spawning aggregations									●	●					
	Extraction — herbivores							●		●	●					●
	Extraction — lower order predators							●		●	●					●
	Extraction — lower trophic orders							●		●	●					●
	Extraction — top order predators							●		●	●					●
	Illegal fishing and poaching									●	●					
	Marine debris*		●		●	●	●		●	●	●	●	●			
	Noise pollution		●			●	●							●	●	●
	Outbreak of disease*		●				●									
	Outbreak or bloom of other species*		●											●		
	Physical damage — fishing									●						
	Physical damage — other							●	●	●	●	●	●	●	●	●
	Physical damage — ship grounding								●	●				●		
	Spill – large chemical				●	●	●							●		
	Spill – large oil					●	●							●		
	Spill – small chemical or oil				●	●	●		●	●	●	●	●	●		
	Vessel strike on wildlife						●		●	●	●	●	●	●	●	
	Waste discharge from vessels							●	●	●	●	●	●	●	●	●
	Wildlife disturbance						●		●	●	●	●	●	●	●	

## 6.3 Legacy impacts

### Direct legacy impacts

Some activities previously undertaken within what is now the Great Barrier Reef Region and on the islands have had severe and long-lasting impacts on the values of the Region (Table 6.3). The legacy of these past activities is still affecting the Reef ecosystem and contributing to cumulative impacts in the Region.

Of the past activities directly undertaken in the Region, the most significant legacy impacts have resulted from large-scale commercial harvesting, especially of long-lived species. Past commercial harvesting of dugongs, green turtles, crocodiles and humpback whales has had major effects on the populations of these species. Historic reductions in dugong, green turtle and hawksbill turtle populations have substantially affected those species' ability to bounce back from more recent impacts.<sup>3,4,5</sup>

The east Australian humpback whale population is now showing strong recovery more than 50 years after harvesting stopped, and it is estimated that the population may be 50 to 75 per cent of its pre-whaling size.<sup>6,7</sup>

**Table 6.3 Past activities resulting in direct legacy impacts on the Region's values**

Activity		Time period
Commercial harvesting	Harvesting humpback whales	1952–1962 <sup>8</sup>
	Harvesting dugong — meat, bones, hide, oil	1847–1969 <sup>8</sup>
	Harvesting hawksbill turtles — tortoise shell	1871–1940s <sup>5,8</sup>
	Harvesting green turtles	1867–1962 <sup>4,8</sup>
	Farming green and hawksbill turtles	1970–1979 <sup>8</sup>
	Harvesting estuarine crocodiles	1945–1974
	Trawling without bycatch reduction devices	1950s–2001
	Trawling sensitive lagoon floor communities <sup>9</sup>	1960s–2000
Physical destruction of coral	Mining coral for limestone	1900–1940 <sup>8</sup>
	Dynamiting coral reefs <sup>8</sup>	Unknown, but pre-1970s
	Constructing transit channels across reef flats <sup>8</sup>	Unspecified
Collecting (commercial and recreational)	Unrestricted collecting of coral	Ceased in 1933 <sup>10</sup>
	Collecting shells — pearl shell, trochus	From late 1800s <sup>8</sup>
	Collecting seabird eggs	Unspecified
	Collecting black teatfish	Ceased in 1999 <sup>11</sup>
Islands	Deliberate introduction of exotic plants and animals on islands, e.g. goats for hunting and food <sup>8</sup>	Unknown, but pre-1980s
Mining and exploration	Mining guano and rock phosphate	1860–1940 <sup>8</sup>
Defence	Widespread activities during World War II and subsequent bombing and shelling during military target practice	1943–1965 <sup>8</sup>

Dugongs were commercially harvested from 1847 until 1969 in the Moreton Bay area (south of the Region) and for shorter periods at other locations on the Queensland coast, including sites within the Region.<sup>12</sup> In the first 50 years, the number taken was so high that the industry was initially closed in 1890 due to a scarcity of dugong.<sup>12</sup> This, combined with large harvests especially between the 1930s and 1969, has increased the vulnerability of dugongs to subsequent impacts.<sup>3</sup> Populations have subsequently failed to recover due to a combination of conservative life history traits and exposure to multiple impacts either directly and through habitat degradation.

Commercial use of green turtles began in the 1860s and continued until 1962. The harvest was primarily to supply turtle meat and soup for export. It was focused in the Capricorn Bunker group, but occurred as far north as Raine Island.<sup>12</sup> It is estimated thousands of mature females were taken during this period.<sup>12</sup> Given green turtles are long-lived, late maturing and display a high degree of fidelity to their foraging grounds and nesting areas, this species is highly vulnerable to overharvesting.<sup>4</sup> Long-term monitoring of the green turtle nesting population in the southern Great Barrier Reef shows this stock is now recovering, increasing at an average of about 3.8 per cent per year for the past four decades.<sup>13</sup> However, the northern Great Barrier Reef stock is at best stable and may be showing

early signs of a population in decline.<sup>4</sup> The northern stock is mainly affected by harvesting of adult and near-adult turtles throughout much of the foraging range, as well as climate and habitat-related loss of hatchling production.<sup>4</sup>

Hawksbill turtles were also harvested commercially for many decades, primarily as a source of tortoiseshell. This activity was concentrated in the northern Great Barrier Reef where hawksbill turtles were abundant. By 1900, the hawksbill turtle had been already heavily exploited by Europeans; harvesting has had lasting impacts on the species.<sup>5,12</sup> Current population trends for the Torres Strait–northern Great Barrier Reef hawksbill turtle stock suggests the stock has been declining at three to four per cent each year since at least 1990, which projects to an expected reduction of greater than 80 per cent in the breeding female population in less than one generation.<sup>5</sup> The most obvious large source of mortality is the continuing harvest of hawksbill turtles for food and tortoiseshell in the broader Coral Sea region.<sup>14</sup>

Otter trawling in the Region dates back to the 1950s. The effect of high intensity prawn trawling may be substantial, locally removing about 70 to 90 per cent of seabed animals.<sup>15</sup> Although very few areas of the Region were fished so intensively, scientific evidence shows historical patterns and the amount of trawl fishing effort resulted in substantial effects and changes to seabed habitats and species at a Great Barrier Reef-wide scale.<sup>9,15,16</sup> Science also shows that over the last decade these impacts have been arrested and reversed by a sequence of management interventions implemented by the Queensland Government and the Authority.<sup>17</sup>

Historically, otter trawling also caused marine turtles to drown in the trawl nets. This activity was one of the major causes of marine turtle mortality in the Region and contributed to their population decline (especially for the endangered loggerhead turtle).<sup>18</sup> The mandatory use of turtle excluder devices since the early 2000s has largely mitigated this impact.<sup>1,17,18,19</sup>

### **Indirect legacy impacts**

Past broadscale land clearing, principally in the southern two-thirds of the Great Barrier Reef catchment, has indirectly caused major changes to the Region's environment, affecting habitats and species. Clearing began in the 1870s and was undertaken to allow more intensive agricultural use. It further increased when intensive cropping on the coastal floodplain began in the early 1900s and again in the 1930s and 1940s when heavy machinery made clearing easier. The rate of clearing continued to increase until the late 1990s when controlling legislation was introduced.<sup>20</sup>

The resultant loss and modification of habitats has led to significant increases in pollutants, principally nutrients and sediments, entering the Great Barrier Reef lagoon.<sup>20</sup> Modelling and ongoing monitoring of habitats indicates a reduction in the ecosystem's ability to bounce back after impacts (resilience), especially in southern inshore areas.<sup>21,22</sup> In addition, freshwater coastal habitats are important to numerous marine species, including the freshwater sawfish which is now threatened, in part, due to habitat loss.<sup>23</sup>

The previous construction of dams (for example, Fairbairn Dam in the 1970s and Burdekin Falls Dam in the 1980s) has led to lasting changes in the hydrological processes in the Great Barrier Reef catchment, in turn affecting the Region's values and reducing connectivity between coastal and marine environments.<sup>24</sup> Levees, bunds, drains and other impediments to flow have also had negative legacy effects on habitats, species, connectivity and hydrology.<sup>24</sup>

## **6.4 Impacts on biodiversity values**

This section contains an analysis of how each impact identified in Table 6.2 affects the Region's biodiversity values as set out in Chapter 4. It is based on the best available information, including input from stakeholders and Traditional Owners. It concludes with an assessment of the relative effect of each impact on the Region's biodiversity values.

### **6.4.1 Climate change impacts**

#### **Altered ocean currents**

Ocean currents play an important part in regulating heat, connecting biodiversity and supporting productivity.<sup>25</sup> They vary naturally and are now also being influenced by climate change.<sup>26</sup> The effects of such changes could be Reef-wide.

There is increasing evidence of intensified flow and accelerated warming in the East Australian Current adjacent to the Region's southern coast.<sup>27</sup> This current is transporting greater volumes of warmer water southward. There is little information about the Hiri Current travelling north along the coast in northern Great Barrier Reef waters.<sup>25,27,28</sup>

Changes to ocean currents have the potential to affect entire marine food webs from corals and sponges to top predators such as sharks. Altered ocean circulation patterns may affect the transport of eggs and larvae within and among coral reefs and other Great Barrier Reef habitats. In addition, the changes may affect the movement of hatchling marine turtles away from nesting beaches. Pelagic foraging seabirds are also highly vulnerable to changes in ocean currents, as many search for food in pelagic communities that follow upwellings, eddies and other sea surface temperature gradients.<sup>29</sup> Changes may also result in species being more widely distributed or occurring in different areas.




At a local scale, currents can be altered by coastal infrastructure (for example, groynes and marinas), and other coastal works (for example, coastal reclamation, beach replenishment, dredging and dumping of dredge material).<sup>30</sup> Historically, the development and location of coastal works and infrastructure has affected coastal water movement which has resulted in negative effects on beaches and islands.<sup>31</sup>

### Cyclone activity

Cyclones are a natural part of the weather cycle in tropical areas. However, the global climate system is now warmer and moister than it was 50 years ago, and this is increasing the chances of more extreme weather events.<sup>32</sup> As cyclone activity is influenced by environmental conditions such as sea surface temperature, climate change predictions suggest an increase in the intensity of cyclones in Queensland and an extension of their southern range.<sup>33</sup> However, the overall frequency of all categories of cyclones is predicted to decrease.<sup>34</sup> Projections for the period 2051 to 2090, relative to 1971 to 2000, suggest an approximate 50 per cent decrease in frequency, a small decrease in duration (0.3 days), and a 100 kilometre southward movement in where they occur.<sup>34</sup>

Over the last decade, particularly between 2005 and 2012, there have been a number of severe cyclones in the Region (Figure 6.2).




**Figure 6.2 Cyclone winds (2005–2011) and flood plume exposure (1991–2011) in the Great Barrier Reef Region**

The cluster of cyclones and flood events in recent years has seriously affected the condition of many Great Barrier Reef habitats and species. The likelihood of flood plume exposure (brown areas) is a cumulative assessment of multiple flood plumes based on remotely sensed conditions at the sea surface. The flood plume extent for 2010–11 (brown line) indicates the distribution of the flood plume as a result of the extreme weather events experienced over that summer.

Severe tropical cyclones can damage the Region's habitats such as coral reefs on a regional scale.<sup>35</sup> It is estimated that they have been responsible for about half of the total coral cover loss since 1985.<sup>36</sup> One effect has been high intensity waves damaging underwater habitats and coastlines (Figure 6.3). Most of the Region is likely to have experienced damaging waves at least once during the last decade; the area between Cairns and Townsville has been particularly affected (Figure 6.3).

Tropical habitats such as coral reefs, seagrass meadows and islands have a natural resilience to physical disturbances such as cyclones. Severely damaged reefs have shown strong signs of recovery within about five to 15 years if there are few other stresses to impede regrowth, recruitment and survivorship of corals.<sup>37,38,39</sup> It is thought the effects of cyclones on vulnerable habitats are becoming more severe and persistent due to a reduction in the resilience of the Reef.<sup>40,41</sup> This has flow-on effects to other parts of the ecosystem.




**Figure 6.3 Cyclone wave exposure in the Region, 2000–2011<sup>42</sup>**

The extent and severity of cyclone wave damage to corals and other biota (e.g. seagrasses) depends on the size and duration of cyclone generated waves, exposure of organisms to those waves, and the vulnerability of the organisms to wave action.<sup>43</sup> Therefore, the significant wave height (the average size of the largest one-third of the waves in a given sea) needed to damage coral or other biota varies by species and area.<sup>44</sup> For the Region, a significant wave height of four metres is likely to damage most species, and is used here. The map represents the probability that four metre significant wave heights formed for at least an hour in a given year over the period 2000 to 2011. Gradings are scaled between 0 and 1, based on a maximum yearly probability of 31.7 per cent: low (less than 11 per cent chance), medium (11 to 22 per cent chance) and high (22 to 31.7 per cent chance).

### Increased sea temperature

In line with global ocean trends, surface waters in the Coral Sea, including the Region, have warmed substantially over the last century<sup>33,45</sup> and will likely continue to rise over the coming century.<sup>46</sup> The average summer (December to February) sea surface temperature anomalies (when the temperature is above the 1982 to 2000 summer maximum) in the Region for the 10 year period to 2011 for non-reef areas are presented in Figure 6.4.



**Figure 6.4 Summer sea surface temperature anomalies in the Region, 2001–2011<sup>47</sup>**

The map was developed by scaling the sums of the average annual positive sea surface temperature anomalies for summer periods (December to February) between 2001 and 2011, for non-reef areas. Values are scaled from 0 to 1, based on the maximum value of 68.79 degree heating days.

Average ocean temperatures around Australia have warmed by 0.68 degrees Celsius since the period 1910 to 1929.<sup>46</sup> Within the 102-year instrumental record, 15 of the 20 warmest years have been in the past 20 years.<sup>46</sup> For many areas of tropical Australia, a new high for recorded sea surface temperatures occurred during October to December 2010.<sup>46</sup> Further records were set in the summer of 2012–13 when the hottest sea surface temperatures for the Australian region were recorded.<sup>32</sup> Whatever climate scenario is used, it is predicted that by 2035 the average sea surface temperature will be warmer than any previously recorded, and by 2100 sea temperatures off north-eastern Australia could be about 2.5 degrees Celsius warmer than the present average.<sup>46</sup>

Increasing sea temperatures are already affecting the Great Barrier Reef.<sup>48,49,50,51,52,53,54,55</sup> Further increases pose significant risks over the coming decades across the whole Region, influencing a range of physical, chemical and biological conditions and processes and, hence, many different habitats and species.<sup>48</sup> As temperatures rise past the levels at which an organism's metabolic processes perform best, physiological efficiency decreases. This can cause stress and increase susceptibility to disease and predation. Higher temperatures also reduce the amount of



oxygen available within the water. Projected increases in ocean stratification — the vertical layers in the water column — are predicted to affect the supply of nutrients and oxygen into deeper pelagic and benthic (seafloor) ecosystems with implications for their biodiversity.<sup>49</sup>

There have been a number of widespread coral bleaching events in the Great Barrier Reef. The two most severe coral bleaching events spanned the summers of 1997–98 and 2001–02<sup>52,53,56</sup>, resulting from prolonged elevated sea temperatures. During both events, more than 50 per cent of reefs were affected to some degree by bleaching, with lasting damage to an estimated five per cent of reefs.<sup>53</sup> In 1997–98, coral bleaching affected reefs around the world. This coincided with observations of 500-year-old corals dying and, what was (at the time), the highest sea surface temperatures ever recorded on the Great Barrier Reef.<sup>52</sup> Another localised, but severe, bleaching event occurred in 2006 in the Keppel Bay region in the southern Great Barrier Reef. During this event, 40 per cent of corals died, but some recovery has since been observed.<sup>38</sup>

A future predicted increase of two degrees Celsius in the average sea temperature will likely lead to annual bleaching, with up to 97 per cent of reefs affected and almost certain regular large-scale coral mortality.<sup>55,57</sup>

### Ocean acidification


A decline of 0.1 units in the pH of Great Barrier Reef waters has already been recorded, and further declines are expected. From a current pH of 8.2 (alkaline), it is predicted that the ocean could fall to a pH of about 7.6 by 2100, with slight regional variation.<sup>49</sup> While rates of decline are greater in southern areas of the Region, it is predicted that the ecosystem will be affected on a Reef-wide scale.

The effects of global warming and ocean acidification may magnify each other<sup>58</sup> and may not occur uniformly from place to place and over time.<sup>59</sup> Regardless of the rate of change, even relatively small increases in ocean acidity reduce the capacity of corals to build skeletons, which in turn reduces their capacity to create habitat for reef biodiversity in general.<sup>60</sup> Field observations at natural carbon dioxide seeps have found more acidic oceanic conditions (a decline in pH from 8.1 to 7.8) do not necessarily affect coral cover but reduce species diversity and structural complexity, and increase macroalgae and seagrass cover.<sup>61</sup>

The most immediate effects of ocean acidification may be on crustose coralline algae, a species vital to reef building. Increasing acidity is likely to reduce the capacity of coralline algae to cement reef debris into solid limestone, and in turn affect coral recruitment and establishment.<sup>62</sup> Other species such as molluscs, phytoplankton and foraminifera will also be affected by acidification.

### Rising sea level

Sea levels are rising because of a combination of thermal expansion of the ocean and the addition of water volume to the ocean from melting glaciers and the ice sheets of Greenland and Antarctica. Globally, the sea level has risen more than 200 millimetres since 1880, and the rate of rise is increasing<sup>63</sup> (Figure 6.5). Sea level rose at a globally-averaged rate of about 1.7 millimetres per year during the 20th century as a whole and about 3 millimetres per year between 1993 and 2011.<sup>63</sup> Around Australia, the fastest rates of sea level rise are in northern Australia (Figure 6.6). In the Region, it continues to rise by about 3.1 millimetres per year on average.



**Figure 6.5 Global average sea level, 1880–2011<sup>64</sup>**

Dark blue line shows sea level records collected from tide gauges with light blue shading that provides an indication of accuracy.

Future responses of sea level to temperature increases are difficult to accurately predict. The Intergovernmental Panel on Climate Change predicts a rise of between 18 and 79 centimetres by 2095, from a 1990 baseline.<sup>65</sup> Sea levels are currently rising at near the upper end of these projections.<sup>63</sup>

Even modest rises in sea level may have substantial consequences throughout the Region when combined with natural variability arising from the El Niño Southern Oscillation and the Pacific Decadal Oscillation.



Rises in sea level are important for the Great Barrier Reef ecosystem as some habitats are shallow and strongly influenced by sea level. In particular, because much of the land adjacent to the Great Barrier Reef is low lying, small changes in sea level will mean increased erosion and land inundation, causing significant changes in tidal habitats such as mangroves, and saltwater intrusion into low lying freshwater habitats.<sup>1,66</sup> Brackish saltmarsh habitats are being displaced by mangroves.<sup>67</sup> Turtle nesting beaches are particularly vulnerable to rising sea levels which exacerbate beach erosion<sup>68</sup> and inundate nests. Seabird nesting and shorebird roosting sites are also at risk.<sup>69</sup>




Figure 6.6 Rates of sea level rise in Australia<sup>64</sup>

#### 6.4.2 Catchment run-off

##### Increased freshwater inflow

Freshwater inflow to the marine environment from heavy or prolonged rainfall and flood events can reduce salinity and have a range of effects on marine life on a regional scale. Freshwater input is generally higher in the region south of Cooktown, corresponding with the larger catchments. Recent La Niña conditions resulted in increased average annual rainfall and flood events between 2008 and 2012 (Figure 6.7). Much of the inshore areas of the Region have been exposed to freshwater over the last decade (Figure 6.8).





Figure 6.7 Annual freshwater discharge from the major rivers of the Great Barrier Reef catchment, 2002–2012

Much greater volumes of freshwater entered the Great Barrier Reef lagoon between 2008 and 2012 compared to previous years. Higher than average annual freshwater discharges were recorded for many of the major rivers, especially in southern catchments. In 2011, discharge volumes in the Fitzroy and Proserpine Rivers were the largest ever recorded. In the Herbert River, the volume was equal to the biggest ever recorded, while the Burdekin River experienced the third biggest. For each year, the inflow is for the 12-month period starting in October of the previous year. Data for 2012 is up to August of that year. (Figure compiled from data provided by the Australian Institute of Marine Science<sup>70</sup>).



**Figure 6.8 Freshwater exposure in the Region, 2001–2011<sup>47</sup>**

Observed frequency of freshwater plumes between 2001 and 2011 modelled from remotely sensed concentrations of dissolved organic matter (a proxy for freshwater), provided by CSIRO Land and Water. Gradings are based on the number of times a freshwater plume with a measured concentration of salinity less than 30 (+/- 4) parts per thousand was observed in any given year over the 10 year period. Gradings were expressed as: low (1–3 events), medium (4–6 events) and high (7–10 events) where the maximum frequency of events observed was 10.

Across Australia, heavy rainfall events are likely to become more frequent as the temperature increases<sup>32</sup>, with a tendency for more abnormally large freshwater inflows to the marine environment.

In a natural system, freshwater wetlands and the forested floodplain capture and slow freshwater flows. Changes to catchment ecosystems, such as infilling wetlands and constructing levee banks and drainage, have had major effects on the quantities of freshwater entering coastal seas.<sup>71</sup> At the same time, the clearing of forests and woodlands, and their replacement with grasslands for grazing, have substantially affected overland and groundwater flow.

Activities such as agriculture, urbanisation and industrial development can increase the magnitude and timing of freshwater inflows into the Region. For example, roads, stormwater drains and other urban infrastructure can increase the volume and speed of freshwater inflow, compared to natural vegetation and soil which retard water movement.<sup>72</sup> Stormwater drains release large quantities of fresh water into the sea that would normally percolate through the groundwater along shorelines and in riparian areas.

Generally, natural freshwater flows enhance estuarine productivity, provide connectivity between freshwater habitats and the sea, and improve fish recruitment and growth.<sup>73</sup> Maintaining natural freshwater flows can have positive effects, for example on breeding and recruitment of estuarine and marine fish of commercial and recreational value including barramundi and king threadfin.<sup>74,75</sup> Abnormally large freshwater inflows can have negative effects, for example low salinity bleaching and mortality in corals<sup>76</sup> or widespread damage to seagrass meadows<sup>41</sup>.

Increased freshwater inflow during flood events also carries with it pulses of nutrients, sediments, pesticides and other pollutants from catchment run-off, which has significant effects on Great Barrier Reef habitats and species<sup>77</sup> (see relevant impacts in this chapter).

### **Nutrients in catchment run-off**

The majority of coral reef habitats in the Great Barrier Reef evolved over thousands of years in nutrient-poor waters. The symbiosis between corals and the algae that live inside them evolved as a response to this lack of nutrients, allowing corals to harvest energy from photosynthesis by the algae.

The changes in land use since European settlement, including clearing and the application of fertilizers, have significantly altered nutrient concentrations in the inshore waters of the Great Barrier Reef lagoon to the extent that it is one of the most significant factors affecting Reef health.<sup>1</sup>


Dissolved inorganic nitrogen and phosphorous continue to enter the Great Barrier Reef ecosystem at greatly enhanced levels compared to those prior to European settlement. Estimates suggest the total nitrogen discharge into the Great Barrier Reef ecosystem has increased from 14,000 tonnes annually before European settlement (late 1800s) to 80,000 tonnes annually in the early 2000s — an almost six-fold increase, ranging from a doubling in some catchments to an almost twenty-fold increase in others.<sup>78</sup> Similarly, it is estimated that almost nine times more phosphorus is entering the ecosystem compared to pre-European settlement, with some catchments as high as 30 times more.<sup>78</sup>

Sources for nutrients into the Great Barrier Reef system include river discharges<sup>79</sup>, urban run-off<sup>80,81</sup>, atmospheric input<sup>82</sup>, nitrogen fixation by marine organisms<sup>83</sup>, deep ocean supply from upwellings<sup>84</sup>, deposition of dust from storms and wind<sup>85</sup>, and resuspension of nearshore sediments<sup>86,87</sup>. Of these, the single largest source is river discharges<sup>88</sup>, largely driven by the application of fertilisers and the subsequent loss of dissolved inorganic nutrients into the water column. Nutrients are also transported as part of the sediment load bound to particulates (particulate nutrients).<sup>89</sup> More than 90 per cent of these river discharges occur during the wet season.<sup>81,90,91</sup>

While the coarser-grained sediments and their bound particulate nutrients initially settle out of the water column close to river mouths, nutrients bound to clay and sediment fractions travel much further and dissolved nutrients disperse widely in the Great Barrier Reef lagoon.<sup>89,92</sup>

Changes to catchment ecosystems from human activities can also contribute to nutrients in catchment run-off through altering processes such as nutrient and chemical cycling.

Nutrients in the marine environment can be assessed by measuring chlorophyll concentrations, as the amount of planktonic algae containing chlorophyll in the water column is proportional to nutrient concentrations. Monitoring and modelling indicate that chlorophyll concentrations in up to 10 to 15 per cent of the Region have exceeded the Water Quality Guidelines for the Great Barrier Reef Marine Park in the last decade.<sup>93,94</sup> For much of the southern nearshore environment, concentrations are frequently above the annual guidelines (Figure 6.9), with some areas more than double.



**Figure 6.9 Years that chlorophyll concentrations exceeded the guidelines in the Region, 2003–2012<sup>103</sup>**

The water quality guidelines for the Great Barrier Reef Marine Park use chlorophyll concentration as an indicator for nutrient concentrations in open waters. The guideline trigger value is an annual mean of 0.45 micrograms of chlorophyll per litre. The map shows the number of years that the guideline value was exceeded between 2003 and 2012.

Once dissolved inorganic nutrients enter the marine system, they are taken up by phytoplankton, bacteria and benthic organisms such as macroalgae and seagrasses. This promotes growth in these organisms which in turn affects a variety of other species, habitats and processes. Imbalances in the nutrient cycle are having severe consequences for many of the Region's values. For example:


- Nutrients in flood discharges result in extensive, observable phytoplankton blooms and likely shifts in the species composition of phytoplankton.<sup>89,95</sup>
- There is strong scientific evidence that an increase in the frequency of crown-of-thorns starfish outbreaks is linked to increased nutrients in the open water. More nutrients lead to greater concentrations of phytoplankton, the food source of the planktonic larval stage of the starfish (Figure 6.10)<sup>96,97</sup>
- The complex relationship between corals and macroalgae is affected by nutrient status. Higher nutrient concentrations may contribute to an increase in macroalgal abundance and a decrease in coral diversity. This drastically affects the overall resilience of the ecosystem, as a dominance of macroalgae reduces the chance for hard corals to recruit, establish and grow, which perpetuates the shift towards an algae-dominated ecosystem.<sup>98,99</sup>



- There is evidence that elevated nutrient concentrations may make corals more sensitive to temperature stress.<sup>100,101</sup>
- While seagrasses may benefit from small increases in nutrients in the water column, elevated concentrations can be detrimental to seagrass health and resilience. Lowered ambient light levels, as a result of increased growth of phytoplankton, macroalgae and algal epiphytes that compete for light, can reduce photosynthesis in seagrass, particularly in deeper waters.<sup>102</sup>

A key target of the *Reef Water Quality Protection Plan 2013*<sup>104</sup> (Reef Plan) is to achieve a 50 per cent reduction in nutrient loads entering the Great Barrier Reef by 2018. Total fertiliser use on farming lands in the catchment has been reduced in recent years (see Figure 5.10), and recent monitoring and modelling show current initiatives are successfully reducing nutrient concentrations in catchment run-off. The Reef Plan Report Card for 2011 reports that, through implementation of best management practices in sugar, grazing and horticulture industries, over the previous two years there has been a reduction in the sediment load of six per cent, the total nitrogen load of seven per cent and the pesticide load of 15 per cent entering the Great Barrier Reef.<sup>105</sup> Early evidence shows reductions in the nutrient load entering the marine system has resulted in reduced nutrient concentrations in open waters.<sup>106,107</sup>

Elevated levels of another nutrient, dissolved organic carbon, can accelerate the growth of microbes on corals, which can cause coral death by oxygen depletion, accumulation of poisons and/or microbial predation on weakened coral.<sup>108</sup> In fact, it has been suggested that dissolved organic carbon causes more coral death than other more routinely measured water quality components (for example, nitrogen, phosphorus and ammonia).<sup>108</sup>



**Figure 6.10 Potential role of nutrients in the population dynamics of crown-of-thorns starfish**

Crown-of-thorns starfish are a major cause of loss of coral cover. One line of evidence suggests that their populations are significantly affected by the concentration of nutrients and, therefore, the amount of phytoplankton in Great Barrier Reef waters.

### Outbreak of crown-of-thorns starfish


Crown-of-thorns starfish are a major predator of coral. Under natural conditions, it is likely that their populations increase to outbreak concentrations in a 50 to 80 year cycle.<sup>97</sup> However, there have been three major population outbreaks of crown-of-thorns starfish over the past 50 years.<sup>97</sup> These occurred from 1962 to 1976, 1978 to 1990, and 1993 to 2005.<sup>96</sup> An emerging outbreak was detected in 2011. An outbreak of crown-of-thorns starfish is considered to be occurring when they are at densities greater than about 30 starfish per hectare.<sup>109,110</sup>

Outbreaks generally appear to initiate in the area between Lizard Island and Cairns, and gradually progress south over several years,<sup>111</sup> although independent outbreaks have been observed in the Swains Reefs in the far south (Figure 6.11). Each of these outbreaks has resulted in severe reductions in coral cover on a regional scale, particularly in the central area of the Region.<sup>112</sup> A crown-of-thorns starfish can consume up to 478 square centimetres of coral each day.<sup>113</sup>

Outbreaks of crown-of-thorns starfish have been one of the major causes of coral death and reef damage on the Great Barrier Reef since surveys began in the 1980s. Analysis of long-term monitoring data has estimated a decline in coral cover of more than 50 per cent in the past 27 years — 42 per cent of which has been attributed to crown-of-thorns starfish outbreaks<sup>36</sup> (see Section 7.1.2).


There is scientific evidence that, while occasional outbreaks are to some extent natural, human impacts have increased their frequency and severity.<sup>97</sup> There are indications that increased nutrient loads lead to crown-of-thorns starfish outbreaks due to increased food supply for their larvae<sup>97,114,115</sup> (Figure 6.10). The reduction in populations of fish that prey on crown-of-thorns starfish may also have some effect, although this is still a matter of conjecture.<sup>116,117</sup> Importantly, the increased frequency of outbreaks, combined with other stresses on corals, means coral populations are unable to fully recover before the next outbreak occurs.

During 2009 and 2010, increases in local populations of crown-of-thorns starfish were beginning to be observed. By late 2011, tourism operators, fishers, rangers and researchers were reporting signs of an emerging outbreak of crown-of-thorns starfish in the northern Great Barrier Reef between Lizard Island and just south of Cairns. In June 2012, the Australian Government invested \$1.43 million for targeted control of starfish in this area. The Authority contracted the Association of Marine Park Tourism Operators to undertake the control operations, with priority given to protecting sites of high value to the Reef tourism industry and, where possible, conducting broader scale control. Between August 2012 and May 2013, 63 reefs were visited out of 330 reefs in the area and 77,679 starfish were removed. A summary of the number of starfish removed to September 2012 is presented in Figure 6.12. In addition, the Queensland Government has invested \$1 million to control crown-of-thorns starfish through the Skilling Queensland program. Work placement participants assist in small-scale control of crown-of-thorns starfish on selected coral reefs.



**Figure 6.11 Extent of observed crown-of-thorns starfish outbreaks in the Region, 1986–2011<sup>47</sup>**

The map shows areas where evidence of an outbreak has been recorded. This includes active outbreaks, incipient outbreaks or recovery from an outbreak.



**Figure 6.12 Removal of crown-of-thorns starfish, offshore Cairns, August 2012 – September 2013**

Reefs where high numbers of starfish have been reported are targeted for repeat visits in the starfish control program. Effort is focused on one or more sites at each reef. The map shows the number of starfish removed from each reef over 39 10-day voyages. (Data provided by the Association of Marine Park Tourism Operators).



### **Pesticides from catchment run-off**

Pesticides, including herbicides, insecticides and fungicides, are used to kill or control pests in agricultural and urban environments and would have been absent from the Region's environment prior to European settlement.<sup>118</sup> It is estimated that at least 28,000 kilograms of herbicides are now introduced into the World Heritage Area each year.<sup>78</sup> The total pesticide load is not accurately known as not all pesticides used in the catchment and draining into its waterways are currently monitored and not all locations can be easily sampled.<sup>78,118</sup> Recently, more systematic monitoring of pesticide residues<sup>119</sup> has shown widespread contamination by a range of pesticides in rivers, streams and estuaries draining to the Region<sup>118</sup>, particularly areas south of Port Douglas. This includes frequent exceedances of the Australian and New Zealand Water Quality Guidelines for fresh waters<sup>120</sup> (often 10 to 50 times), for example atrazine and diuron.<sup>121</sup>

Elevated herbicide concentrations have been particularly linked with sugarcane cultivation in the adjacent catchment.<sup>122,123,124</sup> Irrigation shortly after herbicide application is a major mechanism of herbicide loss from farms.<sup>125</sup> The sugarcane industry has taken initiatives, many funded through the Reef Rescue program, to reduce the run-off of herbicides including precision application (band spray) of herbicides.<sup>126</sup>

Pesticides have been detected in inshore areas of the Region (Figure 6.13) and are of concern as they can have a negative impact on marine plants and animals.<sup>121,122</sup> Herbicide concentrations in flood plumes that extend into the marine environment can exceed concentrations shown to have negative effects on certain species of coral, seagrass and microalgae and present risks to marine mammals.<sup>127,128,129,130,131,132,133</sup> Estuarine areas and freshwater wetlands are more exposed to pesticides as they are closer to the source.

The risk from pesticide exposure is of particular concern for inshore areas in the Mackay Whitsunday Region, Bowling Green and Keppel bays and waters between Port Douglas and Cardwell<sup>132</sup> (Figure 6.13). Other nearshore areas are of low risk, and further offshore the risk might be considered insignificant or zero.

### **Sediments from catchment run-off**

Sediments from the catchment are transported to the Region via coastal rivers and floodplains during the wet season.<sup>134,135</sup>

It is estimated that since European settlement of the catchment in the 1850s, the average annual suspended sediment load entering the Great Barrier Reef has increased by 5.5 times from 3100 to 17,000 kilotonnes per year, with some catchments as high as 14 times more.<sup>78</sup> This is mainly due to increased soil erosion where areas have been cleared of native vegetation to establish pasture grasses for grazing. It has been exacerbated by poor land management practices. In addition, hardened surfaces and straightened channels, as a result of urban and industrial development and agriculture, mean run-off has more erosive power, exacerbating stream bank erosion.


Much of the inshore southern area of the Region is now frequently affected by increased suspended solids that often exceed water quality guidelines (Figure 6.14). Most sediment is confined to the inner shelf and settles out of the water column within five to 15 kilometres of the coastline<sup>134,136,137</sup> where it may be later resuspended by wind-generated waves and currents. However, during flood events, suspended sediment may be carried further offshore. For example, during the 2010–11 wet season, when the Burdekin River had highly elevated discharge over 200 days, most sediment initially settled within approximately 10 kilometres of the river mouth, but some fine silt and clay was carried as far as 100 kilometres northward.<sup>138</sup> These fine colloidal sediments also carry nutrients and other contaminants further into the Region.<sup>139</sup>

Increasing sediment loads have far-reaching effects on the Great Barrier Reef values. For example:

- The heavier erosion sediments, which flow more slowly through the system, are filling freshwater stream beds and deep waterholes. This degrades these habitats, affecting the distribution, abundance and recruitment of many freshwater species and some marine-related species such as sawfish.<sup>140</sup>
- Increases in suspended sediment significantly alter light regimes — lower light levels reduce primary production, both pelagic and benthic (including in the coral-algae symbiosis).
- As the increased sediment load settles, it can smother benthic organisms such as seagrass and corals, making it harder or impossible for them to grow, survive and reproduce.<sup>141,142</sup> This has significant flow-on effects to organisms and animals dependent on these habitats.
- The resuspension of sediments increases the turbidity of open waters and releases additional nutrients previously bound up or buried in sediments.<sup>95,143,144</sup>


Tourism operators voluntarily monitor the health of their tourism sites as part of the Eye on the Reef program. Six years of data collected from tourism sites (Figure 6.15) has found water clarity at some sites is diminishing, particularly in the southern Great Barrier Reef, which is consistent with results presented in Figure 6.14.






**Figure 6.13 Areas of exposure to PSII herbicide residues modelled in the Great Barrier Reef Region<sup>145</sup>**

The model calculated additive photosystem II inhibiting (PSII) herbicide concentrations using end-of-river monitoring data and applied the established relationship of concentration of dissolved organic matter to salinity to corresponding satellite images of flood plumes to predict the additive PSII concentrations. Conservative mixing processes in the Great Barrier Reef lagoon were assumed. Consequence categories are based on known toxicity data to coral and seagrass species and these thresholds, where they were reached, were mapped using the modelled concentrations.



**Figure 6.14 Years that total suspended solids exceeded the guidelines in the Region, 2003–2012<sup>103</sup>**

The map shows the number of years that the relevant water quality guidelines value was exceeded between 2003 and 2012. The exceedances in offshore areas, particularly the Mackay-Fitzroy region (indicated by the dashed line), are under investigation. They may be due to an overestimate of mean concentrations or may indicate a need to revise the boundaries and concentrations in the guidelines.



**Figure 6.15 Water clarity at three tourism sites**

Secchi disc depth data collected voluntarily by tourism operators as part of the Authority's Eye on the Reef monitoring program. The six year data series shows water clarity has declined slightly in the northern Great Barrier Reef. In the south, the rate of decline has been greater.

### Urban and industrial discharge

Urban and industrial discharge refers to all pollutants that enter the Region from the land via pipes and drains. It includes sewage, industrial waste, stormwater and wastewater discharge. The contribution of key pollutants from urban sources is relatively small when compared to diffuse pollutant sources. These impacts can however be locally significant.<sup>146</sup>

There have been significant improvements in the management of sewage in major urban centres of the catchment (see Section 5.3.12) over the last decade. Large investments have been made in upgrading sewage treatment plants at the major coastal population centres of Cairns, Townsville, Mackay and Bundaberg to a tertiary treatment standard. Urban populations with fewer than about 3000 people generally use individual septic tanks at dwellings. The rapidly growing population in most coastal centres is placing increased pressure on local governments to maintain and improve sewage management.

Wastewater is treated at a treatment station before it is discharged to creeks or rivers or reused over land, and some (but not all) contaminants are removed. Stormwater run-off receives no treatment (other than gross pollutant traps) and therefore any chemicals or rubbish it contains can flow directly into creeks or rivers and into the marine environment. To improve stormwater management, local councils are working with their communities and Queensland Government agencies to manage rubbish disposal and the use of chemicals, while documenting environmental values for their local rivers and streams.

Industrial discharge is subject to strict environmental controls. The discharge can release chemicals such as metals, metalloids and non-metallic inorganics in wastewater. There are national, state and Great Barrier Reef-

specific water quality guidelines that identify trigger levels on the discharge of these chemicals.<sup>1</sup> However, many facilities were built decades ago, with long-term permits containing a variety of conditions.

Historically, there have been extensive small-scale mining operations through much of the catchment, including gold, tin, nickel and uranium mines.<sup>147</sup> Some operations have released toxicants which have had severe effects, at least locally, in streams and creeks, for example arsenic associated with tin mining near Herberton.<sup>147</sup> Sediment cores from inshore areas near Townsville have shown a spike in mercury of 25 times the background levels (before European settlement) that coincides with a period of intense gold mining in the adjacent catchment area (between 1870 and the early 1900s) when mercury was used in gold processing.<sup>148</sup>

Extreme flooding events can affect mines within the catchment. Recently, after widespread flooding, more than 30 temporary environmental programs were permitted in central Queensland to address mine flooding. The Queensland Flood Commission of Inquiry proposed a number of changes to the state's Regulations and recommended actions to address mining and water management issues.<sup>149</sup> These included further legislative changes regarding temporary environmental programs and associated permissions. A pilot program of allowing mine wastewater releases during flood conditions was implemented in the 2013 wet season.<sup>150</sup> The results of monitoring associated with these releases will improve understanding of the extent of any effects on water quality, including within the Region.

## 6.4.3 Degradation of coastal ecosystems

### Acid sulphate soils

Potential acid sulphate soils often form where seawater (containing sulphate ions) mixes with land sediments that contain iron oxides and organic matter in a waterlogged condition without oxygen. In these conditions, the iron sulfides are stable and the surrounding soil is often weakly acid to weakly alkaline. However, when this soil is exposed to air (termed 'actual acid sulphate soils'), these soils produce sulphuric acid, often releasing toxic quantities of iron, aluminium and heavy metals.<sup>151</sup> Potential acid sulphate soils are often found in mangroves, salt marshes, floodplains, swamps, wetlands, estuaries, and brackish or tidal lakes.<sup>152</sup> Typically, they may be disturbed through coastal development (specifically the removal of coastal vegetation) and some activities associated with agricultural practices such as drain construction.<sup>153</sup> In drained wetlands, run-off from acid sulphate soils is known to travel through groundwater to estuaries during floods.<sup>154</sup> Modifying floodplains and wetlands can expose acid sulphate soils, resulting in mobilisation of large quantities of iron and aluminium through surface and groundwater, which has been linked to algal blooms such as the toxic *Lyngbya* species.<sup>155</sup>

Potential acid sulphate soils are found extensively in Great Barrier Reef coastal areas and islands.<sup>156,157,158</sup> There has been historical disturbance to these soils<sup>159</sup> and once disturbed, if not treated, acidic water and heavy metals can be released during rain events for decades to come. The combination of acid waters (often with a pH of between two and four) and high concentrations of heavy metals (usually toxic in trace amounts) will affect many species, both immediately and as accumulators in food chains. The impact of acid sulphate soils can affect a range of values such as beaches and coastlines, mangroves, seagrass meadows, invertebrates and fish at a local scale. The effects are often long-term and difficult to reverse.<sup>160,161</sup>

### Artificial barriers to riverine and estuarine flow

Artificial barriers, dams and other modifications to waterways, such as the straightening of channels, affect the natural hydrology in the catchment and may affect Great Barrier Reef species that move between terrestrial and freshwater habitats and the sea.

Barriers to riverine and estuarine flow, such as weirs, dams, barrages, gates, levees, ponded pastures and weeds are widespread in the catchment. For example, 41 impediments to natural environmental flows have been identified as affecting the internationally listed Bowling Green Bay wetland<sup>162</sup>, while in the Burdekin Dry Tropics natural resource management region there are estimated to be more than 1000 obstructions to fish passage.<sup>163</sup>

The interface between the estuarine and freshwater systems is a very productive component of the food chain. Estuaries and coastal waterways are also critical corridors for movement between ecosystems.<sup>20</sup> As many as 78 Great Barrier Reef marine and estuarine fish species use the freshwater systems for part of their life cycle<sup>164</sup> and can be affected by changes in water flow and the presence of artificial barriers.

The loss and modification of estuarine and brackish water habitats, largely saltmarshes and mud flats<sup>20</sup>, is likely to have had effects on shorebirds and fish productivity and survival, particularly through the disruption of tidal systems. Nine per cent of estuarine habitats have been lost in the Great Barrier Reef catchment since European settlement, while up to 30 per cent of the saltmarsh habitats have been modified.<sup>20</sup>

Dams and weirs on rivers reduce freshwater flows in the wet season and often result in increased flows in the dry when the stored water is released.<sup>147</sup> The released water usually comes from deep in the dam, and its temperature and chemistry (for example concentration of dissolved oxygen) is significantly different to natural river flows. Dams also act as traps for sediments (mainly the coarse fraction). For example, the Burdekin Falls Dam captures 60 to 80 per cent of the sediment of the Burdekin River's upstream tributaries.<sup>165</sup>

Hardened surfaces and straightened channels, as a result of urban and industrial development and agriculture, mean there is greater run-off and less water retention on the land. Levees and other structures used to channel waters off the land have significantly modified hydrological processes in many parts of the catchment.<sup>20,166,167,168</sup>

Heavy, salty water sinks are a driver of ocean currents. Water extraction, plus barriers to river and groundwater flow, can increase the salinity of seawater in bays, modifying ocean currents.<sup>169</sup>



## Atmospheric pollution

In general terms, atmospheric pollution refers to that from domestic, industrial and business activities. For the purposes of this report, this potential impact does not include the release of carbon dioxide and other greenhouse gases into the atmosphere — the effects of such pollution are covered in the impacts related to climate change (Section 6.4.1) such as increasing sea temperature.

Atmospheric pollution is not a major issue for the Region's environment. Urban and industrial development is relatively minor and localised in the Great Barrier Reef catchment; however, future projections suggest an increase in both. If not properly managed, this could result in increased atmospheric pollution. Atmospheric pollution has the potential to affect the health of species living in the Region on a local scale, including birds and island plants. It could also affect aesthetic values.

Air pollution from coal dust can occur anywhere coal is handled, conveyed or open to erosion by the wind.<sup>170</sup> Coal dust tends to float<sup>171</sup> and can therefore remain on the water's surface. Although the risk is low, there is a potential it may reduce the amount of sunlight reaching seagrasses and corals and be captured by filter-feeding organisms. Coal dust particles can also aggregate and settle, potentially smothering benthic habitats. While trace elements can leach from coal particles into seawater — some of major concern include arsenic, mercury, lead, sulphur and boron<sup>172</sup> — Australian coal has relatively low trace element concentrations.<sup>170</sup>

## Coastal reclamation

Coastal reclamation refers to the process of creating new land where there was ocean, wetlands, or other water bodies by filling the area with 'land fill' or infrastructure such as groynes and jetties. Reclamation projects can be for public open space, housing or commercial and industrial developments.

The total area reclaimed within the Region is approximately eight square kilometres, the majority of which is in the Gladstone area (approximately 5.5 square kilometres). The largest coastal reclamation projects have been for the development of ports, for example in Gladstone and Townsville.<sup>1</sup>

For some of the major ports in the last few years, the dredge material has been largely used for reclamation works in ports areas. For example, 14 million cubic metres of dredge material has been disposed to the Fisherman's Landing reclamation area in Gladstone Harbour as part of the development of Gladstone's port facilities. Land disposal of dredge material including reclamation can have localised effects on habitats critical to the Great Barrier Reef's health.

Sand replenishment is a form of reclamation used to address beach erosion in order to protect coastal properties and preserve areas for recreational use.

Coastal reclamation can have local effects on the Region's environment, for example removing coastal habitats, covering marine habitats (such as seagrass meadows), altering small-scale local currents, impeding natural drainage from the catchments, altering groundwater levels, and diminishing local aesthetic values. If not properly managed, reclamation works can affect water quality in the adjacent waters and potentially expose acid sulphate soils. Coastal reclamation may also alter the mainland boundary of the World Heritage Area — defined by mean low water.

## Light impacts (artificial)


The presence of artificial light in those localised areas of the Region's coast where there are ports, industrial developments, urban areas and resorts can affect some species. Clearing vegetation and flattening dune systems can also reduce natural light cues and increase impacts from artificial light.<sup>173</sup>

Altered light regimes can disrupt animal behaviour, for example the nocturnal orientation of both adult marine turtles and their hatchlings.<sup>18,174</sup> Artificial lighting can disorient nesting females and turtle hatchlings by reducing the effect of natural lighting and altering topography horizons which are used as guidance mechanisms.<sup>175,176</sup> Seabird fledglings have been found to be attracted to artificial light, causing them to land and stay in urban areas.<sup>177</sup> Some fish and marine invertebrates are attracted to light. Pelagic fish have been shown to be deterred by artificial light, making them disperse and migrate to deeper waters.<sup>178</sup> This response may also lead to consequential impacts on inshore food webs.

## Modifying supporting terrestrial habitats


Since European settlement, there have been significant losses of and modifications to terrestrial habitats that support the Great Barrier Reef ecosystem (Figure 6.16 and Figure 6.17). For the purposes of this report, modifying supporting terrestrial habitats refer to changes that have direct effects on reef species, and environmental processes. It does not include changes to the hydrology or degradation of water quality as those impacts are covered under catchment run-off (Section 6.4.2) or artificial barriers to flow (in this section).

Overall, approximately 60 per cent of pre-clear vegetation — classified as remnant vegetation — remains intact in the Great Barrier Reef catchment.<sup>20</sup> However, the status of coastal ecosystems varies greatly across regions and basins. Generally, human development and the modification of supporting terrestrial habitats intensifies from north to south in the catchment and closer to the coast. For example, much of Cape York's catchment ecosystems remain intact and these make up around one-third of the total area of currently intact ecosystems within the Great Barrier Reef catchment.<sup>20</sup> By contrast, in the Wet Tropics natural resource management region there has been substantial loss of native grasslands and freshwater wetlands across the entire region (greater than 55 per cent) and a high loss of forests and woodlands in several basins, especially on the coastal floodplain.<sup>20</sup>



The loss of functional ecosystems (such as forested floodplains and estuarine habitats) close to the Reef is having significant effects on the feeding and reproduction of many marine species, as well as diminishing dry season refuges. For example, where forested floodplains have been lost through changes in land use, the areas no longer provide nesting habitat or roosts for waterbirds and shady migratory pathways for aquatic species with connections to the Great Barrier Reef.<sup>20</sup> Another example is the replacement of coastal grasslands with intensive agriculture or urban settlements, reducing breeding habitat for many bird and reptile species, including saltwater crocodiles.<sup>20</sup>

Some coastal habitats such as sand dunes are highly sensitive to trampling associated with recreational activities and damage from recreational vehicle use. Damaged coastal vegetation and compacted dune sands influence soil moisture, run-off, erosion, vegetation and micro-organisms. Some animals may also be affected by these activities. For example, shorebirds are highly sensitive to intensive use of beaches, and frequent disturbance results in reduced feeding times, lower survival of chicks, and ultimately population declines.



**Figure 6.17 Coastal ecosystems of the Great Barrier Reef catchment as they are today<sup>20</sup>**

The majority of vegetation in the catchment is classed as 'non-remnant', that is it has been modified to the extent that it is no longer recognised as a functioning habitat. Much has been changed from forest to grassland for grazing purposes.

#### 6.4.4 Direct use

##### Dredging

Dredging involves the extraction of parts of the seafloor (predominantly sand and fine silt, but also harder substrate such as rock) to deepen an area and allow increased access for navigation. It is usually associated with ports, shipping channels, marinas and boat ramps. The term 'capital' dredging is used to refer to dredging that is undertaken to create, lengthen, widen or deepen channels, berth areas, swing basins, marinas and harbour areas. 'Maintenance' dredging refers to dredging which is undertaken to ensure that previously dredged depths are maintained (that is, removing accumulated silt from the channel). The disposal or 'dumping' of dredge material from the dredge site to a disposal site is dealt with separately in the section below.

Both capital and maintenance dredging is undertaken at the majority of trading ports (see Figure 5.11) and at a number of marinas within and adjacent to the Region.

Proposals currently under assessment (as at May 2013) would involve an estimated maximum of approximately 54 million cubic metres of dredging in the World Heritage Area.<sup>179</sup> Decisions are yet to be made in relation to these applications and it is important to note that a change in economic circumstances may result in applications being withdrawn or modified. While it is difficult to predict whether all pending applications will proceed through the full assessment process, projected increases in economic and population growth over the next 25 years (see



Chapter 5) clearly demonstrate that there will be a need for Great Barrier Reef ports to handle increasing volumes of exports and imports and therefore a future need for both capital and maintenance dredging.

Consideration of dredging impacts requires an understanding of: values or attributes likely to be affected by dredging activities, including their current condition; the scale (zone of influence), duration and frequency of dredging activities; non-dredging impacts affecting values or attributes within the dredge zone of influence and the likely combined (or cumulative) effects of dredging and non-dredging impacts on values and attributes; and ecosystem thresholds for health.

The effects of dredging activities are well documented and include: seabed disturbance<sup>180</sup>; removal or modification of habitats<sup>181,182</sup>; loss of species, including benthic organisms<sup>183</sup> and injury or mortality to species of conservation concern<sup>180,184,185</sup>; changes in species behaviour<sup>186</sup>; degradation of water quality<sup>181,187</sup> including increased turbidity levels<sup>182</sup>; changes to hydrodynamics and coastal hydrology<sup>182</sup>; increased underwater noise<sup>188</sup>; and an increased risk of oil spills<sup>186</sup>. However, there is an acute lack of region-specific knowledge and more research and monitoring is required to establish adequate baselines and region-specific impacts.

### **Dumping and resuspension of dredge material**

Once material is extracted from the seafloor during dredging, it requires disposal. Disposal sites may include ocean disposal sites, near-shore reclamation areas and land-based receiving facilities. The National Assessment Guidelines for Dredging<sup>189</sup> require an evaluation of alternatives to ocean disposal including the environmental, social and economic impacts of each disposal option. When disposal sites are ocean-based, the disposal is referred to as 'sea dumping'.<sup>183</sup>

This potential impact considers sea dumping — land disposal is considered under impacts related to degradation of coastal ecosystems (see modifying supporting terrestrial habitats and coastal reclamation in Section 6.4.3).

Between 2000 and 2012, the total volume of dredge material (from both capital and maintenance dredging) dumped in the Great Barrier Reef World Heritage Area was approximately 26 million cubic metres (Figure 6.18). An additional 13 million cubic metres has been approved for disposal within the Region, but has not yet been disposed (Figure 6.18).

The largest quantity of dredge material disposed in the Marine Park in a single campaign was 8.6 million cubic metres associated with the Port of Hay Point in 2006. As noted above, if all dredge and disposal proposals currently under assessment were to be approved, up to approximately 54 million cubic metres of dredge material would need to be disposed, either on land or at sea, over coming years.

Potential effects of sea dumping depend on a number of factors including the: volume and composition of the dredge material; oceanographic conditions in and around the disposal location; proximity of the disposal location to sensitive species and habitats; and the timing and frequency of dredge material disposal.


Major impacts include the burial or smothering of benthic fauna and flora<sup>181,187</sup>, degradation of water quality<sup>183</sup>, and losses and modification of habitat<sup>182</sup>. As at January 2012, the combined area of dredge spoil disposal grounds in the Great Barrier Reef World Heritage Area where localised effects are concentrated was 66 square kilometres.

Less well understood are the broader regional and cumulative effects of sea dumping on inshore biodiversity and more work is required to understand these effects, and their consequences for management. There is evidence that material disposed at existing dredge spoil disposal grounds does not remain within the defined disposal area<sup>190</sup> and that previous modelling of predicted sediment plumes may have significantly underestimated the dispersal and direction of sediments<sup>191</sup> and thus the full extent and potential magnitude of potential impacts.

While dredging and sea dumping do not introduce additional loads of sediments or nutrients to the Great Barrier Reef World Heritage Area, these activities remobilise, redistribute and resuspend sediments and nutrients. It is the effect of the redistribution and resuspension of significant quantities of fine sediments which is a key concern. Increases in turbidity result in decreases in light penetration, affecting seagrass habitats and species such as dugongs and marine turtles which rely on seagrass as a food source.<sup>182</sup> This is particularly significant if these effects happen during periods critical for seagrass survival, growth and reproduction. Increased turbidity also affects coral growth, structure and survival.<sup>187,192</sup>

Increases in turbidity are predominantly caused by fine sediments. Fine sediments can potentially travel large distances (more than 100 kilometres)<sup>138</sup> affecting the light available for photosynthesis of marine plants over significant areas of the Region's inshore waters. Monitoring and modelling of turbidity due to river plumes show that suspended materials can travel over large distances, although this in part reflects the buoyant nature of lower salinity water in the plume<sup>138,143</sup>





**Figure 6.18 Dredge material disposed to date, approved for disposal or being assessed, by port**


Actual dumping volume of dredge material originating from ports within the Great Barrier Reef World Heritage Area. The graphs indicate the amount of dredge material dumped between 2000 and 2012 (red bar), the remaining dredge material yet to be dumped under current permits (blue bar) and the amount of dredge material requested to be dumped in future pending a permit assessment process. Data for Rosslyn Bay (previously dumped: 107,023 ; currently permitted: 63,130; under assessment: 40,000 cubic metres) and Port Douglas marina (previously dumped: 22,000; currently permitted: 0; under assessment: 50,000 cubic metres) was not plotted due to the relatively low amounts compared with other ports. Data derived from the International Maritime Organization sea dumping reports, referrals under the *Environment Protection and Biodiversity Conservation Act* and Marine Park permit applications.

The exact fraction of finer sediment in dredge material will vary. On average, about 30 per cent of capital dredge material is known to be finer sediments such as silts and clays,<sup>193</sup> while riverine inputs typically contain a higher proportion of finer sediments (approximately 70 per cent)<sup>138</sup>.


A recent modelling study, *Improved dredge material management for the Great Barrier Reef Region*, has found dredge material disposed at sea has the potential to migrate over greater spatial and temporal scales than previously understood, due in part to repeated resuspension and deposition.<sup>193</sup> The study was a screening-level sensitivity analysis of the relative merits, if any, of potential alternative material disposal areas and was intended for comparative purposes only. It was the first to incorporate the effects of regional oceanic currents in modelling dredge material and found they are a key factor in the migration or resuspension of dredge material over the long term (12 months). The study also highlighted inter-annual variations of large-scale currents at each of the five ports examined (Figure 6.19), which in turn would influence sediment migration patterns. While the study had a number of limitations (for example, modelled plumes were not field validated and plumes were modelled for an energetic year) and its outcomes cannot be used in project-specific assessments, it highlighted the need for future modelling to take into consideration large-scale currents (and their inter-annual variability) and greater temporal and geographic scales to better predict the extent of dredge material dispersion.

In order to improve scientific understanding of the effects of dredging and offshore dredge material disposal on the Great Barrier Reef, the Authority, in partnership with the Australian Institute of Marine Science, is convening an expert panel to develop and publish a synthesis statement outlining what is known, what is scientifically contentious, and what are the key gaps in our knowledge, focusing on the biophysical effects on the environment. The synthesis statement should enhance the foundation for the ongoing development of policy and guidelines for best practice management and assessment of dredging and dredge material disposal.


### Port of Cairns




### Port of Townsville




### Port of Abbot Point



### Port of Hay Point



### Port of Gladstone



**Figure 6.19 Surface large-scale currents for El Niño (2004), neutral (2007) and La Niña (2011) conditions at major ports**

The rose diagrams show the variability of large-scale currents around the major ports, both in any one year and across years. Data was generated closest to the existing material placement sites.<sup>193</sup>

### Exotic species and diseases

Exotic species and diseases can be introduced through vessel ballast discharge, hull fouling, imported bait and aquaculture operations; in cargo; and by intentional releases (for example, unwanted aquarium fish, dogs and cats). They can travel on all types and sizes of vessels from yachts to cargo ships. Species may be transferred on external and internal surfaces of vessel hulls and on equipment which makes contact with the water (for example, propellers, ropes, chains and intake grates). It is estimated that about 7000 different marine species, including viruses, bacteria and small marine invertebrates, are transported around the world in ballast water every day<sup>194</sup> and more than 150 million tonnes of ballast water is discharged into Australia's major ports each year<sup>194</sup>. More than 250 introduced marine species have been documented in Australia. While most have been relatively harmless, some, such as the northern Pacific seastar, have become aggressive pests in temperate waters.


Introduced marine species have been found in ports along the Great Barrier Reef coastline (for example, the Asian green mussel and Caribbean tubeworm in the Cairns port), although none have been recorded beyond these ports. Any introductions in the future are likely to have regional effects on the ecosystem — the nature of those effects would depend on the species introduced.

Land-based introduced species, such as rats and dogs, affect seabird and marine turtle nesting. Feral foxes and pigs are known to dig up turtle nests — foxes have been implicated as one of the significant factors in the recent decline of loggerhead turtle numbers in Queensland<sup>195</sup>. Feral cats are known predators of green turtle hatchlings in the Seychelles.<sup>196</sup> Although no reports are known for Queensland, feral cats are known to occur on turtle nesting beaches (for example on The Strand, Townsville).

On some Great Barrier Reef islands, insect invasions have caused declines in *Pisonia* forests which are major nesting sites for several seabird species.<sup>1</sup> Weed species have been introduced to islands within the Region.<sup>197</sup>

### Extraction — death of discarded catch


A range of marine species are captured but not retained during fishing activities in the Region. Species include those targeted by the fishery that are discarded due to size or catch restrictions, low market value, 'catch and release' practices, or unwanted species that have been unintentionally captured (incidental catch) and discarded. The largest amount of discarded (non-retained) catch is in the commercial sector (Figure 6.20) and results mainly from trawling activities (Figure 6.21).



**Figure 6.20 Retained and non-retained fisheries catch, Great Barrier Reef, 2007<sup>198</sup>**

A high proportion of the weight of fisheries resources caught in the Great Barrier Reef each year is discarded (non-retained). The survival rates of the discarded species vary and are poorly understood. In the recreational and charter sectors catch and release is increasingly popular.





**Figure 6.21 A breakdown of commercial fisheries non-retained catch, Great Barrier Reef, 2007<sup>198</sup>**

Trawling is responsible for most of the commercial non-retained catch. The continuing interactions between the net fishery and species of conservation concern are of ecological concern. The species and groups of species listed in the figure are those most commonly caught but not retained. Species of conservation concern are in bold type. Syngnathids includes seahorses and pipefish. There is considerable uncertainty in estimates of non-retained catch for fisheries operating in the Great Barrier Reef World Heritage Area, and no contemporary estimates of total extraction (retained and non-retained catch).

Discarded catch in the trawl fishery includes several groups of species of conservation concern, such as sea snakes, marine turtles, seahorses, and sharks and rays.<sup>199</sup> Hundreds of other non-target marine species, most of which are infrequently caught, may be injured or killed by trawling activities. Management requirements in the trawl sector, such as the requirement to install bycatch reduction devices and turtle excluder devices, combined with lower trawl fishing effort have reduced the death of discarded catch in the fishery.<sup>200,201,202</sup> The current risks from trawling to target and non-target species and to the broader environmental values and integrity of the World Heritage Area have been assessed as low or intermediate-low,<sup>199</sup> with a few remaining high risks to species of conservation concern — skates, rays and sea snakes.<sup>199,202</sup> There is no certainty that the level of risk for other species will not increase in the future, as there is potential for trawl fishing effort to increase substantially in the Region under current management arrangements.

The commercial inshore net fishery can unintentionally catch a number of species of conservation concern that are injured or killed in the nets (for example, marine turtles, dugongs, inshore dolphins, sharks and sawfish), many of which are matters of national environmental significance. While strict rules for the use of nets are in place, and Marine Park zoning and Dugong Protection Areas assist to reduce the risks, there is still concern about continuing interactions with these species. Even low levels of mortality may cause population declines and compromise the ability of depleted populations to recover.

Some species may be affected by activities in several fisheries, for example interactions with marine turtles occur in three of the main fisheries (Figure 6.21).

A wide range of species are unintentionally captured through the Queensland Shark Control Program, including some sharks, rays, sawfish, dugongs, whales, dolphins and marine turtles (Table 6.4). Some, such as grey nurse sharks, whale sharks, marine turtles and humpback whales, are listed threatened and listed migratory species under the *Environment Protection and Biodiversity Conservation Act, 1999* (EPBC Act). Not all bycatch are identified to a species level so the actual effects on some species of conservation concern are unknown.

**Table 6.4 Marine species captured in the Queensland Shark Control Program, 1993–2003<sup>203</sup>**

List compiled using catch data from the former Queensland Department of Primary Industries and Fisheries for 1992–2003.

† species is a 'target shark species' and euthanased upon capture

\* common name is inclusive of several species

Sharks		Rays, sawfishes, stingrays	
Blind shark	Common blacktip whaler	Australian cownose ray	Narrow sawfish
Grey nurse shark	Creek whaler	Devil ray*	Queensland sawfish*
Great hammerhead †	Dusky whaler †	Manta ray	Shark ray
Scalloped hammerhead †	Graceful whaler	Eagle ray*	White-spotted guitar ray
Smooth hammerhead †	Grey reef whaler	Bull ray	Shovelnosed ray*
Winged hammerhead †	Hardnose whaler	White-spotted eagle ray	Eastern shovelnosed ray
Port Jackson shark	Long nose whaler †	Electric ray*	Giant shovelnosed ray
School shark	Mangrove whaler	Ray*	Reticulate whipray
Grey carpet shark	Milk shark	Sawfish*	
Great white shark †	Pigeye whaler †	Other marine species	
Mako shark †	Sandbar whaler †	Australian fur seal	Lobster*
Tawny shark	Sharptooth shark †	Barramundi	Mackerel*
Tropical sawshark*	Silky whaler †	Black kingfish	Marlin*
Fossil shark	Sliteye shark	Blue groper	Mud crab*
Weasel shark	Spot-tail shark	Bonita*	Puffer fish*
Whale shark	Spot-tail whaler	Catfish*	Queenfish
Australian blacktip shark	Tiger shark †	Cobia	Queensland groper
Australian sharpnose shark	Whaler shark*	Cod*	Sand crab
Bignose whaler	Whitetip reef shark	Conga eel*	Snapper
Blacktip reef whaler	Tasselled wobbegong	Crayfish*	Swordfish
Blue shark	Zebra shark	Estuarine crocodile*	Toad fish
Bronze whaler	<i>Unknown shark*</i>	Dugong	Tuna*
Bull whaler †		Fish*	Yellowtail kingfish
Whales and dolphins		Marine turtles	
Bottlenose dolphin	Spinner dolphin	Flatback turtle	Leatherback turtle
Common dolphin	Unknown dolphin*	Green turtle	Loggerhead turtle
Snubfin dolphin	Whale*	Hawksbill turtle	Olive ridley turtle
Indo-Pacific humpback dolphin	Humpback whale		

**Extraction — fishing in spawning aggregations**

A fish spawning aggregation is a repeated concentration of fishes, gathered for the purpose of spawning, that is predictable in time and space.<sup>204,205</sup> The density or number of individuals participating in a spawning aggregation is at least four times that found outside the aggregation.<sup>205</sup> This makes the fish easier to catch. Such aggregations are in decline globally, with many decreasing or wiped out due to overfishing.<sup>205</sup> Loss of fish spawning aggregations leads to declines in fish populations with negative ecological consequences.<sup>206</sup> Fish spawning aggregations are classed by the International Union for Conservation of Nature as 'wildlife spectacles'.<sup>206</sup>

Numerous bony fish species in reef and inshore habitats are known to form transient spawning aggregations in the Region. Spanish mackerel are known to aggregate in large numbers to spawn in the Region. During the 1970s, aggregations of spawning Spanish mackerel were reported between Lizard Island and Townsville. In recent years, these aggregations have been reported over a much smaller area on several reefs east of Ingham.<sup>207</sup> A central Great Barrier Reef spawning aggregation site for flowery cod and camouflage grouper has

occasionally been targeted by commercial fishers in the past, and the current status of this aggregation is unknown.<sup>208,209</sup> Monitoring shows a coral trout spawning aggregation site near Cairns has persisted for at least 20 years.<sup>209</sup>

Some fish spawning is protected by Queensland legislation, for example through the Coral Reef Fin Fish Fishery spawning closures and the protection of barramundi during its main spawning season. The protection of representative examples of habitat types in the Marine Park, while not a direct aggregation protection strategy, includes six known fish spawning aggregation sites for coral trout, Spanish mackerel and javelin fish (barred grunter) within no-take areas.<sup>210</sup>

### **Extraction — herbivores**

Herbivory (feeding on plants) is a major ecological process in the Great Barrier Reef ecosystem, performed by a number of species, including many fishes, dugongs and green turtles. Herbivory is a key process in structuring plant communities on coral reefs.<sup>211</sup> Removing herbivores from the ecosystem can lead to unintended ecological consequences for these communities.

On coral reefs around the world, regional reductions in herbivores have underpinned shifts from coral to macroalgae dominance — increases in macroalgae suppress coral growth by competing for space, thus affecting coral recruitment and growth.<sup>98</sup> Extraction of key herbivorous fish can lead to deleterious effects on coral reefs that are hard to reverse<sup>211,212</sup> and it is predicted these effects would be extremely serious should climate change impacts lead to more frequent reef disturbances (such as mass bleaching). Such disturbances would result in widespread algal overgrowth, which a diminished herbivore population may not be able to remove, resulting in limited space for corals to grow and recover.<sup>212,213,214,215</sup>

Currently, there is a relatively small take of herbivorous fish in the Region, and there are a range of management controls that limit the capacity for it to significantly expand. However, any increased demand for herbivorous fish in the future may affect the amount of take and possibly lead to effects higher up in the food chain, and may affect the process of herbivory on coral reefs.

Historically, there has been substantial extraction of the large herbivores — dugongs and green turtles — in the Great Barrier Reef ecosystem (Section 6.3). The effects on seagrass ecosystems in particular from the reduced grazing pressure are unknown. The resulting reduction in turtle and dugong numbers has reduced the capacity of these species to withstand the current multiple impacts on them.<sup>12</sup>

Traditional Use of Marine Resources Agreements, which are developed by Traditional Owner groups in partnership with the Authority and based on the best available science and expert opinion, set limits to the traditional take of dugongs and turtles. Many Traditional Owner groups have taken steps to reduce or stop traditional hunting of turtle and dugong in the wake of recent extreme weather events which heightened concerns for these species (see Section 5.4.1).

### **Extraction — lower order predators**

Lower order predators are species that occupy the middle ranks of the food chain. These species prey upon other species (for example, small fishes and crustaceans) but are also eaten by top order predators (for example sharks). In the Great Barrier Reef ecosystem, lower order predators include predatory fishes such as coral trout, king threadfin, snappers and emperors, groupers and cods, small sharks and stingrays.

Many of these lower order predators are targeted in the Region's fisheries (Figure 6.22) and the abundance of some target fishes is lower in fished areas.<sup>117,216</sup> Some lower order predators may live for a long time and are thus more vulnerable to pressures (for example, the flowery cod can live for up to 42 years<sup>217</sup>) and several could be vulnerable to fishing in inshore or reef areas.<sup>217,218</sup> Loss of predatory reef fish could change the types of species present in groups of juvenile fish<sup>219</sup> which would affect overall reef fish populations. Long-term monitoring has found reduced numbers of predatory fishes may leave coral reefs more susceptible to outbreaks of crown-of-thorns starfish.<sup>116</sup> Changes in predator abundance and behaviour causes changes throughout trophic levels which modify food webs, indirectly acting upon herbivore populations that maintain coral substrates.<sup>213</sup>


Some stingrays and groupers may help to physically shape marine habitats and create micro-habitats for other species.<sup>220,221</sup> Declines in these species could affect habitat formation.

### **Extraction — lower trophic orders**

Lower trophic orders include omnivores, particle feeders and detritivores, for example some fish species, scallops, prawns, lobsters, crabs, sea cucumbers and corals. Herbivores are considered separately in the section above.

The commercial fisheries that extract lower trophic order species for seafood or to supply aquariums include trawl, net, pot, tropical rock lobster, beche-de-mer (sea cucumber), marine aquarium fish and coral collection. Recreational and traditional fishers also take some lower trophic order species such as crabs, prawns and lobsters. Particle feeders and scavengers taken in the trawl and pot fisheries made up more than half of the retained commercial catch from the Great Barrier Reef in 2007 (Figure 6.22).<sup>1</sup> These include valuable seafood species such as eastern king prawns, tiger prawns, endeavour prawns and mud crabs.

The hawksbill turtle is omnivorous, and historic commercial harvesting (Section 6.3) has had lasting effects on this species. Australia supports an internationally significant hawksbill nesting population.<sup>5</sup>



**Figure 6.22 Ecological groups retained by commercial fishing, Great Barrier Reef, 2007<sup>198</sup>**

Most species caught are carnivores (top predators and predators) or particle feeders. Herbivores are not generally targeted on the Great Barrier Reef.

### Extraction — top order predators

Top predators are species that live near or at the top of the food web. In the Great Barrier Reef ecosystem, these species include sharks, very large fishes such as marlin, swordfish, potato cod and Queensland groper, some seabirds, whales, dolphins and the estuarine crocodile.

Many top order predator species are extracted during fishing operations in the Region, either as targeted species (Figure 6.22) or as non-target catch. Some, such as the grey reef shark, whitetip reef shark and several species of sawfishes, have declined.<sup>222,223,224,225</sup> While many top predators are not directly targeted by fishing activities in the Great Barrier Reef, some are unintentionally caught and retained, and several sharks and rays are at high risk from fishing.<sup>218</sup> Many top predators are extracted as part of Queensland's Shark Control Program (Table 6.4). In the last decade, between 521 and 716 sharks have been removed each year throughout Queensland as a result of the program.<sup>226</sup>

Predators play a key role in maintaining a healthy ecosystem and reductions in their populations can have long-term effects<sup>227</sup>, including direct and indirect effects on the food chain.<sup>213</sup>

### Illegal fishing and poaching

Illegal fishing and poaching activities occur Reef-wide and include fishing in no-take zones, the use of fishing equipment or methods where they are not permitted, the incursion of foreign fishing vessels and hunting contrary to legislative requirements. In 2011–12, a total of 305 possible offences involving fishing or collecting (commercial and recreational) and hunting were reported across the World Heritage Area.

Under the Field Management Program's (see Chapter 3 for a description of this program) coordinated risk-based compliance framework for the World Heritage Area, priority is placed on high impact offences such as illegal coastal mesh netting and its incidental impacts on vulnerable species, trawling and commercial reef line fishing. Recently there has been a significant increase in the detection of recreational fishing offences, with 250 recorded between July and December 2012, in part due to the increase of compliance effort on this activity.<sup>228</sup> Incident information shows some fishers are targeting no take (green) zones which has effects on the Region's biodiversity and resilience as well on benefits to other fishers.

The wide geographical range of high priority compliance issues and the growing use of surveillance avoidance tactics by commercial and recreational fishers increases concerns about illegal activity in the Region.

Illegal hunting of dugongs and marine turtles (poaching) in the Region is known to occur, and all reports received are investigated. In recent years, most reports have been found to be legal hunting activities undertaken by Traditional Owners or by people from Indigenous communities hunting with Traditional Owners.

### Mining

Whilst mining of the Great Barrier Reef was halted 40 years ago and subsequent coral growth has concealed most of these impacts on the coral reefs, mining islands for guano-based phosphate rock has had a long-lasting impact. In addition, many important beach ridge sequences have been seen (and still are) as a source of building materials by developers. Daley and Griggs (2006)<sup>229</sup> identified the following locations in the Region that have been mined in the past:

- Guano removal: Raine Island, Michaelmas Cay, Upolu Cay, Oyster Cay, Holbourne Island, Tryon Island, North-west Island, North Fairfax Island, Lady Musgrave Island, Lady Elliott Island. In some cases the geomorphology and vegetation was completely changed and bird nesting sites have been destroyed.
- Coral mining: Snapper Island, Alexandra Reef, Oyster Cay, Upolu Cay, Green Island, Sudbury Cay, Jessie Island, Hutchinson Island, King Reef, Beaver Reef, unnamed cay off Lucinda, Sandpiper Reef.



Sand mining and the destruction of beach ridges is common along the Queensland coast. Beach ridges not only have aesthetic value but can contain important geomorphological information about the last 6000 years. Two of the most important beach ridge sequences are the 100 ridges south of Cape Cleveland near Townsville and the 604 ridges at Cowley Beach near Innisfail.


### Marine debris

There is a massive amount of anthropogenic material accidentally or deliberately released into the marine environment. Common items found within the Region are plastic bags, discarded fishing gear, plastic and glass bottles, rubber thongs, aerosols and drink cans.<sup>230</sup> The annual global demand for plastics has continually grown over recent years. In 2011, it was estimated to be 245 million tonnes.<sup>231</sup> Plastic is the most prevalent type of debris found on beaches worldwide, comprising between 50 to 90 per cent by number of all debris items recorded.<sup>232,233,234</sup> Plastic debris poses a significant threat to wildlife which can choke on the material and starve, absorb chemicals, or ingest items that have degraded into microplastics.<sup>235</sup> In Australia, plastic waste, including discarded fishing gear (nets, lines and ropes), is potentially one of the most harmful types of debris to marine wildlife because of ingestion and entanglement.<sup>236,237,238,239</sup> Marine debris is identified as a key threatening process under the EPBC Act and can affect species and habitats throughout the Region. Marine debris from the catchment appears to accumulate and remain confined within the lagoon system of the Reef but with a northward movement.<sup>230</sup> At the southern end of the Reef, debris appears to be more ocean-sourced.<sup>230</sup>

### Noise pollution

Greater shipping and boating activity, the use of sonar, activities associated with coastal development including pile driving, and defence activities all contribute to increased underwater noise on a local scale. Sound is extremely important to many marine animals, playing a role in communication, navigation, feeding, orientation and the detection of predators.<sup>240</sup> Concerns about the impacts of man-made sound on marine animals has grown over recent decades and is now considered a significant stressor on marine life worldwide.<sup>241</sup> Sounds can have a range of effects, depending on the acoustic frequency animals are able to detect and produce (Figure 6.23) and their proximity to the source. Effects to marine life range from detection with no adverse impacts, to significant behavioural changes, to hearing loss, physical injury and mortality.<sup>240</sup>

While there is a national policy addressing the acoustic impacts of seismic surveys on whales<sup>242</sup>, there are no specific standards for the range of noise pollution affecting Great Barrier Reef species. Given the increases in man-made underwater noise and the observed effects on marine life around the world<sup>241</sup>, there is an urgent need for a greater understanding of the ecological impacts of noise within the Region and for guidance on measures to avoid or mitigate these impacts.



**Figure 6.23 Man-made noise sources and the hearing ranges of marine mammals and fish<sup>243</sup>**

There has been no comprehensive study of the effect of man-made noise on Great Barrier Reef species. This research from North America demonstrates that, for many groups of species, their hearing range overlaps with noises introduced into the environment by human activity. The dotted lines show the range of human hearing in air.

### Outbreak of disease

Whether natural or introduced, disease outbreaks are a clear indicator of stress. They have affected a range of the Region's species in recent years, including corals, green turtles and the Queensland groper. In other countries, coral reef ecosystems that have become much degraded are characterised by a high incidence of diseases. For example, mass coral bleaching in Florida and the Caribbean are usually accompanied by disease outbreaks, probably resulting from low energy reserves in surviving corals.<sup>244</sup> This phenomenon is being

increasingly observed on the Great Barrier Reef and is likely to cause major impacts in the future.<sup>245,246</sup> For the naturally occurring white syndrome disease, major outbreaks have been recorded after especially warm years on reefs with high coral cover, indicating a potential link between coral disease and increasing sea temperatures as a result of climate change.<sup>245,247</sup> Coral disease has been identified as a key indicator of coral reef resilience due to its prevalence in disturbed areas,<sup>248</sup> such as those exposed to flood events,<sup>249</sup> and higher levels of turbidity and sedimentation<sup>250</sup>.

Outbreaks of disease in species of conservation concern are of particular concern. Green turtle fibropapillomatosis was first reported in Australia more than 40 years ago<sup>251</sup> and the frequency of recorded cases increased up to the early 1990s<sup>252</sup>. In the Queensland population, fibropapillomas are rare on green turtles from offshore reefal environments, but prevalent from turtles in semi-enclosed bays.<sup>4</sup> There is evidence of a link to land use practices.<sup>253</sup> The overall effect on the Region's population from this disease currently appears to be low,<sup>254</sup> and a few green turtles have been shown to recover naturally<sup>4,255</sup>.

Necropsies conducted on deceased dugongs reported within the Queensland Marine Wildlife Strandings and Mortality Database indicate disease was the cause of death for between 20 and 25 per cent of the 298 animals examined between 1996 and 2010 for which the cause of death was determined.<sup>256</sup> In 2011, after extreme weather, 30 dugongs died of disease or ill health in Queensland. Twelve dugongs died after extended ill health and had poor body condition. Pneumonia was associated with the deaths of three dugongs, and a further 15 died of unidentified disease.<sup>257</sup>

There is little monitoring of disease in other species.

### **Outbreak or bloom of other species**

An outbreak or bloom of a species refers to a rapid increase in abundance, biomass or population of naturally occurring marine plants and animals. Outbreaks of crown-of-thorns starfish and exotic species are examined above. Outbreaks and blooms can be harmful or lethal to other marine species as they can compete for resources such as food, sunlight and oxygen.

In 2002, a bloom of *Lyngbya majuscula*, a toxic cyanobacteria that can smother seagrass, was observed in Shoalwater Bay, an important feeding ground for green turtles. While the bloom appeared to have no immediate effect on turtle body condition, their diet is likely to have been affected and they were potentially exposed to tumour-promoting compounds produced by the cyanobacteria.<sup>258</sup>

*Trichodesmium* is a cyanobacteria found in nutrient-poor tropical waters. It appears as slicks on the water's surface and can be distinctly pungent. It was first described by Captain Cook, and though it occurs naturally, blooms in the central Great Barrier Reef are thought to have increased, possibly due to nutrients in catchment run-off, in particular phosphorus, iron and organic material.<sup>259,260</sup> The blooms have been implicated in directly smothering corals and increasing the bioavailability of heavy metals.<sup>261</sup>

*Drupella* are marine snails that occur naturally in the Indo-Pacific and are known to damage corals in high densities. Outbreaks have been reported in Ningaloo Reef, Japan and the northern Red Sea.<sup>262</sup> To date, no population outbreaks of *Drupella* have been reported on the Great Barrier Reef, although some tourism operators are permitted to implement control measures for this species. Numbers are monitored daily at some locations in the Region through the Eye on the Reef monitoring program.

### **Physical damage — fishing**

Trawling causes most of the fishing-related physical damage to the Region's marine habitats. It is only allowed in 34 per cent of the Marine Park and generally occurs more than once per year in about seven per cent of the Marine Park.<sup>9,263</sup>

The trawl fishery uses trawl nets designed to operate on or near the seabed, mainly on muddy, sandy or silty habitats. Some trawling occurs in areas with patchy algae (including small amounts of *Halimeda*), patchy seagrass and very small amounts of structural habitat elements such as whip, sponge and gorgonian gardens. Trawl gear can cause direct physical impacts on habitats and remove or damage seabed plants and animals.<sup>15,16,264,265,266,267</sup> Other potential physical impacts include altering the vertical relief of seabed features and redistributing sediments. Although damage to more sensitive lagoon floor communities from trawling has occurred in the past,<sup>9</sup> ongoing physical impacts in most regularly trawled areas are likely to be low, as these areas are generally muddy, silty or sandy and likely to be regularly disturbed naturally.<sup>9,199</sup>

Changes in management arrangements, gear improvements, improved understanding and reduced trawl fishing effort have combined to lower the ecological risk in recent years.<sup>199</sup> However, an upper continental slope habitat (90 to 300 metres deep) in the south-east of the Region that includes deepwater eastern king prawn fishing grounds is considered at high risk from trawling.

Other types of commercial and recreational fishing activities may also cause physical damage to marine habitats, such as damage to live coral tissue and coral colonies from line fishing gear. Such physical impacts may have very localised effects on the health of marine habitats and species at popular fishing locations.

The commercial coral collection fishery also causes some physical damage to habitats, however the total harvest in the fishery is very small relative to the amount of coral on the Great Barrier Reef. The potential impacts are very localised.<sup>268</sup> Appropriate management and practices are in place, including an industry Stewardship Action

Plan<sup>269</sup>. This has included a voluntary moratorium on coral collecting around the Keppel Islands after reef degradation from flooding and coral bleaching.

### **Physical damage — other**


Diving, snorkelling and boating activities can cause local damage to habitats on the seafloor and the plants and animals that live there. Such habitats include seagrass meadows, coral reefs and the lagoon floor. In this report, anchoring of vessels of any size and grounding of vessels less than 50 metres are included under this potential impact along with diving and snorkelling. Physical damage from fishing and the grounding of ships are considered separately.

The effect of anchoring depends on the size of the vessel, its anchoring equipment and the substrate of the seafloor. For example, anchoring has minimal effect on soft muddy bottoms, but is capable of breaking multiple coral colonies as a result of dropping the anchor or the movement of the anchor and chain across the seafloor. Anchoring can also scar seagrass beds. Management arrangements are in place to minimise anchor damage in areas of highest use, for example moorings and no-anchoring reef protection markers. Areas considered most at risk are fragile fringing reefs around popular inshore island destinations.

Disturbance from the anchoring of ships is a localised but chronic impact which is expected to become more frequent close to ports.<sup>270</sup> For many of the areas where ships anchor, the biodiversity values are considered to be relatively low and there is only minor concern about any effects. In contrast, the inner anchorage of Gladstone port supports higher biodiversity than other anchorages, and therefore the overall risk to the biodiversity from anchoring is considered to be higher.<sup>270</sup>

The effects of snorkelling and diving are considered to be low and well managed, especially as the majority of Great Barrier Reef tourism operators use a range of measures including briefings and close supervision to minimise diver and snorkeller interactions with coral reefs. First time divers and those taking photographs are the most likely to damage coral.<sup>271</sup> Best environmental practices for diving and snorkelling are available for tourists and recreational users.

Grounding a vessel can damage coral reefs and other fragile habitats. Weather events and operator error and negligence have caused a number of groundings of small vessels in the Region. While most have had negligible consequences, more of these types of incidents have occurred in the last decade because of an increase in severe weather events. The location of all reported vessel groundings in the Region from 1987 to 2012 is presented in Figure 6.24.



**Figure 6.24 Reported vessel groundings in the Region, 1987–2012**

The locations with a very high number of groundings over the period are those associated with cyclonic events.

### Ship grounding

The nature of the Region's environment, with its thousands of coral reefs and islands, increases the navigational hazards for all ships including cargo, large tourist and recreational vessels. The grounding of a ship can have significant and long-lasting environmental effects on a local area. The grounding of smaller vessels (less than 50 metres in overall length) generally has less effect and is considered in the section on physical damage — other (see above).

Potential impacts from the grounding of a ship include direct physical damage to coral reef or other habitats at the grounding site, leaving visible scars; scraping off and releasing into the local environment toxic substances including anti-fouling paint, copper, zinc, and tributyl tin (also known as TBT, an anti-fouling chemical that targets marine organisms); and cargo and oil spills which can have damaging or lethal effects on marine life.

Despite more than 4000 ship calls to ports adjacent to the Great Barrier Reef each year, there have only been a small number of collisions and groundings.<sup>272</sup> Since 1996, 22 ship groundings and collisions have been recorded in the World Heritage Area.<sup>273</sup>

In April 2010, a Chinese registered coal carrier *Shen Neng I* ran aground on Douglas shoal in the Great Barrier Reef. The vessel was grounded for nine days, severely damaging an estimated 80,000 to 400,000 square metres of reef.<sup>274</sup> It is the largest ship grounding scar on the Great Barrier Reef. At best, it is expected the site of impact will take decades to recover.<sup>274</sup> Anti-fouling chemicals from this and other ship groundings, particularly the *Bunga Terati Satu*, *Doric Chariot* and *Peacock*, will have affected marine life. These chemicals combine with pulverised



reef to damage corals at the grounding sites and surrounding areas where currents transport the paint flakes and pulverised particles.

### Spill — large chemical

For the purposes of this report, large chemical spills are those that trigger a national or regional response, are generally more than 10 tonnes, or are likely to cause considerable environmental effects.

Chemicals commonly transported through the Region include: cement, fertilisers, ammonium nitrate, sugar and molasses, liquefied natural gas, coal, sulphuric acid and caustic soda. However, the full breadth of chemicals being transported is not well understood, particularly when it is transported within shipping containers as there is often mislabelling of the container.

Although there has never been a large chemical spill reported in the Region, greater shipping activity and industrial development along the Great Barrier Reef coast is increasing its likelihood. While a large chemical spill would not necessarily be visible, it could have regional and long-lasting effects on Great Barrier Reef values. Apart from the physical smothering of plants and animals, a chemical's toxicity and reactions with water could result in persistent effects on the health, growth, reproduction, development and survival of a range of marine plants and animals for several years, for example:

- Sulphuric acid would reduce the pH of seawater, affecting calcium deposition in animals with skeletons. It could also coat marine animals such as turtles and cause damage through inhalation.<sup>275</sup>
- Sugar and molasses dissolves and dissipates over large areas. It causes oxygen depletion which can result in mass deaths of fish and other organisms and increased dissolved organic carbon causing microbial changes.<sup>108</sup>
- Liquefied natural gas vaporises after release and can affect air-breathing animals. It is flammable and may also cause thermal shock.<sup>276</sup>

### Spill — large oil

A large oil spill is considered to be any spill of oil that has triggered a regional or national response and/or is more than 10 tonnes (Tier 2 and Tier 3 of the National Marine Oil Spill Contingency Plan<sup>277</sup>).

Much of the oil carried in the Region is refined, such as petrol, diesel and heavy fuel oils, rather than crude unrefined oils. Individual tankers carry up to 60,000 tonnes of oil through the inner shipping route. Large bulk carriers operating to Hay Point can carry up to 4000 tonnes of heavy fuel oil.<sup>278</sup> The volumes carried and the potential effects mean a large oil spill is viewed as one of the greatest risks from ships transiting through the Great Barrier Reef.<sup>278</sup>

The largest recorded oil spill in the World Heritage Area occurred in January 2006 when approximately 25 tonnes of heavy fuel oil was spilt from the *Global Peace* while berthing at a coal loading facility in Gladstone Harbour (Table 6.5). While some of the oil was recovered quickly, much deposited on the coastline. There were local effects on coastal species including mangroves, crabs and sediment-dwelling species.<sup>279,280</sup> In December 2002, a spill off Border Island in the Whitsundays resulted in a slick of at least 70 kilometres long in the Region.

**Table 6.5 Large oils spills that have affected the values of the Great Barrier Reef**

\* Although the Torres Strait is outside the Region, many of the Region's values, including large mobile species such as marine turtles and dugongs, were affected including through loss of connectivity to breeding and feeding grounds.

Date	Vessel	Location	Oil amount
March 1970	<i>Oceanic Grandeur</i>	Torres Strait*	1100 tonnes
December 2002	<i>Pacific Quest</i>	Border Island, Whitsundays	>70 km slick
January 2006	<i>Global Peace</i>	Gladstone Harbour	25 tonnes

In 1970, Australia's second largest oil spill occurred in the Torres Strait when the *Oceanic Grandeur* struck an uncharted rock, releasing an estimated 1100 tonnes of oils, including crude oil from the cargo tanks and a lighter diesoline fuel oil. Despite being outside the Region, it is likely the spill affected the Region's values. At the time, the event highlighted Australia's lack of preparedness to respond to a major oil spill and accelerated the development of the National Contingency Plan.

In 2009, two major incidents involving the release of oil occurred in waters adjacent to the Region, one in the Torres Strait only 20 kilometres from the Region's northern boundary, and the other off Moreton Bay near Brisbane (*Pacific Adventurer*), to the south of the Great Barrier Reef.<sup>1</sup> Another major spill involving 12,000 litres of diesel occurred in January 2013 after the grounding of a refuelling barge in the Burnett River just north of Mon Repos, an important turtle nesting beach.

At a global scale, these spills in or adjacent to the Region are relatively small. The top 10 global oil spills recorded range from 140,000 tonnes to 1.6 million tonnes.

Despite increases in shipping traffic, improvements in shipping safety management have resulted in fewer major shipping incidents in the past 10 years.<sup>273</sup> However, a recent survey of coastal pilots indicates there are a high

number of near misses that are not reported.<sup>281</sup> These may increase with the growing shipping traffic, combined with factors such as fatigue.<sup>282</sup>

Apart from the physical smothering of plants and animals, oil toxicity and its chemical reactions with water mean a large spill is likely to have persistent effects on the health, growth, reproduction, development and survival of a range of marine plants and animals for several years.<sup>278</sup>

### Spill — small chemical or oil

Small chemical and oil spills (defined as less than 10 tonnes) are likely to have had localised effects on environmental and social values. Such spills often happen during refuelling, during port construction and operational activities, while transporting chemicals or oils, and through illegal discharges of bilge waste.<sup>278</sup>

Since 1987, more than 700 reports of spills and discharges have been recorded in the Authority's marine incident database (Figure 6.25). However, most of these reports involve small spills of petroleum products (for example, from bilge pump outs, sinking vessels or vessel groundings, and discharges of hydraulic fluids from construction equipment).<sup>278</sup> Some of these reports are unsubstantiated.

The environmental consequences of small spills will depend on factors such as the type of oil or chemical, the amount spilled, weather conditions and the surrounding habitat. Potential direct effects on the environment include toxicity to marine biota; coating of animals, plants or habitats which can affect respiration and insulation (particularly for seabirds); as well as burns, poisoning and in some cases, mortality.




Figure 6.25 Reported spills and discharges in and adjacent to the Region, 1987–2012

### **Vessel strike on wildlife**

Vessel strikes typically affect surface-breathing marine animals such as marine turtles, dugongs, dolphins and whales, often resulting in injury or death. Wildlife may be struck by any moving vessel (for example, jet skis, yachts, dinghies and cargo ships), but the chances and consequences are far greater for high-speed vessels.<sup>283</sup> The likelihood of a strike increases where high-speed vessels overlap with key shallow water habitats (for example seagrass meadows) or movement corridors for vulnerable animals. It also increases in areas adjacent to large regional communities where visits to the Region are more frequent and there are larger volumes of commercial traffic.<sup>284</sup> Dugongs do not appear to swim away from passing vessels, making them at high risk of vessel strike, especially from high speed vessels.<sup>285</sup>

Current published information on vessel strikes on dugongs, whales and dolphins suggests its impact on these species has not increased substantially within the Region and is a relatively infrequent occurrence on these species.<sup>256,286</sup>

Recently, the waters offshore from Gladstone have had an elevated number of marine turtle strandings. In 2011, there were eight times more interactions between marine turtles and vessels compared to 1996 when strandings data began to be recorded.<sup>287</sup>

Voluntary go slow areas and transit lanes have been declared in some areas where there is high vessel traffic and large populations of marine turtles or dugongs, such as near Hinchinbrook Island. However, compliance to these voluntary measures remains low, especially among frequent users of the Hinchinbrook area.<sup>288</sup> The increase in boat traffic and the consequent risks to dugongs was of great concern to Traditional Owners prior to adoption of the management arrangements.<sup>289</sup>

### **Waste discharge from a vessel**

The types of waste discharged from vessels into the sea include food waste, litter, grey water (that is, from showers, laundry) and sewage. There are specific requirements under the International Convention for the Prevention of Pollution from Ships, as well as Queensland and Commonwealth legislation (including Great Barrier Reef Marine Park Regulations), regarding how this waste should be treated and/or macerated prior to discharge. There is the potential for animals, such as planktivores and filter feeders, feeding on vessel-generated waste to suffer ill health from an unnatural diet.<sup>290</sup>

Waste discharge from vessels also increases nutrients in the water column, particularly sewage discharge which has high concentrations of nitrogen and phosphorous. Total sewage discharge into the Marine Park only accounts for about six per cent<sup>291</sup> of the nitrogen load, and vessel discharge contributes only a small portion of this.

### **Wildlife disturbance**

Activities such as diving, snorkelling, fish feeding, fishing, reef walking, scenic flights, boating, walking and driving on beaches, and visiting islands can disrupt wildlife on a local scale, particularly in sensitive or heavily visited areas. These activities have the potential to disrupt animal behaviour, reduce their fitness and their ability to feed, breed, nest and migrate.<sup>241,285,292,293,294,295,296,297,298,299,300,301,302,303,304,305</sup>

Attitudes towards interacting with wildlife have changed over the decades. While visitors may have previously ridden on, touched and chased wildlife, they are now much more aware of their activities and the best environmental practices for viewing these animals. At the same time, management arrangements have addressed many interactions (such as approaching whales, visiting important seabird rookeries or approaching critical migratory shorebird roosts).

While there is evidence that the presence of divers can attract or deter marine species, these effects may only be temporary<sup>306</sup>. Feeding to attract marine wildlife can potentially affect some species.<sup>307</sup> Tourism operators must have a permit from the Authority to feed fish as part of their tourist program and must abide by guidelines relating to the activity.

### 6.4.5 Assessment of impacts on biodiversity

A detailed assessment of the effects the above impacts have had or are currently having on the key biodiversity values identified in Chapter 4 is presented in Table 6.6 and Table 6.7.

The assessment is of past and present effects on indicators, with an indication of future trend. It is recognised that this assessment is generalised and that effects on individual values are not evenly distributed across all the components that make up a value or across the Region.

Understanding Table 6.6 and Table 6.7	
Grading statements	
	<b>No effect:</b> No known interaction or the interaction is considered insignificant.
	<b>Data deficient:</b> Effects are unknown.
	<b>Very low effect:</b> Any effects attributable to the impact are minor or localised, with no observable effects on the values.
	<b>Low effect:</b> The effects of the impact are observable in some locations or to some species, but only to the extent that limited additional intervention would be required to maintain the values.
	<b>High effect:</b> The effects of the impact are obvious in many locations or for many species to the extent that significant additional intervention would be required to maintain the values.
	<b>Very high effect:</b> The effects of the impact are widespread to the extent that the values are severely compromised.
Trend	
↑	<b>Increasing:</b> The intensity and/or spatial extent of the impact is increasing.
↔	<b>Stable:</b> The intensity and/or spatial extent of the impact is staying about the same.
↓	<b>Decreasing:</b> The intensity and/or spatial extent of the impact is decreasing.

This process is used as the first step to identify the impacts having the most effect on the Region's values. It is these impacts that are the focus of exploring cumulative impacts.

The future risk that these impacts present to the Region's values is assessed in Chapter 10.



**Table 6.6 Past and present effects of impacts on Great Barrier Reef habitats**

The assessments presented are based on the information presented in Section 6.4 above. The references cited there also apply to this assessment. \* indicates impact is likely to be the result of a variety of activities.

	Impacts	Future trend of the impact	Islands	Beaches and coastline	Mangrove forests	Seagrass meadows	Coral reefs (<30m)	Deeper reefs	Lagoon floor	Shoals	Halimeda banks	Open waters	Continental slope
Climate change	Altered ocean currents	↑											
	Cyclone activity	↑											
	Increased sea temperature	↑											
	Ocean acidification	↑											
	Rising sea level	↑											
Catchment run-off	Increased freshwater inflow	↑											
	Nutrients from catchment run-off	↔											
	Outbreak of crown-of-thorns starfish	↑											
	Pesticides from catchment run-off	↔											
	Sediments from catchment run-off	↔											
	Urban and industrial discharge	↑											
Degradation of coastal ecosystems	Acid sulphate soils	↑											
	Artificial barriers to flow	↑											
	Atmospheric pollution	↑											
	Coastal reclamation	↑											
	Light impacts (artificial)	↑											
	Modifying supporting terrestrial habitats	↑											
Direct use	Dredging	↑											
	Dumping and resuspension of dredge material	↑											
	Exotic species and diseases *	↑											
	Extraction — death of discarded species	↔											
	Extraction — fishing spawning aggregations	↔											
	Extraction — herbivores	↔											
	Extraction — lower order predators	↔											
	Extraction — lower trophic orders	↔											
	Extraction — top order predators	↔											
	Illegal fishing and poaching	↑											
	Marine debris *	↑											
	Noise pollution	↑											
	Outbreak of disease *	↑											
	Outbreak or bloom of other species *	↑											
	Physical damage — fishing	↔											
	Physical damage — other	↑											
	Physical damage — ship grounding	↑											
	Spill — large chemical	↑											
	Spill — large oil	↑											
	Spill — small chemical or oil	↑											
	Vessel strike on wildlife	↑											
	Waste discharge from vessels	↑											
	Wildlife disturbance	↑											

**Table 6.7 Past and present effects of impacts on Great Barrier Reef species and groups of species**

The assessments presented are based on the information presented in Section 6.4 above. The references cited there also apply to this assessment. \* indicates impact is likely to be the result of a variety of activities.

	Impacts	Future trend of the impact	Mangroves	Seagrasses	Macroalgae	Benthic microalgae	Corals	Other invertebrates	Plankton and microbes	Bony fish	Sharks and rays	Sea snakes	Marine turtles	Estuarine crocodiles	Seabirds	Shorebirds	Whales	Dolphins	Dugongs
Climate change	Altered ocean currents	↑																	
	Cyclone activity	↑	Orange	Red	Orange	Orange	Dark Purple	Red	Grey	Orange	Orange		Orange	Orange	Yellow	Yellow			Orange
	Increased sea temperature	↑		Orange			Red	Orange	Yellow	Orange			Orange	Orange	Red	Orange			
	Ocean acidification	↑					Yellow	Yellow	Yellow										
	Rising sea level	↑											Yellow						
Catchment run-off	Increased freshwater inflow	↑	Yellow	Orange	Yellow	Yellow	Red	Yellow	Yellow				Yellow	Orange					Yellow
	Nutrients from catchment run-off	↔	Grey	Red	Red	Red	Dark Purple	Yellow	Red				Orange						Orange
	Outbreak of crown-of-thorns starfish	↑					Dark Purple	Orange	Grey	Orange									
	Pesticides from catchment run-off	↔	Orange	Red	Red	Red	Red	Orange	Orange	Yellow	Yellow	Yellow	Orange	Yellow	Yellow	Yellow	Yellow	Yellow	Orange
	Sediments from catchment run-off	↔	Yellow	Red	Red	Orange	Dark Purple	Orange	Orange				Orange						Orange
	Urban and industrial discharge	↑	Yellow	Yellow	Yellow	Yellow	Yellow	Grey					Yellow		Yellow	Yellow			Orange
Degradation of coastal ecosystems	Acid sulphate soils	↑	Orange	Yellow	Grey	Grey	Yellow	Orange	Yellow	Orange	Orange			Yellow		Yellow			
	Artificial barriers to flow	↑	Yellow	Yellow				Yellow		Orange	Yellow			Orange		Red			
	Atmospheric pollution	↑	Grey	Grey					Grey										
	Coastal reclamation	↑	Orange	Orange		Yellow	Yellow	Grey	Grey				Grey	Yellow		Yellow			Grey
	Light impacts (artificial)	↑						Yellow		Yellow	Yellow		Orange		Yellow	Yellow			
	Modifying supporting terrestrial habitats	↑	Red	Grey	Grey	Grey	Grey	Orange	Grey	Red	Red		Orange	Orange	Orange	Orange			
Direct use	Dredging	↑	Yellow	Orange	Yellow	Yellow	Yellow	Orange	Grey	Yellow	Yellow	Grey	Orange	Yellow		Yellow	Grey	Orange	Yellow
	Dumping and resuspension of dredge material	↑	Yellow	Red	Yellow	Orange	Red	Orange	Grey	Yellow	Grey	Grey	Orange	Grey		Yellow			Orange
	Exotic species and diseases*	↑						Yellow	Grey										
	Extraction — death of discarded species	↔						Orange		Orange	Red	Red	Red	Yellow	Yellow		Yellow	Red	Red
	Extraction — fishing spawning aggregations	↔								Red									
	Extraction — herbivores	↔								Yellow			Orange						Red
	Extraction — lower order predators	↔								Orange	Orange		Orange				Red		
	Extraction — lower trophic orders	↔					Yellow	Orange		Yellow									
	Extraction — top order predators	↔								Orange	Red		Orange						
	Illegal fishing and poaching	↑								Orange	Red		Red						Red
	Marine debris*	↑						Yellow		Yellow	Yellow	Yellow	Orange	Orange	Orange	Orange	Orange	Orange	Orange
	Noise pollution	↑						Yellow		Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Orange	Orange	Orange
	Outbreak of disease*	↑					Orange		Grey	Orange			Orange						Orange
	Outbreak or bloom of other species*	↑		Yellow			Yellow	Yellow	Yellow				Yellow						Yellow
	Physical damage — fishing	↔		Orange			Yellow												
	Physical damage — other	↑		Orange	Yellow	Yellow	Orange	Orange											
	Physical damage — ship grounding	↑		Yellow	Yellow	Yellow	Orange	Orange	Yellow	Orange	Yellow	Yellow	Yellow	Yellow	Yellow				
	Spill — large chemical	↑																	
	Spill — large oil	↑	Orange	Orange	Orange	Orange	Yellow	Orange	Orange	Yellow	Yellow	Yellow	Yellow		Orange	Yellow		Yellow	Yellow
	Spill — small chemical or oil	↑	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow
	Vessel strike on wildlife	↑											Orange				Yellow	Yellow	Orange
	Waste discharge from a vessel	↑			Yellow	Yellow	Yellow	Yellow											
	Wildlife disturbance	↑						Yellow		Yellow	Yellow	Yellow	Orange	Yellow	Orange	Orange	Orange	Yellow	Yellow

## 6.5 Impacts on geomorphological features

A detailed assessment of the effects the identified impacts have had or are currently having on geomorphological features of the Region (as identified in Chapter 4) is presented in Table 6.8.

The assessment is of past and present effects on the values, with an indication of future trend. As with biodiversity above, the assessment is generalised — effects on individual values may not be evenly distributed across all the components that make up a value or across the Region, and there will be spatial variation.

It can be assumed that many of the impacts affecting the condition of biodiversity values (Section 6.4) may also be affecting the geomorphological features of the Region as the geomorphology and ecology of the Great Barrier Reef are strongly interdependent.<sup>308</sup> However, there is likely to be a lag in the response of geomorphological features to impacts compared to the ecological system. For example, coral species and coral reefs can degrade or die as a result of rises in sea temperature; however, the death of coral may not affect the overall reef structure until much later or until the dead coral begins to erode.<sup>309</sup>

Understanding Table 6.8	
Grading statements	
	<b>No effect:</b> No known interaction or the interaction is considered insignificant.
	<b>Data deficient:</b> Effects are unknown.
	<b>Very low effect:</b> Any effects attributable to the impact are minor or localised, with no observable effects on the values.
	<b>Low effect:</b> The effects of the impact are observable in some locations or to some geomorphological features, but only to the extent that limited additional intervention would be required to maintain the values.
	<b>High effect:</b> The effects of the impact are obvious in many locations or for many geomorphological features to the extent that significant additional intervention would be required to maintain the values.
	<b>Very high effect:</b> The effects of the impact are widespread to the extent that the values are severely compromised.
Trend	
↑	<b>Increasing:</b> The intensity and/or spatial extent of the impact is increasing.
↔	<b>Stable:</b> The intensity and/or spatial extent of the impact is staying about the same.
↓	<b>Decreasing:</b> The intensity and/or spatial extent of the impact is decreasing.

The assessment presented in Table 6.8 demonstrates no high or very high effects on the Region's geomorphological features. Impacts which have had the most effect on geomorphological features are cyclone activity, modifying supporting terrestrial habitats, sediment in catchment run-off, increased freshwater inflow, dredging and dumping of dredge material.

- **Cyclones** have produced destructive effects on geomorphological features throughout the Region. Cyclone-induced waves and currents carrying sand and rubble can cause erosion of reefs and reworking of pre-existing sediment deposits.<sup>309,310</sup> Cyclones can also result in large areas of seagrass being ripped up,<sup>310</sup> leading to further instability of sediment in the area.<sup>309</sup> Recent cyclones have also caused the formation of new rubble cays, loss of some sandy cays and erosion of some shorelines by waves. Shoreline and shallow subtidal features are likely to be affected the most, whereas deeper features experience fewer effects.<sup>309</sup> Features such as continental islands and islands with some fringing vegetation and reefs are also likely to be better protected.<sup>309,311</sup> Cyclones are also likely to have rearranged some river deltas, however these are naturally dynamic systems.
- **Artificial barriers to flow** such as weirs and dams; **modifying supporting terrestrial habitats; dumping and resuspension of dredge material;** and increased **sediment from catchment run-off** affect the amount of suspended sediment within the Region. Suspended sediment can affect the health of some corals and increase their susceptibility to erosion.<sup>309</sup> It also affects the health of seagrass meadows and can potentially infill submerged palaeochannels.<sup>309</sup> Changes in sediment loads can also affect the shape and size of river deltas.
- Deepwater geomorphological features (90 to 300 metres deep) off the southern Great Barrier Reef may have been affected by **physical damage from fishing activities** (deep sea trawling), however there is limited knowledge about the effects.<sup>309</sup>
- Alterations to channels in river deltas and paleochannels from **dredging** have the potential to remove or rearrange sediment deposits that make up the geomorphological feature.<sup>309</sup>
- **Increased freshwater inflow**, particularly during flood events, has had an effect on islands and shorelines and river deltas within the Region. It also affects the health of coral reefs and seagrass meadows. Extreme weather events in the summer of 2010–11 resulted in increased freshwater inflow.

The worst effects were on some inshore reefs close to the mouths of major rivers and on the mainland sides of islands.<sup>310</sup>

- **Coastal reclamation** has affected some shorelines and river deltas by changing the way sediment is eroded, transported and deposited.

**Table 6.8 Past and present effects of impacts on Great Barrier Reef geomorphological features**

\* indicates impact is likely to be the result of a variety of activities

	Impacts	Future trend of the impact	Coral reefs	Islands and shorelines	Channels and canyons	River deltas	Halimeda banks	Seagrass beds
Climate change	Altered ocean currents	↑						
	Cyclone activity	↑						
	Increased sea temperature	↑						
	Ocean acidification	↑						
	Rising sea level	↑						
Catchment run-off	Increased freshwater inflow	↑						
	Nutrients from catchment run-off	↔						
	Outbreak of crown-of-thorns starfish	↑						
	Pesticides from catchment run-off	↔						
	Sediments from catchment run-off	↔						
	Urban and industrial discharge	↔						
Degradation of coastal ecosystems	Acid sulphate soils	↑						
	Artificial barriers to flow	↑						
	Atmospheric pollution	↑						
	Coastal reclamation	↑						
	Light impacts (artificial)	↑						
	Modifying supporting terrestrial habitats	↑						
Direct use	Dredging	↑						
	Dumping and resuspension of dredge material	↑						
	Exotic species and diseases*	↑						
	Extraction — death of discarded species	↔						
	Extraction — fishing spawning aggregations	↔						
	Extraction — herbivores	↔						
	Extraction — lower order predators	↔						
	Extraction — lower trophic orders	↔						
	Extraction — top order predators	↔						
	Illegal fishing and poaching	↑						
	Marine debris*	↑						
	Noise pollution	↑						
	Outbreak of disease*	↑						
	Outbreak or bloom of other species*	↑						
	Physical damage — fishing	↔						
	Physical damage — other	↑						
	Physical damage — ship grounding	↔						
	Spill — large chemical	↑						
	Spill — large oil	↑						
	Spill — small chemical or oil	↑						
	Vessel strike on wildlife	↑						
	Waste discharge from a vessel	↑						
	Wildlife disturbance	↑						



Submarine canyons and turbidite deposits occur in deep waters along the continental shelf. Due to the relative inaccessibility, they are presumed to be largely unaffected by the impacts assessed.

## 6.6 Impacts on Indigenous and historic heritage values

### 6.6.1 Indigenous heritage values

As demonstrated in Chapter 4, the Indigenous heritage values of the Region are interconnected in a cultural landscape. The natural environment is fundamental to Traditional Owner connections to their land and sea country. Therefore, when considering likely impacts on Indigenous heritage values, the first step is to recognise that all impacts identified as affecting natural values in Sections 6.4 and 6.5 are likely to be equally affecting Indigenous heritage values.

Added to this are the effects on the other aspects of Indigenous heritage such as cultural practices, sacred sites, sites of particular significance, stories, songlines, totems, language, technology, tools and archaeology. The following is a series of examples of the ways in which the identified impacts have been and continue to affect these values since European settlement. A structured assessment of the impacts is presented in Table 6.9.

Traditional Owners report their ability to catch food has been affected by the **death of discarded species**, resulting in a break with cultural practice, lore and custom, and interference with sites of particular cultural significance. For example, if all the fish are gone from an area, Traditional Owners may not be able to go there anymore. Indigenous communities in Cape York have reported to the Authority's staff large hauls of queenfish being discarded by commercial netters in Princess Charlotte Bay.<sup>312</sup> Indigenous heritage values are also severely affected by the loss of dugong, a cultural keystone species that can be incidentally caught and killed in nets.<sup>313</sup>

Although a legacy activity, past **commercial harvesting of herbivores**, such as marine turtle and dugong, has ongoing effects on Indigenous cultural values and has significantly changed cultural practices. The low population size, in part due to the previous commercial harvests, limits the number of animals available for hunting, as well as changing where animals occur.<sup>313</sup>

Indigenous heritage values can be affected by the **extraction of lower order predators**. For example, a key food source by Injinoo Traditional Owners, the black jewfish, is now found in much lower numbers and in much smaller sizes, altering their ability to maintain their cultural practices and customs. This decline has been attributed to fishing pressure.<sup>289</sup>

The **extraction of lower trophic orders** has affected Indigenous heritage values in some areas. For example, Gooreng Gooreng Traditional Owners from the Gladstone region report a decline in seafoods such as oysters and crabs.<sup>289</sup> As well as being an important food source, the cultural practice tied to collecting these shellfish (usually by women) is being lost as the resource becomes increasingly scarce.<sup>289</sup>

The **illegal take** of culturally significant species including dugong, marine turtles, sea snakes, sharks, rays, crayfish, mullet, oysters, pipi, clamshells, whiting and bream can have direct effects on Indigenous heritage values such as cultural practices, observances, lore, stories, songlines and sites. Taken at the wrong time of year, in the wrong numbers or at a culturally sensitive site reduces the numbers available for use in traditional ways and forces Traditional Owners to change their customs and practice.<sup>289</sup>

Species most affected by **vessel strike on wildlife** are also species culturally significant for many Great Barrier Reef Traditional Owners. Marine turtles, dugongs and whales are totem animals and are part of cultural practice, observances, tradition and stories and songlines. A reduction in these species has directly affected Indigenous heritage values. For example, they are not available as a food source and, therefore, cultural practices such as hunting cannot be undertaken. In turn, observances, lore and customs cannot be continued.

The increasing amount of **marine debris** is affecting cultural keystone species for Traditional Owners. For example, plastics that float in the open ocean affect seabirds<sup>314</sup> — many of which are totem animals and intimately linked with storylines.

The **modification of supporting terrestrial habitats** since European settlement has changed Traditional Owners' country and diminished cultural and heritage values. For example, timbers that were once abundantly available to make tools are no longer found along many stretches of coastline.<sup>315</sup> Also, in Dharumbal country, coastal pandanus was and still is used for making baskets and matting. The distribution and abundance of the pandanus tree has been greatly reduced since European settlement.<sup>316</sup> Clearing and modifying coastal areas has also reduced the distribution and abundance of the corkwood tree, a well-used and important tree for Wulgurukaba Traditional Owners.<sup>317</sup> A traditional burial site was disturbed in 1998–99 when a dam was built at Francis Creek on the north side of Palm Island; more than forty remains were disturbed during sand removal.<sup>318</sup> Another example is along the shores of Cleveland Bay and Ross River, which have significance as 'Dreaming places' and as important fishing and hunting grounds. Traditional Owners report that traces of the 'big Murri camps' are 'long gone' as a result of modern disturbance to the landscape.<sup>319</sup>

Historically, some **coastal reclamation** has been undertaken without proper engagement or consultation with Traditional Owners, resulting in effects on Indigenous heritage values. Any changes to land and seascapes are changes to Traditional Owners' country and, therefore, diminish culture and heritage values such as story places, songlines and sacred sites. For example, in the Nelly Bay harbour development on Magnetic Island there was an

incident that involved digging up of remains.<sup>320</sup> Bindal Traditional Owners have also reported that Ross Creek fish traps have disappeared due to reclamation, meaning it is no longer possible to pass on this traditional practice to future generations.<sup>312</sup> Another example is at Clump Point near Mission Beach — a culturally important story place. Part of the story involves the shape of the bay and headland; changes due to development mean the storyline is broken and the significant cultural site has been affected.<sup>321</sup>

Traditional Owners have adapted to the changing environment as the current Great Barrier Reef was formed. When sea level was much lower, Traditional Owners inhabited what is now the sea floor of the Great Barrier Reef. Any disturbance of the sea floor is considered to impact Indigenous cultural and heritage values. There can be sea burial sites, sacred sites and sites of other cultural significance in the areas where **dredging** is undertaken and, previously, inadequate consultation with Traditional Owners has meant some of these values have been affected.<sup>322</sup> For example, places where access channels are dredged for island resorts or communities are usually also common access points for Traditional Owners, identified and used for thousands of years. In addition, dredging activities can disturb cultural practices and sacred sites. For example, Traditional Owners raised concerns about dredging impacts on local plants and animals in Cleveland Bay.<sup>319</sup> Furthermore, it is reported that any prehistoric archaeological remains in that port area would likely have been destroyed by the prolonged and repeated dredging.<sup>319</sup>

For Traditional Owners, the reefs of the Great Barrier Reef have many stories associated with them. Cultural practices and lore linked with story places are ‘broken’ or affected if there is a **ship grounding** on those reefs. Particular examples include Piper Reef where the *Peacock* ran aground in 1996<sup>323</sup> and the *Doric Chariot* in 2002<sup>324</sup>. Piper Reef is an important story place for its Traditional Owners and these groundings are likely to have affected the cultural heritage of the site. Sudbury Reef, where the *Bungu Terati Satu* ran aground in 2000,<sup>325</sup> is not only a story place but an important cultural place where young men go for traditional rite of passage<sup>326</sup>.

**Cyclones** change land and seascapes, changing places of significance to Traditional Owners’ and affecting Indigenous structures. Subsistence limits on marine resources are affected, potentially resulting in moratoria on Traditional Owner hunting and temporary changes to cultural practices.<sup>327</sup> Cyclones have also had effects on important cultural sites. For example, cyclone Charlie (1988) and cyclone Aivu (1989) destroyed about 50 per cent of the archaeological sites recorded at Upstart Bay in 1987.<sup>328</sup> The recent increases in **freshwater inflow** have affected areas that Traditional Owners use for subsistence. The effects on seagrass meadows and cultural keystone species, such as marine turtles and dugongs, have resulted in Traditional Owners deciding to apply moratoria on hunting while the habitats recover or while the animals are in poor condition.<sup>327,329</sup> That, in turn, means places important for cultural tradition may not be visited, or stories and songlines might not be practiced or passed down to younger generations, because those aspects are directly related to fishing, collecting or hunting activities.

Traditional Owners have observed impacts on Indigenous heritage values from **rising sea levels**. For example, the fish traps in Giringun country in the Cardwell area are being affected.<sup>330</sup> Once culturally significant sites are affected, then stories and songlines are compromised and customary practice has to be changed.

Understanding Table 6.9	
Grading statements	
	<b>No effect:</b> No known interaction or the interaction is considered insignificant.
	<b>Data deficient:</b> Effects are unknown.
	<b>Very low effect:</b> Any effects attributable to the impact are minor or localised, with no observable effects on the values.
	<b>Low effect:</b> The effects of the impact are observable in some locations or to some heritage sites or values, but only to the extent that limited additional intervention would be required to maintain the values.
	<b>High effect:</b> The effects of the impact are obvious in many locations or for many heritage sites or values to the extent that significant additional intervention would be required to maintain the values.
	<b>Very high effect:</b> The effects of the impact are widespread to the extent that the values are severely compromised.
Trend	
↑	<b>Increasing:</b> The intensity and/or spatial extent of the impact is increasing.
↔	<b>Stable:</b> The intensity and/or spatial extent of the impact is staying about the same.
↓	<b>Decreasing:</b> The intensity and/or spatial extent of the impact is decreasing.

**Table 6.9 Past and present effect of impacts on Indigenous and historic heritage values**

The assessment of effects on Indigenous heritage takes into account the interconnectedness of nature and culture for Indigenous people. It reflects effects on the Region's natural environment and other aspects of Indigenous heritage such as cultural practices, stories and language. \* indicates impact is likely to be the result of a variety of activities.

	Impacts	Future trend of the impact	Cultural practices, observances, customs and lore	Sacred sites, sites of particular significance, places important for cultural tradition	Stories, songlines, totems and languages	Indigenous structures, technology, tools and archaeology	Places of historic significance — historic shipwrecks	Places of historic significance — World War II features and sites	Places of historic significance — lightstations	Places of historic significance — other	Places of scientific significance	Places of social significance — iconic sites
Climate change	Altered ocean currents	↑										
	Cyclone activity	↑										
	Increased sea temperature	↑										
	Ocean acidification	↑										
	Rising sea level	↑										
Catchment run-off	Increased freshwater inflow	↑										
	Nutrients from catchment run-off	↔										
	Outbreak of crown-of-thorns starfish	↑										
	Pesticides from catchment run-off	↔										
	Sediments from catchment run-off	↔										
Degradation of coastal ecosystems	Urban and industrial discharge	↔										
	Acid sulphate soils	↑										
	Artificial barriers to flow	↑										
	Atmospheric pollution	↑										
	Coastal reclamation	↑										
Direct use	Light impacts (artificial)	↑										
	Modifying supporting terrestrial habitats	↑										
	Dredging	↑										
	Dumping and resuspension of dredge material	↑										
	Exotic species and diseases*	↑										
	Extraction — death of discarded species	↔										
	Extraction — fishing spawning aggregations	↔										
	Extraction — herbivores	↔										
	Extraction — lower order predators	↔										
	Extraction — lower trophic orders	↔										
	Extraction — top order predators	↔										
	Illegal fishing and poaching	↑										
	Marine debris*	↑										
	Noise pollution	↑										
	Outbreak of disease*	↑										
	Outbreak or bloom of other species*	↑										
	Physical damage — fishing	↔										
	Physical damage — other	↑										
	Physical damage — ship grounding	↔										
	Spill — large chemical	↑										
	Spill — large oil	↑										
	Spill — small chemical and oil	↑										
	Vessel strike on wildlife	↑										
	Waste discharge from a vessel	↑										
	Wildlife disturbance	↑										

The effect of **pesticides** on the natural environment has a flow-on effect on the cultural values of Traditional Owners. Marine animals and plants affected by pesticides are likely to be linked to a range of Indigenous heritage values.

Cultural practices have been affected by **urban and industrial discharge** including heavy metal contaminants. High levels of heavy metals have been detected in the livers of turtles<sup>331</sup> and dugongs,<sup>128</sup> which could pose health risks to Indigenous people. In the Torres Strait, there is a program designed to encourage Traditional Owners not to eat the liver of dugong or turtle for health reasons. In the Gladstone region, some Traditional Owners consider the health of turtles to be so poor they are not prepared to take animals for cultural reasons.<sup>322</sup>

Traditional Owners report that **physical damage** to reefs, cays and the seabed has affected sacred sites and sites of cultural significance. For example, boats often anchor in bays near Hinchinbrook Island where the Haven fish traps are located. These fish traps are of extremely high cultural significance to Giringun Traditional Owners.<sup>289</sup> In turn, this can have an effect on cultural observances, songlines and storylines.

Traditional Owners have expressed concern about reductions in seagrass due to **physical damage from fishing** such as trawling in shallow water and its effect on their access to marine resources.<sup>289</sup>

## 6.6.2 Historic heritage values

The historic heritage of the Region includes its lightstations, wrecks, places of significance and iconic sites as described in Chapter 4. A structured assessment of the effects of past and present impacts on this heritage is presented in Table 6.9.

Many of the impacts considered are assessed to have not affected the Region's historic heritage. Of those considered to be affecting historic heritage, the effects have been assessed as either: minor or localised with no observable effects on the values; or observable in some locations, but only to the extent that limited additional intervention would be required to maintain the values.

**Cyclones** have affected historic shipwrecks, World War II features and sites, and lightstations. A recent example is the effect of cyclone Yasi on the wreck of the *SS Yongala* in 2011, including moving some parts and 'sandblasting' its exterior. Throughout historical times, cyclones have damaged historic structures such as resorts.<sup>310</sup>

The connections between the natural environment and the scientific activities historically undertaken in the Region means the heritage value of places of scientific significance is affected to some extent by the impacts affecting the places' natural values.

## 6.7 Impacts on community benefits of the environment

The benefits that Reef-dependent communities and industries derive from the Region's environment are founded on its biodiversity, geomorphological features, and Indigenous and historic heritage values. The quality and magnitude of community benefits, including its aesthetic value, are therefore affected by impacts that diminish the condition of those values.

The depth and extent of the flow-on effects to community benefits as a result of impacts on biodiversity, geomorphology and heritage values are difficult to assess. However, it is recognised that the economic and employment benefits derived by Reef-dependant industries (such as tourism, commercial fishing and some aspects of recreational use) are at highest risk and stand to be the most affected by a declining ecosystem.

Because everyone's interaction with the Reef is different, effects on community benefits derived from the environment will depend very much on each Reef user's individual perspective.

In broad terms:

- Income and employment benefits are diminished by impacts that affect the productivity and attractiveness of the environment, such as those related to catchment run-off, loss or modification of coastal habitats, climate change, declines in iconic species and a reduction in the aesthetic qualities of the Region. These qualities are essential for Reef-dependent industries like tourism, fishing, recreational use and traditional use of marine resources.
- The benefits derived by people understanding, appreciating, enjoying and admiring the Region's environment are most affected by those impacts that significantly affect key values — in particular coral reef habitats and species, the quality and clarity of the Reef's water, the land and seascapes and iconic species such as dugongs, marine turtles, sharks, whales and dolphins.
- The ability for people to access the Reef's resources is affected by the impacts that diminish those resources, such as those that affect coral reefs as a place to visit and populations of targeted fish species. In this regard, declines in top predators could affect the commercial and recreational fisheries that target these species, and tourism (particularly dive tourism) for which big sharks and fishes are important natural attractions.<sup>332</sup> A high density of vessels at anchor at a ship anchorage area can interfere with other user's ability to effectively use that area (for example, tourism and commercial, recreational and traditional fisheries).<sup>270</sup>



- People's personal connection with the Reef's environment is diminished by the range of impacts that degrade the quality of its key habitats, iconic locations and iconic species such as those related to catchment run-off, climate change and declines in species of conservation concern. In particular, Traditional Owner connections are affected by any impacts that affect their sea country and their ability to maintain their culture in relation to it.
- The health benefits people derive from the Region are diminished by those impacts that make the Reef a less attractive and fulfilling place to visit, and by those that reduce the quality and availability of its wild-caught food resources.
- The aesthetic values of the Region may be diminished by development activities. For example the building of structures, industrial and port developments, and coastal reclamation could affect the natural scenic values of the coastal areas. Marine debris, along with oil and chemical spills, also affects the aesthetic value of seascapes and islands. Coastal reclamation may affect aesthetic qualities of the landscape and seascape.<sup>333</sup> Increased turbidity diminishes the Region's underwater aesthetic values.<sup>333</sup> Artificial light and noise pollution associated with coastal development and increased shipping activities and anchorage areas<sup>270</sup> may diminish aesthetic attributes such as tranquillity, solitude and remoteness.<sup>333</sup>

The concept of 'shifting baselines' (see Section 7.1) is relevant in considering the extent to which the community benefits derived are affected by changes in the condition of the Region's biodiversity, geomorphological features, and Indigenous and historic heritage values. Over time, many users have adapted their activities and expectations in response to the changing condition of the Reef environment.

## 6.8 Cumulative impacts

The examination of the effect of impacts on individual values presented in the previous sections identifies the impacts having the most significant effects on key values relevant to matters of national environmental significance in the Region. However, this one-on-one assessment does not take into account cumulative impacts — the successive and combined effects of impacts that overlap in time or space on the environment. In this section, both qualitative models and spatial analysis techniques are applied to investigate the complex interactions between some of the key impacts and values.

### 6.8.1 Qualitative models

As part of the Sustainable Regional Development Program project to develop a resilience framework to inform decision making in the Great Barrier Reef coastal zone<sup>334</sup>, qualitative models were developed for coral reef and seagrass ecosystems, and dugong (a species dependent on seagrass). The models examine the relationships of drivers, activities, impacts and values, and were developed through workshops with experts in coral reef and seagrass biology and ecology

They broadly detail the main variables and effects — excluding minor species groups and weak effects. The models outlined below are only preliminary and will require further refinement and validation before implementation.

A basic feature of qualitative models is the development and analysis of sign-directed graphs, or signed digraphs. These are used to describe the main interacting variables within a system, linking them to their surrounding ecosystem and also to the drivers, activities and impacts of concern. Analysis of the structured lists (for example Table 6.2) and value-impact matrices (for example Table 6.6 to Table 6.9) helps inform assessment of the relative importance of drivers, activities and impacts on values, and how they affect the system. While model links are qualitative — they represent only the 'sign' of the effects (that is, positive, negative or nil) — they, nonetheless, provide a useful means to formally assess a system's dynamics and its response to disturbances.

## Modelling approach

The use of models for the strategic assessment process is guided by a strategy of model building that ultimately seeks to combine different modelling approaches to better understand, predict and intervene in complex ecological and socio-economic systems.<sup>335,336</sup> The underlying premise is that there are three desired properties of models: *generality*, *realism* and *precision*. A model that attempts to maximise all three properties, however, quickly becomes impractical to apply and difficult to understand. There needs to be a trade-off — a choice to emphasise two properties over a third, leading to three alternative modelling approaches.

*Quantitative process models* — Models that emphasise precision and realism over generality result in precise predictions for highly specified details of a system's biological and ecological components, processes and relationships. Such models produce precise and testable predictions, and are favoured by managers who ask 'how much do I have to spend on X to get this amount of Y.' Developing these models, however, typically requires extensive amounts of data. They are tuned to the specific context from which their data were drawn (that is, they lack generality), and thus are not easily transferred to new applications without additional data gathering, tuning of parameters and calibration.

*Statistical models* — Models that emphasise precision and generality over realism result in precise testable predictions based on correlations among system variables. They are useful for describing general patterns with measured confidence and are more easily exported to new situations, but again, they come with a burden of data acquisition and they provide less understanding of the underlying processes, as correlation is not synonymous with causation.


*Qualitative process models* — Models that emphasise realism and generality over precision are free from the constraint of extensive and expensive data collection, and while predictions are not precise, they are nonetheless rigorous in their derivation and testable. They are based on the qualitative interpretation of processes and do not require exact specification of functions or parameter values. The process of building a qualitative process model requires answering the general questions of: 'does a variable have a positive or negative effect on another variable?' and 'is a function increasing or decreasing, above or below a threshold?' Predictions are expressed in terms of directional change in a variable (that is, increase, decrease, no change) or as inequalities that highlight key relationships or parameters in the system.

Each approach has its own inherent strengths, and each can provide useful results. Models based on each approach will necessarily be derived from different simplifying assumptions, and each purposefully leaves something out about how the world works. The dilemma is not about which approach is better — each is incomplete but serves the common goals of understanding, prediction and intervention. Rather, the aim is to confront the problem of complexity with an array of alternative models drawn from each approach. Then, if the different models arrive at similar results, even though they are based on different assumptions, a robust concept of the system has been developed that is relatively independent of model details.<sup>336</sup>

While there will always be a need to address management questions with precision, the scale, complexity and pace of change affecting the Great Barrier Reef Region presents significant limitations to the widespread application of quantitative models. However, many — if not most — of the management objectives for the Region can be meaningfully interpreted and approached qualitatively. The challenge is to match the relative scale and complexity of the management problem with the appropriate set of analytical tools and to deliver practical results in a timely manner.

## Coral reef ecosystem model

The coral reef model (Figure 6.26) provides a generalised depiction of some of or the key processes that enhance or diminish corals, and the role that coral plays in supporting biodiversity. Corals are strongly dependent on successful coral recruitment, compete for space with macroalgae, and provide critical habitat and resources for fishes and invertebrates. Crown-of-thorns starfish can strongly affect corals during outbreaks, and it is hypothesised that such outbreaks are enhanced by nutrients. Drivers, activities and impacts on coral reefs do not stand alone, but are intertwined in a complex web of synergistic or cumulative impacts. The model shows some cause-and-effect relationships are relatively simple, such as an increase in ocean warming increasing the frequency of coral bleaching events, which then leads to a reduction in coral cover. However, an increase in catchment run-off from agriculture leads to increases in four impacts (toxins, nutrients, turbidity and sedimentation) that affect a total of seven ecosystem variables (predatory fish, herbivorous fish, crown-of-thorns starfish, fish and invertebrates, macroalgae, crustose coralline algae and coral recruitment, and coral cover).



**Figure 6.26 Qualitative model of how coral reefs are affected by drivers, activities and impacts**


The model illustrates how coral reefs are affected by multiple impacts that result from various direct drivers and activities. It broadly details the main variables and effects, and excludes minor (rare) species groups and weak effects. Links describe direct positive or negative effects of one variable on another. There are a number of links that are uncertain or contentious. These are represented by dashed lines and provide the basis to consider alternative model structures in subsequent analyses (see Chapter 11).

### Seagrass meadows model and dugongs

The seagrass model (Figure 6.27) represents the multiple pressures on seagrass meadows. These meadows can be composed of fast-growing species of seagrass favoured by frequent disturbances and high grazing pressure (known as 'r' species) or slow-growing species favoured by conditions of low disturbance and low levels of grazing (known as 'K' species). One of the variables in the model is the ratio of 'r' and 'K' species, representing the relative dominance of these two types of species. In the model, some variables such as dugong populations or storms can act to shift the relative balance between these two types of seagrass species. Other factors that regulate seagrass include epiphytic algae which grow on the surface of seagrass and can inhibit their growth through shading.

Similar to the model for coral reef ecosystems, the seagrass meadows model shows some cause-and-effect relationships are relatively simple, such as an increase in turbidity and sedimentation suppressing seagrass distribution and abundance, which then leads to a reduction in dugongs. An increase in catchment run-off from agriculture, however, leads to the increase of four impacts (toxins, nutrients, turbidity and sedimentation) that affect a total of five ecosystem variables (scrapers (prawns and fishes), marine turtles, dugongs, epiphytes and seagrass distribution and abundance). There are both positive and negative influences on the relationship between catchment run-off and seagrass distribution and abundance. There are also thresholds for which positive influences can become negative. For example, seagrasses are generally considered nutrient-limited and thrive under conditions of increased nutrients. However, after a certain threshold, they can deteriorate due to light reduction from an increase in epiphytes, macroalgae or phytoplankton.<sup>337</sup>





**Figure 6.27 Qualitative model of how seagrass meadows are affected by drivers, activities and impacts**

The model illustrates how seagrass meadows are affected by multiple impacts that result from various drivers and activities. It broadly details the main variables and effects and excludes minor (rare) species groups and weak effects. Links describe direct positive or negative effects of one variable on another. There were a number of links that were uncertain or contentious. These are represented by dashed lines and provide the basis to consider alternative model structures in subsequent analyses (see Chapter 11).

## 6.8.2 Spatial approaches

Mapping approaches show the spatial distribution of interactions between values and impacts. As an initial step to understanding combined or cumulative impacts across the Region, standard geographic information system analysis, where individual impact gradients are standardised, has been undertaken for both coral reefs and water quality as described in Brodie *et al.* 2013.<sup>338</sup>


The coral reef analysis examines the amount the reefs throughout the Region have been exposed to identified key impacts. The water quality analysis examines the risk that the ecosystem has faced as a result of key water quality impacts, taking into account the likely vulnerability of various ecosystem components.

### Coral reefs

As outlined in Chapter 7, the primary drivers of coral decline in the Region have been crown-of-thorns starfish, cyclones and coral bleaching events. Coral bleaching is typically caused by elevated temperatures but can also be caused by freshwater exposure. Figure 6.28 presents the exposure of reefs to each of these key impacts for 2001 to 2011. These individual impacts are then combined and scaled in a similar way for the cumulative exposure map (Figure 6.29).




It is important to note this analysis does not represent all past and current impacts on coral reefs. In particular, many notable impacts to coral reef systems occurred before the decade presented — notably, major crown-of-thorns starfish outbreaks in the 1980s and 1990s, and the 1998 mass coral bleaching event. The 2001–2011 period was chosen based on the availability of remote sensing data and because the decade covers a range of El Niño and La Niña conditions.



**Figure 6.28 Exposure of coral reefs to key impacts, 2001–2011**

The key impacts analysed are (a) damaging waves from cyclones, (b) crown-of-thorns starfish outbreaks, (c) thermal stress events and (d) freshwater exposure. Individual impacts are scaled between zero and one, with the most exposed reef areas being scored at one.



**Figure 6.29 Cumulative exposure of coral reefs to key impacts, 2001–2011<sup>339</sup>**

The map presents the modelled cumulative exposure of coral reefs to cyclone-induced waves, crown-of-thorns starfish outbreaks, elevated sea surface temperatures and freshwater inflow. The magnitude of the impacts has been normalised between zero and one, where the highest score was 2.40 out of a maximum possible of 4.

## Water quality

Freshwater influxes from catchment run-off, and the load of nutrients, sediments, pesticides and other pollutants they carry, affect the marine ecosystems in the Region, particularly inshore Great Barrier Reef habitats and species.<sup>77</sup> Individual habitats and species will be affected by these impacts in different ways, depending on how much it is exposed and how sensitive it is. A relative risk assessment of degraded water quality on Great Barrier Reef ecosystems has been conducted to inform implementation of the Reef Plan and Reef Rescue programs, and has substantially improved understanding of the combined effects of water quality.<sup>340</sup> The outcomes of this risk assessment are reported below.

The exposure of a species or habitat to an impact is typically a function of the intensity of the impact (the concentration or load of a pollutant) and the length of time it is exposed to the impact. For example, a seagrass meadow may be exposed to a high intensity impact for a short period of time, or to lower intensities for longer periods. When quantifying exposure, it is important to determine the threshold concentrations — the concentration which potentially leads to damage or mortality within hours or days — for each habitat or species, as well as determining long-term average concentrations and the duration of exposure.<sup>340</sup> This complicates the description of exposure thresholds given their values may change by one to two orders of magnitude between days, seasons and years.<sup>340</sup> Hence, some key water quality variables, such as suspended sediments, are divided into different thresholds based on ecological responses, periods of exposure and the potential severity of the impact on habitats.<sup>340</sup> Due to limitations in the spatial and temporal availability of data, this assessment defines relative risk as the total area of coral reefs and seagrass meadows in the highest severity classes for several water quality variables.

As outlined in Section 6.4.2, the key impacts in relation to water quality in the Region are nutrients, sediments and pesticides in catchment run-off. Based on these three broad impacts, the key variables examined in the combined impact analysis are:

- Concentrations of **total suspended solids above two milligrams per litre** (Figure 6.30a). This variable correlates strongly with declines in ecosystem condition such as increased growth of macroalgae and declining diversity.<sup>341</sup> Two milligrams per litre is the average annual threshold in the Great Barrier Reef water quality guidelines.<sup>127</sup> The data is extracted from daily remote sensing imagery (at a resolution of one square kilometre) and processed using methods described in Brando *et al.* (2013).<sup>103</sup> The map shows the frequency of valid daily observations where the total suspended solids exceeded two milligrams per litre in the 10 year period from 1 November 2002 to 30 April 2012.
- Concentrations of **total suspended solids above 6.6 milligrams per litre** (Figure 6.30b). This concentration of suspended solids is equivalent to a turbidity of five nephelometric turbidity units.<sup>340</sup> It has been used in the analysis as it has been shown to have various ecosystem effects including coral stress,<sup>342</sup> declines in seagrass cover,<sup>343</sup> fish habitat choice,<sup>344</sup> home range movement<sup>345</sup> and (above 7.5 nephelometric turbidity units) foraging and predator-prey relationships<sup>346,347</sup>. The data is extracted from daily remote sensing imagery (at a resolution of one square kilometre) and processed using methods described in Brando *et al.* (2013).<sup>103</sup> The map shows the frequency of valid daily observations where the total suspended solid concentration exceeded the threshold of 6.6 milligrams per litre (or five nephelometric turbidity units) in the 10 year period from 1 November 2002 to 30 April 2012.
- **Surface exposure to total suspended solids** (Figure 6.30c). An assessment of the frequency and extent of the influence of flood plumes containing differing concentrations of total suspended solids provides an estimation of the extent of surface exposure of habitats during wet season conditions. In the map, surface exposure is modelled using an assessment of plume frequency from satellite imagery and monitored end-of-catchment loads in each wet season (November to May) from 2007 to 2011, as detailed in Devlin *et al.* (2013).<sup>145</sup> The mean of the five annual distributions was selected as a way of factoring in inter-annual variability in river discharge, although it is recognised that this period was characterised by several extreme rainfall events.
- **Chlorophyll** concentrations (Figure 6.30d). Chlorophyll is an indicator of nutrient enrichment in marine waters. Death and Fabricius (2008)<sup>93</sup> identified 0.45 micrograms per litre as an important ecological threshold for macroalgal cover, hard coral species richness and octocoral species richness. This concentration is identified as the annual average threshold for chlorophyll in the Great Barrier Reef water quality guidelines.<sup>127</sup> Significant benefits for the ecological status of reefs in the Region are likely if mean annual chlorophyll concentrations remain below this concentration. The data is extracted from daily remote sensing imagery (one square kilometre resolution) and processed using methods described in Brando *et al.* (2013).<sup>103</sup> The map shows the frequency of valid daily observations where the chlorophyll concentration exceeded the threshold of 0.45 micrograms per litre in the 10 year period from 1 November 2002 to 30 April 2012.




- **Surface exposure to dissolved inorganic nitrogen** (Figure 6.31e). This variable is a direct indicator of nutrient enrichment. High concentrations of dissolved inorganic nitrogen can reduce coral calcification and increase abundance of macroalgae.<sup>348</sup> There is typically more dissolved inorganic nitrogen in waters adjacent to catchments with intensive agriculture. Surface exposure is modelled annually using plume frequency data from satellite imagery and monitored end-of-catchment loads in each wet season as described in Devlin *et al.* (2013).<sup>145</sup> The mean of five annual distributions in the period 2007 to 2011 was selected as a way to factor in inter-annual variability in river discharge, although it is recognised that this period was characterised by several extreme rainfall events.
- **Photosystem II inhibiting (PSII) herbicides** such as diuron, atrazine, simazine, ametryn, hexazinone and tebuthiuron (Figure 6.31f). These chemicals can disrupt photosynthetic activity in marine plants including mangroves, seagrass, coral zooxanthellae and crustose coralline algae. The map is based on an estimate of the relationship between PSII herbicide concentrations and coloured dissolved organic matter (a proxy for salinity) in flood plume conditions in 2010–2011. Risk categories are based on thresholds of diuron for coral and seagrass species: no risk (less than 0.025 micrograms per litre), insignificant (0.025 to 0.1 micrograms per litre), minor (0.1 to 0.5 micrograms per litre), moderate (0.5 to 2.3 micrograms per litre), and major (2.3 to 10 micrograms per litre). This is likely to be an underestimate of the actual risk due to limitations of the method, as outlined in Brodie *et al.* (2013).<sup>338</sup>
- **The crown-of-thorns starfish initiation zone** (Figure 6.31g). This zone, between Cairns and Lizard Island, has been identified as the area where there is the highest likelihood that a primary outbreak of crown-of-thorns starfish will begin. The four primary outbreaks originating in this zone since 1960 all began two to five years after wet seasons when aggregate discharges early in the season (November to February) from the Burdekin to Daintree Rivers exceeded 10 million megalitres.<sup>349</sup> The area defined to be at highest risk in initiating crown-of-thorns outbreaks is between latitudes 14.5 degrees south and 17 degrees south and is described in Furnas *et al.* (2013).<sup>349</sup>

For most of the maps of the individual variables presented in Figure 6.30 and Figure 6.31, the data is scaled on a five-point scale. The maps of exposure to total suspended solids and dissolved inorganic nutrients are based on 2007–2011 data and represented on a three-point scale. The crown-of-thorns initiation zone is simply mapped.

The combined effect of water quality in the Region has been examined spatially by combining the individual variables and normalising the outcomes on a five-point scale (Figure 6.32).

In preparing the analysis, it has been recognised there remains many issues, complexities, variability, uncertainties and unknowns surrounding the understanding of all elements and the extent and effects of catchment run-off.






**Figure 6.30 Key water quality exposures in the Region (part 1)**

(from Brodie *et al.* (2013)<sup>338</sup>)

Key water quality parameters analysed are:

part 1 — (a) total suspended solids, 2 milligrams per litre exceedance (2002–2012); (b) total suspended solids, 6 milligrams per litre exceedance (2002–2012); (c) total suspended solids exposure (2007–2011); (d) chlorophyll concentrations, 0.45 micrograms per litre exceedance (2002–2012);

part 2 — (e) dissolved inorganic nitrogen exposure (2007–2011); (f) photosystem II inhibiting (PSII) herbicide exposure (2010–2011); and (g) crown-of-thorns starfish initiation zone.




**Figure 6.31 Key water quality exposures in the Region (part 2)**

(from Brodie *et al.* (2013)<sup>338</sup>)

Key water quality parameters analysed are:

part 1 — (a) total suspended solids, 2 milligrams per litre exceedance (2002–2012); (b) total suspended solids, 6 milligrams per litre exceedance (2002–2012); (c) total suspended solids exposure (2007–2011); (d) chlorophyll concentrations, 0.45 micrograms per litre exceedance (2002–2012);  
part 2 — (e) dissolved inorganic nitrogen exposure (2007–2011); (f) photosystem II inhibiting (PSII) herbicide exposure (2010–2011); and (g) crown-of-thorns starfish initiation zone.



**Figure 6.32 Combined key water quality exposures**

(from Brodie *et al.* (2013)<sup>338</sup>)

Combined assessment of: total suspended sediments (exceedance of 2 milligrams per litre and 6 milligrams per litre thresholds, 2002–2012, and average annual surface exposure, 2007–2011); nutrients (chlorophyll exceedance of 0.45 micrograms per litre thresholds, 2002–2012, and dissolved inorganic nitrogen average annual surface exposure, 2007–2011), PSII herbicide exposure, 2010–2011, and crown-of-thorns starfish initiation zone.



### 6.8.3 Assessment of outcomes arising from cumulative impacts

#### Cumulative impacts on the Great Barrier Reef

The preliminary qualitative models for coral reefs and seagrass meadows show some cause-and-effect relationships are relatively simple, while others are far more complex involving a large number of network pathways and feedbacks. While it is recognised that the models are preliminary, they demonstrate the ways in which different components are likely to interact and the consequences that changes in the system are likely to have on some of the Region's values. It is however acknowledged that these models require validation through further monitoring and experimental studies before they provide a complete picture of cumulative impacts.

The spatial analysis of the key impacts on coral reefs and water quality shows the areas of highest exposure to cumulative impacts are inshore, particularly in the southern two-thirds of the Region. The results illustrate the accumulating nature of major pressures on reef and non-reef areas in the Region over the last decade. They provide greater understanding of the scale at which these pressures affect the Region's values.

Over the past 10 years, coral reefs between Cairns and Townsville have experienced the highest cumulative impacts from cyclones, thermal stress, crown-of-thorns starfish outbreaks and increased freshwater inflow. The spatial assessment also shows almost all coral reefs in the Region have been exposed to one or more of these threats over the last decade (Figure 6.29).

Inshore areas, particularly those south of Cairns, are most at risk of poor water quality. Areas along the coast in the vicinity of Hinchinbrook Island, Townsville and Gladstone showed a very high combined water quality risk, whereas the water quality risk in areas north of Cairns and in offshore areas is distinctly lower (Figure 6.32).

Recognising that the inshore waters of the Region are already under significant stress from the decline in water quality driven by increased land-based run-off of suspended solids, nutrients and pesticides<sup>78</sup>, the uncertainty regarding the additional effects of sea dumping of dredge material is a key concern, particularly given the potential for large volumes to be dumped.

The frequency and timing of dredge material disposal can also significantly affect for the cumulative impacts on ecosystems. Catchment inputs occur as strong but highly variable pulses during wet seasons, and are ongoing (decadal scale). Dredge inputs involve major pulses during periods of capital dredging, with ongoing maintenance dredging typically occurring in the dry season. Emerging evidence shows turbidity effects of flood plumes may persist for months after major wet seasons.<sup>78,138</sup> If dredging activity results in increased turbidity during otherwise natural clear periods of the year, the combined impacts on ecosystems may be considerably greater than either impact alone. Significant volumes of sea dumping over the next five to 10 years has the potential to add further pressure to already declining inshore ecosystems.

It is important to note the spatial analyses presented in Section 6.8.2 only cover the 10-year period for which accurate and comparable records of these impacts are available. In addition, the analysis does not include many other relevant impacts, all of which combine to affect the natural systems of the Region. Research is currently underway to build a more complete, dynamic understanding of cumulative impacts to guide future management actions to support the resilience of the Reef.

#### Cumulative impact assessment methods


The analyses provided in this chapter demonstrate that effective cumulative impact assessment requires a suite of tools and approaches to address the challenges of complexity, scale and pace of change. It highlights the need for the systematic application of the tools and approaches to improve understanding of the cause-and-effect relationships of multiple and compounding impacts, and the spatial and temporal scales at which processes and impacts are operating.

It demonstrates the importance of understanding the zones of influence of different activities (the sources of impact) and the need to be particularly vigilant in areas where zones of influence overlap and cumulative impact intensity is highest. Determination of zones of influence requires the use of geographic information systems to spatially identify, map and model activities and impacts and their likelihood of occurring. It also requires 'breaking up' of impacts from point and non-point sources into intensity categories. For example, in Figure 6.33 impact gradients from a point-source discharge from an urban development, and more diffuse impacts such as water quality changes associated with agricultural run-off from catchments, have been divided into high, moderate and low intensities. Their overlapping zones of influence create a mosaic of impact intensities that, combined, have an effect on the values of the area.

Importantly impacts vary in both time and space and with differing levels of severity. For example, major storm and flood events may only occur once every two decades, yet their impact in a flooding year can be severe with a legacy that persists for decades. Hence, the determination of the zone of influence of individual and cumulative impacts requires an understanding of both the spatial and temporal dimensions of the activity or event and its associated effects.

The assessment of impacts in this report also highlights the need for point-source impacts to be assessed at a scale that takes into consideration the influence of other human-related activities and ecological processes operating in the Region.





**Figure 6.33 Zones of influence of different activities**

Zones of influence of different activities (sources) overlap to create a mosaic of impact intensities.

Because quantitative process models need large amounts of data, they have only had limited use in relation to the Region. A staged and complementary approach allows qualitative process models to be used for initial assessments of cumulative impacts, and for them to be employed where there are critical management questions and sufficient data. Knowledge gained from analysis and testing of qualitative models can be used to better focus the application of and inform the construction of quantitative process models.

The spatial modelling examples demonstrate that this approach is feasible. However, a strong understanding of the vulnerability, extent and significance of habitats, as well as their exposure to particular impacts, is required. A basic requirement is systematic habitat classification and mapping. Spatial modelling of cumulative impacts also depends upon an understanding of the extent, linkages and importance of ecosystem processes. These processes act as vectors that redistribute influences or pollutants from a source to a wider zone (for example, currents distribute coral and fish larvae and may also disperse crown-of-thorns starfish larvae and dredge spoil over similar scales and locations). Mapping these influences is crucial to understanding the extent of their impacts.

While significant investment in monitoring and research has increased our understanding of the effects of the pollutants and their interactions, effects on a wide variety of species or habitats are still not fully understood. Each of the numerous inshore species has its own tolerance limit, including at different life stages. Interactions between the organisms and pollutants also add to the complexity. Significant progress has been made, in particular, in identifying thresholds and interactive effects of multiple stressors on seagrass. The Reef Rescue marine monitoring program science providers have developed metrics that include biological indicators of health that have potential for inclusion in revised guidelines.<sup>350</sup>

The water quality spatial analysis has highlighted a paucity of data relating to the toxicity of many contaminants to primary producers — the basis of all food webs — in tropical marine ecosystems, as well as a lack of a comprehensive understanding of biogeochemical cycling.

## 6.9 Effects on matters of national environmental significance

Based on the assessment of impacts on key values and attributes relevant to matters of national environmental significance, presented above, the following is a summary of the effects on each matter.

### 6.9.1 World heritage properties

The past and present effects on the outstanding universal value of the Great Barrier Reef World Heritage Area are summarised in Table 6.10. It is structured around the four world heritage criteria relevant to the property and considers those key values and attributes identified in Table 4.8, in Chapter 4.

The impacts assessed have also had some effect on the integrity of the World Heritage Area. While all elements necessary to express its outstanding universal value remain largely intact and the area continues to be an

adequate size to ensure complete representation of its features and processes, there have been adverse effects on the property, such as a 50 per cent average decline in coral cover since 1987<sup>36</sup>. The legacy impacts of activities in the Region, combined with impacts arising externally, have meant that the condition of some values is in decline.

**Table 6.10 Summary of impacts affecting outstanding universal value**

Short title of world heritage criteria	Summary of most severe impacts
<b>Major stages of the Earth's evolutionary history</b>	The assessment of impacts on geomorphological features shows the main impacts affecting them are cyclones and increased sediments. Cyclones have the ability to alter and weaken the structure of the geomorphological features, while sediments affect the features that are reliant upon photosynthesis (such as coral reefs and seagrass meadows) and can infill palaeochannels.
<b>Ecological and biological processes</b>	Modifying supporting terrestrial habitats has changed hydrological processes in many parts of the catchment. This has had downstream impacts on ecological and biological processes, reducing connectivity and the health of the World Heritage Area. The strong ongoing links between Aboriginal and Torres Strait Islanders and their land and sea country have been and continue to be affected by those impacts which are diminishing the natural environment of the Region.
<b>Natural beauty and phenomena</b>	Combined effects from cyclones, declining water quality and increasing sea temperature are affecting on the natural beauty and phenomena of the Great Barrier Reef ecosystem. Coastal habitat degradation and highly visible activities, such as coastal development, dredging and shipping, may have also diminished natural scenic values, especially in coastal areas. Transient impacts to the aesthetic experience of a place can include large numbers of tourists, passing ships or turbidity due to weather. Natural phenomena such as nesting marine turtles have been affected by previous commercial harvesting which has increased the species' vulnerability to subsequent impacts. The most severe current impacts affecting nesting marine turtles include death of discarded species and illegal fishing and poaching.
<b>Habitats for conservation of biodiversity</b>	Cyclones, increasing sea temperature, crown-of-thorns starfish outbreaks and increased freshwater inflow are threatening habitats and species. Almost all areas of the Region have been exposed to one or more of these threats over the last decade. The assessment shows the greatest impacts to seagrass and coral habitats arise from the adjacent catchment (specifically increased freshwater inflow, nutrients, pesticides and sediments) and from extreme weather.

## 6.9.2 Great Barrier Reef Marine Park

The impacts having the greatest effect on the Great Barrier Reef World Heritage Area also apply to the Great Barrier Reef Marine Park. Freshwater influxes from catchment run-off, and the load of nutrients, sediments, pesticides and other pollutants they carry, are affecting the marine ecosystems of the Marine Park. Cumulatively these impacts, along with the other listed severe impacts such as cyclones and those associated with coastal development, mean the southern two-thirds of the Marine Park is experiencing the greatest impacts

The benefits that Reef-dependent communities, industries and Traditional Owners derive from the Region's environment are based on its natural and heritage values. The quality and magnitude of benefits derived are also diminished as a result of the declining condition of the underlying values. Impacts on coral cover (from crown-of-thorns starfish, cyclones and declining water quality) have meant there are fewer opportunities for people to experience and enjoy healthy coral reefs. Illegal fishing and poaching and death of discarded catch have reduced numbers of targeted fish and iconic fauna (for example, marine turtles and dugong), further affecting catch rates and recreational experiences. Poorer water quality (from impacts related to catchment run-off) has changed underwater visibility, affecting appreciation of some coral reefs.

Given the attributes of the biological and geomorphological features of the Marine Park contribute to its aesthetic value, effects on aesthetic value are implicit.<sup>333</sup> Transient impacts to the aesthetic experience of a place can include large numbers of tourists, passing ships or turbidity due to weather. More significant and long-lasting impacts directly affecting tangible attributes of aesthetic values include modifying supporting terrestrial habitats and coastal development.<sup>333</sup>

## 6.9.3 National heritage places

As the Great Barrier Reef was listed as a national heritage place because of its status as a world heritage property, the summary of impacts affecting outstanding universal value (above) are also relevant to this matter of national environmental significance.

#### 6.9.4 Commonwealth marine area

The assessment summary for the condition and trend of the Great Barrier Reef Marine Park above is also relevant to the Commonwealth marine area.

#### 6.9.5 Listed migratory and listed threatened species

The most severe impacts to listed migratory and threatened species include climate-related impacts (increased sea temperature and cyclone activity), impacts that affect nesting, feeding and breeding habitats such as clearing and modifying supporting terrestrial habitats (which may also increase light and noise impacts) and direct impacts from activities that pose a threat to survival (for example, marine debris, chemical and oil spills and wildlife disturbance).

A wide range of listed species are unintentionally captured during the Queensland Shark Control Program, including the grey nurse shark, whale shark, marine turtles and humpback whale.<sup>226</sup> Death of discarded species as a result of fishing practices has high effects on listed sharks, marine turtles and dugong. Trawling remains a risk to some listed species including sharks. The commercial inshore net fishery unintentionally catches a number of species which may be injured or killed in the nets (for example, marine turtles, dugongs, inshore dolphins, sharks and sawfish). Even low levels of mortality may compromise the ability of depleted populations to recover. Extraction of top order predators has a very high effect on listed sharks and, through the legacy impacts of commercial harvesting, has had a high effect on crocodiles. While many top predators are not directly targeted by fishing activities in the Great Barrier Reef, some are unintentionally caught and retained, and several sharks are at high risk from fishing.

Recent severe tropical cyclones have damaged wide areas of marine habitat on which dugong and marine turtles rely. In addition, seagrass mortality as a result of floods, especially in early 2011, is thought to be responsible for severe effects on dugong populations.

Pelagic-foraging seabirds are also highly vulnerable to changes in ocean currents and sea temperature as many search for food in upwellings, eddies and other sea surface temperature gradients.<sup>29</sup>

Listed shorebirds are affected by artificial barriers to flow and clearing and modification of coastal habitats. Previous broadscale land clearing, principally in the southern two-thirds of the Great Barrier Reef catchment, has indirectly caused major changes to the Region's nearshore environment. This has, in turn, affected the shorebirds that use these habitats. In addition, changes to catchment ecosystems, such as infilling wetlands, and constructing levee banks and drainage, have affected connectivity and the habitats for listed shorebirds.

#### 6.9.6 Wetlands of international importance

The Shoalwater and Corio Bay Area is predominantly within a defence training area. The restricted access provisions applying to the defence area reduce the direct human-related impacts (for example, fishing and coastal reclamation). As a defence training area, some impacts from these activities are likely in the marine component of the area (for example noise, boat and aircraft presence) and there is a risk of marine environmental incidents such as oil spills. However, mitigation strategies are in place. Environmental monitoring has shown activities, such as low-flying aircraft, are not posing significant biological threats to the area and the species which rely upon it (for example roosting seabirds).<sup>351</sup> However, the area is not immune to climate-related impacts, with changes in sea temperature and ocean currents likely to affect many species that migrate to the area.<sup>352</sup>

### 6.10 Key information gaps

There is a tendency to consider the impacts on biodiversity and heritage values as distinct entities, but it is the interaction between drivers, activities and, in particular, their cumulative impacts, that are often of greatest concern. Further, complex and cumulative interactions between biodiversity, ecosystem health, and social, cultural and economic values determine the resilience of the natural and human systems — a concept which is fundamental to the protection of the matters of national environmental significance.

An understanding of impacts relies on knowledge of the value being affected, the activity causing the impact (as well as what is driving it) and how they interact. There is a developing understanding of cause-and-effect relationships for many values and impacts in the Region, but the quantitative nature of these interactions is not well known in most cases. Indirect ecosystem effects of impacts are even less well understood. For instance, it is known that fishing removes a portion of top order predators and that this has the effect of reducing predator populations in fished areas, but little is known about the effect on the rest of the ecosystem. Likewise, the introduction of pesticides has direct effects on plants and animals, however the nature of many of these interactions is not well quantified and indirect impacts are even less understood.

Quantitative understanding of cause-and-effect relationships, both now and under a range of future scenarios, will greatly improve understanding of impacts and their cumulative effects — this currently constitutes a major information gap.

### 6.10.1 Biodiversity

The impacts of most concern that have been identified for biodiversity are categorised into the broad areas of: climate change, catchment run-off, degradation of coastal ecosystems and direct use. Greater understanding is required for these and other impacts relevant to the Region, particularly:

#### Climate change

- the likely long-term environmental changes resulting from climate change impacts
- the direct and indirect effects of climate change impacts on the function and behaviour of species and ecosystem processes
- the effect of a changing climate on cyclone activity and ocean currents
- the effects of impacts related to climate change on seabird feeding grounds and nesting areas.

#### Catchment run-off

- the fate of nutrients, sediments and pesticides entering the Region, including where they go and what impacts they have on an individual catchment basis
- the time lags between changes in agricultural practices and improvements in the Region's water quality
- the volume and composition of sediment from various sources, including from dumped dredge material and sediment from catchment run-off, compared with natural levels
- the concentration of nutrients in marine sediments, including the rates and relative contribution to the pool of re-mineralised particulate nutrients
- the trigger levels for pollutants that are appropriate for maintaining the health and function of non-Reef ecosystems within the Region
- pollutants from urban and industrial discharge and their relative contribution to pollutant loads in the Region
- the concentrations of heavy metal and implications of legacy mining activities on water quality
- the extent of pollutant delivery to the Region through groundwater
- the exact locations and numbers of crown-of-thorns starfish outbreaks
- a more operationally efficient technique to control crown-of-thorns starfish outbreaks
- the most effective changes to land management practices for reducing the probability and/or severity of future crown-of-thorns starfish outbreaks
- the status of reef health, particularly in areas affected by the current crown-of-thorns starfish outbreak and by tropical cyclone Yasi, to inform management priorities (such as crown-of-thorns starfish control).

#### Degradation of coastal ecosystems

- changes to natural water movement in the catchment and its effect on the condition of the Region's habitats and species
- identification of catchment sources of bio-available nitrogen and phosphorus constituents in the Region, through improved understanding of nutrient dynamics in riverine, estuarine and marine environments.

#### Direct use

- the risks to the Region's ecosystems from commercial fisheries (that is, ecological risk assessments, such as that done for the trawl fishery<sup>17</sup>)
- for some targeted fish species, stock assessments and harvest strategies are required
- the amounts of retained and discarded catch (including the proportion that dies after being discarded) from the recreational fishing sector
- the amount, species composition and post-release survival rates of discarded catch in commercial fisheries
- information about the status of populations of targeted species where there is limited or no data available, including fishery independent data
- the ecological consequences of localised depletion of populations targeted through fisheries
- development of further technologies to minimise or mitigate fishing impacts
- the impacts resulting from illegal activities in the Region and the ecological benefits of compliance and enforcement
- the effects of illegal fishing on species of conservation concern including dugong, inshore dolphins, marine turtles, sharks and rays
- the fate of dredge material dumped at sea including the full extent of dispersal and the recovery time of any affected benthic communities
- the direct and indirect effects of noise on species within the Region, including from shipping and port development
- the direct and indirect effects of dredging on the surrounding habitats and species



- the risks and key drivers of marine pest introductions, especially the ability of potential marine pests to colonise and affect coral reef environments
- the source, type, quantity and fate of marine debris, including microplastics, within the Region.

### **6.10.2 Geomorphological features**

There is limited information about past and present impacts that have had an effect on geomorphological features. Key information gaps include:

- effects of sediment from catchment run-off and dumping and resuspension of dredge material on geomorphological features such as submerged reefs, palaeochannels and karstic channels
- effects of deep sea trawling on deepwater geomorphological features
- effects of ocean acidification on geomorphic features
- effects of altered ocean currents on geomorphological features such as islands and shorelines.

### **6.10.3 Indigenous and historic heritage values**

The limited knowledge of Indigenous and historic heritage values of the Region means it is difficult to understand the effects of various impacts on those values. Consequently, improving baseline knowledge about the values is an important first step to better management of impacts on those values. Aspects include:

- improved identification of Indigenous and historic heritage values relevant to the Region, including consultation and identification of Indigenous heritage in partnership with relevant Traditional Owners
- a centralised information management system for Indigenous and historic heritage values, including spatial mapping
- improved understanding of the effects of impacts on Indigenous and historic heritage values and potential avenues to avoid, mitigate and offset impacts.

### **6.10.4 Community benefits of the environment**

Because the community benefits derived from the Region's environment are heavily dependent on the quality of that environment, the knowledge gaps described above are also relevant to community benefits. In addition, there is a need to better understand the connection between the state of the environment and the benefits derived from it, including:

- positive and negative impacts of use of the Region on community benefits
- impacts of ports and shipping on community benefits including enjoyment
- positive and negative impacts of urban and industrial development on community benefits
- quantifying impacts on aesthetic values of the Region.

### **6.10.5 Cumulative impact assessment**

Assessment of cumulative impacts is an emerging field with regard to management of the Region and protection of its matters of national environmental significance, including outstanding universal value. As highlighted earlier, there is a need to improve our understanding of:

- the relationships between Great Barrier Reef ecological and human systems and the benefits derived from these systems
- the cause-and-effect relationships of multiple or compounding impacts on values underpinning matters of national environmental significance
- the multiple spatial and temporal scales at which impacts and processes occur and interact with ecological and human systems
- models that account for uncertainty in the range of environmental conditions that will occur over the rest of this century.

The systematic application of tools and approaches outlined in this chapter will greatly assist the Authority to address these information gaps.

The various approaches to cumulative impact assessment adopted as part of this strategic assessment process have also highlighted some significant gaps in the way that information required to inform cumulative impact assessments is collected and applied. Improvements in the following areas would enhance the assessments and further accelerate our understanding of issues highlighted above:

- an accepted set of indicators for the values that underpin matters of national environmental significance that can be used to measure the effects of impacts on the state of ecological and human systems and the benefits derived from these systems
- a comprehensive, dynamic assessment of the key zones of influence within the Region
- establishment of a set of Great Barrier Reef monitoring standards, allowing results to be applied at a range of scales

- integration of monitoring results into management systems to enable impacts from various activities to be jointly considered
- systems to provide improved access (discoverability) to monitoring data (point-source and ambient data) across management agencies and for the public
- improved adaptive management mechanisms and processes, based on greater monitoring and evaluation, to enhance the effectiveness of management actions.

## 6.10.6 Processes to address information gaps

### Scientific knowledge

To guide delivery of the science needed to protect and manage the Great Barrier Reef World Heritage Area, the Authority outlined its scientific information needs in 2001 and 2005. The Outlook Report 2009<sup>1</sup> provided a fresh basis for identifying the current information gaps and scientific needs for management. These were published after the report and will be updated every five years in line with the outlook report cycle.

These identified science information needs guide future research investment. They are intended to open the dialogue between managing agencies and research and monitoring providers. For example, the key research questions used to guide investment in the Great Barrier Reef Node of the Tropical Ecosystems Hub of the National Environment Research Program were formulated with reference to the Authority's published needs.

Targeted and effective research and monitoring programs for addressing management needs require strong partnerships between research providers and the relevant management agencies. Governance structures and engagement activities are required to align monitoring and research activities with management needs. Effective governance structures create implicit information networks that can provide managers with regular updates about emerging issues, as well as trends and changes detected through monitoring activities that can be incorporated into management decisions and actions.

Research providers include:

- research institutions where research and monitoring is carried out by scientists and technicians highly trained in the relevant fields. Major institutions include the Australian Institute of Marine Sciences, James Cook University, the University of Queensland, the University of Central Queensland, the University of the Sunshine Coast, CSIRO, and government agencies such as the Authority and Queensland Government departments
- Reef-based industries where monitoring may be conducted by individuals with varying levels of training. It may be a voluntary contribution to management, undertaken to meet a permit condition or a compulsory reporting obligation
- members of the community, typically on a voluntary basis, and by people associated with a recognised 'citizen science' monitoring program
- Traditional Owners, through Traditional Use of Marine Resources Agreements and through natural resource management initiatives.

As part of its management activities, the Authority is working in partnership with CSIRO to establish a social and economic long-term monitoring program for the Great Barrier Reef.

Investment in Great Barrier Reef science is delivered through a range of programs and government initiatives, as well as through co-investment from the private sector. Apart from the ongoing funding of the Australian Institute of Marine Science, none of the major monitoring programs for the Great Barrier Reef currently have secure ongoing funding. Most rely, at least partly, on time-limited funding programs such as the National Environment Research Program and the Australian Government Caring for our Country Reef Rescue initiative for their continuation. In parallel with the strategic assessment, an integrated monitoring framework is being developed to provide the basis for reviewing monitoring priorities and activities and coordinating monitoring efforts and outputs. The framework will contain recommendations for long-term funding and governance mechanisms for monitoring (see Chapter 13).

Scientific programs are becoming increasingly spatial in focus and output, generating a variety of spatial datasets about the Region's values, its use and impacts. As a consequence, the process of gathering, synthesising, interpreting and delivering these datasets is becoming increasingly important. Continued collaboration between the Authority and its partners will help to identify and address gaps in spatial data and opportunities to share data and make it more accessible by others.

The Authority helps facilitate the delivery of the science needed to protect and manage the World Heritage Area through:

- articulating problems
- framing specific questions
- assisting in project design
- providing letters of support for grant applications
- advising on the most useful form and timing of delivery
- synthesising spatial datasets and delivering outputs to science partners and the community

- helping interpret science for government and the public
- facilitating policy and operational outcomes from science.

### Community knowledge

The Authority aims to have better outcomes for management through partnerships with Traditional Owners, industry and the community — building and sharing knowledge to support biodiversity and heritage conservation at local and regional scales.

Indigenous people are the primary source of information on the value of their heritage and how it is best conserved. The Authority is conscious of the need for Indigenous people to control intellectual property and other information relating to their heritage.

Traditional ecological knowledge is held by the many Traditional Owner clan groups with links to Great Barrier Reef sea country and is gradually being transferred to managers, especially through the establishment of Traditional Use of Marine Resources Agreements along the length of the Great Barrier Reef.

Strong relationships with stakeholders, such as tourism operators and commercial fishers, and their associations help the Authority understand use of the Region and allow information about the Reef, its condition and matters of interest to be relayed to the Authority. A range of voluntary programs formalise and streamline the provision of such information, in particular the Eye on the Reef monitoring program. The Authority also works with the community to gather knowledge and understanding of the importance and extent of values and processes.

The Authority's Reef Guardian and High Standard Tourism programs encourage participants to share their knowledge and experiences about stewardship activities which benefit the Reef.

The Authority receives a range of invaluable information from members of its Reef Advisory Committees and Local Marine Advisory Committees and during its day-to-day interactions with community members.

## 6.11 Summary of conclusions

- **The Great Barrier Reef Region is a large and complex natural system** which is being affected by a wide range of past and present impacts, including direct, indirect, consequential and cumulative impacts. A broad list of impacts has been consolidated into 40 separate impacts, grouped into four categories: climate change; catchment run-off; degradation of coastal ecosystems; and direct use.
- **The impacts affect many of the values** relevant to the matters of national environmental significance in the Region. Attributes of the outstanding universal value have been affected, including those relating to biodiversity, geomorphological features, ecosystem processes, natural beauty and man's interaction with the natural environment. The impacts assessed have also had some effect on the integrity of the World Heritage Area.
- **Twenty-five impacts are assessed as having high or very high effects** on the Region's values. The most severe past and present impacts include climate change impacts (cyclones, sea temperature increase); catchment run-off impacts (freshwater inflow, nutrients, crown-of-thorns starfish outbreaks, pesticides and sediments); modifying supporting terrestrial habitats; and several impacts of direct use (death of discarded species, dredging, dumping and resuspension of dredge material, and illegal fishing and poaching).
- **Impacts are compounding, focused inshore** in the southern two-thirds of the Region. The impacts affecting the Great Barrier Reef do not act in isolation. Many of those of most concern are land-based in origin, arising from activities undertaken in the southern two-thirds of the catchment. Combined with inshore activities, it means the zone of most intensive impact is the inshore area in the southern two-thirds of the Region. This corresponds with the declining state of values found in central and southern inshore areas. Cumulative impacts are particularly challenging to quantify, assess and manage. These impacts are beginning to be better understood through descriptive qualitative models and spatial mapping tools.
- **Some of the Region's values have been particularly affected** by the impacts assessed, with many rated as experiencing a number of high and very high effects. The combined effects of increased nutrients, more sediments and extreme weather events have seriously affected key habitats such as coral reefs and seagrass meadows. Species of conservation concern continue to be at risk, including dugongs, inshore dolphins, sawfish and some sharks, primarily due to degradation of inshore and estuarine habitats, loss of connectivity, legacy impacts and some past and present fishing activities (both legal and illegal).
- **Indigenous heritage values have been severely affected**, and these effects are intensified by the closeness of Traditional Owners' relationship to the environment. Their intimate association with their sea country is made up of tangible and intangible values, stretching over tens of thousands of years. Traditional Owners and their ancestors have witnessed profound effects relevant to all aspects of their culture and heritage, both in the past and now. For example, there have been significant and widespread effects on cultural practices, customs and lore, sacred sites, sites of particular significance, songlines, totems, languages and burial sites from a wide range of impacts. Traditional Owners emphasise that, for them, these changes in country have impacted on their culture.

- **The vast majority of impacts are assessed as increasing into the future**, driven mainly by climate change, economic growth and population growth. Some, related to direct use of the Region, are assessed as likely to be stable (such as extraction — death of discarded species, extraction — top order predators and physical damage — fishing). Improvements in shipping management mean that despite projected increases in shipping traffic a stable trend is also likely for impacts from ship groundings. The success of recent initiatives to improve the quality of catchment run-off entering the Region means related impacts are likely to stabilise in the future.
- **The future impacts of climate change are predicted to be very serious.** As greenhouse gas concentrations continue to rise at unprecedented rates, effects on the Great Barrier Reef become more severe. The concentration of carbon dioxide has increased from a pre-Industrial Revolution concentration of 280 parts per million to 396 parts per million in March 2013. Already coral bleaching as a result of elevated sea temperature has been a major cause of coral decline. If carbon dioxide levels are allowed to reach 450 parts per million, scientists predict reefs will be in rapid and terminal decline worldwide from multiple impacts including mass bleaching and ocean acidification.
- **Recent extreme weather has had significant effects.** There has been a cluster of floods, temperature extremes and cyclones over the last decade which has significantly affected the Region's values.
- **High concentrations of nutrients, sediments and pesticides continue to have long-lasting effects.** Targeted investment in halting and reversing the decline in water quality is producing results. However, there is likely to be a significant lag time before measurable water quality improvements are achieved in the marine environment, with sediments and nutrients projected to continue affecting biodiversity for many years. With continued pesticide use in the catchment, it is almost certain that pesticides will continue to be a component of catchment run-off.
- **Crown-of-thorns starfish outbreaks are a major and more frequent impact on coral reefs.** Rather than experiencing outbreaks in a natural cycle of about every 50 to 80 years, the Reef has been affected by three in the past 50 years and the beginning of another outbreak has been detected. Crown-of-thorns starfish have been a major cause of coral loss in recent decades. There is evidence of a link between outbreaks and increased nutrient concentrations and possibly loss of predators in fished areas.
- **The effects of dredge disposal may be more widespread than previously understood.** The operation of ports and further port development will require capital and maintenance dredging, potentially involving much larger volumes. Recent research indicates re-suspended dredge material may move over much greater distances from disposal sites than previously assumed. While the full extent of any effects on the Region's values is not well understood, uncertainty regarding the additional effects of sea dumping is a key concern, particularly given the potential for large volumes of proposed dredge material to be dumped and resuspended in areas of the Region already in poor condition.
- **Growth in the catchment is intensifying impacts.** Queensland's strong economic growth at rates faster than the Australian and Organisation for Economic Co-operation and Development averages is substantially the result of activities in the Great Barrier Reef catchment. This economic growth and its accompanying population growth are, in turn, driving changes in activities that affect the Region, as well as the terrestrial habitats and processes that support it.
- **The effects of legacy impacts continue.** It is important to recognise that the Region is not a pristine natural system. The conditions and trends evident today are a product of the past 200 years. Past activities in and adjacent to the Region continue to significantly influence components of the environment and reduce its ability to bounce back from current impacts. Legacy impacts are also assessed as significantly affecting many Indigenous values in the Region; for example the past commercial harvesting of dugongs and green turtles continues to have a widespread effect on cultural practices, places important for cultural tradition, and stories.
- **Past and some current impacts from fishing are affecting values.** Over the past two decades, significant progress has been made in addressing the environmental impacts from fishing. A number of high risks remain and continue to affect the Region's values.
- **Marine debris is a global problem affecting the Region's values.** Around the world, marine debris, particularly plastic, is an increasing problem as it accumulates in the ocean and on beaches. As a result of ingestion and entanglement, it is affecting a range of the Reef's wildlife.
- **The majority of popular destinations are inshore in the south of the Region.** Some of the most nationally and internationally recognised tourism destinations are nearshore islands along the southern two-thirds of the Region — the area experiencing the most intensive cumulative impacts. Such tourism destinations are highly dependent on a healthy Reef ecosystem. If not addressed, continuing declines in biodiversity and water quality are expected to reduce the attractiveness of these destinations and their viability as tourist destinations. Given the proximity of these areas to major population centres, declines in the natural environment also significantly affect recreational use of the Region. Combined, there are significant flow-on impacts on the economies and employment opportunities of adjacent Great Barrier Reef catchment communities.

The overall effect that each of the identified impacts has had on the Region's biodiversity, geomorphology and Indigenous and historic heritage values is summarised in Figure 6.11, based on the number of values affected and the severity of the effect.



**Table 6.11 Summary of the past and present effects of impacts on the Region's values**

This table is a summary of the past and present effects of impacts on attributes as shown in Table 6.6, Table 6.7, Table 6.8 and Table 6.9. Where the effects are different across attributes, the highest effect is shown.

Effect on values	Impacts affecting the Region's values			
	Biodiversity (habitats and species)	Geomorphological features	Indigenous heritage	Historic heritage
Unknown	<ul style="list-style-type: none"> <li>Atmospheric pollution</li> </ul>		<ul style="list-style-type: none"> <li>Atmospheric pollution</li> <li>Ocean acidification</li> <li>Outbreak or bloom of other species</li> <li>Spill — large chemical</li> </ul>	
Very low	<ul style="list-style-type: none"> <li>Light impacts (artificial)</li> <li>Ocean acidification</li> <li>Outbreak or bloom of other species</li> <li>Rising sea level</li> <li>Vessel strike on wildlife</li> <li>Waste discharge from a vessel</li> <li>Spill — small chemical and oil</li> </ul>	<ul style="list-style-type: none"> <li>Acid sulphate soils</li> <li>Coastal reclamation</li> <li>Ocean acidification</li> <li>Pesticides from catchment run-off</li> <li>Physical damage — fishing</li> <li>Physical damage — other</li> <li>Physical damage — ship grounding</li> </ul>	<ul style="list-style-type: none"> <li>Exotic species and diseases</li> <li>Increased sea temperature</li> <li>Light impacts (artificial)</li> <li>Spill — large oil</li> <li>Spill — small chemical and oil</li> <li>Waste discharge from a vessel</li> </ul>	<ul style="list-style-type: none"> <li>Atmospheric pollution</li> <li>Coastal reclamation</li> <li>Extraction — herbivores</li> <li>Extraction — lower order predators</li> <li>Extraction — lower trophic orders</li> <li>Extraction — top order predators</li> <li>Modifying supporting terrestrial habitats</li> <li>Outbreak of disease</li> <li>Outbreak or bloom of other species</li> <li>Physical damage — ship grounding</li> <li>Waste discharge from a vessel</li> </ul>
Low	<ul style="list-style-type: none"> <li>Acid sulphate soils</li> <li>Coastal reclamation</li> <li>Dredging</li> <li>Extraction — lower trophic orders</li> <li>Noise pollution</li> <li>Outbreak of disease</li> <li>Physical damage — other</li> <li>Physical damage — ship grounding</li> <li>Spill — large oil</li> <li>Urban and industrial discharge</li> <li>Wildlife disturbance</li> </ul>	<ul style="list-style-type: none"> <li>Artificial barriers to flow</li> <li>Cyclone activity</li> <li>Dredging</li> <li>Dumping and resuspension of dredge material</li> <li>Increased freshwater flow</li> <li>Increased sea temperature</li> <li>Modifying supporting terrestrial habitats</li> <li>Nutrients from catchment run-off</li> <li>Outbreak of crown-of-thorns starfish</li> <li>Sediments from catchment run-off</li> </ul>	<ul style="list-style-type: none"> <li>Acid sulphate soils</li> <li>Altered ocean currents</li> <li>Artificial barriers to flow</li> <li>Dumping and resuspension of dredge material</li> <li>Extraction — fishing spawning aggregations</li> <li>Extraction — top order predators</li> <li>Increased freshwater inflow</li> <li>Marine debris</li> <li>Noise pollution</li> <li>Nutrients from catchment run-off</li> <li>Outbreak of crown-of-thorns starfish</li> <li>Outbreak of disease</li> <li>Pesticides from catchment run-off</li> <li>Sediments from catchment run-off</li> <li>Urban and industrial discharge</li> <li>Wildlife disturbance</li> </ul>	<ul style="list-style-type: none"> <li>Cyclone activity</li> <li>Dumping and resuspension of dredge material</li> <li>Exotic species and diseases</li> <li>Illegal fishing and poaching</li> <li>Increased freshwater flow</li> <li>Marine debris</li> <li>Noise pollution</li> <li>Nutrients from catchment run-off</li> <li>Outbreak of crown-of-thorns starfish</li> <li>Pesticides from catchment run-off</li> <li>Physical damage — other</li> <li>Sediments from catchment run-off</li> <li>Spill — large oil</li> <li>Spill — small chemical and oil</li> <li>Urban and industrial discharge</li> <li>Wildlife disturbance</li> </ul>

Effect on values	Impacts affecting the Region's values				
	Biodiversity (habitats and species)	Geomorphological features	Indigenous heritage	Historic heritage	
High	<ul style="list-style-type: none"><li>• Artificial barriers to flow</li><li>• Dumping and resuspension of dredge material</li><li>• Exotic species and diseases</li><li>• Extraction — death of discarded species</li><li>• Extraction — fishing spawning aggregations</li><li>• Extraction — herbivores</li><li>• Extraction — lower order predators</li><li>• Extraction — top order predators</li><li>• Illegal fishing and poaching</li><li>• Increased freshwater inflow</li><li>• Increased sea temperature</li><li>• Marine debris</li><li>• Modifying supporting terrestrial habitats</li><li>• Pesticides from catchment run-off</li><li>• Physical damage — fishing</li></ul>		<ul style="list-style-type: none"><li>• Coastal reclamation</li><li>• Cyclone activity</li><li>• Extraction — death of discarded species</li><li>• Modifying supporting terrestrial habitats</li><li>• Physical damage — fishing</li><li>• Physical damage — other</li><li>• Rising sea level</li><li>• Vessel strike on wildlife</li></ul>		
Very high	<ul style="list-style-type: none"><li>• Cyclone activity</li><li>• Nutrients from catchment run-off</li><li>• Outbreak of crown-of-thorns starfish</li><li>• Sediments from catchment run-off</li></ul>		<ul style="list-style-type: none"><li>• Dredging</li><li>• Extraction — herbivores</li><li>• Extraction — lower order predators</li><li>• Extraction — lower trophic orders</li><li>• Illegal fishing and poaching</li><li>• Physical damage — ship grounding</li></ul>		
Effect on value	Unknown	Very low	Low	High	Very high

## References

- Great Barrier Reef Marine Park Authority 2009a, *Great Barrier Reef Outlook Report 2009*, GBRMPA, Townsville.
- Department of Sustainability, Environment, Water, Population and Communities 2012, *A guide to undertaking strategic assessments: Environment Protection and Biodiversity Conservation Act 1999*, DSEWPC, Canberra.
- Marsh, H., De'ath, G., Gribble, N.A. and Lane, B. 2005, Historical marine population estimates: triggers or targets for conservation? The dugong case study, *Ecological Applications* 15(2): 481-492.
- Limpus, C.J. 2008a, *A biological review of Australian marine turtle species, 2 Green turtle, Chelonia mydas (Linnaeus)*, Environmental Protection Agency, Brisbane.
- Limpus, C.J. 2009, *A biological review of Australian marine turtle species, 3 Hawksbill turtle, Eretmochelys imbricata*, Environmental Protection Agency, Brisbane.
- Noad, M.J., Dunlop, R.A., Paton, D. and Kniest, H. 2011, *Abundance estimates of the east Australian humpback whale population: 2010 survey and update*, University of Queensland, Brisbane.
- Noad, M.J., Dunlop, R.A., Cato, D. and Paton, D. 2008, *Abundance estimates of the east Australian humpback whale population: Final report for the Australian Department of the Environment, Water, Heritage and the Arts*, University of Queensland, Brisbane.
- Daley, B. 2005, *Changes in the Great Barrier Reef since European settlement: Implications for contemporary management*, PhD thesis, James Cook University, Townsville.
- Pitcher, C.R., Doherty, P., Arnold, P., Hooper, J., Gribble, N.A., Bartlett, C., Browne, M., Campbell, N., Cannard, T., Cappel, M., Carini, G., Chalmeres, S., Cheers, S., Chetwynd, D., Colegax, A., Coles, R., Cook, S., Davie, P., De'ath, G., Devereux, D., Done, B., Donovan, T., Ehrke, B., Ellis, N., Ericson, G., Fellegara, I., Forcey, K., Furey, M., Gledhill, D.,

- Good, N., Gordon, S., Haywood, M., Hendricks, M., Jacobsen, I., Johnson, J., Jones, M., Kininmonth, S., Kistler, S., Last, P., Leite, A., Amrks, S., McLeod, I., Oczkowicz, S., Robinson, M., Rose, C., Seabright, D., Sheils, J., Sherlock, M., Skelton, P., Smith, D., Smith, G., Speare, P., Stowar, M., Strickland, C., Van der Geest, C., Venables, W., Walsh, C., Wassenberg, T.J., Welna, A. and Yearsley, G. 2007, *Seabed biodiversity on the continental shelf of the Great Barrier Reef World Heritage Area*. AIMS/CSIRO/QM/QDPI CRC Reef Research Task final report, CSIRO Marine and Atmospheric Research, Hobart.
10. Daley, B. and Griggs, P. 2008, 'Loved to Death': Coral collecting in the Great Barrier Reef, Australia, 1770–1970, *Environment and History* 14(1): 89–119.
11. Roelofs, A. 2004, *Ecological assessment of the Queensland's east coast bêche-de-mer fishery*, Department of Primary Industries, Brisbane.
12. Daley, B., Griggs, P. and Marsh, H. 2008, Exploiting marine wildlife in Queensland: the commercial dugong and marine turtle fisheries 1847–1969, *Australian Economic History Review* 48(3): 227–265.
13. Chaloupka, M.Y., Bjørndal, K.A., Balazs, G.H., Bolten, A.B., Ehrhart, L.M., Limpus, C.J., Suganuma, H., Tröeng, S. and Yamaguchi, M. 2008, Encouraging outlook for recovery of a once severely exploited marine megaherbivore, *Global Ecology and Biogeography* 17(2): 297–304.
14. Limpus, C.J. and Miller, J.D. 2008, *Australian hawksbill turtle population dynamics project*, Environmental Protection Agency, Brisbane.
15. Burrridge, C.Y., Pitcher, C.R., Wassenberg, T.J., Poiner, I.R. and Hill, B.J. 2003, Measurement of the rate of depletion of benthic fauna by prawn (shrimp) otter trawls: An experiment in the Great Barrier Reef, Australia, *Fisheries Research* 60(2–3): 237–253.
16. Poiner, I.R., Glaister, J., Pitcher, C.R., Burrridge, C.Y., Wassenberg, T.J., Gribble, N.A., Hill, B.J., Blaber, S.J.M., Milton, D., Brewer, D. and Ellis, N. 1998, *Environmental effects of prawn trawling in the far northern section of the Great Barrier Reef: 1991–96*, CSIRO Division of Marine Research, Cleveland.
17. Pears, R.J., Morison, A.K., Jebreen, E.J., Dunning, M.C., Pitcher, C.R., Courtney, A.J., Houlden, B. and Jacobsen, I.P. 2012a, *Ecological risk assessment of the East Coast Otter Trawl Fishery in the Great Barrier Reef Marine Park: Technical report*, Great Barrier Reef Marine Park Authority, Townsville.
18. Limpus, C.J. 2008b, *A biological review of Australian marine turtle species, 1 Loggerhead turtle, Caretta caretta (Linnaeus)*, Environmental Protection Agency, Brisbane.
19. Brewer, D., Heales, D., Milton, D., Dell, Q., Fry, G., Venables, B. and Jones, P. 2006, The impact of turtle excluder devices and bycatch reduction devices on diverse tropical marine communities in Australia's northern prawn trawl fishery, *Fisheries Research* 81(2–3): 176–188.
20. Great Barrier Reef Marine Park Authority 2012a, *Informing the outlook for Great Barrier Reef coastal ecosystems*, GBRMPA, Townsville.
21. Wooldridge, S.A. 2009, Water quality and coral bleaching thresholds: formalising the linkage for the inshore reefs of the Great Barrier Reef, Australia, *Marine Pollution Bulletin* 58(5): 745–751.
22. Thompson, A., Costello, P., Davidson, J., Schaffelke, B., Uthicke, S. and Liddy, M. 2013a, *Reef Rescue Marine Monitoring Program. Report of AIMS activities: inshore coral reef monitoring 2012. Report for Great Barrier Reef Marine Park Authority*, Australian Institute of Marine Science, Townsville.
23. Kyne, P.M., Carlson, J. & Smith, K. 2013, *Pristis pristis (Indo-West Pacific subpopulation): IUCN Red List of Threatened Species Version 2013.1*, International Union for Conservation of Nature, Cambridge, UK, <<http://www.iucnredlist.org/details/43508905/0>>.
24. Great Barrier Reef Marine Park Authority 2013a, *Haughton Basin Assessment: Burdekin Dry Tropics Natural Resource Management Region*, GBRMPA, Townsville.
25. Steinberg, C. 2007, Impacts of climate change on the physical oceanography of the Great Barrier Reef, in *Climate change and the Great Barrier Reef: a vulnerability assessment*, eds J.E. Johnson and P.A. Marshall, Great Barrier Reef Marine Park Authority and Australian Greenhouse Office, Townsville, pp. 51–74.
26. Hoegh-Guldberg, O. and Bruno, J.F. 2010, The impact of climate change on the world's marine ecosystems, *Science* 328: 1523–1528.
27. Ridgway, K. and Hill, K. 2012, East Australian Current, in *A marine climate change impacts and adaptation report card for Australia 2012*, eds E.S. Poloczanska, A.J. Hobday and A.J. Richardson, CSIRO, Canberra, pp. 47–60.
28. Williams, K.J., Dunlop, M., Bustamante, R.H., Murphy, H.T., Ferrier, S., Wise, R.M., Liedloff, A., Skewes, T.D., Harwood, T.D., Kroon, F., Williams, R.J., Joehnk, K., Crimp, S., Smith, M.S., James, C. and Booth, T. 2012, *Queensland's biodiversity under climate change: Impacts and adaptation – synthesis report*, CSIRO Climate Adaptation Flagship, Canberra.
29. Chambers, L.E., Devney, C.A., Congdon, B.C., Dunlop, N., Woehler, E.J. and Dann, P. 2011, Observed and predicted effects of climate on Australian seabirds, *Emu* 111(3): 235–251.
30. Environment Protection Agency 2002, *State Coastal Management Plan: Queensland's Coastal Management Policy*, EPA, Brisbane.
31. Australian State of the Environment Committee 2002, *Australia: state of the environment 2001*, CSIRO Publishing, Collingwood.
32. Steffen, W., Hughes, L. and Karoly, D. 2013, *The critical decade: Extreme weather*, Climate Commission Secretariat, Canberra.
33. Williams, K.J. and Crimp, S. 2012, *Queensland's biodiversity under climate change: An overview of climate change in Queensland, Climate Adaptation Flagship Working Paper No 12A*, CSIRO Climate Adaptation Flagship, Canberra.
34. Abbs, D. 2012, *The impact of climate change on the climatology of tropical cyclones in the Australia region, Working Paper No. 11*, CSIRO Climate Adaptation Flagship.
35. Great Barrier Reef Marine Park Authority 2011a, *Impacts of tropical cyclone Yasi on the Great Barrier Reef: A report on the findings of a rapid ecological impact assessment*, GBRMPA, Townsville.
36. De'ath, G., Fabricius, K.E., Sweatman, H. and Puotinen, M. 2012, The 27-year decline of coral cover on the Great Barrier Reef and its causes, *Proceedings of the National Academy of Sciences* 109(44): 17995–17999.
37. Fabricius, K.E., De'ath, G., Puotinen, M.L., Done, T.J., Cooper, T.F. and Burgess, S.C. 2008, Disturbance gradients on inshore and offshore coral reefs caused by a severe tropical cyclone, *Limnology and Oceanography* 53(2): 690–704.
38. Diaz-Pulido, G., McCook, L.J., Dove, S., Berkelmans, R., Roff, G., Kline, D.I., Weeks, S., Evans, R.D., Williamson, D.H. and Hoegh-Guldberg, O. 2009, Doom and boom on a resilient reef: Climate change, algal overgrowth and coral recovery, *PLoS ONE* 4(4): e5239.
39. Hughes, T.P. and Connell, J.H. 1999, Multiple stressors on coral reefs: a long-term perspective, *Limnology and Oceanography* 44(3): 932–940.
40. Roff, G., Clark, T.R., Raymond, C.E., Zhao, J., Feng, Y., McCook, L.J., Done, T.J. and Pandolfi, J.M. 2013, Palaeoecological evidence of a historical collapse of corals at Pelorus Island, inshore Great Barrier Reef, following European settlement, *Proceedings of the Royal Society B: Biological Sciences* 280(1750): 2012–2100.
41. McKenzie, L.J., Collier, C. and Waycott, M. 2012, *Reef Rescue Marine Monitoring Program: nearshore seagrass, annual report for the sampling period 1st July 2010 - 31st May 2011*, Fisheries Queensland, Cairns.
42. Puotinen, M., Maynard, J., Johnson, J., Johnson, J., Beeden, R., Devlin, M., Dryden, J. and Marshall, P. (in preparation), *Development of a cyclone damage predictive model for the Great Barrier Reef: Technical report*, Great Barrier Reef Marine Park Authority, Townsville.
43. Done, T. 1992, Effects of tropical cyclone waves on ecological and geomorphological structures on the Great Barrier Reef, *Continental Shelf Research* 12(7–8): 859–872.
44. Madin, J. and Connelly, S. 2006, Ecological consequences of major hydrodynamic disturbances on coral reefs, *Nature* 444: 477–480.

45. Lough, J.M. 2009, A changing climate: Evidences and consequences, *Microbiology Australia* 30(2): 58-60.
46. Lough, J., Gupta, A.S. and Hobday, A.J. 2012, Temperature, in *A marine climate change: impacts and adaptation report card for Australia 2012*, eds E.S. Poloczanska, A.J. Hobday and A.J. Richardson, CSIRO, Canberra.
47. Maynard, J.A., Johnson, J.C., Beeden, R., Puotinen, M., Devlin, M., Dryden, J. and Marshall, P. 2014, *Historic and projected future exposure of habitats in the Great Barrier Reef Marine Park to disturbances. Draft Report*, Great Barrier Reef Marine Park Authority, Townsville.
48. Lough, J. 2007, Climate and climate change on the Great Barrier Reef, in *Climate change and the Great Barrier Reef: A Vulnerability Assessment*, eds J.E. Johnson and P.A. Marshall, Great Barrier Reef Marine Park Authority and Australian Greenhouse Office, Townsville, pp. 15-50.
49. Hobday, A.J. and Lough, J.M. 2011, Projected climate change in Australian marine and freshwater environments, *Marine and Freshwater Research* 62(9): 1000-1014.
50. Glynn, P.W. 1993, Coral bleaching: Ecological perspectives, *Coral Reefs* 12(1): 1-17.
51. Glynn, P.W. 1984, Widespread coral mortality and the 1982-83 El Niño warming event, *Environmental Conservation* 11(2): 133-146.
52. Lough, J.M. 2000, 1997-98: Unprecedented thermal stress to coral reefs? *Geophysical Research Letters* 27(23): 3901-3904.
53. Berkelmans, R., De'ath, G., Kininmonth, S. and Skirving, W.J. 2004, A comparison of the 1998 and 2002 coral bleaching events on the Great Barrier Reef: Spatial correlation, patterns and predictions, *Coral Reefs* 23(1): 74-83.
54. Ward, S., Harrison, P. and Hoegh-Guldberg, O. 2000, Coral bleaching reduces reproduction of scleractinian corals and increases susceptibility to future stress, in *Proceedings of the 9th International Coral Reef Symposium*, eds M. K. Moosa and et al, Ministry of Environment, Indonesia Institute of Sciences, Bali, Indonesia, pp.1123-1128.
55. Veron, J.E.N., Hoegh-Guldberg, O., Lenton, T.M., Lough, J.M., Obura, D.O., Pearce-Kelly, P., Sheppard, C.R., Spalding, M., Stafford-Smith, M.G. and Rogers, A.D. 2009, The coral reef crisis: The critical importance of <350 ppm CO<sub>2</sub><SUB>2</SUB>, *Marine Pollution Bulletin* 58(10): 1428-1436.
56. Hoegh-Guldberg, O., Berkelmans, R. and Oliver, J.K. 1999, *Coral bleaching: Implications for the Great Barrier Reef Marine Park*, CRC Reef Research, Townsville.
57. Hoegh-Guldberg, O., Mumby, P.J., Hooten, A.J., Steneck, R.S., Greenfield, P., Gomez, E., Harvell, C.D., Sale, P.F., Edwards, A.J., Caldeira, K., Knowlton, N., Eakin, C.M., Iglesias-Prieto, R., Muthiga, N., Bradbury, R.H., Dubi, A. and Hatzioiols, M.E. 2007, Coral reefs under rapid climate change and ocean acidification, *Science* 318: 1737-1742.
58. Anthony, K.R.N., Kline, D.I., Diaz-Pulido, G., Dove, S. and Hoegh-Guldberg, O. 2008, Ocean acidification causes bleaching and productivity loss in coral reef builders, *Proceedings of the National Academy of Sciences* 105(45): 17442-17446.
59. Pandolfi, J.M., Connolly, S.R., Marshall, D.J. and Cohen, A.L. 2011, Projecting coral reef futures under global warming and ocean acidification, *Science* 333(6041): 418-422.
60. Anthony, K.R.N. and Marshall, P. 2012, Coral Reefs, in *A marine climate change impacts and adaptation report card Australia 2012*, eds E.S. Poloczanska, A.J. Hobday and A.J. Richardson, CSIRO, Canberra.
61. Fabricius, K.E., Langdon, C., Uthicke, S., Humphrey, C., Noonan, S., De'ath, G., Okazaki, R., Muehllehner, N., Glas, M. and Lough, J.M. 2011, Losers and winners in coral reefs acclimatized to elevated carbon dioxide concentrations, *Nature Climate Change* 1: 165-169.
62. Doropoulos, C., Ward, S., Diaz-Pulido, G., Hoegh-Guldberg, O. and Mumby, P.J. 2012, Ocean acidification reduces coral recruitment by disrupting intimate larval-algal settlement interactions, *Ecology Letters* 15(4): 338-346.
63. Church, J.A., White, N.J., Hunter, J.R. and McInnes, K.L. 2012, Sea level, in *A marine climate change impacts and adaptation report card for Australia 2012*, eds E.S. Poloczanska, A.J. Hobday and A.J. Richardson, CSIRO, Canberra, pp. 27-46.
64. CSIRO and Bureau of Meteorology 2012, *State of the Climate 2012*, CSIRO Publishing, Collingwood.
65. Church, J.A., White, N.J., Hunter, J.R. and Lambeck, K. 2008, *Briefing: a post-IPCC AR4 update on sea-level rise*, The Antarctic Climate and Ecosystems Cooperative Research Centre, Hobart.
66. Department of Environment and Resource Management 2011, *Queensland coastal processes and climate change*, Queensland Climate Change Centre of Excellence, Brisbane.
67. Rogers, K., Saintilan, N. and Copeland, C. 2012, Modelling wetland surface elevation dynamics and it's application to forecasting the effects of sea level rise on estuarine wetlands, *Ecological Modelling* 244: 148-157.
68. Fuentes, M.M.P.B., Limpus, C.J., Hamann, M. and Dawson, J. 2010, Potential impacts of projected sea-level rise on sea turtle rookeries, *Aquatic Conservation: Marine and Freshwater Ecosystems* 20(2): 132-139.
69. Chambers, L.E., Dann, P., Devney, C., Dunlop, N. and Woehler, E.J. 2012, Seabirds, in *Marine Climate Change Impacts and Adaptation Report Card for Australia 2012*, eds E.S. Poloczanska, A.J. Hobday and A.J. Richardson, 2012 Report Card edn, CSIRO, Canberra.
70. Schaffelke, B., Carleton, J., Costello, P., Davidson, J., Doyle, J., Furnas, M., Gunn, K., Skuza, M., Wright, M. and Zagorskis, I. 2012, *Reef Rescue Marine Monitoring Program. Final report of AIMS activities 2011/12: Inshore water quality monitoring. Produced for the Great Barrier Reef Marine Park Authority*, Australian Institute of Marine Science, Townsville.
71. Devlin, M., Brodie, J.E. and Waterhouse, J. 2001, Community and connectivity : summary of a community based monitoring program set up to assess the movement of nutrients and sediments into the Great Barrier Reef during high flow events, *Water Science and Technology* 43(9): 121-131.
72. Wong, T., Breen, P. and Lloyd, S. 2000, *Water sensitive road design: Design options for improving stormwater quality of road run-off*, CRC for Catchment Hydrology, Canberra.
73. Halliday, I. and Robins, J. 2007, *Environmental flows for sub-tropical estuaries: understanding the freshwater needs of estuaries for sustainable fisheries production and assessing the impacts of water regulation*, Final Report FRDC Project No. 2001/022 , Fisheries Research and Development Corporation, Canberra.
74. Staunton-Smith, J., Robins, J.B., Mayer, D.G., Sellin, M.J. and Halliday, I.A. 2004, Does the quantity and timing of fresh water flowing into a dry tropical estuary affect year-class strength of Barramundi (*Lates calcarifer*)? *Marine and Freshwater Research* 55(8): 787-797.
75. Robins, J.B., Halliday, I., Staunton-Smith, J., Mayer, D.G. and Sellin, M.J. 2005, Freshwater-flow requirements of estuarine fisheries in tropical Australia: a review of the state of knowledge and application of a suggested approach, *Marine and Freshwater Research* 56(3): 343-360.
76. Berkelmans, R., Jones, A.M. and Schaffelke, B. 2012, Salinity thresholds of *Acropora* spp. on the Great Barrier Reef, *Coral Reefs* 31: 1103-1110.
77. Brodie, J., Binney, J., Fabricius, K., Gordon, I., Hoegh-Guldberg, O., Hunter, H., O'Reagain, P., Pearson, R., Quirk, M., Thorburn, P., Waterhouse, J., Webster, I. and Wilkinson, S. 2008, *Scientific consensus statement on water quality in the Great Barrier Reef*, Reef Water Quality Protection Plan Secretariat, Brisbane.
78. Kroon, F.J., Kuhnert, P.M., Henderson, B.L., Wilkinson, S.N., Kinsey-Henderson, A., Abbott, B., Brodie, J.E. and Turner, R.D.R. 2012, River loads of suspended solids, nitrogen, phosphorus and herbicides delivered to the Great Barrier Reef lagoon, *Marine Pollution Bulletin* 65(4-9): 167-181.
79. Mitchell, A.W., Bramley, R.G.V. and Johnson, A.K.L. 1997, Export of nutrients and suspended sediment during a cyclone-mediated flood event in the Herbert River catchment, Australia, *Marine and Freshwater Research* 48: 79-88.
80. Brodie, J.E. 2000, The problem of nutrients and eutrophication in the Australian marine environment, in *The Marine Environment: The State of the Marine Environment Report for Australia, Technical Annex 2: Pollution*, eds L. Zann



and D.C. Sutton, Department of the Environment, Sport and Territories, Ocean Rescue, Canberra, pp. 1-29.

81. Mitchell, A.W. and Furnas, M.J. 1997, Terrestrial inputs of nutrients and suspended sediments to the Great Barrier Reef lagoon, in *Proceedings of the Great Barrier Reef, Science, Use and Management: A National Conference: Vol 1*, eds. N. Turia and C. Dalliston, Great Barrier Reef Marine Park Authority, Townsville, pp.59-71.

82. Furnas, M., Mitchell, A.W. and Skuza, M. 1995, *Nitrogen and Phosphorus Budgets for the Central Great Barrier Reef Shelf*, Great Barrier Reef Marine Park Authority, Townsville.

83. Furnas, M. and Brodie, J. 1996, Current status of nutrient levels and other water quality parameters in the Great Barrier Reef, in *Downstream effects of landuse*, eds H.M. Hunter, A.G. Eyles and G.E. Rayment, Department of Natural Resources, Brisbane, pp. 9-21.

84. Furnas, M. and Mitchell, A. 1986, Oceanographic conditions on the north Queensland shelf after passage of cyclone Winifred, in *Proceedings of the Workshop on the offshore effects of cyclone Winifred, 20 June 1986, Townsville*, eds. I. M. Dutton, Great Barrier Reef Marine Park Authority, Townsville, .

85. Shaw, C.M., Lam, P.K.S. and Muller, J.F. 2008, Photosystem II herbicide pollution in Hong Kong and its potential photosynthetic effects on corals, *Marine pollution bulletin* 57(6): 473-478.

86. Gagan, M.K., Sandstrom, M.K. and Chivas, A.R. 1987, Restricted terrestrial carbon input to the continental shelf during Cyclone Winifred: implications for terrestrial runoff to the Great Barrier Reef province, *Coral Reefs* 6(2): 113-119.

87. Walker, T.A. and O'Donnell, G. 1981, Observations on Nitrate, Phosphate and Silicate in Cleveland Bay, Northern Queensland, *Australian Journal of Marine and Freshwater Research* 32: 877-887.

88. Furnas, M.J., Mitchell, A. and Skuza, M. 1997, Shelf scale nitrogen and phosphorus budgets for the Central Great Barrier Reef (16 - 19 S), in *Proceedings of the 8th International Coral Reef Symposium, Panama, 24-29 June 1996*, eds. H. A. Lessios and I. G. Macintyre, Smithsonian Tropical Marine Research Institute, Balboa, Republic of Panama, pp.809-814.

89. Brodie, J.E., Devlin, M., Haynes, D. and Waterhouse, J. 2011, Assessment of the eutrophication status of the Great Barrier Reef lagoon (Australia), *Biogeochemistry* 106(2): 281-302.

90. Packett, R., Dougall, C., Rohde, K. and Noble, R. 2009, Agricultural lands are hot-spots for annual runoff polluting the southern Great Barrier Reef lagoon, *Marine Pollution Bulletin* 58(7): 976-986.

91. Mitchell, C., Brodie, J.E. and White, I. 2005, Sediments, nutrients and pesticide residues in event flow conditions in streams of the Mackay Whitsunday Region, Australia, *Marine Pollution Bulletin* 51(1-4): 23-36.

92. Lewis, S.E., Brodie, J., Ledee, E. and Alewijnse, M. 2006, *The spatial extent of delivery of terrestrial materials from the Burdekin region of the Great Barrier Reef lagoon. ACTFR Report No. 06/02*, Australian Centre for Tropical Freshwater Research, James Cook University, Townsville.

93. De'ath, G. and Fabricius, K.E. 2008, *Water quality of the Great Barrier Reef: Distributions, effects on reef biota and trigger values for the protection of ecosystem health*, Great Barrier Reef Marine Park Authority, Townsville.

94. Great Barrier Reef Marine Park Authority 2010, *Water quality guidelines for the Great Barrier Reef Marine Park*, GBRMPA, Townsville.

95. Furnas, M., Mitchell, A., Skuza, M. and Brodie, J.E. 2005, In the other 90%: Phytoplankton responses to enhanced nutrient availability in the Great Barrier Reef lagoon, *Marine Pollution Bulletin* 51: 253-265.

96. Brodie, J., Fabricius, K.E., De'ath, G. and Okaji, K. 2005, Are increased nutrient inputs responsible for more outbreaks of crown-of-thorns starfish? An appraisal of the evidence, *Marine Pollution Bulletin* 51(1-4): 266-278.

97. Fabricius, K.E., Okaji, K. and De'ath, G. 2010, Three lines of evidence to link outbreaks of the crown-of-thorns seastar *Acanthaster planci* to the release of larval food limitation, *Coral Reefs* 29: 593-605.

98. McCook, L.J., Folke, C., Hughes, T., Nyström, M., Obura, D. and Salm, R. 2007, Ecological resilience, climate change and the Great Barrier Reef, in *Climate Change and the Great Barrier Reef: A vulnerability assessment*, eds J.E. Johnson and P.A. Marshall, Great Barrier Reef Marine Park Authority and Australian Greenhouse Office, Townsville, pp. 75-96.

99. McCook, L.J. 1999, Macroalgae, nutrients and phase shifts on coral reefs: scientific issues and management consequences for the Great Barrier Reef, *Coral Reefs* 18: 357-367.

100. Wooldridge, S.A. and Done, T.J. 2009, Improved water quality can ameliorate effects of climate change on corals, *Ecological Applications* 19: 1492-1499.

101. Wiedenmann, J., D'Angelo, C., Smith, E.G., Hunt, A.N., Legiret, F., Postle, A.D. and Achterberg, E.P. 2013, Nutrient enrichment can increase the susceptibility of reef corals to bleaching, *Nature Climate Change* 3: 160-164.

102. Walker, D., Dennison, W.C. and Edgar, G.J. 1999, Status of Australian seagrass research and knowledge, in *Seagrass in Australia: Strategic Review and Development of an R & D Plan*, eds A.J. Butler and P. Jernakoff, CSIRO Publishing, Collingwood, pp. 1-25.

103. Brando, V.E., Schroeder, T., Dekker, A.G. and Clementson, L. 2014, *Reef Rescue Marine Monitoring Program: Assessment of terrestrial run-off entering the Reef and inshore marine water quality monitoring using Earth observation data, Final Report for 2011/2012 activities*, Unpublished report.

104. Department of the Premier and Cabinet 2013a, *Reef Water Quality Protection Plan 2013*, Reef Water Quality Protection Plan Secretariat, Brisbane.

105. Department of the Premier and Cabinet 2013b, *Great Barrier Reef Report Card 2011, Reef Water Quality Protection Plan*, Reef Water Quality Protection Plan Secretariat, Brisbane.

106. Schaffelke, B., Carleton, J., Doyle, J., Furnas, M., Gunn, K., Skuza, M. and Zagorskis, I. 2011, *Reef Rescue Marine Monitoring Program, Final report of AIMS Activities 2010/11: Inshore water quality monitoring, Report for the Great Barrier Reef Marine Park Authority*, Australian Institute of Marine Science, Townsville.

107. Schaffelke, B., Carleton, J., Doyle, J., Furnas, M., Gunn, K., Skuza, M., Wright, M. and Zagorskis, I. 2010, *Reef Rescue Marine Monitoring Program: Final report of AIMS activities 2009/10: Inshore water quality monitoring*, Australian Institute of Marine Science, Townsville.

108. Kline, D.I., Kuntz, N.M., Breitbart, M., Knowlton, N. and Rohwer, F. 2006, Role of elevated organic carbon levels and microbial activity in coral mortality, *Marine Ecology Progress Series* 314: 119-125.

109. Engelhardt, U., Miller, I., Lassig, B.R., Sweatman, H.P.A. and Bass, D. 1997, Crown-of-thorns starfish (*Acanthaster planci*) populations in the Great Barrier Reef World Heritage Area: Status report 1995-96, in *Proceedings of the State of the Great Barrier Reef World Heritage Area Workshop*, eds. D. Wachenfeld, J. Oliver and K. Davis, Great Barrier Reef Marine Park Authority, Townsville, pp.158-187.

110. Sweatman, H., Cheal, A.J., Coleman, G.J., Emslie, M.J., Johns, K., Jonker, M., Miller, I.R. and Osborne, K. 2008, *Long-term monitoring of the Great Barrier Reef: Status Report 8*, Australian Institute of Marine Science, Townsville.

111. Kenchington, R.A. 1977, Growth and recruitment of *Acanthaster planci* (L.) on the Great Barrier Reef, *Biological Conservation* 11: 103-118.

112. Brodie, J. and Waterhouse, J. 2012, A critical review of environmental management of the 'not so Great' Barrier Reef, *Estuarine, Coastal and Shelf Science* 104-105: 1-22.

113. Keesing, J.K. and Lucas, J.S. 1992, Field measurement of feeding and movement rates of the crown-of-thorns starfish *Acanthaster planci* (L.), *Journal of Experimental Marine Biology and Ecology* (156): 89-104.

114. Brodie, J.E. 1992, Enhancement of larval and juvenile survival and recruitment in *Acanthaster planci* from the effects of terrestrial runoff: a review, *Australian Journal of Marine and Freshwater Research* 43: 539-554.

115. Brodie, J.E. and Mitchell, A.W. 2005, Nutrients in Australian tropical rivers: Changes with agricultural development

and implications for receiving environments, *Marine and Freshwater Research* 56(3): 279-302.

116. Sweatman, H. 2008, No-take reserves protect coral reefs from predatory starfish, *Current Biology* 18(14): R598-R599.

117. McCook, L.J., Ayling, T., Cappo, M., Choat, J.H., Evans, R.D., DeFreitas, D.M., Heupel, M., Hughes, T.P., Jones, G.P., Mapstone, B., Marsh, H., Mills, M., Molloy, F.J., Pitcher, C.R., Pressey, R.L., Russ, G.R., Sutton, S., Sweatman, H., Tobin, R., Wachenfeld, D. and Williamson, D. 2010, Adaptive management of the Great Barrier Reef: A globally significant demonstration of the benefits of networks of marine reserves, *Proceedings of the National Academy of Sciences* 107(43): 18278-18285.

118. Brodie, J.E., Kroon, F.J., Schaffelke, B., Wolanski, E.C., Lewis, S.E., Devlin, M.J., Bohnet, I.C., Bainbridge, Z.T., Waterhouse, J. and Davis, A.M. 2012, Terrestrial pollutant runoff to the Great Barrier Reef: An update of issues, priorities and management responses, *Marine Pollution Bulletin* 65: 81-100.

119. Carroll, C., Waters, D., Vardy, S., Silburn, D.M., Attard, S., Thorburn, P.J., Davis, A.M., Halpin, N., Schmidt, M. and Wilson, B. 2012, A Paddock to reef monitoring and modelling framework for the Great Barrier Reef: Paddock and catchment component, *Marine Pollution Bulletin* 65(4-9): 136-149.

120. Australian and New Zealand Environment and Conservation Council and Agriculture and Resource Management Council of Australia and New Zealand 2000, *Australian and New Zealand guidelines for fresh and marine water quality, National Water Quality Management Strategy Paper No 4*, ANZECC and ARMCANZ, Canberra.

121. Smith, R., Middlebrook, R., Turner, R., Huggins, R., Vardy, S. and Warne, M. 2012, Large-scale pesticide monitoring across Great Barrier Reef catchments: Paddock to Reef Integrated Monitoring, Modelling and Reporting Program, *Marine Pollution Bulletin* 65(4-9): 117-127.

122. Lewis, S.E., Brodie, J.E., Bainbridge, Z.T., Rohde, K.W., Davis, A.M., Masters, B.L., Maughan, M., Devlin, M.J., Mueller, J.F. and Schaffelke, B. 2009, Herbicides: A new threat to the Great Barrier Reef, *Environmental Pollution* 157(8-9): 2470-2484.

123. Davis, A., Lewis, S., Bainbridge, Z., Brodie, J. and Shannon, E. 2008, Pesticide residues in waterways of the lower Burdekin region: challenges in ecotoxicological interpretation of monitoring data, *Australasian Journal of Ecotoxicology* 14(2-3): 89-108.

124. Bainbridge, Z.T., Brodie, J., Faithful, J.W., Sydes, D.A. and Lewis, S.E. 2009, Identifying the land-based sources of suspended sediments, nutrients and pesticides discharged to the Great Barrier Reef from the Tully-Murray Basin, Queensland, Australia, *Marine and Freshwater Research* 60(11): 1081-1090.

125. Davis, A.M., Thorburn, P.J., Lewis, S.E., Bainbridge, Z.T., Attard, S.J., Milla, R. and Brodie, J.E. 2013, Environmental impacts of irrigated sugarcane production: Herbicide run-off dynamics from farms and associated drainage systems, *Agriculture, Ecosystems and Environment* 180(1): 123-135.

126. Silburn, M., Rojas-Ponce, S., Fillols, E., Olsen, D., McHugh, J., Baillie, C., Lewis, S. and Davis, A. 2013, *Precision application (band spray) of herbicides on sugarcane in the Burdekin region*, State of Queensland, Brisbane.

127. Great Barrier Reef Marine Park Authority 2009b, *Water quality guidelines for the Great Barrier Reef Marine Park*, GBRMPA, Townsville.

128. Haynes, D., Carter, S., Gaus, C., Muller, J. and Dennison, W. 2005, Organochlorine and heavy metal concentrations in blubber and liver tissue collected from Queensland (Australia) dugong (*Dugong dugon*), *Marine Pollution Bulletin* 51(1-4): 361-369.

129. Gaus, C., O'Donohue, M. and Connell, D. 2004, An assessment of risks to dugongs from exposure to dioxins, in *Proceedings of the Catchment to Reef: Water Quality Issues in the Great Barrier Reef Region Conference, 9-11 March 2004*, Townsville, eds. D. Haynes and B. Schaffelke, CRC Reef Research Centre, Townsville, .

130. Gaus, C., O'Donohue, M., Connell, D., Muller, J., Haynes, D. and Papke, O. 2004, Exposure and potential risks of

dioxins to the marine mammal dugong, *Organohalogen Compounds* 66: 1559-1566.

131. McMahon, K., Nash, S.B., Eaglesham, G., Muller, J.F., Duke, N.C. and Winderlich, S. 2005, Herbicide contamination and the potential impact to seagrass meadows in Hervey Bay, Queensland, Australia, *Marine Pollution Bulletin* 51(1-4): 325-334.

132. Lewis, S.E., Schaffelke, B., Shaw, M., Bainbridge, Z.T., Rohde, K.W., Kennedy, K., Davis, A.M., Masters, B.L., Devlin, M.J. and Mueller, J.F. 2012, Assessing the additive risks of PSII herbicide exposure to the Great Barrier Reef, *Marine Pollution Bulletin* 65(4-9): 280-291.

133. Devlin, M., McKinna, L. and Harkness, P. 2010, *Reef Rescue Marine Monitoring Program: terrestrial runoff in the Great Barrier Reef (3.7.2b): Flood plume monitoring for 2009/10 annual report*, Australian Centre for Tropical and Freshwater Research, Townsville.

134. Devlin, M.J. and Brodie, J. 2005, Terrestrial discharge into the Great Barrier Reef lagoon: Nutrient behaviour in coastal waters, *Marine Pollution Bulletin* 51(1-4): 9-22.

135. Devlin, M.J., McKinna, L.I.W., Alvarez-Romero, J.G., Abbott, B., Harkness, P. and Brodie, J. 2012, Mapping the pollutants in surface river plume waters in the Great Barrier Reef, Australia, *Marine Pollution Bulletin* 65: 224-235.

136. Wolanski, E., Fabricius, K.E., Cooper, T.E. and Humphrey, C. 2008, Wet season fine sediment dynamics on the inner shelf of the Great Barrier Reef, *Estuarine, Coastal and Shelf Science* 77(4): 755-762.

137. Wolanski, E., Jones, M. and Williams, W.T. 1981, Physical properties of Great Barrier Reef lagoon waters near Townsville. II. Seasonal variations, *Australian Journal of Marine and Freshwater Research* 32: 321-334.

138. Bainbridge, Z.T., Wolanski, E., Alvarez-Romero, J.G., Lewis, S.E. and Brodie, J.E. 2012, Fine sediment and nutrient dynamics related to particle size and floc formation in a Burdekin River flood plume, Australia, *Marine Pollution Bulletin* 65(4): 236-248.

139. Weber, M., Lott, C. and Fabricius, K.E. 2006, Sedimentation stress in a scleractinian coral exposed to terrestrial and marine sediments with contrasting physical, organic and geochemical properties, *Journal of Experimental Marine Biology and Ecology* 336: 18-32.

140. Great Barrier Reef Marine Park Authority 2012b, *A vulnerability assessment for the Great Barrier Reef: Sawfish*, GBRMPA, Townsville.

141. Baird, A.H., Babcock, R.C. and Mundy, C.P. 2003, Habitat selection of larvae influences the depth distribution of six common coral species, *Marine Ecology Progress Series* 252: 289-293.

142. Fabricius, K.E., De'ath, G., McCook, L.J., Turak, E. and Williams, D.M. 2005, Changes in algal, coral and fish assemblages along water quality gradients on the inshore Great Barrier Reef, *Marine Pollution Bulletin* 51(1-4): 384-398.

143. Fabricius, K.E., De'ath, G., Humphrey, C., Zagorskis, I. and Schaffelke, B. 2013, Intra-annual variation in turbidity in response to terrestrial runoff on near-shore coral reefs of the Great Barrier Reef, *Estuarine Coastal and Shelf Science* 116: 57-65.

144. Walker, T.A. 1981, Dependence of phytoplankton chlorophyll on bottom resuspension in Cleveland Bay, northern Queensland, *Australian Journal of Marine and Freshwater Research* 32: 981-986.

145. Devlin, M., Petus, C., da Silva, E., Alvarez-Romero, J., Zeh, D., Waterhouse, J. and Brodie, J. 2013, Chapter 5. Mapping of exposure to flood plumes, water types and exposure to pollutants (DIN, TSS) in the Great Barrier Reef: toward the production of operational risk maps for the World's most iconic marine ecosystem. in *Assessment of the relative risk of water quality to ecosystems of the Great Barrier Reef: supporting studies. A report to the Department of the Environment and Heritage Protection, Queensland Government*, ed. J. Waterhouse, Centre for Tropical Water & Aquatic Ecosystem Research, James Cook University, Townsville.

146. Brodie, J., Waterhouse, J., Schaffelke, B., Kroon, F., Thorburn, P., Rolfe, J., Johnson, J., Fabricius, K., Lewis, S., Devlin, M., Warne, M. and McKenzie, L.J. 2013a, *2013 Scientific*

*Consensus Statement: Land use impacts on Great Barrier Reef water quality and ecosystem conditions*, Reef Water Quality Protection Plan Secretariat, Brisbane.

147. Brodie, J., Pearson, R., Lewis, S.E., Bainbridge, Z.T., Waterhouse, J. and Prange, J. 2009, *Water quality research: baseline synthesis and year 1 summary*, Report to the Marine and Tropical Sciences Research Facility, Reef and Rainforest Research Centre, Cairns.

148. Walker, G.S. and Brunskill, G.J. 1997, A history of anthropogenic mercury input into the Great Barrier Reef lagoon, Australia, in *Proceedings of the 8th International Coral Reef Symposium Vol. 2*, Smithsonian Tropical Research Institute, Panama, eds. H. A. Lessios and I. G. Macintyre, Smithsonian Tropical Research Institute, Panama, pp.1889-1892.

149. Queensland Floods Commission of Inquiry 2012, *Final Report*, Queensland Government, Brisbane.

150. Department of Environment and Heritage Protection 2013, *Central Queensland mines water releases 2012: Pilot program*, DEHP, Brisbane, <<http://www.ehp.qld.gov.au/land/mining/water-release-pilot.html>>.

151. Sammut, J., Callinan, R. and Fraser, G. 1996, An overview of the ecological impacts of acid sulfate soils in Australia, in *Proceedings of the 2nd National Conference on Acid Sulfate Soils, Coffs Harbour, 4-6 Sept, 1996*, eds. Anonymous, NSW Agriculture, Wollongbar, NSW, pp.140-145.

152. National Working Party on Acid Sulfate Soils 2000, *National strategy for the management of coastal acid sulfate soils*, Agricultural and Resource Management Council of Australia and New Zealand, Canberra.

153. Russell, D.J. and Helmke, S.A. 2002, Impacts of acid leachate on water quality and fisheries resources in northern Australia, *Marine and Freshwater Research* 53(1): 19.

154. De Weys, J., Santos, I.R. and Eyre, B.D. 2011, Linking groundwater discharge to severe estuarine acidification during a flood in a modified wetland, *Environmental Science and Technology* 45(8): 3310-3316.

155. Ahern, K.S., Udy, J.W. and Pointon, S.M. 2006, Investigating the potential for groundwater from different vegetation, soil and landuses to stimulate blooms of the cyanobacterium, *Lyngbya majuscula*, in coastal waters, *Marine and Freshwater Research* 57: 177-186.

156. Murtha, G.G., Cannon, M.G. and Smith, C.D. 1996, *Soils in the Babinda-Cairns area, North Queensland*. Divisional Report No 123, CSIRO Division of Soils, South Australia.

157. Smith, C.D. and Ahern, C.D. 1998, *Report on acid sulfate soil investigation Ingham to Tully drainage schemes*, Department of Natural Resources, Brisbane.

158. Ross, D.J. 2002, *Acid sulfate soils, Tannan Sands to St Laurence, Central Queensland Coast*, Department of Natural Resources and Mines, Rockhampton, Queensland.

159. Environmental Protection Agency 2008, *State of the Environment Queensland 2007*, EPA, Brisbane.

160. Sammut, J., White, I., Melville, M.D. and Melville, M.D. 1996, *Coastal in-stream impacts and assessment of acid sulfate soils*, Research report 63, Water Research Foundation of Australia, Canberra.

161. Powell, B. and Martens, M. 2005, A review of acid sulfate soil impacts, actions and policies that impact on water quality in Great Barrier Reef catchments, including a case study on remediation at East Trinity, *Marine Pollution Bulletin* 51: 149-164.

162. Queensland Department of Employment, Economic Development and Innovation 2011a, *Inventory of Instream Structures Impacting on Ramsar Wetlands*, DEEDI, Brisbane.

163. Carter, J., Tait, J., Kapitzke, R. and Corfield, J. 2007, *Burdekin Dry Tropics NRM region fish passage study*, Alluvium Consulting, Townsville.

164. Veitch, V. and Sawynok, B. 2005, *Freshwater wetlands and fish: importance of freshwater wetlands to marine fisheries resources in the Great Barrier Reef*, Great Barrier Reef Marine Park Authority, Townsville.

165. McKergrow, L.A., Prosser, I.P., Hughes, A.O. and Brodie, J. 2005, Sources of sediment to the Great Barrier Reef World Heritage Area, *Marine Pollution Bulletin* 51: 200.

166. Hyland, S.J. 2002, *An investigation of the impacts of ponded pastures on barramundi and other finfish populations in tropical coastal wetlands: Final project report Q002005*, Fisheries Queensland, Brisbane.

167. Lawrence, M.L., Sully, D.W., Couchman, D. and Beumer, J.P. 2011, *Inventory of instream structures impacting on Ramsar wetlands*, Report to the Queensland Wetlands Program, Fisheries Queensland, Department of Employment, Economic Development and Innovation, Brisbane.

168. Resource Assessment Commission 1993, *coastal zone Inquiry, Final Report*, RAC, Canberra.

169. Gräwe, U., Wolff, J.O. and Ribbe, J. 2010, Impact of climate variability on an east Australian Bay, *Estuarine, Coastal and Shelf Science* 86(2): 247-257.

170. Toki, B., Baheerathan, R., Richardson, D. and Bothelho, D. 2012, *Abbot Point Cumulative Impact Assessment Technical Report: Assessment of potential impacts of coal dust on the marine environment at the Port of Abbot Point*, Abbot Point Working Group, BMT WBM Pty Ltd, Brisbane.

171. Johnson, R. and Bustin, R.M. 2006, Coal dust dispersal around a marine coal terminal (1977-1999), British Colombia: the fate of coal dust in the marine environment, *International Journal of Coal Geology* 68: 57-69.

172. Lucas, S.A. and Planner, J. 2012, Grounded or submerged bulk carrier: The potential for leaching of coal trace elements to seawater, *Marine Pollution Bulletin* 64: 1012-1017.

173. Salmon, M., Wyneken, J. and Fritz, E. and Lucas, M. 1992, Seafinding by hatchling sea turtles: role of brightness, silhouette and beach slope as orientation cues. *Behaviour* 122: 56-77.

174. Kamrowski, R.L., Limpus, C.J., Moloney, J. and Hamann, M. 2012, Coastal light pollution and marine turtles: Assessing the magnitude of the problem, *Endangered Species Research* 19: 85-98.

175. Witherington, B.E. 1992, Behavioral responses of nesting sea turtles to artificial lighting, *Herpetologica* 48(1): 31-39.

176. Lohmann, K.J. and Lohmann, C.M.F. 1998, Migratory guidance mechanisms in marine turtles, *Journal of Avian Biology* 29: 585-596.

177. Rodriguez, A. and Rodriguez, B. 2009, Attraction of petrels to artificial light in the Canary Islands: effects of the moon phase and age class, *Ibis* 151(2): 299-310.

178. Levenez, J.J., Gerlotto, F. and Petit, D. 1990, Reaction of tropical coastal pelagic species to artificial lighting and implications for the assessment of abundance by echo integration, in *Proceedings of the International Symposium on Fisheries Acoustics*, eds. Anonymous, Seattle, WA, USA, .

179. Great Barrier Reef Marine Park Authority and Department of Sustainability Environment Water Populations and Communities 2013, *Information on applications for Marine Park permits, Sea Dumping permits and Environment Protection and Biodiversity Conservation referrals*.

180. Foster, T., Corcoran, E., Erftemeijer, P.L., Fletcher, C., Peirs, K., Dolmans, C., Smith, A. and Yamamoto, H. 2010, *Dredging and port construction around coral reefs*, *Envicom Report 108*, PIANC Secretariat, Belgium.

181. Smith, R., Boyd, S.E., Rees, H.L., Dearnaley, M.P. and Stevenson, J. 2006, Effects of dredging activity on epifaunal communities - surveys following cessation of dredging, *Estuarine, Coastal and Shelf Science* 70(1-2): 207-223.

182. Erftemeijer, P.L. and Lewis, R.R.R. 2006, Environmental impacts of dredging on seagrass: a review, *Marine Pollution Bulletin* 52(12): 1553-1572.

183. Brunner, R.W.J. 2011, *Sustainability in the design of bulk coal ports: an Australian case study*, North Queensland Bulk Ports Corporation, Mackay, Queensland.

184. Biddle, T.M. and Limpus, C.J. 2011, Marine wildlife stranding and mortality database annual report 2005-2010, Marine turtles, *Conservation and Technical Data Report 2011(1)*: 1-124.

185. Erftemeijer, P.L., Riegl, B., Hoeksema, B.W. and Todd, P.A. 2012, Environmental impacts of dredging and other sediment disturbances on corals: a review, *Marine Pollution Bulletin* 64(9): 1737-1765.



186. Organisation for Economic Co-operation and Development 2011, *Environmental impacts of international shipping: The role of ports*, OECD Publishing, Paris, France.
187. Rasheed, K. and Balchand, A.N. 2007, Environmental studies on impacts of dredging. *International Journal of Environmental Studies* 58(6): 703-725.
188. Ellison, W.T., Southall, B.L., Clark, C.W. and Frankel, A.S. 2011, A new context-based approach to assess marine mammal behavioral responses to anthropogenic sounds, *Conservation Biology* 26(1): 21-28.
189. Department of the Environment, Water, Heritage and the Arts 2009, *National assessment guidelines for dredging*, Department of the Environment, Water, Heritage and the Arts, Canberra.
190. Islam, A., Wang, L., Smith, C., Reddy, S., Lewis, A. and Smith, A. 2007, Evaluation of satellite remote sensing for operational monitoring of sediment plumes produced by dredging at Hay Point, Queensland, Australia, *Journal of Applied Remote Sensing* 1(1): 1-15.
191. Trimarchi, S. and Keane, J. 2007, *Port of Hay Point apron areas and departure path capital dredging project: environmental review*, Ports Corporation of Queensland Limited, Brisbane.
192. Smith, A., Foster, T., Corcoran, E. and Monkivitch, J. 2007, Dredging and material relocation in sensitive coral environments, in *Proceedings of the 18th World Dredging Conference, 1 June, 2007, Florida*, eds. Anonymous, Great Barrier Reef Marine Park Authority, Townsville, pp.945-955.
193. Sinclair Knight Merz Pty Ltd and Asia-Pacific Applied Science Associates 2013, *Improved dredge material management for the Great Barrier Reef Region: Synthesis report*, Great Barrier Reef Marine Park Authority, Townsville.
194. Goggin, L. 2004, *Introduced species in tropical waters*, CRC Reef Research Centre, Townsville.
195. Limpus, C.J. and Reimer, D. 1994, The loggerhead turtle, *Caretta caretta*, in Queensland: a population in decline, in *Proceedings of the Australian marine turtle conservation workshop*, eds. R. James, Department of Environment and Heritage and Australian Nature Conservation Agency, Canberra, pp.39-59.
196. Seabrook, W. 1994, Feral cat predation on hatchling green turtles. in *Proceedings of the Australian Marine Turtle Conservation Workshop*, eds. R. James, Department of Environment and Heritage and the Australian Nature Conservation Agency, Canberra, pp.191.
197. Batianoff, G.N., Naylor, G.C., Olds, J. and Neldner, V.J. 2009, Distribution patterns, weed incursions and origins of terrestrial flora at the Capricorn-Bunker Islands, Great Barrier Reef, Australia, *Cunninghamia* 11(1): 107-121.
198. Queensland Department of Primary Industries and Fisheries 2008, 'Unpublished data'.
199. Pears, R.J., Morison, A.K., Jebreen, E.J., Dunning, M.C., Pitcher, C.R., Courtney, A.J., Houlden, B. and Jacobsen, I.P. 2012b, *Ecological risk assessment of the East Coast Otter Trawl Fishery in the Great Barrier Reef Marine Park: summary report*, Great Barrier Reef Marine Park Authority, Townsville.
200. Brewer, D., Rawlinson, N., Eayrs, S. and Burridge, C. 1998, An assessment of bycatch reduction devices in a tropical Australian prawn trawl fishery, *Fisheries Research* 36(2-3): 195-215.
201. Courtney, A.J., Haddy, J.A., Campbell, v., Roy, D.P., Tonks, M.L., Gaddes, S.W., Chilcott, K.E., O'Neill, M.F., Brown, I.W., McLennan, M., Jebreen, E.J., van der Geest, C., Rose, C., Kistle, S., Turnbull, C.T., Kyne, P.M., Bennett, M.B. and Taylor, J. 2007, *Bycatch weight, composition and preliminary estimates of the impact of bycatch reduction devices in Queensland's trawl fishery*, Department of Primary Industries and Fisheries, Brisbane.
202. Courtney, A.J., Schemel, B.I., Wallace, R.M., Campbell, M.J. and Mayer, D.J. 2010, *Reducing the impact of Queensland's trawl fisheries on protected sea snakes*, Fisheries Research and Development Corporation and Department of Employment, Economic Development and Innovation, Brisbane.
203. Department of Sustainability, Environment, Water, Population and Communities 2009, *Death or injury to marine species following capture in beach meshing (nets) and drum lines used in Shark Control Programs*, DSEWPA, Canberra, <<http://www.environment.gov.au/biodiversity/threatened/ktp/shark-control-programs.html#attachment>>.
204. Domeier, M.L. and Colin, P.L. 1997, Tropical reef fish spawning aggregations: defined and reviewed, *Bulletin of Marine Science* 60(3): 698-726.
205. Sadovy de Mitcheson, Y. and Colin, P.L. (eds) 2012, *Reef fish spawning aggregations: Biology, research and management*, *Fish and Fisheries Series, Vol 35*, Springer Science and Business Media, New York, USA.
206. Russell, M. and Pears, R.J. 2007, *Management and science of fish spawning aggregations in the Great Barrier Reef Marine Park, summary of workshop held 17-20 July 2007*, *Museum of Tropical Queensland*, Great Barrier Reef Marine Park Authority, Townsville.
207. Queensland Department of Employment, Economic Development and Innovation 2011b, *Annual status report 2010: East Coast Spanish Mackerel Fishery*, Department of Employment, Economic Development and Innovation, Brisbane.
208. Pears, R., J., Choat, J., H., Mapstone, B., D. and Begg, G., A. 2007, Reproductive biology of a large, aggregation-spawning serranid, *Epinephelus fuscoguttatus*: management implications, *Journal of Fish Biology* 71: 795-817.
209. Pears, R., Nemeth, R.S., Ferreira, B.P., Hostim-Silva, M., Bertocini, A.A., Coleman, F.C., Koenig, C.C., Rhodes, K.L., Sadovy de Mitcheson, Y., Heppell, S.A., Colin, P.L., Samoilys, M.A., Sakaue, J., Akino, H., Ida, H., Jackson, G., Warner, R.R., Hamilton, R.J., Choat, H., Etpison, M.T., Collins, P., Clavijo, I., J. and Hillmann Kitalong, A. 2012c, Species case studies, in *Reef fish spawning aggregations: Biology, research and management*, Vol 35, eds Y. Sadovy de Mitcheson and P.L. Colin, Springer Science + Business Media, New York, USA, pp. 405.
210. Russell, M.W., Luckhurst, B.E. and Lindeman, K.C. 2012, Management of spawning aggregations, in *Reef fish spawning aggregations: Biology, research and management*, Vol 35, eds Y. Sadovy de Mitcheson and P.L. Colin, Springer Science and Business Media, New York, USA, pp. 371-404.
211. Hoey, A.S. and Bellwood, D.R. 2009, Limited functional redundancy in a high diversity system: single species dominates key ecological process on coral reefs, *Ecosystems* 12(8): 1316-1328.
212. Bellwood, D.R., Hoey, A.S. and Choat, J.H. 2003, Limited functional redundancy in high diversity systems: resilience and ecosystem function on coral reefs, *Ecology Letters* 6(4): 281-285.
213. Hughes, T.P., Rodrigues, M.J., Bellwood, D.R., Ceccarelli, D., Hoegh-Guldberg, O., McCook, L.J., Moltschanivskyj, N.A., Pratchett, M.S., Steneck, R.S. and Willis, B. 2007, Phase shifts, herbivory, and the resilience of coral reefs to climate change, *Current Biology* 17(4): 360-365.
214. Jompa, J. and McCook, L.J. 2002, The effects of nutrients and herbivory on competition between a hard coral (*Porites cylindrica*) and a brown alga (*Lobophora variegata*), *Limnology and Oceanography* 47: 527-534.
215. Hoey, A. and Bellwood, D.R. 2011, Suppression of herbivory by macroalgal density: a critical feedback on coral reefs? *Ecology Letters* 14: 267.
216. Williamson, D.H., Russ, G.R. and Ayling, A.M. 2004, No-take marine reserves increase abundance and biomass of reef fish on inshore fringing reefs of the Great Barrier Reef, *Environmental Conservation* 31(2): 149-159.
217. Pears, R., J., Choat, J., H., Mapstone, B., D. and Begg, G., A. 2006, Demography of a large grouper *Epinephelus fuscoguttatus*, from the Great Barrier Reef: implications for harvest limits and size regulations, *Marine Ecology Progress Series* 307: 259-271.
218. Tobin, A.J., Simpfendorfer, C.A., Mapleston, A., Currey, L., Harry, A.V., Welch, D.J., Ballagh, A.C., Chin, A., Szczecinski, N., Schlaff, A., White, J. and Moore, B. 2010, *A quantitative ecological risk assessment of sharks and finfish of the Great Barrier Reef World Heritage Area inshore waters: a tool for fisheries and marine park managers. Identifying species at risk and potential mitigation strategies*, Marine and Tropical Sciences Research Facility, Cairns.



219. Almany, G.R. and Webster, M.S. 2004, Odd species out as predators reduce diversity of coral-reef fishes, *Ecology* 85(11): 2933-2937.
220. O'Shea, O.R., Thums, M., van Keulen, M. and Meekan, M. 2012, Bioturbation by stingrays at Ningaloo Reef, Western Australia, *Marine and Freshwater Research* 63(3): 189-197.
221. Coleman, F.C. and Williams, S.L. 2002, Overexploiting marine ecosystem engineers: potential consequences for biodiversity, *Trends in Ecology and Evolution* 17(1): 40-44.
222. Cavanagh, R.D., Kyne, P.M., Fowler, S.L., Musick, J.A. and Bennett, M.B. 2003, *The conservation status of Australian chondrichthyans: report of the IUCN Shark Specialist Group Australia and Oceania Regional Red List Workshop*, School of Biomedical Sciences, University of Queensland, Brisbane.
223. Heupel, M.R., Williams, A.J., Welch, D.J., Ballagh, A., Mapstone, B.D., Carlos, G., Davies, C. and Simpfendorfer, C.A. 2009, Effects of fishing on tropical reef associated shark populations on the Great Barrier Reef, *Fisheries Research* 95(2-3): 350-361.
224. Hisano, M., Connolly, S.R. and Robbins, W. 2011, Population growth rates of reef sharks with and without fishing on the Great Barrier Reef, *PLoS ONE* 6(9): e25028.
225. Robbins, W.D., Hisano, M., Connolly, S.R. and Choat, J.H. 2006, Ongoing collapse of coral reef shark populations, *Current Biology* 16: 2314-2319.
226. Office of Economic and Statistical Research 2012, *Shark control program: shark caught by type, Queensland, 2001 to 2002 until 2011 to 2012*, Queensland Treasury and Trade, Brisbane, <<http://www.oesr.qld.gov.au/products/tables/shark-control-program-caught-type/index.php>>.
227. Stevens, J.D., Bonfil, R., Dulvy, N.K. and Walker, P.A. 2000, The effects of fishing on sharks, rays and chimaeras (chondrichthyans), and the implications for marine ecosystems, *ICES Journal of Marine Science* 57: 476-494.
228. Great Barrier Reef Marine Park Authority 2012c, *Field management quarterly report (IN-CONFIDENCE)*, GBRMPA, Townsville.
229. Daley, B. and Griggs, P. 2006, Mining the reefs and cays: Coral, guano and rock phosphate extraction in the Great Barrier Reef Australia, 1844-1940. *Environment and History*, 12: 395-433.
230. Taylor, H. and Smith, W. 2009, *2009 Far north Queensland marine debris project report*, Tangaroa Blue Ocean Care Society, Port Douglas.
231. Andrady, A.L. 2011, Microplastics in the marine environment, *Marine pollution bulletin* 62(8): 1596-1605.
232. Simmonds, M.P. 2012, Cetaceans and Marine Debris: The Great Unknown, *Journal of Marine Biology* 2012: 684279.
233. Coe, J.M. and Rogers, A.B. (eds) 1997, *Marine Debris. Source, Impacts and Solutions*, Springer-Verlag, New York.
234. United National Environment Program 2005, *Marine Litter: An analytical overview*, UNEP, Kenya.
235. Barnes, D.K., Galgani, F., Thompson, R.C. and Barlaz, M. 2009, Accumulation and fragmentation of plastic debris in global environments, *Philosophical Transactions of the Royal Society B: Biological Sciences* 364(1526): 1985-1998.
236. Pemberton, D., Brothers, N. and Kirkwood, R. 1992, Entanglement of Australian fur seals in man-made debris in Tasmanian waters, *Wildlife Research* 19: 151-159.
237. Slip, D.J. and Burton, H.R. 1991, Accumulation of fishing debris, plastic litter, and other artifacts on Heard and Macquarie Islands in the Southern Ocean, *Environmental Conservation* 18(3): 249-254.
238. Chatto, R. 1995, Sea turtles killed by flotsam in northern Australia, *Marine Turtle Newsletter* 69(April): 17-18.
239. Laist, D. 1996, Marine debris entanglement and ghost fishing: A cryptic and significant type of bycatch? in *Proceedings of the Solving Bycatch Workshop: Considerations for today and tomorrow*, Report No 96-03, 25-27 September 1995, Seattle, WA, eds. Alaska Sea Grant, Alaska Sea Grant College Program, Fairbanks, AK, pp.33-39.
240. Richardson, W.J., Greene, C.R., Malme, C.I. and Thomson, D.H. 1995, *Marine mammals and noise*, Academic Press, San Diego.
241. United Nations Environment Programme 2012, Scientific synthesis on the impacts of underwater noise on marine and coastal biodiversity and habitats, in *Proceedings of the 16th Meeting of the Subsidiary Body on Scientific, Technical and Technological Advice, 30 April - 5 May 2012, Montreal, Canada*, eds. UNEP, UNEP, Nairobi, pp.1-93.
242. Department of the Environment, Water, Heritage and the Arts 2008, *EPBC Act Policy Statement 2.1- Interaction between offshore seismic exploration and whales*, Department of the Environment, Water, Heritage and the Arts, Canberra.
243. Slabbekoorn, H., Bouton, N., van Opzeeland, I., Coers, A., ten Cate, C. and Popper, A.N. 2010, A noisy spring: The impact of globally rising underwater sound levels on fish, *Trends in Ecology and Evolution* 25(7): 419-427.
244. Brandt, M.E. and McManus, J.W. 2009, Disease incidence is related to bleaching extent in reef-building corals, *Ecology* 90(10): 2859-2867.
245. Bruno, J.F., Selig, E.R., Casey, K.S., Page, C.A., Willis, B., Harvell, C.D., Sweatman, H. and Melendy, A.M. 2007, Thermal stress and coral cover as drivers of coral disease outbreaks, *PLoS Biology* 5(6): e124.
246. Heron, S.F., Willis, B.L., Skirving, W.J., Eakin, C.M., Page, C.A. and Miller, I.R. 2010, Summer hot snaps and winter conditions: modelling white syndrome outbreaks on Great Barrier Reef corals, *PLoS One* 5(8): e12210.
247. Harvell, C.D., Jordan-Dahlgren, E., Merkel, S., Rosenberg, E., Raymundo, L., Smith, G., Weil, G. and Willis, B. 2007, Coral disease, environmental drivers and the balance between coral and microbial associates, *Oceanography* 20(1): 36-59.
248. McClanahan, T.R., Donner, S.D., Maynard, J.A., MacNeil, M.A., Graham, N.A.J., Maina, J., Baker, A.C., Alemu, J.B., Beger, M., Campbell, S.J., Darling, E.S., Eakin, C.M., Heron, S.F., Jupiter, S.D., Lundquist, C.J., McLeod, E., Mumby, P.J., Paddock, M.J., Selig, E.R. and van Woessik, R. 2012, Prioritizing key resilience indicators to support coral reef management in a changing climate, *PLoS ONE* 7(8): e42884.
249. Thompson, A., Costello, P., Davidson, J., Schaffelke, B., Uthicke, S. and Liddy, M. 2013b, *Reef Rescue Marine Monitoring Program, Report of AIMS Activities – Inshore coral reef monitoring 201, Report for Great Barrier Reef Marine Park Authority*, Australian Institute of Marine Science, Townsville.
250. Pollock, J. 2013, *Personal Communication*.
251. Flint, M.J., Limpus, C.J., Patterson-Kane, J.C., Murray, P.J. and Mills, P.C. 2010a, Corneal fibropapillomatosis in green sea turtles (*Chelonia mydas*) in Australia, *Journal of Comparative Pathology* 142: 341-346.
252. Herbst, L.H. 1994, Fibropapillomatosis of marine turtles, *Annual Review of Fish Diseases* 4: 389-425.
253. van Houtan, K.S., Hargrove, S.K. and Balazs, G.H. 2010, Land use, macroalgae, and a tumour-forming disease in marine turtles, *PLoS ONE* 5(9): e12900.
254. Flint, M., Patterson-Kane, J.C., Limpus, C.J. and Mills, P.C. 2010b, Health surveillance of stranded green turtles in southern Queensland, Australia (2006-2009): an epidemiological analysis of causes of disease and mortality, *EcoHealth* 7(1): 135-145.
255. Chaloupka, M.Y., Balazs, G.H. and Work, T.M. 2009, Rise and fall over 26 years of a marine epizootic in Hawaiian green sea turtles, *Journal of Wildlife Diseases* 45(4): 1138-1142.
256. Biddle, T.M., Boyle, M. and Limpus, C.J. 2011, *Marine wildlife stranding and mortality database annual report 2009 and 2010: Dugong*, Department of Environment and Resource Management, Brisbane.
257. Meager, J.J. and Limpus, C.J. 2012a, Marine wildlife stranding and mortality database annual report 2011, I Dugong, *Conservation and Technical Data Report* 2011(1): 1-30.
258. Arthur, K.E., Limpus, C.J., Roelfsema, C.M., Udy, J.W. and Shaw, G.R. 2006, A bloom of *Lyngbya majusculina* Shoalwater Bay, Queensland, Australia: an important feeding

ground for the green turtle (*Chelonia mydas*), *Harmful Algae* 5(3): 251-265.

259. Bell, P.R.F., Elmetri, I. and Uwins, P. 1999, Nitrogen fixation by *Trichodesmium* spp. in the central and northern Great Barrier Reef lagoon: relative importance of the fixed-nitrogen load, *Marine Ecology Progress Series* 186: 119-126.

260. Bell, P. 1991, Must GBR pollution become chronic before management reacts? *Search* 22(4): 117-119.

261. Jones, G.B. 1992, Effect of trichodesmium blooms on water quality in the Great Barrier Reef lagoon, in *Marine pelagic cyanobacteria: trichodesmium and other diazotrophs*, eds E.J. Carpenter, D.G. Capone and J.G. Reuter, Kluwer Academic Press, Dordrecht, pp. 273-287.

262. Cumming, R. 2009, *Case study: impact of Drupella spp. on reef-building corals of the Great Barrier Reef*, Great Barrier Reef Marine Park Authority, Townsville.

263. Grech, A. and Coles, R. 2011, Interactions between a trawl fishery and spatial closures for biodiversity conservation in the Great Barrier Reef World Heritage Area, Australia, *PLoS ONE* 6(6): e21094.

264. Pitcher, C.R., Burrige, C.Y., Wassenberg, T.J., Hill, B.J. and Poiner, I.R. 2009, A large scale BACI experiment to test the effects of prawn trawling on seabed biota in a closed area of the Great Barrier Reef Marine Park, Australia, *Fisheries Research* 99(3): 168-183.

265. Pitcher, C.R., Wassenberg, T.J., Cappo, M.C., Smith, G.P., Austin, M., Gordon, S.R., Bustamante, R.H., Moeseneder, C.H., Speare, P.J., Kennedy, J.A., Doherty, P.J. and Hooper, J.N.A. 2004, *Dynamics of large sessile seabed fauna, important for structural fisheries habitat and biodiversity of marine ecosystems and the use of these habitats by key finfish species, Final report to the FRDC on project 1997/205*, CSIRO Marine Research, Cleveland, Qld.

266. Pitcher, C.R., Poiner, I.R., Hill, B.J. and Burrige, C.Y. 2000, Implications of the effects of trawling on sessile megazoobenthos on a tropical shelf in northeastern Australia, *ICES Journal of Marine Science* 57(5): 1359-1368.

267. Burrige, C.Y., Pitcher, C.R., Hill, B.J., Wassenberg, T.J. and Poiner, I.R. 2006, A comparison of demersal communities in an area closed to trawling with those in adjacent areas open to trawling: A study in the Great Barrier Reef Marine Park, Australia, *Fisheries Research* 79(1-2): 64-74.

268. Harriott, V.J. 2001, *The sustainability of Queensland's coral harvest fishery*, CRC Reef Research Centre, Townsville.

269. Donnelly, R. 2009, *Pro-vision reef: Stewardship action plan, a statement of operational standards and climate change contingency planning*, Great Barrier Reef Marine Park Authority, Townsville.

270. GHD Pty Ltd 2013, *Ship anchorage management in the Great Barrier Reef World Heritage Area*, Great Barrier Reef Marine Park Authority, Townsville.

271. Barker, N.H.L. and Roberts, C.M. 2004, Scuba diver behaviour and the management of diving impacts on coral reefs, *Biological Conservation* 120(4): 481-489.

272. Australian Maritime Safety Authority 2010, *Improving Safe Navigation in the Great Barrier Reef*, AMSA, Canberra.

273. Great Barrier Reef Marine Park Authority 2013b, *Maritime incident database*, GBRMPA, Townsville.

274. Great Barrier Reef Marine Park Authority 2011b, *Grounding of the Shen Neng 1 on Douglas Shoal, April 2010: impact assessment report*, GBRMPA, Townsville.

275. Asia-Pacific ASA 2012a, *Maritime Chemical Spill Response Card Sulphuric Acid*, Great Barrier Reef Marine Park Authority, Townsville.

276. Asia-Pacific ASA 2012b, *Maritime Chemical Spill Response Card Liquefied Natural Gas (LNG) - Methane*, Great Barrier Reef Marine Park Authority, Townsville.

277. Australian Maritime Safety Authority 2011, *National marine oil spill contingency plan*, AMSA, Canberra.

278. Aston, J. 2006, Shipping and oils spills, in *The State of the Great Barrier Reef Online*, ed. A. Chin, Great Barrier Reef Marine Park Authority, Townsville.

279. Melville, F., Andersen, L.E. and Jolley, D.F. 2009, The Gladstone (Australia) oil spill-Impacts on intertidal areas:

Baseline and six months post-spill, *Marine Pollution Bulletin* 58(2): 263-271.

280. Anderson, L.E., Melville, F. and Jolley, D.F. 2008, An assessment of an oil spill in Gladstone, Australia - impacts on intertidal areas at one month post-spill, *Marine Pollution Bulletin* 58: 263-271.

281. Australian Transport Safety Bureau 2011, *Independent investigation into the grounding of the Chinese registered bulk carrier Shen Neng on Douglas Shoal, Queensland 3 April 2010*, ATSB, Canberra.

282. Australian Transport Safety Bureau 2012, *Independent safety issue investigation into Queensland Coastal Pilotage, Report No. 282-MI-2010-011*, ATSB, Canberra.

283. Hazel, J., Lawler, I.R., Marsh, H. and Robson, S. 2007, Vessel speed increases collision risk for the green turtle *Chelonia mydas*, *Endangered Species Research* 3(2): 105-113.

284. Limpus, C.J., Currie, K.J. and Haines, J. 2004, *Marine wildlife stranding and mortality database annual report 2003, I Dugong*, Environmental Protection Agency, Brisbane.

285. Hodgson, A.J. and Marsh, H. 2007, Response of dugongs to boat traffic: The risk of disturbance and displacement, *Journal of Experimental Marine Biology and Ecology* 340(1): 50-61.

286. Greenland, J.A. and Limpus, C.J. 2008, *Marine wildlife stranding and mortality database annual report 2007, II Cetacean and Pinniped*, Environmental Protection Agency, Brisbane.

287. Meager, J.J. and Limpus, C.J. 2012b, Marine wildlife stranding and mortality database annual report 2011, III Marine turtles, *Conservation and Technical Data Report 3*: 1-46.

288. Andersson, M. 2008, *Hinchinbrook Area Boaters' opinions of and compliance with dugong conservation initiatives* Independent Study Projects (ISP), Independent Study Projects (ISP), Study Abroad, Cairns.

289. Ross, H., Innes, J., George, M. and Gorman, K. 2004, *Traditional owner aspirations towards cooperative management of the Great Barrier Reef World Heritage Area: Community case studies*, CRC Reef Research Centre, Townsville.

290. Wu, R.S.S. 1999, Eutrophication, Water Borne Pathogens and Xenobiotic Compounds: Environmental Risks and Challenges, *Marine Pollution Bulletin* 39(1-12): 11-22.

291. Productivity Commission 2003, *Industries, land use and water quality in the Great Barrier Reef catchment: research report*, Productivity Commission, Canberra.

292. Rogers, D., Hassall, C. and Lewis, J. 2006, *Shorebird disturbance on the beaches of Roebuck Bay, 2005-2006: conservation implications and recommendations: Report by Broome Bird Observatory for the WA Department of Conservation and Land Management*, Department of Conservation and Land Management, Perth.

293. Williams, A.J., Ward, V.L. and Underhill, L.G. 2004, Waders respond quickly and positively to the banning of off-road vehicles from beaches in South Africa, *Wader Study Group Bulletin* 104: 79-81.

294. Banks, P.B. and Bryant, J.V. 2007, Four legged friend or foe? Dog walking displaces native birds from natural areas, *Biology Letters* 3(6): 611-613.

295. Bejder, L., Samuels, A., Whitehead, H., Gales, N., Mann, J., Connor, R., Heithaus, M.R., Watson-Capps, J., Flaherty, C. and Krützen, M. 2006, Decline in relative abundance of bottlenose dolphins exposed to long-term disturbance, *Conservation Biology* 20(6): 1791-1798.

296. Parra, G.J., Corkeron, P.J. and Marsh, H. 2006, Population sizes, site fidelity and residence patterns of Australian snubfin and Indo-Pacific humpback dolphins: Implications for conservation, *Biological Conservation* 129(2): 167-180.

297. Mattson, M.C., Thomas, J.A. and Aubin, D.S. 2005, Effects of boat activity on the behavior of bottlenose dolphins (*Tursiops truncatus*) in waters surrounding Hilton Head Island, South Carolina, *Aquatic Mammals* 31(1): 133-140.

298. Lukoschek, V. and Chilvers, B.L. 2008, A robust baseline for bottlenose dolphin abundance in coastal Moreton Bay: a large carnivore living in a region of escalating anthropogenic impacts, *Wildlife Research* 35(7): 593-605.



299. Corkeron, P.J. 1995, Humpback whales (*Megaptera novaeangliae*) in Hervey Bay, Queensland: behaviour and responses to whale-watching boats, *Canadian Journal of Zoology/Revue Canadien de Zoologie* 73(7): 1290-1299.
300. Preen, T. 2001, *Dugongs, boats, dolphins and turtles in the Townsville-Cardwell region and recommendations for a boat traffic management plan for the Hinchinbrook Dugong Protection Area*, Great Barrier Reef Marine Park Authority, Townsville.
301. Stamation, K.A., Croft, D.B., Shaughnessy, P.D., Waples, K.A. and Briggs, S.V. 2010, Behavioral responses of humpback whales (*Megaptera novaeangliae*) to whale-watching vessels on the southeastern coast of Australia, *Marine Mammal Science* 26(1): 98-122.
302. Lusseau, D. and Bejder, L. 2007, The long-term consequences of short-term responses to disturbance experiences from whalewatching impact assessment. *International Journal of Comparative Psychology* 20(2-3): 228-236.
303. Devney, C.A., Short, M. and Congdon, B.C. 2009, Cyclonic and anthropogenic influences on tern populations, *Wildlife Research* 36: 368-378.
304. Hulsman, K., O'Neill, P.O. and Stokes, T. 1997, *Current status and trends of seabirds on the Great Barrier Reef*, Great Barrier Reef Marine Park Authority, Townsville.
305. Clemens, R.S., Haslem, A., Oldland, J., Shelley, L., Weston, M.A. and Abdullah Abu Diyan, M. 2008, *Identification of significant shorebird areas in Australia: mapping, thresholds and criteria*, Birds Australia, Carlton.
306. Chapman, C.J. and Atkinson, R.J.A. 1986, Fish behaviour in relation to divers, *Progress of Underwater Science* 11: 1-14.
307. Fitzpatrick, R., Abrantes, K.G., Seymour, J. and Barnett, A. 2011, Variation in depth of whitetip reef sharks: does provisioning ecotourism change their behaviour? *Coral Reefs* 30: 569-577.
308. Smithers, S.G., Harvey, N., Hopley, D. and Woodroffe, C.D. 2007, Vulnerability of geomorphological features in the Great Barrier Reef to climate change, in *Climate change and the Great Barrier Reef: a vulnerability assessment*, eds J.E. Johnson and P.A. Marshall, Great Barrier Reef Marine Park Authority and Australian Greenhouse Office, Townsville, pp. 667-716.
309. Whiteway, T., Smithers, S., Potter, A. and Brooke, B. 2013, *Geological and geomorphological features of Outstanding Universal Value in the Great Barrier Reef World Heritage Area: Technical report prepared for the Department of Sustainability, Environment, Water, Population and Communities*, Geoscience Australia and James Cook University, Townsville.
310. Great Barrier Reef Marine Park Authority 2011c, *Extreme weather and the Great Barrier Reef*, GBRMPA, Townsville.
311. Hopley, D., Smithers, S.G. and Parnell, K.E. 2007, *The geomorphology of the Great Barrier Reef: Development, diversity and change*, Cambridge University Press, Cambridge, UK.
312. Tapim, J. 2013, *Personal communication*, Manager, Indigenous Partnerships Group, Great Barrier Reef Marine Park Authority.
313. Ross, A., Herman, K.P., Snodgrass, J.G., Delcore, H.D. and Sherman, R. 2011, *Indigenous peoples and the collaborative stewardship of nature*, Left Coast Press, Walnut Creek, California, USA.
314. Azzarello, M.Y. and van Vleet, E.S. 1987, Marine birds and plastic pollution, *Marine Ecology Progress Series* 37: 295-303.
315. Kamminga, J. 2002, *Australian Aboriginal Timber Quick Search*, Australian Institute of Aboriginal and Torres Strait Islander Studies, Canberra.
316. Mann, M. 2013, *Personal Communication*, Dharumbal Traditional Owner.
317. Allan, R. and Barnett, B. 2010, *Caring for our Coast Community Resource Book*, North Queensland Dry Tropics, Townsville.
318. Department of Sustainability, Environment, Water, Population and Communities 2013, *Commonwealth marine reserves*, Department of the Environment, Canberra, <<http://www.environment.gov.au/marinereserves/management.html>>.
319. Bird, M. and Heijm, N. 2009, Appendix R1: Indigenous cultural heritage report, in *Port Expansion Project EIS Port of Townsville*, Townsville, pp. 79-84.
320. Townsville Bulletin 2011, *Forensics examines skeleton*, November 30, News Limited, Townsville.
321. Pentecost, P. 2007, *Indigenous cultural significance assessment, Mission Beach: An initiative of FNQ NRM Ltd*, Giringun Aboriginal Corporation, Tully.
322. Blackman, C. 2012, *Personal Communication*, Gurang Traditional Owner.
323. Australian Transport Safety Bureau 1997, *Departmental investigation into the grounding of the Panamanian flag refrigerated cargo vessel Peacock on Piper Reef, in the Great Barrier Reef, on 18 July 1996*, ATSB, Canberra.
324. Marshall, P.A. 2002, *Impact assessment report: grounding of the MVDoric Chariot on Piper Reef*, Great Barrier Reef Marine Park Authority, Townsville.
325. Australian Transport Safety Bureau 2001, *Independent investigation into the grounding of the Malaysian flag container ship: Bunga Teratai Satu*, ATSB, Canberra.
326. Tapim, J. 2012, *Gungandji Traditional Owners Discussion with Manager, Indigenous Partnerships Group*, Great Barrier Reef Marine Park Authority.
327. Fairfax Media 2011, *Clans agree to stop dugong hunting*, September 9, Sydney Morning Herald.
328. Bird, M.K. 1992, The impact of tropical cyclones on the archaeological record: An Australian example, *Archaeology in Oceania* 27(2): 75-86.
329. Pickrell, J. 2011, *Aboriginal groups impose dugong hunting ban*, September 13 edn, Australian Geographic Society, Sydney.
330. Rist, P. 2011, *Students learn first hand impacts of climate change on Indigenous communities, Media release*, Great Barrier Reef Marine Park Authority, Townsville, <<http://www.gbrmpa.gov.au/archive/archive-media-releases/media-releases-2009/students-learn-first-hand-impacts-on-climate-change-on-indigenous-communities>>.
331. Harris, A.N.M., Kwan, D. and Williams, G. 2000, *Torres Strait Turtles*, Australian Fisheries Management Authority, Canberra.
332. Whatmough, S., VanPutten, I. and Chin, A. 2011, From hunters to nature observers: a record of 53 years of diver attitudes towards sharks and rays and marine protected areas. *Marine and Freshwater Research* 62: 755-762.
333. Context Pty Ltd 2013, *Defining the aesthetic values of the Great Barrier Reef*, Department of Sustainability, Environment, Water, Population and Communities, Canberra.
334. Anthony, K., Dambacher, J.M., Walsh, T. and Beeden, R. 2013, *A framework for understanding cumulative impacts, supporting environmental decisions and informing resilience-based management of the Great Barrier Reef World Heritage Area*, Great Barrier Reef Marine Park Authority, Canberra.
335. Levins, R. 1998, Qualitative mathematics for understanding, prediction, and intervention in complex systems, in *Ecosystem Health*, eds D. Rapport, R. Costanza, P.R. Epstein, C. Gaudet and R. Levins, Blackwell Science, Malden, Massachusetts, USA, pp. 178-204.
336. Levins, R. 1966, The Strategy of Model Building in Population Biology, *American Scientist* 54: 421.
337. Burkholder, J.M., Tomasko, D.A. and Touchette, B.W. 2007, Seagrasses and eutrophication, *Journal of experimental marine biology and ecology* 350(1-2): 46-72.
338. Brodie, J., Waterhouse, J., Maynard, J., Bennett, J., Furnas, M., Devlin, M., Lewis, S., Collier, C., Schaffelke, B., Fabricius, K., Petus, C., da Silva, E., Zeh, D., Randall, L., Brando, V., McKenzie, L.J., O'Brien, D., Smith, R., Warne, M.S.J., Brinkman, R., Tonin, H., Bainbridge, Z., Bartley, R., Negri, A., Turner, R.D.R., Davis, A., Bentley, C., Mueller, J., Alvarez-

Romero, J.G., Henry, N., Waters, D., Yorkston, H. and Tracey, D. 2013b, *Assessment of the relative risk of water quality to ecosystems of the Great Barrier Reef: A report to the Department of the Environment and Heritage Protection*, Centre for Tropical Water & Aquatic Ecosystem Research, James Cook University, Townsville.

339. Johnson, J.E., Maynard, J.A., Devlin, M.J., Wilkinson, S., Anthony, K.R.N., Yorkston, H., Heron, S.F., Puotinen, M.L. and van Hooijdonk, R. 2013, Resilience of Great Barrier Reef marine ecosystems and drivers of change, in *Synthesis of evidence to support the Reef Water Quality Scientific Consensus Statement 2013*, Department of the Premier and Cabinet, Brisbane.

340. Waterhouse, J., Maynard, J., Brodie, J., Zeh, D., Randall, L., Lewis, S., Petus, C., Devlin, M., da Silva, E., Furnas, M., Schaffelke, B., Fabricius, K., Brando, V., McKensie, L., Collier, C., Warne, M.S.J., Smith, J., Henry, N., Yorkston, H. and Tracey, D. 2013, Assessment of the risk of pollutants to ecosystems of the GBR including differential risk between sediments, nutrients and pesticides and among NRM Regions, in *Assessment of the relative risk of water quality to ecosystems of the Great Barrier Reef: A report to the Department of the Environment and Heritage Protection*, eds J. Brodie, J. Waterhouse, J. Maynard and et al., Centre for Tropical Water & Aquatic Ecosystem Research, James Cook University, Townsville, pp. 18-89.

341. De'ath, G. and Fabricius, K.E. 2010, Water quality as a regional driver of coral biodiversity and macroalgae on the Great Barrier Reef, *Ecological Applications* 20(3): 840-850.

342. Cooper, T.F., Ridd, P.V., Ulstrup, K.E., Humphrey, C., Slivkoff, M. and Fabricius, K.E. 2008, Temporal dynamics in coral bioindicators for water quality on coastal coral reefs of the Great Barrier Reef, *Marine and Freshwater Research* 59: 703-716.

343. Collier, C.J., Waycott, M. and McKenzie, L.J. 2012, Light thresholds derived from seagrass loss in the coastal zone of the northern Great Barrier Reef, Australia, *Ecological Indicators* 23: 211-219.

344. Wenger, A.S., Johansen, J.L. and Jones, G.P. 2011, Suspended sediment impairs habitat choice and chemosensory discrimination in two coral reef fishes, *Coral Reefs* 30: 879-887.

345. Wenger, A.S. and McCormick, M.I. 2013, Determining trigger values of suspended sediment for behavioral changes in a coral reef fish, *Marine Pollution Bulletin* 70(12): 73-80.

346. Wenger, A.S., Johansen, J.L. and Jones, G.P. 2012, Increasing suspended sediment reduces foraging, growth and condition of a planktivorous damselfish, *Journal of Experimental Marine Biology and Ecology* 428: 43-48.

347. Wenger, A.S., McCormick, M., McLeod, I.M. and Jones, G.P. 2013, Suspended sediment alters predator-prey interactions between two coral reef fish, *Coral Reefs* 32(2): 369-374.

348. Brodie, J. and Fabricius, K. 2008, Terrestrial runoff to the Great Barrier Reef and the implications for its long term ecological status, in *The Great Barrier Reef: Biology, Environment and Management*, eds P.A. Hutchings, M.J. Kingsford and O. Hoegh-Guldberg, CSIRO, Collingwood, pp. 108-113.

349. Furnas, M., Brinkman, R., Fabricius, K., Tonin, H. and Schaffelke, B. 2013, Chapter 1: Linkages between river runoff, phytoplankton blooms and primary outbreaks of crown-of-thorns-seastars in the northern GBR, in *Assessment of the relative risk of degraded water quality to ecosystems of the Great Barrier Reef: supporting studies. A report to the Department of Environment and Heritage Protection, Queensland Government*, ed. J. Waterhouse, Centre for Tropical Water & Aquatic Ecosystem Research, James Cook University, Townsville.

350. Great Barrier Reef Marine Park Authority 2013c, *Reef rescue marine monitoring program quality assurance and quality control manual 2011/2012*, GBRMPA, Townsville.

351. O'Neill, P. 2009, *State of the Environment Report for Shoalwater Bay Training Area 2008*, Department of Defence, Canberra.

352. Turner, M. and Batianoff, G.N. 2007, Vulnerability of island flora and fauna of the Great Barrier Reef to climate change, in *Climate change and the Great Barrier Reef: a vulnerability assessment*, eds J.E. Johnson and P.A. Marshall, Great Barrier Reef Marine Park Authority and Australian Greenhouse Office, Townsville, pp. 621-666.





## Chapter 7

### Current condition and trend







<b>7.5</b>	<b>Environmental processes .....</b>	<b>7–29</b>
<b>7.6</b>	<b>Condition and trend of matters of national environmental significance .....</b>	<b>7–33</b>
7.6.1	World heritage properties .....	7–33
7.6.2	Great Barrier Reef Marine Park .....	7–41
7.6.3	National heritage places.....	7–41
7.6.4	Commonwealth marine area .....	7–42
7.6.5	Listed migratory and listed threatened species.....	7–42
7.6.6	Wetlands of international importance .....	7–43
<b>7.7</b>	<b>Key information gaps .....</b>	<b>7–43</b>
7.7.1	Biodiversity — habitats and species .....	7–43
7.7.2	Geomorphological features .....	7–44
7.7.3	Indigenous and historic heritage values .....	7–44
7.7.4	Community benefits of the environment .....	7–45
7.7.5	Environmental processes.....	7–45
7.7.6	Processes to address information gaps .....	7–46
<b>7.8</b>	<b>Summary of conclusions .....</b>	<b>7–46</b>
<b>References.....</b>		<b>7–47</b>

## Table of contents

<b>7.1</b>	<b>Biodiversity .....</b>	<b>7–3</b>
7.1.1	Summary of overall condition of biodiversity.....	7–13
7.1.2	Coral reefs .....	7–13
7.1.3	Seagrass meadows .....	7–15
7.1.4	Marine turtles .....	7–16
7.1.5	Inshore dolphins .....	7–17
7.1.6	Dugongs .....	7–19
7.1.7	Seabirds.....	7–19
7.1.8	Targeted bony fish species.....	7–19
7.1.9	Sharks and rays.....	7–19
<b>7.2</b>	<b>Geomorphological features.....</b>	<b>7–20</b>
<b>7.3</b>	<b>Indigenous and historic heritage values .....</b>	<b>7–22</b>
7.3.1	Traditional Owner connections .....	7–24
7.3.2	Cultural keystone species.....	7–24
7.3.3	Identifying Indigenous heritage values .....	7–25
7.3.4	Historic heritage values .....	7–25
<b>7.4</b>	<b>Community benefits of the environment.....</b>	<b>7–25</b>
7.4.1	Economic benefit .....	7–27
7.4.2	Visitation .....	7–28
7.4.3	Understanding and appreciation.....	7–28
7.4.4	Personal connections .....	7–29
7.4.5	Knowledge about community benefits .....	7–29

Cover page image: Seabirds on Michaelmas Cay, with anchored yacht.

## **Extract from Great Barrier Reef Region Strategic Assessment terms of reference**

### **2.2 Condition and trend of matters of national environmental significance**

- a) *describe the current condition and trend of key indicators of the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area*
- b) *for world heritage values, benchmark the current condition of key indicators of outstanding universal value against the retrospective Statement of Outstanding Universal Value which describes the state of the Great Barrier Reef World Heritage Area at the time of listing in 1981*
- c) *identify key information gaps and processes to address critical information needs.*

This chapter contains an assessment of the current condition and trend of the key indicators of matters of national environmental significance in the Great Barrier Reef Region (the Region). Each key indicator identified in Chapter 4 is assessed against a standard set of grading statements in four categories: biodiversity, geomorphological features, Indigenous and historic heritage values, and community benefits derived from the environment (Sections 7.1 to 7.4). The current condition of the key indicators of environmental processes relevant to the matters of national environmental significance is also assessed against a standard set of grading statements (Section 7.5).

A summary of the condition and trend of each matter of national environmental significance, including the Reef's outstanding universal value, is presented in Section 7.6 and the chapter concludes with a summary of key information gaps concerning the Region's values.

## **7.1 Biodiversity**

Outcomes of an assessment of the condition and recent trend of key biodiversity values and attributes (see Chapter 4) are presented below:

- Table 7.1 — Great Barrier Reef habitats (11 elements)
- Table 7.2 — terrestrial habitats that support the Great Barrier Reef (nine elements)
- Table 7.3 — species (17 elements).

A discussion of the key findings about the condition and trend of some biodiversity indicators follows the tables.

While Indigenous heritage values are considered separately in Section 7.3, it is recognised that the interconnection between the Great Barrier Reef's natural environment and its Indigenous heritage means the condition of biodiversity values is intrinsically linked to the condition of Indigenous heritage values. Similarly, the community benefits derived from the environment (Section 7.4) are linked to the condition of the Reef's biodiversity. Likewise, the aesthetic values encompassed under criterion (vii) of the World Heritage Convention for attributes which 'contain superlative natural phenomena or areas of exceptional natural beauty and aesthetic importance' are closely linked to the condition of biodiversity within the property.

## Shifting baselines: How much has the Reef changed?

When we look at the Great Barrier Reef, we tend to compare it with our previous experiences — we have a ‘baseline’ of what we think the Reef is like naturally and, therefore, how much it has changed. However, such a baseline can move with time as gradual changes accumulate, often without being noticed. Older generations may remember when there were more and bigger fish, but younger generations may consider the current abundance as normal or healthy. This is particularly an issue in marine environments where humans, as terrestrial beings, have only recently developed the technology to undertake surveys and research, although we have been exploiting the oceans for much, much longer. In fact, research has been largely conducted in marine systems that are already degraded to some extent, and there is little understanding of how these systems operated in the absence of human activity.<sup>1</sup> A lack of marine ecosystem knowledge and understanding, combined with our preconceptions of what comprises a pristine ecosystem<sup>2</sup>, has resulted in a ‘shifting baseline’ syndrome, where chronic, imperceptible decline has influenced our collective awareness of what the system used to be.

On the Great Barrier Reef, most scientific research and monitoring began in the 1970s and 1980s. Only a few years before proclamation of the *Great Barrier Reef Marine Park Act 1975* and the 1981 world heritage listing, two of the Reef’s principal research institutions were established — James Cook University in 1978 and the Australian Institute of Marine Science in 1972. But there is increasing evidence of profound changes in the Reef well before this time, some stretching back over the past 200 years (Figure 7.1). The lack of rigorous scientific data before the 1980s presents a significant challenge for assessing the true condition and long-term trend of the ecosystem, including the risk of using a shifted baseline to make the assessment. For example, the first systematic surveys of subtidal reefs in the late 1960s began after a severe outbreak of crown-of-thorns starfish had impacted coral populations along much of the Great Barrier Reef.<sup>1</sup> Our understanding of the significance of recent declines in coral reefs<sup>3,4</sup> depends critically on the context of those declines — if they follow significant undocumented declines prior to the commencement of monitoring, then we have an even greater cause for concern.

Evidence for shifted baselines in the Great Barrier Reef has come from older people who remember how conditions were different<sup>5</sup> or from observations recorded in images, journals and ship’s logs. Historical photographs of inshore coral reefs have been especially powerful in illustrating long-term changes in these habitats (Figure 7.1).<sup>6</sup> Comparison of historical photographs showing abundant and diverse corals with more recent ones suggests, in many cases, there has been a long-term decline in condition in many areas. There are some scientific challenges in interpreting these photographs<sup>7</sup> but, importantly, it is now simply not possible to find healthy reef communities at many sites that had such communities historically. This strongly suggests the changes represent long-term degradation.


Traditional Owners and many older people in the broader community consider that fish stocks and other marine resources have declined from the very considerable early bounty that was available on the Reef. This conclusion is supported by fish surveys on inshore reefs. The results of these surveys suggest coral trout stocks on these reefs in general were markedly depleted by as early as 1984, before widespread monitoring began.<sup>8</sup>

Recent research provides rigorous evidence of major changes in inshore reefs, before extensive monitoring was developed in the 1980s. This supports the realisation that our current knowledge of the condition of inshore reefs is actually a shifted baseline. The coral community composition on an inshore island (Pelorus Island in the Palm Islands, north of Townsville) had remained largely stable for many hundreds of years, until the mid-1900s when there were major, unprecedented changes, either to different coral types or to communities no longer dominated by corals.<sup>9</sup> These changes coincided with major development of the coastal zone and catchments in the area. Importantly, the results suggest that what we think of as the ‘natural’ community on these reefs may well be a result of much more turbid, disturbed water conditions, all occurring before the research and monitoring of recent decades.

A shifting baseline is also an issue for understanding trends in environmental conditions on the Reef, such as water quality. Again, there is very little data prior to the 1980s. One of the few pieces of evidence comes from the Low Isles expedition of 1928 when measurements of water turbidity were taken. These suggest the water on the inshore reefs near Port Douglas was relatively clear compared to current conditions. However, water clarity is notoriously variable in time and space, and there is insufficient replication of measurements since 1928 to know if those conditions were typical. Major changes in the activities that cause increased sediments and nutrients in Reef waters suggest the current monitoring is set in the context of a dramatically changed baseline.

This emerging picture of how much the Reef may have changed presents a significant challenge for management of the Great Barrier Reef Region, since it makes it difficult to benchmark progress. It is clear that key aspects of the ecosystem, especially in the inshore southern areas, were already significantly degraded at the time of world heritage listing in 1981. This means we may need to aim for improvements to 1981 conditions, rather than just maintaining or restoring conditions present at the time of listing.





**Figure 7.1 Shifting baselines on inshore coral reefs, Stone Island, offshore Bowen**

Comparisons of historical photographs provide an illustration of the changes in inshore coral reefs over the last century. This series of photographs can be accurately compared using the skyline in the background. The changes largely took place before monitoring programs commenced. Although the condition of the Reef has been fairly stable for the past 20 years or so, it cannot be assumed to be in a natural or healthy condition — in fact, the Reef has degraded and its current, stable condition is a shifted baseline. (2012 photograph courtesy of the University of Queensland)

**Table 7.1 Current condition and trend of biodiversity values — Great Barrier Reef habitats**

Understanding the table			
Very good	Good	Poor	Very poor
<b>Very good:</b> All major habitats are essentially structurally and functionally intact and able to support all dependent species.	<b>Good:</b> There is some habitat loss, degradation or alteration in some small areas, leading to minimal degradation but no persistent, substantial effects on populations of dependent species.	<b>Poor:</b> Habitat loss, degradation or alteration has occurred in a number of areas leading to persistent substantial effects on populations of some dependent species.	<b>Very poor:</b> There is widespread habitat loss, degradation or alteration leading to persistent, substantial effects on many populations of dependent species.
Area (See Chapter 2, Figure 2.3)		Trend	Confidence in condition and trend
<b>N.I.</b> Northern inshore		↑ Improving	● Adequate high-quality evidence and high level of consensus
<b>N.O.</b> Northern offshore		↔ Stable	◐ Limited evidence or limited consensus
<b>S.I.</b> Southern inshore		↓ Deteriorating	○ Very limited evidence, assessment based on anecdotal information
<b>S.O.</b> Southern offshore		— No clear trend	

	Area	Condition and trend			
		Very good	Good	Poor	Very poor
<b>Overview:</b> Information on condition and trend of habitats is highly variable with some well known (e.g. shallow coral reefs) and others poorly known, particularly habitats in remote areas or deep waters (e.g. <i>Halimeda</i> banks). Habitats such as coral reefs, seagrass meadows and open waters which are critical in supporting a range of other habitats and species have deteriorated in the southern two-thirds of the Region, especially inshore. This has resulted in an overall grade of poor for that area. The habitats of the northern third of the Great Barrier Reef are believed to remain in good or very good condition, and are able to support dependent species.	N.I.	↔			
	N.O.	↔			
	S.I.			↓	
	S.O.		↔		

	Area	Condition and trend				Confidence	
		Very good	Good	Poor	Very poor	Condition	Trend
<b>Islands:</b> There are about 1050 islands within the World Heritage Area, including continental islands, coral cays and mangrove islands. There is limited monitoring of the condition of most islands. Most are likely to be in good condition, but there is increasing pressure from recreational use, coastal development and climate change. <b>References:</b> <sup>10,11,12,13,14,15</sup>	N.I.		↔			●	●
	N.O.		↔			●	●
	S.I.		↓			●	●
	S.O.		↔			●	●
<b>Beaches and coastlines:</b> Beaches and coastlines are important habitats for migratory shorebirds, seabirds and marine turtles. In the remote north, they remain relatively undisturbed, except for marine debris. Structures near urban centres and ports have extensively modified some coastline habitats and affected coastal processes. Artificial barriers to freshwater flow have disrupted sediment supply to beaches, and increased fine sediments have resulted in mangrove forests replacing beaches. <b>References:</b> <sup>16,17,18,19</sup>	N.I.	↔				●	●
	S.I.		↓			●	●
<b>Mangrove forests:</b> The Region includes an estimated 2070 square kilometres of mangrove habitat. Mangrove forests are a dynamic habitat, with some localised declines and some expansions. The overall condition of mangrove forests is relatively stable and abundance is being maintained. <b>References:</b> <sup>17,20,21,22,23,24,20,25</sup>	N.I.	↔				●	●
	S.I.		↔			●	●
<b>Seagrass meadows:</b> Intertidal seagrass meadows are in poor condition with serious declines reported over the past four years, especially those in the paths of cyclones and those exposed to flooding. Little is known about the abundance and condition of subtidal and deepwater seagrass meadows. The limited information available suggests these habitats can be affected by severe cyclones. Fewer impacts mean northern area meadows are likely to be in very good condition. <b>References:</b> <sup>26,27,28,29,30,31,32,33,34</sup>	N.I.	↔				●	●
	N.O.	↔				○	○
	S.I.				↓	●	●
	S.O.			↓		○	○
<b>Coral reefs (&lt;30 m):</b> Since 1986, average hard coral cover on reefs of the Great Barrier Reef ecosystem is estimated to have declined from 28 to 13.8 per cent and the rate of decline is increasing. The two major causes of loss are cyclones and outbreaks of crown-of-thorns starfish; others include catchment run-off and coral bleaching. The decline is most severe on southern inshore reefs. <b>References:</b> <sup>4,35,36,37,9,38,39,40,41,42,43,44</sup>	N.I.		↔			●	●
	N.O.		↔			●	●
	S.I.				↓	●	●
	S.O.			↓		●	●

	Area	Condition and trend				Confidence	
		Very good	Good	Poor	Very poor	Condition	Trend
<b>Deeper reefs (&gt;30 m):</b> There is no long-term data on the condition of mesophotic and deeper cold water reefs, but for most there is no indication of recent physical damage. The one exception is Myrmidon Reef for which there is information from the 1980s and after cyclone Yasi in 2011. This showed substantial damage to this deepwater reef. Recent studies indicate these reefs have a higher coral diversity than previously thought. <b>References:</b> <sup>45,46,47,48</sup>	N.O.						
	S.O.						
<b>Lagoon floor:</b> The lagoon floor generally consists of soft sand and mud and supports a wide range of species. While a large-scale study of the Region's lagoon floor has provided a comprehensive and extensive snapshot of the habitat, there is no long-term monitoring. The habitat is likely to be in good condition overall. Some areas that have experienced impacts from trawling are likely to be in poor condition, albeit recovering. Recent extreme weather is likely to have damaged lagoon floor habitats, but no assessments have been made. <b>References:</b> <sup>49,50,51,52</sup>	N.I.						
	N.O.						
	S.I.						
	S.O.						
<b>Shoals:</b> Shoals include continental rock, Pleistocene reef edges and submerged coral reefs. They are diverse and variable and provide habitat and structure in an otherwise flat seafloor. They remain in good condition throughout the Region with no evidence that they are significantly threatened. <b>References:</b> <sup>53,54,55,56,57</sup>	N.I.						
	N.O.						
	S.I.						
	S.O.						
<b>Halimeda banks:</b> Large areas of the far northern Great Barrier Reef are dominated by <i>Halimeda</i> , calcareous green algae. The habitat is poorly studied, but is likely to be in very good condition given its isolation from land-based impacts and its level of protection from trawling. Its rate of calcification is likely to be affected by changes in ocean chemistry. <b>References:</b> <sup>49,51,52,58,59,60</sup>	N.O.						
<b>Continental slope:</b> The continental slope makes up about 15 per cent of the Region. The habitat is little studied. Much of the habitat remains undisturbed and is, therefore, likely to be in very good condition. A deepwater trawl fishery operating in the south-east of the Region probably has some physical impacts on the continental slope. <b>References:</b> <sup>45,49,61,62</sup>	N.O.						
	S.O.						
<b>Open waters:</b> Many species depend on open waters. Inshore areas of open water are being degraded, particularly in the southern two-thirds of the Region. There is emerging evidence that an increase in nutrients causes a shift in phytoplankton populations, providing favourable conditions for the development of crown-of-thorns starfish larvae. Offshore and northern areas are considered to be in better condition. <b>References:</b> <sup>63,35,64,65,66,67,68</sup>	N.I.						
	N.O.						
	S.I.						
	S.O.						

**Table 7.2 Current condition and trend of biodiversity values — terrestrial habitats that support the Great Barrier Reef**

This assessment examines the current condition and trend of relevant terrestrial habitats catchments adjacent to the Great Barrier Reef Region. Four broad areas have been examined: northern coastal, northern inland, southern coastal and southern inland.

Understanding the table			
<b>Very good:</b> All major habitats are essentially structurally and functionally intact and able to support all dependent species.	<b>Good:</b> There is some habitat loss, degradation or alteration in some small areas, leading to minimal degradation but no persistent, substantial effects on populations of dependent species.	<b>Poor:</b> Habitat loss, degradation or alteration has occurred in a number of areas leading to persistent substantial effects on populations of some dependent species.	<b>Very poor:</b> There is widespread habitat loss, degradation or alteration leading to persistent, substantial effects on many populations of dependent species.
<b>Area</b> (See Chapter 2, Figure 2.3)		<b>Trend</b>	<b>Confidence in condition and trend</b>
<b>N.C.</b> Northern coastal		↑ Improving	● Adequate high-quality evidence and high level of consensus
<b>N.In.</b> Northern inland		↔ Stable	◐ Limited evidence or limited consensus
<b>S.C.</b> Southern coastal		↓ Deteriorating	○ Very limited evidence, assessment based on anecdotal information
<b>S.In.</b> Southern inland		— No clear trend	

	Condition and trend			
	Very good	Good	Poor	Very poor
<b>Overview:</b> Terrestrial habitats that support the Reef are generally in very good condition in the northern catchment. However, these habitats have been substantially modified in southern areas (south of about Port Douglas), especially saltmarshes, wetlands, woodlands and forests. The functioning of connecting waterbodies has deteriorated, reducing connectivity to the marine environment.				
<b>N.C.</b>	↔			
<b>N.In.</b>	↔			
<b>S.C.</b>			↓	
<b>S.In.</b>			↓	

	Condition and trend				Confidence	
	Very good	Good	Poor	Very poor	Condition	Trend
<b>Saltmarshes:</b> Saltmarshes occur along the length of the Great Barrier Reef coast. They have been significantly modified by coastal development, affecting more than 30 per cent of the habitat in the catchment. The impact is highest in areas with grazing and cropping, urban growth or large communities. <b>References:</b> <sup>66,67,17</sup>						
<b>N.C.</b>	↔				◐	◐
<b>S.C.</b>			↓		◐	◐
<b>Freshwater wetlands:</b> Freshwater wetlands across the catchment are relatively intact, but many are functioning poorly due to a range of factors, including loss of connectivity, sediment and nutrient overload, changes to groundwater and weed infestations. Losses of wetlands are often underestimated, especially for infrequently inundated wetlands on highly developed coastal floodplains. In some coastal floodplain basins, up to 80 per cent of freshwater wetlands have been lost. The rate of wetland loss has slowed in recent years. <b>References:</b> <sup>17</sup>						
<b>N.C.</b>		↔			◐	◐
<b>N.In.</b>		↔			◐	◐
<b>S.C.</b>			↓		◐	◐
<b>S.In.</b>			↓		◐	◐



	Area	Condition and trend				Confidence	
		Very good	Good	Poor	Very poor	Condition	Trend
<b>Forested floodplain:</b> Since European settlement, the area of forested floodplain has been reduced by nearly 50 per cent and much of its remaining extent is subject to grazing. There is 12,700 square kilometres of forested floodplain remaining within the Great Barrier Reef catchment. The habitat is affected by clearing and land modification, changes to overland and groundwater flows, weed and pest invasion, water extraction and reduced connectivity. <a href="#">References:</a> <sup>17</sup>	N.C.	—				●	○
	N.In.	—				●	○
	S.C.			↓		●	●
	S.In.			↓		●	●
<b>Heath and shrublands:</b> Heath and shrublands occur within coastal environments. Approximately 94 per cent of the heath and shrublands in the catchment remains intact, with about 78 per cent afforded protection in national parks, conservation areas and state forests. <a href="#">References:</a> <sup>17</sup>	N.C.	↔				●	●
	N.In.	↔				●	●
	S.C.	↔				●	●
	S.In.	↔				●	●
<b>Grass and sedgelands:</b> Grass and sedgeland habitats occur throughout the catchment. The greatest loss of grasslands has been in the Fitzroy and Burdekin regions where more than 60 and 40 per cent (respectively) of the habitat has been lost. Coastal grasslands have been extensively modified for agricultural production or urban settlements. <a href="#">References:</a> <sup>17</sup>	N.C.	↔				●	●
	N.In.	↔				●	●
	S.C.				—	●	○
	S.In.				—	●	○
<b>Woodlands:</b> The condition of woodlands varies throughout the catchment. There have been significant losses in the Burdekin and Fitzroy regions and an average loss of 39 per cent throughout the catchment. Woodlands are now relatively well protected from broadscale clearing as a result of legislative changes; however, there remains extensive grazing pressure in this habitat. <a href="#">References:</a> <sup>17</sup>	N.C.		↔			●	●
	N.In.		↔			●	●
	S.C.			↔		●	●
	S.In.				↔	●	●
<b>Forests:</b> The condition of forests varies throughout the catchment. There have been losses in all areas, averaging 39 per cent. Forests are now relatively well protected from broadscale clearing. It is estimated that the loss of woody vegetation is due mainly to clearing for agriculture and, to a much lesser extent, urban development. <a href="#">References:</a> <sup>17</sup>	N.C.	↔				●	●
	N.In.		↔			●	●
	S.C.				↔	●	●
	S.In.			↔		●	●
<b>Rainforests:</b> There have been losses in rainforest habitats throughout the catchment, in particular the Wet Tropics, Fitzroy and Burnett–Mary regions. Since pre-European levels, an average 38 per cent has been lost across the catchment. There has been no recorded loss of rainforest in the Wet Tropics since the inscription of the area on the World Heritage List in 1988, at which time logging was completely banned. <a href="#">References:</a> <sup>17</sup>	N.C.	↔				●	●
	N.In.		↔			●	●
	S.C.		↔			●	●
	S.In.		↔			●	●
<b>Connecting waterbodies:</b> Aquatic connections between freshwater and marine environments are still functioning well in northern areas. In contrast, aquatic connectivity has been substantially altered in the south. In southern coastal areas, changes to hydrological flows and the construction of bunds, weirs and other structures have altered the functioning of the waterbodies and decreased connectivity, except in flood events. In the southern upper catchment, the construction of dams is a major barrier. Groundwater connectivity has also been affected. <a href="#">References:</a> <sup>17</sup>	N.C.	↔				●	●
	N.In.	↔				●	●
	S.C.			↓		●	●
	S.In.				↓	●	●

Table 7.3 Current condition and trend of biodiversity values — species

Understanding the table			
<b>Very good:</b> Only a few, if any, species populations have deteriorated as a result of human activities or declining environmental conditions.	<b>Good:</b> Populations of some species (but no species groups) have deteriorated significantly as a result of human activities or declining environmental conditions.	<b>Poor:</b> Populations of many species or some species groups have deteriorated significantly as a result of human activities or declining environmental conditions.	<b>Very poor:</b> Populations of a large number of species have deteriorated significantly.
<b>Area</b> (See Chapter 2, Figure 2.3)	<b>Trend</b>	<b>Confidence in condition and trend</b>	
<b>N.I.</b> Northern inshore	↑ Improving	● Adequate high-quality evidence and high level of consensus	
<b>N.O.</b> Northern offshore	↔ Stable	◐ Limited evidence or limited consensus	
<b>S.I.</b> Southern inshore	↓ Deteriorating	○ Very limited evidence, assessment based on anecdotal information	
<b>S.O.</b> Southern offshore	— No clear trend		

	Area	Condition and trend			
		Very good	Good	Poor	Very poor
<b>Overview:</b> There is only condition and trend information for a limited number of species and species groups; hence, the assessment of some components is highly uncertain and often inferred from the condition of supporting habitats and processes. Of those for which there is information, there have been significant declines in many, especially in the inshore southern two-thirds of the Region, and some iconic and cultural keystone species such as dugongs and some marine turtles. Serious declines have been recorded in most hard corals and seagrasses, some fishes and sharks, and some seabird populations. There are four examples of species showing good recovery after past serious declines: humpback whales, estuarine crocodiles, loggerhead turtles and green turtles (southern stock). However, even these species have not recovered to their original numbers.	N.I.		↔		
	N.O.		↔		
	S.I.			↓	
	S.O.		↔		

	Area	Condition and trend				Confidence	
		Very good	Good	Poor	Very poor	Condition	Trend
<b>Mangroves:</b> The Region's mangrove forests are very diverse, with at least 39 mangrove species and hybrids recorded. The highest diversity is in the far north. In contrast to international trends, the diversity and abundance of mangrove species along the Great Barrier Reef coast are being maintained. <b>References:</b> 69,70,71,22,23	N.I.	↔				◐	◐
	S.I.		↔			◐	◐
<b>Seagrasses:</b> The Great Barrier Reef is maintaining seagrass diversity; however, there have been recent severe declines in abundance and changes in community composition in southern inshore areas. These are mainly due to cyclones and flood events, in addition to the longer term impacts of catchment run-off. Little is known about the condition of offshore seagrass species. <b>References:</b> 27,29,72,30,31	N.I.	↔				◐	◐
	N.O.	↔				○	○
	S.I.				↓	◐	◐
	S.O.			—		○	○
<b>Macroalgae:</b> The biodiversity of macroalgae is being maintained. There is little information about its condition, but it is assumed to be good to very good. Interactions between coral and macroalgae are affected by changes in ocean chemistry and the presence of herbivores. Shifts in the coral-algae balance may influence future coral reef recovery. <b>References:</b> 73,74,75,76,77,78,79,80	N.I.	↔				○	○
	N.O.	↔				○	○
	S.I.		↔			◐	◐
	S.O.	↔				○	○

	Area	Condition and trend				Confidence	
		Very good	Good	Poor	Very poor	Condition	Trend
<b>Benthic microalgae:</b> Benthic microalgae are little studied. It is assumed that they remain undisturbed and in very good condition for much of the Region. <a href="#">References:</a> <sup>81,82</sup>	N.I.	↔				○	○
	N.O.	↔				○	○
	S.I.		↔			○	○
	S.O.	↔				○	○
<b>Corals:</b> Although there is an urgent need for more long-term monitoring of inshore reefs, there is consensus that coral diversity and abundance has substantially decreased on reefs south of Cooktown. Hard coral cover has also declined in southern offshore areas, principally due to cyclone impacts. Coral in the far north of the Region is in better condition, largely due to better water quality and fewer extreme weather events. Little is known about the recently discovered cold water corals in very deep parts of the Region (more than 1000 metres). <a href="#">References:</a> <sup>4,35,36,38,39,40,41,42,44,83,84,85,86,87,88</sup>	N.I.		↔			●	●
	N.O.		↔			●	●
	S.I.				↓	●	●
	S.O.			↓		●	●
<b>Other invertebrates:</b> There are thousands of species of invertebrates in the Region. Some, such as prawns, crabs and sea cucumbers, are important in fisheries. Currently, there are no invertebrate species assessed as overfished in stock status reports. Little is known about the status of most non-commercial species. An outbreak of crown-of-thorns starfish is a serious emerging concern for the health of Great Barrier Reef coral reefs. It can be assumed the changing environment in southern inshore areas has affected dependent invertebrates. <a href="#">References:</a> <sup>49,89,90</sup>	N.I.	↔				○	○
	N.O.	↔				○	○
	S.I.		↓			○	○
	S.O.	↔				○	○
<b>Plankton and microbes:</b> Little is known about the status of plankton and microbes. Changes in water temperature and quality are likely to be altering plankton communities which, in turn, will be affecting higher trophic levels. <a href="#">References:</a> <sup>91,92</sup>	N.I.	↓				○	○
	N.O.	↓				○	○
	S.I.		↓			○	○
	S.O.	↓				○	○
<b>Bony fish:</b> There are about 1600 species of bony fish in the Region. Very little is known about the status of most species. While long-term monitoring of coral reef fish populations does not indicate declines in the species monitored, understanding of fishing activities suggests targeted species are under significantly more pressure in the southern two-thirds of the Region. These are likely to be exacerbated by severe weather events, reduced habitat and declining water quality. There is little evidence of declines in northern populations. <a href="#">References:</a> <sup>90,93,94,95,96,97,98,99,100,101,102,103,104,105,106,107,108,109,110,111</sup>	N.I.	↓				◐	◐
	N.O.	↔				◐	◐
	S.I.		↓			◐	◐
	S.O.		↔			◐	◐
<b>Sharks and rays:</b> There are 136 species of sharks and rays recorded, including some listed threatened and migratory species. Overall, populations of sharks and rays are considered to be in good condition. However, there are significant concerns about a small number of species, including one (the speartooth shark) which may now be extinct in the Region. Their life histories and habitat preferences make them vulnerable to impacts. Inshore and estuarine species are the most affected by cumulative human-induced impacts. <a href="#">References:</a> <sup>112,113,114,115,116,117,118,119,120,121,49,117,122</sup>	N.I.			↓		◐	○
	N.O.		↓			◐	○
	S.I.			↓		◐	○
	S.O.		↓			◐	○
<b>Sea snakes:</b> There are 14 species of sea snakes with permanent breeding populations in the Region. Abundance estimates are only available for a few species or for small areas, and there is little information about population trends. Some species are taken as incidental bycatch in the trawl fishery, but the effects on the populations are poorly understood. <a href="#">References:</a> <sup>49,123,124,125,126</sup>	N.I.			↔		◐	○
	N.O.			↔		◐	○
	S.I.			↔		◐	○
	S.O.			↔		◐	○

	Area	Condition and trend				Confidence	
		Very good	Good	Poor	Very poor	Condition	Trend
<b>Marine turtles:</b> Each of the six species of marine turtle has a different population status and trend. Although the nesting populations of some species are thought to be increasing — for example loggerhead and green turtle (southern stock), other nesting stocks are declining — for example green turtle (northern stock) and hawksbill. Little is known about the foraging components of most populations, but there are concerns for some species particularly due to pressures outside the Region. Declines in seagrass meadows after recent floods and cyclones have placed additional stress on southern green turtle stocks. <b>References:</b> 127,128,129,130,131,132,133,134,135,136,137,138,139	N.I.			↓		●	●
	N.O.			↓		●	●
	S.I.		↔			●	●
	S.O.		↔			●	●
<b>Estuarine crocodiles:</b> Estuarine crocodiles occur in most coastal waters in the Region. They are also regularly reported at mid-shelf and some offshore islands. The species is steadily recovering from previous population declines. <b>References:</b> 140,141,142,143,144	N.I.		↑			○	○
	N.O.		↑			○	○
	S.I.		↑			○	○
	S.O.		↑			○	○
<b>Seabirds:</b> Between 1.4 and 1.7 million seabirds from at least 20 species breed annually on islands and cays in the Region. Declines of up to 70 per cent in nesting seabird populations have been recorded. Although all seabirds are vulnerable to impacts of climate change, it is the offshore and pelagic-foraging seabirds that are most at risk. Some catastrophic nesting failure has already been recorded in the south. <b>References:</b> 12,145,146,147,148,149,150,151,152,153,154,155,156,157,158,159,160,161	N.I.		↔			○	○
	N.O.			↓		○	○
	S.I.		↔			○	○
	S.O.			↓		○	○
<b>Shorebirds:</b> There are no population estimates for the Region's shorebirds. Australia-wide declines of between 70 and 80 per cent have been recorded in the past 24 years. Internationally significant numbers of shorebirds occur at a number of sites within the Region. Changes to the coastline from population growth directly affect the habitats used by shorebirds. <b>References:</b> 162,163,164	N.I.			↓		○	○
	S.I.			↓		○	○
<b>Whales:</b> It is estimated that 15 species of whale visit the Region, and there is no definitive information on the condition of most of these. This assessment is based on knowledge of the humpback whale population, which is recovering strongly after being decimated by whaling. An estimated population of 10,000 in 2007 grew to 14,500 in 2010 and 17,000 in 2012. <b>References:</b> 165,166,167,168,169	N.I.		↑			○	○
	S.I.		↑			○	○
<b>Dolphins:</b> The number of dolphin species in the Region is estimated to be 17. There is limited information about most populations. Two listed inshore species, the Australian snubfin and Indo-Pacific humpback, are considered most at risk and are likely to be in decline. <b>References:</b> 170,171,172,173,174,175,176,177,178,179,180,181,182,183,184,185,186,187,188,189	N.I.		↔			○	○
	N.O.		↔			○	○
	S.I.			↓		○	○
	S.O.		↔			○	○
<b>Dugongs:</b> The northern population of dugong remains healthy with a stable population trend. A substantial decline in dugongs occurred in waters south of Cooktown since the 1960s, but the population was thought to have stabilised in 2009. However, a series of extreme weather events from 2009 to 2011 led to impacts on seagrass meadows and a further recent decline in dugong numbers. <b>References:</b> 190,191,192,193,193,194,195,196,197,198,199,200,201,202,203,204,205,72	N.I.		↔			●	●
	S.I.				↓	●	●



### 7.1.1 Summary of overall condition of biodiversity

The Great Barrier Reef remains one of the world's most unique and biologically diverse ecosystems. Although most habitats and species are assessed to be in good to very good condition at the scale of the whole Great Barrier Reef Region, a lack of accurate or sufficient information on many habitats and species means the assessment is principally based on limited evidence and anecdotal information.

The condition of key habitats, which have adequate certainty, such as coral reefs, seagrass meadows and open waters, all show poor and declining conditions in the southern parts of the Region. Southern mangrove forests are the only habitat with a good and stable condition and trend grading that is supported by adequate high-quality evidence and high level consensus. Corals, turtles and dugongs are the only species associated with an adequate grading of certainty. All of these are in poor or very poor condition and declining in at least one of the four regions. All these species and habitats affect the broader value of the Region, and if these declines are not arrested they are likely to have broadscale impacts.

The habitats and species north of around the Port Douglas–Cooktown area are generally in better condition than those further south. Also, habitats further offshore and in deeper water are typically subject to fewer stresses and are, therefore, presumed to be healthier, including the lagoon floor, shoals, *Halimeda* banks, deeper reefs and the continental slope.

However, unsustainable practices — some dating back more than a century — combined with the last 10 years of extreme weather have affected the ecosystem and reduced its ability to recover from disturbance.<sup>4</sup> This is especially true for inshore areas adjacent to the more developed part of the catchment. For some species, such as dugongs and inshore dolphins, and some habitats, such as coral reefs and seagrass meadows, their condition in this area is assessed as poor or very poor and declining.

The Authority's vulnerability assessments of the elements that make up the Great Barrier Reef's biodiversity<sup>206</sup> identified nine species or groups of species that are highly vulnerable. This included dugong, marine turtles, inshore dolphins (Indo-Pacific humpback and Australian snubfin), sharks and rays including sawfish, sea snakes, seabirds (offshore and pelagic-foraging), king and blue threadfin salmon, and snapper. Many of these at-risk species are listed migratory or threatened species. Many are also cultural keystone species for Great Barrier Reef Traditional Owners.

Vulnerability assessments also identified overwhelming evidence that a range of threats are continuing to affect inshore habitats along the developed coast, and the species that use these habitats.<sup>206</sup> The key ecosystem-level impacts affecting habitats and species and reducing the Great Barrier Reef's resilience are climate change, poor water quality including the effects of catchment run-off, loss of coastal habitats from coastal development, illegal fishing and poaching, and some remaining impacts from lawful fishing.<sup>43</sup> Legacy impacts, some dating back 150 years, include early catchment clearing, historic levels of commercial fishing and unsustainable commercial harvesting of species such as humpback whales, sea cucumber, dugongs and green turtles.<sup>206</sup> These impacts are detailed in Chapter 6.

There are few examples of recovering populations. Those that are recovering are species that declined as a result of human-related impacts which are now eliminated or reduced; for example, commercial whaling for humpback whales and incidental capture of marine turtles in trawl nets. These populations have yet to recover to their original size and, as they tend to be long-lived species, full recovery is likely to take decades.


### 7.1.2 Coral reefs

There are more than 400 species of hard coral and at least 150 species of soft corals, sea fans and sea pens in the Region. Coral reefs are the cornerstone of the Great Barrier Reef ecosystem and their evolutionary history, species diversity, habitat values and natural beauty are major contributors to the Reef's outstanding universal value.

Research indicates coral cover has significantly declined over the past 40 to 50 years through a combination of successive cyclones, crown-of-thorns starfish outbreaks and mass bleaching events<sup>3</sup>, combined with insufficient time for recovery between these disturbances. Since 1986, estimated average hard coral cover on the Great Barrier Reef has declined from 28 to 13.8 per cent (Figure 7.2).<sup>4</sup> The rate of decline has increased substantially in recent years, averaging approximately 1.45 per cent per year since 2006. Two-thirds of the loss has occurred since 1998.<sup>4</sup>

At a Reef-wide scale, the two biggest direct causes of coral cover loss are cyclones<sup>43,44</sup> and crown-of-thorns starfish outbreaks<sup>35,4</sup>.<sup>4</sup> Importantly, such large long-term declines imply corals in some areas are failing to recover fully after disturbances<sup>207</sup>, or their recovery is being interrupted by subsequent disturbances. Catchment run-off is likely to have been a major contributor to such loss of resilience, particularly in inshore areas. At stakeholder meetings during the strategic assessment, participants also raised concerns about the effects on coral reefs of ports and disposal of dredge material.

There is strong evidence of a link between poor water quality and crown-of-thorns starfish outbreaks, through increased nutrients.<sup>35,208</sup> In addition, there is some evidence of a link between these outbreaks and fishing pressures: reefs in fished Habitat Protection Zones (dark blue zones) have almost four times the number of outbreaks as reefs in no-take Marine National Park Zones (green zones)<sup>209</sup>, resulting in reduced coral cover.<sup>8</sup>



**Figure 7.2 Hard coral cover in the Great Barrier Reef, 1986–2012<sup>4</sup>**

Modelling of data from the Australian Institute of Marine Science's long-term monitoring program shows hard coral cover in the Great Barrier Reef has declined significantly since monitoring began in 1986. Declines have been most severe in the central and southern two-thirds of the Region. Crown-of-thorns starfish and cyclones have been responsible for most of the mortality. The solid line represents modelled coral cover, while the dashed lines either side represent the associated standard errors. The inverted bar graphs indicate the relative contributions to annual coral mortality.

The decline in coral cover has been most severe on reefs south of latitude 20 degrees, particularly since 2006. Since that time, coral cover has reduced from about 35 per cent to eight per cent.<sup>4</sup> Over a shorter timeframe and with less available data to assess trends, data from the Authority's marine monitoring program of inshore reefs adjacent to the developed central and southern areas of the catchment, indicates a decline in coral reef health since 2006<sup>210</sup>.

In addition, there is evidence that since European settlement, but prior to most modern monitoring efforts, inshore reefs were particularly affected (Figure 7.1). Research on Pelorus Island employed sophisticated coral core dating techniques of the reef substrate to understand the historical composition of coral assemblages.<sup>9</sup> Results suggest coral communities in the past were remarkably stable and bounced back after disturbance. They were dominated by *Acropora* corals, thought to be characteristic of clearer water. However, between 1920 and 1955, these assemblages collapsed and were replaced either by corals typical of more turbid, muddy waters or by communities with little live coral. Importantly, these changes indicate a reduced capacity for the system to absorb disturbance and bounce back to its previous state.<sup>211</sup> This loss of resilience and degradation coincided with prolonged increases in sediment and nutrient loads in the inshore Great Barrier Reef, due to coastal and catchment development.<sup>9</sup>

Many of the inshore reefs of the southern two-thirds of the Region are similar in condition to those around Pelorus Island, and there is evidence that some of these have undergone similar shifts,<sup>6</sup> suggesting widespread degradation in the condition of these reefs.

It is important to note that the 'very poor' grade for the current condition of southern inshore coral reefs and corals in Table 7.1 and Table 7.3 is based on different criteria to the *Reef Water Quality Protection Plan* report cards and the Reef Rescue marine monitoring program report for 2012 and derives a different outcome. The assessment criteria used in this report are the same for all habitats and species, and are consistent with those used in the Authority's assessment of condition in its outlook reports. It is significant that, despite the different criteria and the difference in grade, neither assessment is suggesting the status of inshore southern coral is 'good' or 'very good'.

In contrast, inshore reefs adjacent to less developed, northern catchments do not appear to have undergone the same loss of resilience and degradation.<sup>212</sup> Coral cover in the far north has not shown similar declines over the past three decades, and is in better condition than in the rest of the Region. This is largely due to better water quality and fewer extreme weather events.<sup>4</sup> The overall health of reefs in this area means they are likely to have greater resilience and are, therefore, better able to recover from impacts — for example, the bleaching event in 2002 which was the cause of the dip in coral cover shown in Figure 7.2.


### 7.1.3 Seagrass meadows

Seagrass meadows are an important component of the Great Barrier Reef ecosystem. They can be found on sandy or muddy bottoms from intertidal areas to depths of 60 metres or more, on reef platforms in individual reef lagoons, and the Great Barrier Reef lagoon.<sup>34</sup> Seagrass meadows are the main food source for dugongs and green turtles — both species of conservation concern. They are also habitat builders and act as nurseries for juvenile prawns, fishes, crabs and tropical lobsters, while also providing high rates of carbon sequestration.

Up until the past five years, seagrass meadows appeared to have been relatively stable for about 20 years.<sup>26</sup> Fluctuations were mainly due to natural cycles, although influenced by catchment run-off. The meadows were also likely to show prompt recovery from declines. However, current evidence from 30 monitored intertidal meadows along the Region's coast suggests seagrass abundance and reproductive effort in and adjacent to large areas in the southern two-thirds of the Region have recently declined (Figure 7.3).<sup>213</sup> Ongoing monitoring shows: 67 per cent of monitoring sites have reduced seagrass abundance; 50 per cent of sites exhibit shrinking meadow area; many sites have limited or no seed production; indications of light limitation at 63 per cent of sites; nutrient enrichment at 33 per cent of sites; and 90 per cent of sites with either high or elevated nitrogen.<sup>34</sup> However, recent data collected through Seagrass-Watch indicates a recovery in the Townsville area since early to mid-2012. Little is known about the abundance and condition of subtidal and deepwater seagrass meadows. The limited information available suggests declines in the abundance of these meadows, and that these habitats can be affected by severe cyclones.


Significant losses of seagrass meadows occurred in 2010–11 in the path of tropical cyclone Yasi.<sup>27</sup> There were also broader scale losses in meadows exposed to flooding. These severe events followed a number of years of decline due to other cyclonic events and major freshwater inflow (see Figure 6.2), as well as declining water quality and extended periods of cloud cover which limits growth periods. Examples of declines include Mourilyan Harbour where seagrass meadows had been consistently present for the past 15 years but have now been lost,<sup>28</sup> as well as substantial reductions in the meadows adjacent to Cairns,<sup>29</sup> Townsville<sup>30</sup> and Gladstone<sup>31</sup>.

The decline in seagrass meadows has had profound effects on species which depend on this habitat.<sup>214,215</sup> The extreme weather events of 2011 led to unprecedented losses of dugongs and green turtles within the Region. About twice as many marine turtles were found stranded on Queensland beaches in that year compared to previous years; the number of strandings remained high in 2012 (Figure 7.4). A detailed analysis of the effects on dugong populations is provided in the dugong demonstration case study presented in Section 9.3.



**Figure 7.3 Average abundance of intertidal seagrass meadows, 1999–2012<sup>27</sup>**

Significant declines in the abundance of intertidal seagrass meadows have been recorded in the southern two-thirds of the Region since about 2009, and are linked to the effects of tropical cyclones, catchment run-off and cloud cover. Intertidal meadows are composed of estuarine, coastal and reefal habitat types. The overall health of seagrass is a combination of abundance, reproductive effort, nutrient status and light availability.



**Figure 7.4 Cumulative monthly marine turtle strandings for the Queensland east coast, 1998–2012<sup>216</sup>**

Marine turtle strandings in 2011 were about double previous years. Higher than normal strandings were also recorded in 2012. The data only includes cases confirmed in the field by a trained person and later verified by an expert.

### 7.1.4 Marine turtles

All marine turtle species in the Region are matters of national environmental significance, being listed as both migratory and threatened species under the *Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act). They are also one of the attributes for listing the Great Barrier Reef as a World Heritage Area. They form a key part of the Marine Park's biodiversity and are listed as a protected species under the Great Barrier Reef Marine Park Regulations 1983. Marine turtle species are also listed under the *Nature Conservation Act 1992* (Qld) and identified as priority species in the Queensland Government's *Back on Track Actions for Biodiversity plans*.<sup>217</sup>

Within the Great Barrier Reef, there are 38 islands which are important nesting sites for marine turtles (see Figure 4.10 in Chapter 4). Of these, Raine Island is highly significant as it supports the world's largest aggregation of nesting green turtles. Other important islands include Milman Island, Wild Duck Island, Peak Island and the cays of the Capricorn Bunker Group.<sup>127,128,129,131</sup> On a global scale, these islands are important for the survival of four species of marine turtle: loggerhead, green, hawksbill and flatback turtles.

Marine turtles, as a group, are exposed to many impacts including climate change, declining water quality from catchment run-off, habitat loss from coastal development, some direct and indirect impacts from commercial and recreational fishing and poaching (illegal hunting), boat strike, disease, feral animals eating eggs and destroying nests, and ingestion of marine debris. There is traditional hunting of green turtles and, based on the best available information, this is considered sustainable. Not all impacts occur on all species or in all parts of the Region. In addition, some impacts occur outside the Region and in the waters and on the beaches of other countries.



Some 1800 marine turtles were reported stranded on the east Queensland coast in 2011 (Figure 7.4).<sup>216</sup> Most were from the southern Great Barrier Reef green turtle stock and were associated with declines in seagrass meadows. Fortunately, the population is expected to withstand this increased mortality as it mainly affected small immature animals rather than productive mature females, and numbers were increasing at the time.<sup>128,135</sup>

Different life histories, habitat preferences and susceptibility to impacts mean the condition and current trend of each of the Region's marine turtle species varies significantly (Table 7.4).

**Table 7.4 Summary of condition and trend for marine turtle species in the Region**

Species	Summary of condition and trend
<b>Green</b> — northern Great Barrier Reef stock	This is the largest remaining green turtle stock in the world. There was an upward trend in the annual nesting population from 1976 to 1996, followed by a downward trend since that time, largely attributed to selective harvesting of adult females throughout their range. There are concerns about juvenile recruitment into the stock. <sup>128</sup>
<b>Green</b> — southern Great Barrier Reef stock	The nesting component of the stock has been increasing at 3.8 per cent per year for the past four decades. There are no indicators of population-level impacts on the nesting beaches used by this stock or its foraging component. <sup>128</sup>
<b>Hawksbill</b>	The primary index nesting site at Milman Island indicates the Torres Strait–northern Great Barrier Reef stock is in decline at about three per cent per year. There is no trend data for the foraging component of the population. <sup>129</sup>
<b>Loggerhead</b>	The nesting population is currently recovering after declining by more than 80 per cent between 1970 and the early 2000s. There are concerns for juvenile recruitment into the foraging population from pressures outside the Region. <sup>127</sup>
<b>Flatback</b>	The nesting component of the eastern Queensland stock is currently stable. There is virtually no data on the foraging population. <sup>131</sup>
<b>Leatherback</b>	A very small number of leatherbacks are known to have nested in the Region, but no nesting has been recorded since 1996. The Region's population is considered to be part of the south-west Pacific genetic stock which is in decline. <sup>132</sup>
<b>Olive Ridley</b>	No nesting occurs in the Region, and there is virtually no data on foraging animals. <sup>130</sup>

### 7.1.5 Inshore dolphins

Two of the Region's dolphin species are considered to be at risk: the Australian snubfin dolphin and the Indo-Pacific humpback dolphin.<sup>176,218</sup> They are listed as migratory species under the EPBC Act and are two of the many species of global conservation significance included in the Statement of Outstanding Universal Value of the Great Barrier Reef World Heritage Area — making them matters of national environmental significance.

The Australian snubfin and Indo-Pacific humpback dolphins occupy similar inshore habitats and have small (less than 350 square kilometres) discrete home ranges along the coast, with little overlap.<sup>219</sup> Each localised population is estimated to have fewer than 100 individuals, meaning there is a very high reliance on mature females within each population.<sup>176,219</sup> These factors make the species particularly susceptible to any human-induced mortality, potentially resulting in localised extinction.

There are currently no overall population estimates for the Australian snubfin or Indo-Pacific humpback dolphins in the Region;<sup>170</sup> however, there are local population estimates for Australian snubfin dolphin aggregations in Cleveland–Halifax Bays<sup>176</sup> and in Keppel Bay–Fitzroy River<sup>219</sup> of about 70 dolphins each. An aggregation has also been recorded at Princess Charlotte Bay–Bathurst Bay on Cape York Peninsula,<sup>220</sup> but no population census has been undertaken. All three areas have either an extensive port expansion program planned (Cleveland Bay) or proposals for new port developments (Fitzroy River delta and Bathurst Bay).

There have been population estimates for Indo-Pacific humpback dolphins in Cleveland Bay (50 or less)<sup>176</sup>; the Capricorn coast (about 64); Keppel Bay (about 107); and Port Curtis (about 85)<sup>221</sup>. Populations are known to inhabit the Great Sandy Strait<sup>172</sup> and Moreton Bay<sup>173</sup>, south of the Region.

The apparent declining trend in populations of these species throughout the Region is of great concern. For populations to remain stable, modelling suggests that the Australian snubfin population in Cleveland–Halifax Bays and Keppel Bay–Fitzroy River can sustain a human-induced death rate of only one animal every four years or one animal every year, respectively.<sup>176,219</sup> The human-induced deaths of two snubfin dolphins in Halifax Bay in 2011 means the long-term viability of this population is at risk. There are similar concerns for Indo-Pacific humpback dolphins in Keppel Bay and Port Curtis where eight died of unknown causes in 2011.<sup>216</sup> Although there are limited reports of mortality of inshore dolphins in gillnets (including nets associated with the Queensland Shark Control Program), the nets are recognised as a serious threat to these species due to the extremely low levels of human-induced mortality that the populations can withstand.<sup>219,176</sup> In the 20-year period from 1991 to 2010, the


Queensland Marine Wildlife Stranding and Mortality Database (StrandNet) reported 35 Australian snubfin and Indo-Pacific humpback dolphins drowned in nets from the Queensland Shark Control Program and from the commercial net fishery.<sup>222</sup> Indirect impacts such as the degradation of inshore habitats and declining water quality are likely to be also affecting inshore dolphins.<sup>186,219,223</sup> The carrying capacity of the home range is directly related to the quality and productivity of its habitat.<sup>224,225</sup> Recognising that known populations of inshore dolphins within the Region are considered to be isolated,<sup>176,219</sup> their survival appears to be closely related to the quality and quantity of their habitat.<sup>219</sup>

Other direct impacts on dolphins include disease and ingestion of or entanglement in marine debris.<sup>183</sup>

### Improving outcomes for inshore dolphins

In recent years, a number of management arrangements have been introduced to better protect the Australian snubfin dolphin and the Indo-Pacific humpback dolphin. Declaration of Dugong Protection Areas in the late 1990s and rezoning of the Great Barrier Reef Marine Park in 2004 mean that 132,500 square kilometres of the inshore waters of eastern Queensland are now closed to large-mesh gill netting. However, core habitats for Australian snubfin and Indo-Pacific humpback dolphins are only partially protected by these arrangements (Figure 7.5).<sup>170,219,226</sup> For example, the majority of the Fitzroy River Australian snubfin dolphin population live in a General Use zone<sup>219</sup> which allows for a wide range of uses such as fishing, aquaculture, shipping and boating.

In 2006, the Queensland Government recommended two devices to reduce interactions between commercial fishers and dolphins: passive acoustic monitoring to enable fishers to detect the presence of dolphins and avoid interaction; and acoustic alarms to deter dolphins from approaching fishing gear. A recent study found these devices may not be as effective as first thought. The dolphins were only vocal about one-third of the time, reducing their chance of detection by fishers, and the likelihood of the dolphins leaving the area did not change significantly when the acoustic alarms were active.<sup>226</sup>



**Figure 7.5 Overlap of protected areas and inshore dolphin populations**

Areas known to be occupied by Indo-Pacific humpback and Australian snubfin dolphins are outlined in purple.  
(a) Hinchinbrook Channel, Halifax and Cleveland Bay, and (b) Keppel Bay and Port Curtis region.

Increasing coastal development is predicted to magnify pressures on inshore dolphins. For example, if approved, proposed developments within the Fitzroy River estuary would directly modify between 14 to 25 per cent of the known Australian snubfin dolphin habitat in that area.<sup>219</sup>

The most critical components likely to contribute to an improvement in the condition and trend of these inshore dolphins are preventing further habitat degradation, enhancing water quality and reducing all impacts causing mortality in the species. Improved management of the species also requires improved knowledge of their abundance and distribution, including a population estimate for the entire World Heritage Area.

### 7.1.6 Dugongs

The Region is home to a globally significant population of dugongs — a listed migratory species under the EPBC Act, and one of the key attributes for listing the Great Barrier Reef as a World Heritage Area. Dugongs are also of great cultural, spiritual and social importance, especially to the Reef's Traditional Owners.

The Great Barrier Reef provides essential habitat and population connectivity between Queensland's two most significant locations for dugong — Torres Strait and Hervey Bay.<sup>227</sup> In northern areas of the Reef, populations are in good condition and the trend over several decades of monitoring is stable. However, dugong populations south of Cooktown have been mostly declining for decades, with an indexed population estimate of only 600 animals between the Daintree River and the southern limit of the Region in 2011,<sup>190,228</sup> compared with an indexed estimate of 2059 from the previous survey in 2005. This is the lowest number recorded in this area since surveys began in 1986, and is considered a response to several years of above average rainfall and the broadscale impacts of cyclone Yasi in February 2011. The decline is likely to be explained mostly by animals moving out of the survey area to seagrass meadows elsewhere, rather than by mortality.

The population status of dugong and the factors that have contributed to this decline are discussed in detail in the dugong demonstration case study (see Section 9.3).

### 7.1.7 Seabirds

The Great Barrier Reef Region supports more than 10 per cent of Australia's breeding seabird population,<sup>146</sup> not counting short-tailed shearwaters which nest in huge numbers in southern Australia. There are significant nesting sites on islands throughout the Region for a range of seabirds. Each year, between 1.4 and 1.7 million seabirds from at least 20 species breed on islands and cays in the Region.<sup>146</sup> This represents more than 25 per cent of Australia's tropical seabirds, more than 50 per cent of offshore foraging black noddies and approximately 25 per cent of wedge-tailed shearwaters, brown and masked boobies and red-tailed tropic birds.<sup>146</sup> The number of non-breeding birds (birds which use the ecosystem for feeding but breed elsewhere) is estimated to be about 425,000, giving a total seabird population that may exceed two million.

Seabirds are relevant to a number of matters of national environmental significance (see Chapter 4). Six of the Region's breeding and non-breeding seabirds are listed threatened species and 23 are listed migratory species under the EPBC Act. In addition, the number of breeding seabird species is included in the Statement of Outstanding Universal Value for the Great Barrier Reef World Heritage Area.

In this time of a changing climate, there is growing concern about seabird species with foraging strategies and prey species that are strongly influenced by climatic conditions, particularly the offshore and pelagic foraging seabirds. In the southern part of the Region, there have already been incidences of serious nesting failure of wedge-tailed shearwaters. These failures are closely correlated with reduced availability of their pelagic prey as a result of the El Niño Southern Oscillation cycles and higher than average increases in sea surface temperature,<sup>155</sup> as well as the interaction between the two.

Fourteen seabird species regularly breed at Raine Island in the Region's north.<sup>12</sup> Significant declines have been reported in the total number of seabirds breeding on the island. The estimated average annual breeding population for 1994–2003 was 70 per cent lower than for 1979–1993. There is no evidence the birds have moved to other breeding sites in the Great Barrier Reef;<sup>149</sup> however, the monitoring design has limited power to determine seasonal peaks in breeding which may contribute to variation in annual breeding survey results.<sup>159</sup> Declines were evident in 13 of the 16 species surveyed including the common noddy, brown booby and lesser frigatebird — the most common species in the area.<sup>149</sup> Between December 1995 and December 2000, a population decline of more than 40 per cent was reported for wedge-tailed shearwaters in the Capricorn cays.<sup>150</sup>

### 7.1.8 Targeted bony fish species

Only a small proportion of the Great Barrier Reef's bony fish species are targeted by fishers. There are concerns for some of these species. The vulnerability of two species of threadfin salmon has been assessed as high.<sup>229</sup> The vulnerability of grey mackerel has been assessed as medium.<sup>230</sup> The stock of snapper has been assessed as overfished.<sup>90</sup> There is evidence of localised depletions of coral trout species, with biomass in the no-take (green) zones being typically two to four times greater than in the fished (blue) zones.<sup>231</sup> The 2012 Queensland Fisheries Stock Status<sup>232</sup> summary reported that coral trout has moved from being considered 'sustainably fished' to 'uncertain' due to depressed catch rates, as well as triggered performance measures.

As with most species of conservation concern, the fundamental biology of some of these fishes, such as being long-lived and late maturing, makes them vulnerable to depletion. In addition, fishing activity, including illegal fishing in no-take zones, has contributed to declines of some targeted fish species. The importance of no-take zones is illustrated in recent research on larval fish connectivity between reefs, which demonstrated that fish productivity in this zone is contributing to the biomass of fish both within the zone and on reefs in surrounding areas.<sup>93</sup>

### 7.1.9 Sharks and rays

There are 136 known shark and ray species that inhabit the Region. Seven shark species and three species of ray are listed as either migratory or threatened species and are, therefore, matters of national environmental significance (see Chapter 4). On an ecosystem level, they are also important values relevant to the World Heritage Area, national heritage place and Marine Park matters of national environmental significance.

Some sharks and rays have more conservative life history traits than other fish, including a slow growth rate, low reproduction rate and late maturation, whereas others are fast growing, have high reproductive output and mature early. Some species display high levels of specificity in their habitat and prey,<sup>112,113,114,115</sup> making them susceptible to overfishing and slow to recover if overfished. Sharks and rays are captured (both as targeted and non-targeted catch) in a number of fisheries (commercial and recreational) within the Region. However, accurate identification of some species taken is still lacking, along with broad confidence in the validity of catch, release and survivorship data.<sup>233</sup> In addition, Queensland's Shark Control Program targets some sharks considered dangerous to bathers; however, the program also unintentionally catches a range of other non-dangerous shark species and many species of ray.<sup>222</sup> There is evidence of a decline in shark numbers and sizes caught under the program.<sup>234</sup> This decline could be a reflection of reduced population numbers rather than improved control methods.











Of the listed species, there have been significant range contractions and population declines for the largetooth (previously called the freshwater), green and dwarf sawfish. More concerning is the possibility that the speartooth shark, listed as critically endangered under the EPBC Act, has now become extinct on the east coast of Australia. The last verified specimen was recorded in 1983 from the Bizant River, which flows into Princess Charlotte Bay.<sup>121</sup> Whale sharks, shortfin makos, longfin makos and porbeagle sharks are pelagic species which are rarely seen within the Region, and there is limited information on their status and trends. Similarly, the white shark and grey nurse shark are temperate species which are also rarely sighted.





Of the remaining species, there is concern for the grey reef shark and whitetip reef shark,<sup>122</sup> and some other sharks such as the coastal Australian blacktip shark, hammerhead sharks and some sharks and rays that interact with the trawl fishery.<sup>112,117,119,122,235,236,237,238</sup> Shark-like batoids (for example, sawfishes, guitarfish and shovelnose rays) may be particularly vulnerable to nets set in inshore waters due to their body shape and preference for inshore habitats.<sup>239</sup>

## 7.2 Geomorphological features

Geomorphology refers to landforms and the processes that shape them. The interplay of all the coastal and marine geomorphic elements of the Region is a major component of the outstanding universal value of the World Heritage Area. The major geomorphological features of the Region are described in Chapter 4.

The current condition and trend of key geomorphological features is presented in Table 7.5.

Table 7.5 Current condition and trend of geomorphological features				
Understanding the table				
				
<b>Very good:</b> All major geomorphological features are structurally intact and there is no evidence of human impacts.		<b>Good:</b> There is minimal degradation or alteration to geomorphological features in some areas, but no substantial effects at a Reef-wide scale.		<b>Poor:</b> Degradation or alteration has occurred in a number of areas affecting the nature and integrity of the geomorphological feature.
<b>Very poor:</b> There is widespread degradation or alteration in geomorphological features at a Reef-wide scale.				
<b>Area</b> (See Chapter 1, Figure 1.1)		<b>Trend</b>		<b>Confidence in condition and trend</b>
<b>GBR</b> Great Barrier Reef Region		 Improving  Stable  Deteriorating  No clear trend		 Adequate high-quality evidence and high level of consensus  Limited evidence or limited consensus  Very limited evidence, assessment based on anecdotal information

	Condition and trend			
	Very good	Good	Poor	Very poor
<b>Overview:</b> Recent advances in mapping technology have provided unprecedented high resolution imagery of the underwater landscapes and geological features of the Great Barrier Reef. Glacio-hydroisostatic processes have resulted in several metres of sea level variation over the past 6500 years. However, features close to the populated coast are likely to have been impacted more due to increased human activity and development in this area. The effects of future climate change impacts on geological features are difficult to predict, but are most likely negative.				
<b>GBR</b>				



	Area	Condition and trend				Confidence	
		Very good	Good	Poor	Very poor	Condition	Trend
<p><b>Coral reefs:</b> Coral reefs are geomorphological features dependent upon the health of reef-building corals and the environmental conditions that promote their growth. Since 1986, average hard coral cover of the Great Barrier Reef ecosystem is estimated to have declined from 28 to 13.8 per cent, and the rate of decline is increasing. Without coral, the reef features become prone to erosion. Some corals are also showing signs of decreased calcification rates which has implications on the future of the structures. There is no long-term data on the condition of deeper submerged reefs but, for most, there is no indication of recent physical damage.</p> <p>References: <sup>4,35,36,37,9,38,39,40,41,42,43,44,45,46,47,48,64,240</sup></p>	GBR						
<p><b>Islands and shorelines:</b> There are about 1050 islands within the World Heritage Area, including continental islands, coral cays and mangrove islands. There is limited monitoring of the condition of most islands. Cyclones and coastal development have impacted the geomorphology of some islands and shorelines. Artificial barriers to flow have disrupted sediment supply to beaches, while increased fine sediments have resulted in mangrove forests replacing beaches.</p> <p>References: <sup>16,17,18,19,10,11,12,13,14,15,240</sup></p>	GBR						
<p><b>Channels and canyons:</b> Although little is known about submarine canyons, karstic channels and blue holes, their depth and distance from shore mean they are likely to be in very good condition and stable into the future. Palaeochannels can be found closer to the coastline and are, therefore, more at risk of human impacts such as sediment from catchment run-off. The impacts of climate change on these features are unknown.</p> <p>References: <sup>240</sup></p>	GBR						
<p><b>River deltas:</b> River deltas within and adjacent to the Region range in condition. They are in relatively good condition in areas north of Port Douglas where there has been relatively limited catchment development. In the central and southern areas, it is likely the geomorphology of river deltas has been affected by coastal development and changed sediment loads from catchment development and artificial barriers to flow. These effects can be positive or negative on the feature.</p> <p>References: <sup>240</sup></p>	GBR						
<p><b>Halimeda banks:</b> Large areas of the far northern Great Barrier Reef are dominated by <i>Halimeda</i>, calcareous green algae. It is likely these banks are in very good condition given their isolation from land-based impacts and level of protection from trawling.</p> <p>References: <sup>58,59,60,240</sup></p>	GBR						
<p><b>Seagrass meadows:</b> Intertidal seagrass meadows are in poor condition with serious declines reported over the past four years, especially those in the paths of cyclones and those exposed to flooding. Little is known about the implications of this loss on geomorphological processes performed by seagrass such as sediment trapping. The abundance and condition of subtidal and deepwater seagrass meadows are also poorly known. Fewer impacts mean northern area meadows are likely to be in very good condition and effectively stabilising and trapping sediment.</p> <p>References: <sup>26,27,28,29,30,31,32,33,240</sup></p>	GBR						

### 7.3 Indigenous and historic heritage values

The Region’s Indigenous and historic heritage values relevant to matters of national environmental significance are outlined in Chapter 4. Outcomes of an assessment of the condition and recent trend of key values are presented below.

They are grouped into:

- Table 7.6 — Indigenous heritage values (four elements)
- Table 7.7 — historic heritage values (six elements).

A discussion of the key findings about the condition and trend of some Indigenous and historic heritage values is presented after the tables.

This report considers Indigenous heritage values as a whole and recognises the interconnectedness of all elements, including the natural values described in Sections 7.1 and 7.2 above.

Table 7.6 Current condition and trend of heritage values — Indigenous			
Understanding the table			
<b>Very good:</b> Places with heritage values have been systematically and comprehensively identified and included in relevant inventories or reserves. Heritage places are in very good condition with identified values retaining a high degree of integrity.	<b>Good:</b> Places with heritage values have been systematically identified and included in relevant inventories or reserves. Heritage places are in good condition with identified values generally retaining their integrity.	<b>Poor:</b> Places with heritage values have not been systematically identified. Heritage places are in poor condition and/or their values lack integrity.	<b>Very poor:</b> Places with heritage values have not been identified. Heritage places are in degraded condition and their values lack integrity.
<b>Area</b> (See Chapter 1, Figure 1.1)	<b>Trend</b>		<b>Confidence in condition and trend</b>
<b>GBR</b> Great Barrier Reef Region	↑ Improving ↔ Stable ↓ Deteriorating — No clear trend		● Adequate high-quality evidence and high level of consensus ◐ Limited evidence or limited consensus ○ Very limited evidence, assessment based on anecdotal information

	Condition and trend			
	Very good	Good	Poor	Very poor
<b>Overview:</b> Traditional Owners with connections to the Great Barrier Reef Region maintain their cultural practices and customs. Places of Indigenous heritage values have not been systematically identified and many have deteriorated, especially around development areas and on islands. Some species of cultural significance are under pressure. Story, language and songlines are being affected by activities in the Region.				
<b>GBR</b>				

	Condition and trend				Confidence	
	Very good	Good	Poor	Very poor	Condition	Trend
<b>Cultural practices, observances, customs and lore:</b> Traditional Owners with connections to the Great Barrier Reef Region maintain their cultural practices and customs. There is evidence of inter-generational transfer throughout the Reef.						
<b>GBR</b>						

	Area	Condition and trend				Confidence	
		Very good	Good	Poor	Very poor	Condition	Trend
<b>Sacred sites, sites of particular significance, places important for cultural tradition:</b> There are many places, especially in coastal systems and on islands, where there is pressure on sacred sites and other sites of cultural significance. This is particularly around areas of high development and those exposed to severe weather events. Other sites are intact and in good condition and are being well managed by Traditional Owners.	GBR						
<b>Stories, songlines, totems and languages:</b> Story, language, songlines and totems are being affected by activities such as shipping, anchoring and dredging. These have a particularly significant impact as they may transgress these traditional systems. Some species of cultural significance, such as whales, dugongs, turtles, rays, sharks and dolphins, and other coastal resources, are under pressure, especially in areas south of Cooktown.	GBR						
<b>Indigenous structures, technology, tools and archaeology:</b> Indigenous structures (for example, fish traps), tools, technologies and archaeology have not been systematically identified. They are under pressure from coastal development and vehicle use. Some specific sites, such as the Hinchinbrook fish traps, are managed by Traditional Owners. The significance of some sites, such as the rock quarry on South Molle Island, is recognised on the National Heritage List.	GBR						

**Table 7.7 Current condition and trend of heritage values — historic heritage**

Understanding the table			
<b>Very good:</b> Places with heritage values have been systematically and comprehensively identified and included in relevant inventories or reserves. Heritage places are in very good condition with identified values retaining a high degree of integrity.	<b>Good:</b> Places with heritage values have been systematically identified and included in relevant inventories or reserves. Heritage places are in good condition with identified values generally retaining their integrity.	<b>Poor:</b> Places with heritage values have not been systematically identified. Heritage places are in poor condition and/or their values lack integrity.	<b>Very poor:</b> Places with heritage values have not been identified. Heritage places are in degraded condition and their values lack integrity.
<b>Area</b> (See Chapter 1, Figure 1.1)  <b>GBR</b> Great Barrier Reef Region	<b>Trends</b> Improving Stable Deteriorating No clear trend	<b>Confidence in condition and trend</b> Adequate high-quality evidence and high level of consensus Limited evidence or limited consensus Very limited evidence, assessment based on anecdotal information	

	Area	Condition and trend			
		Very good	Good	Poor	Very poor
<b>Overview:</b> There is good understanding and recording of some aspects of historic heritage; for example, historic shipwrecks and lightstations. Heritage values are being maintained or restored at some lightstations. Most places of historic significance are poorly recorded and their condition is not well understood.	GBR				

	Area	Condition and trend				Confidence	
		Very good	Good	Poor	Very poor	Condition	Trend
<b>Places of historic significance — historic shipwrecks:</b> There are more than 1300 known historic shipwrecks (i.e. those greater than 75 years) in the Region. Of these, only six are protected. Known wrecks have been systematically recorded as part of the Australian National Shipwrecks Database. However, the majority are poorly recorded or their locations remain unknown. <a href="#">Reference:</a> <sup>241</sup>	GBR						
<b>Places of historic significance — World War II features and sites:</b> The features and sites of significance in relation to World War II include shipwrecks, aircraft wrecks, unexploded ordnances and features on islands. Generally, they have been identified but little is known of their condition.	GBR						
<b>Places of historic significance — lightstations:</b> The locations and values of lightstations, including the lighthouses and ancillary structures, are well known. Heritage values are being maintained or restored at Commonwealth lightstations where there is a permanent presence, such as on Low Isles and Lady Elliot Island. However, certain sites have deteriorated due to poor construction materials.	GBR						
<b>Places of historic significance — other:</b> Other places of historic significance include Endeavour Reef, sites of turtle and dugong factories, and Mrs Watson's cottage on Lizard Island. Most are poorly recorded and their condition is not well understood. Some historic photos have been collected.	GBR						
<b>Places of scientific significance (research stations, expedition sites):</b> The Great Barrier Reef has long been an important site for scientific endeavours, from the Yonge expedition in 1928–29 to modern research stations (for example, Heron, Orpheus and Lizard islands). The location of significant research is generally well recorded in scientific publications. Some sites near research stations are maintained.	GBR						
<b>Places of social significance — iconic sites:</b> The iconic sites of the Great Barrier Reef include Raine Island, Cod Hole, Michaelmas Cay and reef, Green Island and reef, Whitehaven Beach and Hill Inlet. They are well recognised and many are comprehensively managed. As with the overall condition of the Great Barrier Reef, the condition of some iconic sites has declined.	GBR						

### 7.3.1 Traditional Owner connections

Traditional Owner connections integrate nature, heritage and culture, and Indigenous heritage is made up of tangible and intangible elements. Landscapes, seascapes, sites and areas of their country are particularly important to Aboriginal and Torres Strait Islander peoples, intertwined with the more intangible elements such as cultural practices, observances, customs, lore, stories, songlines, totems and languages (see Chapter 4). There are a variety of cultural sites within the sea country of the Great Barrier Reef, including sacred sites, ceremonial sites, burial grounds, rock art sites, middens, fish traps, cultural landscapes and story places. These significant places provide a strong connection to traditional clan areas and form part of Australia's rich heritage.

For Traditional Owners, nature and culture combine to make a living heritage. They are increasingly re-asserting their role in managing their country through active engagement in on-country management and in policy and planning programs.

*"...we've had a long long long association with the Reef. It is one of the seven wonders of the world but we also have a common culture and obligation to it."* <sup>242</sup>

### 7.3.2 Cultural keystone species

As with most cultures, the environment is central to Indigenous heritage values, and includes some plants and animals of particular significance. These plants and animals are referred to as 'cultural keystone species' and they play a fundamental role in Traditional Owner culture, including through diet, materials, medicine, totems and stories. <sup>243</sup> In the Region, populations of many of these cultural keystone species have been significantly reduced — examples include dugongs, green turtles, some sharks and some bony fish (Section 7.1). This, in turn, affects the Region's Indigenous heritage values. For example, the declines in fish, turtle and dugong numbers in the



inshore coastal zone have affected the traditional practice of building platforms out into the sea to hunt and fish for these species.

When Traditional Owner Marie Shipton was asked about seeing marine turtle nesting along her beaches she replied “...no we don't anymore... we used to have a lot of turtle and dugong but it's very few now, but we don't know where they're gone..”<sup>244</sup>

Through their own practices and partnerships with managing agencies, Traditional Owners are working to look after these keystone species. One example is the Traditional Use of Marine Resources Agreements between Traditional Owner groups, the Great Barrier Reef Marine Park Authority (the Authority) and the Queensland Government.

#### **Saltwater people managing 'Gungu' (green turtles) in their sea country**

Ecological monitoring and information exchange between Traditional Owners, scientists and Marine Park managers have helped conserve 'Gungu' (green turtles) in Gudjada sea country around Bowen. The on-ground conservation project undertook a preliminary investigation of the incidence and prevalence of fibropapilloma disease in green turtles in Edgecombe Bay, as well as tagging and monitoring of the green turtle population in Gudjada sea country. The project was a partnership between the Gudjada Reference Group, Girringun Aboriginal Corporation, the Authority, the Queensland Government, WWF-Australia and James Cook University scientists. It provided the opportunity for Traditional Owners to share their cultural lore, custom and Traditional ecological knowledge. It also enabled scientists to share western science and wildlife monitoring techniques with Indigenous communities, building Traditional Owner capacity in monitoring and management of their sea country.

### **7.3.3 Identifying Indigenous heritage values**

One aspect of the statements used to grade the condition of heritage values is the degree to which those values have been recorded and identified. Based on this aspect, many of the Indigenous heritage values of the Great Barrier Reef Region are graded as being in poor condition. There is no systematic and comprehensive inventory of the Region's tangible or intangible Indigenous heritage values. While Traditional Owners have a good understanding of their own country and its values, this has rarely been integrated into the broader information systems of managing agencies. The Authority's website *Story Place* is a reference database that shares information and knowledge about traditional connections to sea.<sup>245</sup>

#### **Girringun Traditional Owners record their important cultural heritage sites**

As part of the implementation plan for their Traditional Use of Marine Resources Agreement, Girringun Traditional Owners have conducted comprehensive cultural heritage surveys to inform management planning. This involves documenting previously unrecorded cultural heritage sites, providing recommendations on their management and protection, and providing baseline information about the natural and cultural landscape so that, for example, damage from cyclones Yasi and Larry may be assessed. Participation of Girringun rangers was fundamental to the activity. It was an opportunity to provide training in cultural and heritage management and to further empower the Traditional Owners of the area.

### **7.3.4 Historic heritage values**

Historic heritage values illustrate the way in which the many cultures of Australian people have modified, shaped and created the cultural environment of the Region. They are assessed as being generally in good condition in the Region, although some historic lightstations and some of the Region's iconic sites are likely to be deteriorating.

While some specific aspects of the Reef's historic heritage have been well documented (for example, shipwrecks and lightstations), knowledge of many historic places or events is limited (for example, World War II features and sites). Even for values that are relatively well recorded, understanding about their full extent in the Region continues to improve as technology improves and visitation to the Region increases. For example, new shipwrecks continue to be found and recent investigations indicate there may be between 160 and 200 aircraft wrecks from World War II within the outer boundaries of the Region.

## **7.4 Community benefits of the environment**

The condition and trend of the Region's community benefits are determined, in large part, by the quality of the environment (its biodiversity, geomorphological features and Indigenous and historic heritage values) from which the benefits are derived. Outcomes of an assessment of the condition and recent trend of community benefits are presented in Table 7.8.

A discussion of the key findings follows the table.

**Table 7.8 Current condition and trend of community benefits of the environment**

Understanding the table			
<b>Very good:</b> There is a significant contribution to the wellbeing of local communities and the nation. The Region contributes strongly to regional and national economies, and is highly valued, well understood, and enjoyed by catchment residents, the nation and the world community.	<b>Good:</b> There is a valuable contribution to the wellbeing of local communities and the nation. The Region contributes to regional and national economies, and is valued, understood and enjoyed by catchment residents, the nation and the world community.	<b>Poor:</b> There is a small contribution to wellbeing of local communities and the nation. The Region makes a minor contribution to regional and national economies and many do not understand or enjoy its values.	<b>Very poor:</b> There is little or no contribution to the wellbeing of local communities or the nation. The Region contributes very little to regional and national economies and most do not understand or enjoy its values.
<b>Area</b> (See Chapter 1, Figure 1.1)  <b>GBR</b> Great Barrier Reef Region	<b>Trend</b> ↑ Improving ↔ Stable ↓ Deteriorating — No clear trend	<b>Confidence in condition and trend</b> ● Adequate high-quality evidence and high level of consensus ◐ Limited evidence or limited consensus ○ Very limited evidence, assessment based on anecdotal information	

	Area	Condition and trend			
		Very good	Good	Poor	Very poor
<b>Overview:</b> The Region's environment provides a range of benefits to catchment communities and the nation. Many catchment residents understand and appreciate the Reef and have a strong personal attachment to it. Much of the natural beauty of the Reef remains, but underwater aesthetic values have declined in some areas. Reef-dependent industries contribute strongly to the Australian economy. In recent years, some community benefits have been adversely affected by the direct and indirect effects of extreme weather.	GBR	<div></div>	<div>↔</div>	<div></div>	<div></div>

	Area	Condition and trend				Confidence	
		Very good	Good	Poor	Very poor	Condition	Trend
<p><b>Income:</b> Much of the economy derived from the Great Barrier Reef is based upon use of its biological resources. The economic contribution generated by tourism, recreation, commercial fishing and scientific research in the Reef catchment and the World Heritage Area in 2012 was about \$5.7 billion. This has been relatively stable over the past five years.</p> <p><b>References:</b> <sup>246,247</sup></p>	GBR	<div></div>	<div>↔</div>	<div></div>	<div></div>	<div>●</div>	<div>●</div>
<p><b>Employment:</b> Tourism, recreation and commercial fishing based on the Great Barrier Reef environment generated almost 69,000 full-time equivalent jobs throughout Australia in 2012, compared to 54,000 in 2007. Scientific research conducted on the Reef generated more than 900 jobs.</p> <p><b>References:</b> <sup>247</sup></p>	GBR	<div></div>	<div>↔</div>	<div></div>	<div></div>	<div>●</div>	<div>●</div>

	Condition and trend				Confidence	
	Very good	Good	Poor	Very poor	Condition	Trend
<p><b>Understanding and appreciation:</b> The Authority, the tourism industry and Reef HQ Aquarium provide information on the Reef, its key issues and actions that can be taken. Students at 293 Reef Guardian schools are receiving key messages about the Reef's protection and actions they can take. Many new residents within the Region may have little knowledge of the Reef and its management.  <a href="#">References:</a> <sup>246</sup></p>	GBR					
<p><b>Enjoyment:</b> Opportunities to experience clear oceans, healthy coral reefs and healthy fish are important to Reef visitors. Visitors generally report that they are satisfied or very satisfied with their visit to the Great Barrier Reef.  <a href="#">References:</a> <sup>246,248</sup></p>	GBR					
<p><b>Access to Reef resources:</b> Millions of people visit the Region each year. It provides a wide range of recreational opportunities such as boating, snorkelling, diving, fishing and nature appreciation. There are also opportunities for commercial fishing, marine tourism and education. In some key locations, management arrangements partition use to allow for a range of activities.  <a href="#">References:</a> <sup>246,249,250</sup></p>	GBR					
<p><b>Personal connection:</b> More than 80 per cent of Reef catchment residents have visited the Great Barrier Reef at least once in their lives, and almost 10 per cent visit the Reef more than 10 times a year. Some residents report that they chose where they live so as to be close to the Reef. Many report a close attachment to particular sites in the Region. Traditional Owners continue to maintain connection to their sea country; for example, through stories and songlines, sites of cultural significance and important saltwater ceremonies. Many people who are employed in the Region show a strong personal attachment to its environment.  <a href="#">References:</a> <sup>246</sup></p>	GBR					
<p><b>Health benefits:</b> Catchment residents often report that they derive health benefits from the Great Barrier Reef environment and its resources. Examples include being able to 'unwind' and 'get away from it all', the release of stress, improved fitness through outdoor activities, and improved diet through access to wild-caught seafood.  <a href="#">References:</a> <sup>246</sup></p>	GBR					
<p><b>Aesthetics:</b> The natural beauty of most of the Region remains intact, especially for offshore coral reefs and aerial vistas, as well as for neighbouring islands (many of which are national parks). The significant loss of coral cover, especially in southern inshore areas, has reduced underwater aesthetic value.  <a href="#">References:</a> <sup>251</sup></p>	GBR					

### 7.4.1 Economic benefit

The Great Barrier Reef provides long-term economic benefits and employment to Queensland coastal communities and the nation. These benefits are based upon use of its biological resources. This includes marine tourism and recreation, which are nature-based and heavily reliant on a healthy and diverse ecosystem; and commercial fishing, which is almost totally dependent on natural ecological systems for productivity and sustainable profit. Exploration and the recovery of minerals or petroleum are banned in the Marine Park.

The Australia-wide economic contribution generated by tourism, recreation, commercial fishing and scientific research in the Reef catchment and the World Heritage Area in 2012 was \$5.7 billion.<sup>247</sup> This supported the equivalent of about 69,000 full-time jobs. These estimates are based on both the direct and indirect contributions of the activities considered, and are driven by just over \$7 billion of expenditure in the catchment (Table 7.9).<sup>247</sup>

Tourism in the catchment and the World Heritage Area represents more than 90 per cent of the direct expenditure of Reef-related activities (Table 7.9).<sup>247</sup> This then flows through the economy, with tourism accounting for 91 per cent and 93 per cent of the value-added and employment contributions to Australia, respectively. Although the Reef provides an impetus for travel to the broader catchment, not all visitors to the catchment visit the Great

Barrier Reef. Analysis of more Reef-specific information indicates direct Reef-related expenditure in 2012 was about \$480 million, which contributed almost \$389 million (value-added) to Australia's economy and generated employment equivalent to more than 4800 full-time jobs. This is based on 1.9 million direct visits to the Marine Park annually and excludes the 2.3 million passengers who transfer through the Marine Park to adjacent islands.<sup>247</sup>

Importantly, the Reef is considered a major driver or incentive for international travellers to visit Australia.

Recreation, which covers household recreational activity by those who live in the catchment area, contributed just over 2700 full-time equivalent jobs. Recreational activity covers trip-related expenditure for fishing, boating, sailing and visiting islands, as well as household expenditure on recreational equipment.

Commercial fishing contributed \$160.3 million and provided the equivalent of 975 full-time jobs in 2012.

**Table 7.9 Economic contributions of Reef-related activities to Australia, 2012** <sup>247</sup>

The economic contribution of tourism, recreation, commercial fishing and scientific research in the Great Barrier Reef catchment and the World Heritage Area are provided. Direct expenditure is the total amount of money (in millions of dollars) spent within each industry. Value-added figures are estimates of the profit (in millions of dollars) that each industry generated during 2012. Full-time equivalent is the equivalent of one employee working full-time.

	Direct expenditure (\$m)	Value-added (\$m)	Employment (full-time equivalent)
Tourism	6410.6	5175.6	64,338
Recreation	332.4	243.9	2785
Commercial fishing	192.5	160.3	975
Scientific research	106.1	98.0	881
<b>Total</b>	<b>7,041.5</b>	<b>5,677.8</b>	<b>68,979</b>

#### 7.4.2 Visitation

There are now more than 1.8 million tourism visitor days spent on the Great Barrier Reef each year (see Figure 5.14 in Chapter 5). The number of tourists visiting the Reef fell between 2006 and 2011 after a peak of almost two million visitor days in 2005; however, there was evidence of growth in 2012. The recent declines can largely be attributed to international financial instability, the strong Australian dollar, extreme weather events, particularly in 2011, and growing competition from alternative international tourism destinations.

In addition to the tourists directly visiting locations within the Marine Park, there are an estimated 2.3 million transfers of visitors, residents and workers to and from islands within the Region.

In the stakeholder workshops held as part of the strategic assessment process (see Appendix 5), participants highlighted favourite Great Barrier Reef places including islands, coral reefs, beaches, estuaries, bays and inlets. There is a spectrum of such locations throughout the Region, supporting recreational activities such as boating, snorkelling, diving, fishing, wildlife watching, nature appreciation, relaxing, walking, camping and generally spending time with family and friends, as well as for commercial fishing, marine tourism and education.

It is estimated there are more than 14 million recreational visits to the Marine Park by Great Barrier Reef catchment residents every year, plus visits by non-paying and independent travellers from outside the catchment.<sup>252</sup> In 2011 and 2012, 87 per cent of coastal residents visited a beach; 73 per cent visited a second beach; 30 per cent visited an island; and 42 per cent went either fishing, boating or sailing in the Great Barrier Reef.<sup>249</sup> Across the Reef catchment, nine per cent of people own a boat. Rates of ownership are higher in coastal areas, with almost 17 per cent of Whitsundays residents owning a boat (see Section 5.4.5).<sup>253</sup>

Results of a 2012 survey show 75 per cent of tourists to coastal parts of the Great Barrier Reef catchment went to a mainland beach, 60 per cent visited an island and 50 per cent went to an offshore reef. Their activities included snorkelling or diving (50 per cent); sailing (25 per cent); private boating (20 per cent); and fishing (20 per cent). These visitors highly valued the opportunities to experience clear oceans, healthy coral reefs and healthy reef fish, and were mostly highly satisfied with these aspects of their visit.<sup>254</sup>

#### 7.4.3 Understanding and appreciation

There are many opportunities for coastal residents and visitors to learn about and help protect the Great Barrier Reef.

A key component of many tourism operations in the Region is presenting and interpreting the Reef environment to their visitors. This is especially the case for the more than 60 per cent of visitors who travel with certified high standard tourism operations,<sup>255</sup> as high standards of presentation and interpretation are key commitments in the certification process.

The Authority's Reef HQ Aquarium, located in Townsville, also provides people of all ages and physical abilities the chance to see and gain an appreciation for a living coral reef ecosystem, what makes it special and why it



needs to be protected. More than 3.4 million local, national and international visitors have visited the facility since it opened in 1987, and visitation continues to increase. In 2012, 94 per cent of respondents to Reef HQ Aquarium's visitor survey believed they had gained an improved understanding of the issues relating to the Great Barrier Reef, while 89 per cent believed they had a better understanding of how they can help protect and conserve the ecosystem.<sup>256</sup> The facility's reef videoconferencing program has also showcased the living reef to more than 680 locations nationally and internationally, directly engaging more than 52,000 people.

Through the Authority's Reef Guardian stewardship program, local stakeholders are encouraged to take hands-on actions to care for the Reef. Participants are encouraged to go beyond what is required by law in their day-to-day activities and become active stewards for the Reef. This includes sharing information about their actions. There are now 293 Reef Guardian schools, involving about 114,900 students working on Reef-related projects. The program was expanded in 2007 to include local councils and, in the past two years, pilot programs were introduced for farmers, graziers and commercial fishers.

Participants in the Authority's Eye on the Reef monitoring program contribute substantially to understanding trends in Reef condition, contributing more than 10,000 surveys from about 500 reefs throughout the Region.

The extent to which people care about the Great Barrier Reef is evident in responses to a 2012 survey where visitors indicated they were more sensitive to changes in the Reef environment — such as oil spills, ship groundings, waste spills from ports, water clarity and rubbish — than to increases in local prices. When asked if they would be willing to pay to help 'fix' various threats to the Reef, visitors indicated they would be prepared to pay \$10 (median response) for each visit to improve water quality. They were also willing to pay \$5 (median response) to protect top predators or reduce the risk of shipping accidents.<sup>254</sup>

#### **7.4.4 Personal connections**

The Great Barrier Reef is a key part of the identity of adjacent coastal communities. It is a major 'point of difference' for these communities and something to be proud of. Traditional Owners are strongly linked to their sea country through their living culture and traditions, including their stories and songlines, sites of cultural significance and important saltwater ceremonies.

Most residents within adjacent coastal communities have visited the Reef environment, and many maintain a close connection, with 10 per cent going to the Reef more than 10 times in a year.<sup>257</sup> Some residents choose to live at their current locations so they can enjoy the benefits of being close to the Great Barrier Reef.<sup>258</sup>

Reef-dependent commercial fishers and tourism operators identify very strongly with their occupations and the places where they live and work. This is highlighted by the fact that few, if any, who were directly affected by cyclone Yasi or the central Queensland floods in 2011, changed their jobs or moved elsewhere after the extreme weather events. This was despite economic imperatives to find alternative income or maintain their wage.<sup>259</sup>

#### **7.4.5 Knowledge about community benefits**

It is widely recognised that communities of the Great Barrier Reef catchment and beyond derive a range of significant benefits from the Region's environment. Supporting evidence specific to the Region is patchy; however, studies from elsewhere show parks and protected areas contribute to the health and wellbeing of communities. They are places of adventure and challenge, exercise, peace and quiet, recreation, gathering and relaxation, and offer a range of recreational activities to bring families and friends together.<sup>260</sup>

It is anticipated that the Great Barrier Reef's role in supporting and sustaining local communities will be better understood in the future when a social and economic long-term monitoring program, designed by CSIRO and James Cook University researchers, is implemented.

The ecosystem services that the Region provides to communities (for example, the role of wetlands, mangroves and seagrass in trapping sediments and therefore maintaining clear water on the Reef) are poorly understood.

### **7.5 Environmental processes**

The key environmental processes relevant to matters of national environmental significance are outlined in Chapter 4. Outcomes of an assessment of the condition and recent trend of these environmental processes are presented in Table 7.10.

It is important to note that while there is a good understanding of the variables that contribute to some of the key physical and chemical processes — such as sedimentation, sea temperature, nutrient cycling and freshwater inflow — there is a poor understanding and almost no monitoring of many others, especially ecological processes.

Table 7.10 Current condition and trend of environmental processes

Understanding the table			
Very good	Good	Poor	Very poor
<b>Very good:</b> There are no significant changes in process as a result of human activities.	<b>Good:</b> There are some significant changes in processes as a result of human activities in some areas, but these are not to the extent that they are significantly affecting ecosystem functions.	<b>Poor:</b> There are substantial changes in processes as a result of human activities, and these are significantly affecting ecosystem functions in some areas.	<b>Very poor:</b> There are substantial changes in processes across a wide area as a result of human activities, and ecosystem functions are seriously affected in much of the area.
Area (See Chapter 2, Figure 2.3)		Trends	Confidence in condition and trend
<b>N.I.</b> Northern inshore		↑ Improving	● Adequate high-quality evidence and high level of consensus
<b>N.O.</b> Northern offshore		↔ Stable	◐ Limited evidence or limited consensus
<b>S.I.</b> Southern inshore		↓ Deteriorating	○ Very limited evidence, assessment based on anecdotal information
<b>S.O.</b> Southern offshore		— No clear trend	

	Area	Condition and trend			
		Very good	Good	Poor	Very poor
<b>Overview:</b> A recent cluster of severe weather events (cyclones and floods), especially in the southern area, has adversely affected the Region's physical processes. Sea temperature and sea level are rising throughout the Region. The acidity of the ocean continues to slowly increase. The processes of sedimentation, including resuspension, and nutrient cycling continue to be affected in inshore areas on the developed coast. Ecological processes remain largely intact in northern areas and any changes are not significantly affecting the ecosystems. Primary production, symbiosis, reef-building and connectivity are in poor condition in the southern inshore area and are deteriorating.	N.I.		↔		
	N.O.		↔		
	S.I.			↓	
	S.O.		↔		

	Area	Condition and trend				Confidence	
		Very good	Good	Poor	Very poor	Condition	Trend
<b>Waves, currents and tides:</b> Ocean currents vary naturally. There is increasing evidence of intensified flow and accelerated warming in the East Australian Current adjacent to the Region's southern coast. This current is transporting greater volumes of warmer water southward, carrying larvae and juveniles with it. There is little information about the Hiri Current travelling north along the coast in northern Great Barrier Reef waters. Severe cyclone activity over the past four years has caused large waves which have damaged benthic habitats. There have been no significant changes to tides within the Region. <b>References:</b> <sup>261,262,263</sup>	N.I.	—				○	○
	N.O.	—				○	○
	S.I.		—			◐	○
	S.O.		—			◐	○
<b>Cyclones:</b> Between 2008 and 2012, there was a cluster of category three and above cyclones that affected the Great Barrier Reef. Impacts have been most severe in the southern half of the Region, causing significant damage to benthic habitats. Cyclones are forecast to become more intense. <b>References:</b> <sup>139,153,264,265,266,267,268,269,270,271,272</sup>	N.I.		↔			●	●
	N.O.		↔			●	●
	S.I.			↓		●	●
	S.O.			↓		●	●

	Area	Condition and trend				Confidence	
		Very good	Good	Poor	Very poor	Condition	Trend
<b>Wind:</b> There is emerging evidence of increases in wind strength Australia-wide, but little information specific to the Region. Changes in the wind patterns may have consequences for inshore ocean turbidity through the resuspension of sediments, for island formation and for the distribution of planktonic larvae. <a href="#">References:</a> <sup>273</sup>	N.I.		↓			○	○
	N.O.		↓			○	○
	S.I.		↓			○	○
	S.O.		↓			○	○
<b>Sedimentation:</b> Exposure of the Great Barrier Reef to terrestrial sediments and resuspended marine sediments has increased, especially in southern inshore areas. Significant investment in land-based management measures has resulted in a six per cent reduction in the sediment load delivered to the Great Barrier Reef since 2009. Dumping of dredge material also affects sedimentation processes with resuspended sediments potentially travelling considerably further than previously thought. There may also be more resuspension in shallow water due to increases in wind in the Region. <a href="#">References:</a> <sup>274,275,276</sup>	N.I.		↔			●	●
	N.O.	—				●	●
	S.I.			↓		●	●
	S.O.		↓			●	●
<b>Sea level:</b> The fastest rates of sea level rise in Australian waters are being recorded in northern areas, with sea level in the Region continuing to rise by about 3.1 mm per year. Even modest rises in sea level may have substantial consequences when combined with natural variability arising from the El Niño Southern Oscillation and the Pacific Decadal Oscillation, for example to turtle nesting success. <a href="#">References:</a> <sup>272,277,278,279,280</sup>	N.I.		↓			●	●
	N.O.		↓			●	●
	S.I.		↓			●	●
	S.O.		↓			●	●
<b>Sea temperature:</b> Average sea surface temperature has risen substantially over the last century, with 15 of the 20 warmest years occurring in the past two decades. This has caused some coral bleaching plus observable effects on growth rates of corals and overall coral cover. Increasing sea temperature provides additional energy to the formation of tropical cyclones. <a href="#">References:</a> <sup>272,281,282,283</sup>	N.I.			↓		●	●
	N.O.			↓		●	●
	S.I.			↓		●	●
	S.O.			↓		●	●
<b>Light:</b> Increased sedimentation may be altering light levels in inshore areas of the southern two-thirds of the Region. On a local scale, there is increasing light pollution from coastal facilities, potentially affecting any turtle nesting and the behaviour of hatchlings in the vicinity. <a href="#">References:</a> <sup>284</sup>	N.I.	↔				●	●
	N.O.	↔				●	●
	S.I.			↓		●	●
	S.O.		↔			●	●
<b>Nutrient cycling:</b> Most inshore areas of the southern two-thirds of the Region are exposed to nutrients, which continue to enter the Great Barrier Reef at greatly enhanced levels. Recent significant investment in land management practices has reduced the nutrient load entering the Region from the adjacent catchment. For example, since 2009 improvements in farming practices has resulted in a seven per cent reduction in the load of total nitrogen delivered to the Great Barrier Reef. <a href="#">References:</a> <sup>285,286,287</sup>	N.I.		↓			●	●
	N.O.		↓			●	●
	S.I.			↓		●	●
	S.O.		↓			●	●
<b>Ocean acidity:</b> A decline of 0.1 units in the pH of Great Barrier Reef waters has already been recorded and further significant declines are expected. Rates of decline are greater in southern areas of the Region. The most immediate effects may be on crustose coralline algae, a species vital to reef building. <a href="#">References:</a> <sup>64,288,289,290</sup>	N.I.		↓			●	●
	N.O.		↓			●	●
	S.I.		↓			●	●
	S.O.		↓			●	●
<b>Freshwater inflow and salinity:</b> After a period of significantly lower than average freshwater flow, significant volumes of freshwater have entered the Region in the past five years, including record flows for some rivers. This freshwater has reached beyond the coastal zone, and has had direct effects on marine species as well as delivering increased loads of sediments, nutrients and pesticides. The salinity of Great Barrier Reef waters is generally stable. <a href="#">References:</a> <sup>291</sup>	N.I.		↓			●	●
	N.O.	↔				●	●
	S.I.			↓		●	●
	S.O.	↔				●	●

	Area	Condition and trend				Confidence	
		Very good	Good	Poor	Very poor	Condition	Trend
<b>Microbial processes:</b> Many marine microbes are sensitive to changes in temperature, ocean acidity, nutrients, salinity and oxygen and respond rapidly to changing environmental conditions. There is little specific information available for the Great Barrier Reef, but it is assumed the condition of microbial processes is a reflection of other ecosystem processes. In recent decades, there has been a global increase in reports of disease in marine organisms, linked to increasing temperatures and thermal stress. <b>References:</b> 92,292,293,294,295,296	N.I.	↔				○	○
	N.O.	↔				○	○
	S.I.		↔			○	○
	S.O.	↔				○	○
<b>Particle feeding:</b> Particle feeding is undertaken by a range of organisms including corals, sponges, crustaceans and holothurians. The decline of coral cover in the southern inshore area is likely to have had a negative impact on this process. Some species of prawns are targeted by the trawl fishery. <b>References:</b> 49,297,298,299,300	N.I.		↔			○	○
	N.O.		↔			○	○
	S.I.		↓			○	○
	S.O.		↓			○	○
<b>Primary production — pelagic:</b> The presence of elevated chlorophyll a, together with intense and extensive phytoplankton blooms following the discharge of nutrient-rich flood waters, suggests pelagic primary production in the southern inshore area is likely to be significantly impacted by elevated nutrient loads. This in turn affects zooplankton populations, such as crown-of-thorns starfish larvae. <b>References:</b> 63,301,302,303	N.I.		↔			●	●
	N.O.		↔			●	●
	S.I.			↓		●	●
	S.O.		↔			●	●
<b>Primary production — benthic:</b> The overall condition of benthic primary production in the Region remains generally good. However, certain primary producers, such as coral and seagrasses, are in decline and have been for many decades, resulting in a loss of primary production and a deterioration of habitat, especially in southern areas. There is evidence of increased primary production by macroalgae, indicating a phase shift from coral to macroalgae domination at some reefs. <b>References:</b> 304,305,306	N.I.		↔			●	●
	N.O.		↔			●	●
	S.I.			↔		●	●
	S.O.			↔		●	●
<b>Herbivory:</b> Populations of herbivorous fish are generally healthy and not under pressure; however, the largest herbivore species, the dugong, has declined significantly in the southern inshore area. The herbivorous green turtle population has increased in the south; however, there are indications that the northern stock is in decline. <b>References:</b> 80,307,308,309,310,311,312,313	N.I.		↔			●	●
	N.O.		↔			●	●
	S.I.			↓		●	●
	S.O.		↔			●	●
<b>Predation:</b> While little is known about trends in the ecological process of predation, the condition of the ecosystem's top predators can provide an indication of its condition. There are 115 species of top predators identified within the Region (for example, sharks, mackerel and coral trout). While several species are declining, threatened or over exploited, the majority are either caught in sustainable numbers or not harvested. <b>References:</b> 314,119,120,235,315	N.I.		↔			○	○
	N.O.		↔			○	○
	S.I.			—		○	○
	S.O.		↔			○	○
<b>Symbiosis:</b> There is a massive number and wide range of symbiotic relationships in the Region including those that are mutually beneficial (mutualism); beneficial to one organism without affecting the other (commensalism); and beneficial to one organism to the detriment of another (parasitism). Very little is known about most; however, based on the overall condition of the ecosystem, it is likely they are in good condition. The most significant symbiosis is between coral and algae, providing coral with more than 90 per cent of its energy. The decline in coral cover will have had a significant effect on this symbiosis which is also known to break down during thermal stress (coral bleaching). Areas affected by poor water quality may also experience higher parasite numbers. <b>References:</b> 316,317,318,319,320,321,322,323,324,325,326,327,328,329	N.I.		↔			○	○
	N.O.		↔			○	○
	S.I.			↓		○	○
	S.O.			↓		○	○



	Area	Condition and trend				Confidence	
		Very good	Good	Poor	Very poor	Condition	Trend
<b>Competition:</b> There is little information about the multitude of competitive interactions in the Great Barrier Reef ecosystem, but it is likely the majority are in good condition. Coral reefs are highly competitive ecosystems; particularly significant is the competitive balance between coral and algae. A phase shift from coral to macroalgae domination has been recorded at some reefs. <b>References:</b> 330,331,332,333,304,73,78	N.I.		↔			○	○
	N.O.		↔			○	○
	S.I.			↓		◐	◐
	S.O.		↔			○	○
<b>Connectivity:</b> Connectivity within the ecosystem has not been comprehensively studied, but marine species and habitats are thought to remain well connected. However, it is likely fragmentation in seabed habitats has reduced connectivity over the lagoon floor. There is reduced aquatic connectivity between marine and freshwater habitats which is affecting dependent species. Continued connectivity between no-take zones and other marine areas is improving overall species abundance. There remains very good connectivity with other areas visited by migratory species; for example, Torres Strait, Coral Sea and Antarctica. <b>References:</b> 48,93,334,335,336,337,338	N.I.	↔				○	○
	N.O.	↔				○	○
	S.I.			↓		○	○
	S.O.		↔			○	○
<b>Recruitment:</b> Recruitment is likely to be in good condition for most of the Region. The apparent lack of recovery of many severely degraded, southern inshore reefs is likely to be partly due to poor recruitment and low juvenile survival. There is also likely to have been poor recruitment in southern inshore seagrass meadows. There is deteriorating recruitment offshore for seabirds and some important green turtle nesting sites. <b>References:</b> 72,288,339,340,341	N.I.		↔			◐	◐
	N.O.		↔			◐	◐
	S.I.			↓		◐	◐
	S.O.		↔			◐	◐
<b>Reef building:</b> Reef building is primarily driven by the formation of calcium carbonate skeletons. How fast a reef is built is the net effect of recruitment, calcification and erosion. Reef building is likely to be in good condition for much of the Region, but has been affected by cyclones and reduced coral cover, especially in the southern area. Increasing sea temperatures and ocean acidification are likely to be reducing calcification rates. <b>References:</b> 64,342,343,344,345,346,347	N.I.		↓			◐	◐
	N.O.		↓			◐	◐
	S.I.			↓		◐	◐
	S.O.			↓		◐	◐

## 7.6 Condition and trend of matters of national environmental significance

### 7.6.1 World heritage properties

The current condition and trend of the outstanding universal value of the Great Barrier Reef World Heritage Area is benchmarked against the retrospective Statement of Outstanding Universal Value as a way of describing changes in the property's condition since inscription (Table 7.11).

Excerpts from the statement relating to each of the four relevant world heritage criteria have been assessed (Table 7.11 (a) to (d)). The grading statements are based on those used by the International Union for Conservation of Nature to assess natural world heritage sites. The title of each grade has been adapted to align with those used in other parts of the strategic assessment. Each assessment is a 'grade of best fit' for the whole property and all matters relating to the excerpt. An overall trend in condition is provided, comparing current condition with the likely condition at the time of inscription. Confidence in the assessment is also indicated.

Like most of the attributes relevant to the matters of national environmental significance, outstanding universal value is distributed across the whole of the Region rather than being found at discrete locations. It is therefore assessed across the entire Region.

An assessment of the property's integrity is presented in Table 7.12. This benchmarking is based on Section II.E of the *Operational Guidelines for the Implementation of the World Heritage Convention*<sup>348</sup> which sets out the requirements for integrity, namely: the extent to which the property includes all elements necessary to express its outstanding universal value; that it is of adequate size to ensure the complete representation of the features and processes which convey the property's significance; and protection from the adverse effects of development and/or neglect.

These two assessments show the outstanding universal value of the World Heritage Area remains largely intact. Many elements that make up the outstanding universal value remain in good condition. The Region remains a

globally outstanding example of an ecosystem that has evolved over the millennia, and almost all geomorphological processes remain in good condition. Examples of all stages of reef development remain, although the overall health of reefs, especially in the southern two-thirds of the Region, has declined significantly. While the condition of some elements, such as humpback whales, has improved others have experienced serious declines since 1981. These include inshore coral reefs in southern areas, southern dugong populations, some species of marine turtle, and offshore and pelagic foraging seabirds. The natural beauty of most of the Region remains intact, however its underwater aesthetic value has declined in southern inshore areas. This, in turn, affects the image and the human experiences of the Reef (for example, Indigenous culture, tourism and recreation). The property continues to meet the requirements for integrity, meaning its natural attributes are whole and intact.

It is important to note that, while most of the excerpts assessed are rated as currently being in very good or good condition, the trend for about two-thirds of these is assessed as deteriorating.

**Table 7.11 Benchmarking the outstanding universal value of the Great Barrier Reef World Heritage Area**

The assessments presented in this table are based on the same information used to assess current condition and trend in Sections 7.1 to 7.5. The references listed there also apply to this assessment.

Understanding the table			
<b>Very good:</b> All elements necessary to maintain the outstanding universal value are essentially intact, and their overall condition is stable or improving. Available evidence indicates only minor, if any, disturbance to this element of outstanding universal value.	<b>Good:</b> Some loss or alteration of the elements necessary to maintain the outstanding universal value has occurred, but their overall condition is not causing persistent or substantial effects on this element of outstanding universal value.	<b>Poor:</b> Loss or alteration of many elements necessary to maintain outstanding universal value has occurred, which is leading to a significant reduction in this element of the outstanding universal value.	<b>Very poor:</b> Loss or alteration of most elements necessary to maintain the outstanding universal value has occurred, causing a major loss of the outstanding universal value.
<b>Area</b> (See Chapter 1, Figure 1.1)	<b>Trends</b>		<b>Confidence in condition and trend</b>
<b>GBR</b> Great Barrier Reef Region	↑ Improving ↔ Stable ↓ Deteriorating — No clear trend		● Adequate high-quality evidence and high level of consensus ◐ Limited evidence or limited consensus ○ Very limited evidence, assessment based on anecdotal information

- a) **Natural beauty and phenomena (previously criterion (iii) now criterion (vii)):** contains unique, rare or superlative natural phenomena, formations or features or areas of exceptional natural beauty, such as superlative examples of the most important ecosystems to man.

	Condition and trend			
	Very good	Good	Poor	Very poor
<b>Overview:</b> The significant loss of coral cover, especially in areas south of about Cooktown, has reduced underwater aesthetic value, as has increasing turbidity in inshore areas. The natural beauty of large areas remains intact, especially for offshore coral reefs in the far north and aerial vistas, as well as for neighbouring islands (many of which are national parks). While many of the natural phenomena remain intact, others are likely to have deteriorated; for example, some turtle nesting locations and coral spawning.				
<b>GBR</b>		↓		

Excerpt from statement	Comment	Area	Condition and trend				Confidence
			Very good	Good	Poor	Very poor	
<i>Superlative natural beauty above and below the water</i>	The natural beauty of most of the Region remains intact, especially for offshore coral reefs and aerial vistas, as well as for neighbouring islands. The significant loss of coral cover has reduced underwater aesthetic value.	GBR					
<i>Some of the most spectacular scenery on Earth</i>	Both above and below the water, the Region's scenery remains spectacular. There have been some declines in the aesthetics of inshore reefs in the southern two-thirds.	GBR					
<i>One of a few living structures visible from space</i>	The Reef remains visible from space and technological advances make these images more accessible.	GBR					
<i>A complex string of reefal structures along Australia's north-east coast</i>	Reefal structures remain intact. Recent estimates vastly increase the extent of coral with the identification of more deepwater reefs.	GBR					
<i>Unparalleled aerial panorama of seascapes comprising diverse shapes and sizes</i>	Aerial vistas remain spectacular, with scenic flights a popular tourism activity.	GBR					
<i>Whitsunday Islands provide a magnificent vista of green vegetated islands and white sandy beaches spread over azure waters</i>	The majority of the Whitsunday Islands are protected and managed as national parks. There have been some changes to island scenery, such as on resort islands.	GBR					
<i>Vast mangrove forests in Hinchinbrook Channel, or the rugged vegetated mountains and lush rainforest gullies</i>	All of Hinchinbrook Island is protected and managed as a national park. Patches of mangrove forests and rainforest were affected by cyclone Yasi.	GBR					
<i>On many of the cays there are spectacular and globally important breeding colonies of seabirds and marine turtles</i>	There have been serious declines in some populations of seabirds and some marine turtle species.	GBR					
<i>Raine Island is the world's largest green turtle breeding area</i>	Long-term data indicates that, since the mid-1970s, green turtle nesting on Raine Island has increased and then plateaued over the past two decades. It is thought to have declined recently.	GBR					
<i>Beneath the ocean surface, there is an abundance and diversity of shapes, sizes and colours... Spectacular coral assemblages of hard and soft corals</i>	Since 1986, average hard coral cover is estimated to have declined from 28 to 13.8 per cent, principally in the southern two-thirds of the Region. This is mainly due to storm damage (48 per cent), crown-of-thorns starfish (42 per cent), and bleaching (10 per cent).	GBR					

Excerpt from statement	Comment	Area	Condition and trend				Confidence
			Very good	Good	Poor	Very poor	
<i>Thousands of species of reef fish provide a myriad of brilliant colours, shapes and sizes</i>	There are about 1500 species of bony fish. Long-term monitoring of about 200 species of coral reef fish has not detected declines in the species monitored. A small number of targeted species are under significant pressure.	GBR	↔				●
<i>The internationally renowned Cod Hole is one of many significant tourist attractions</i>	There is anecdotal evidence of severe declines in the number and condition of potato cod at Cod Hole.	GBR			↓		◐
<i>Superlative natural phenomena include the annual coral spawning, migrating whales, nesting turtles, and significant spawning aggregations of many fish species</i>	The number of migrating humpback whales is increasing. Nesting numbers have declined for at least two of the six species of marine turtle. Protection for fish spawning aggregations has improved, but most sites are unknown.	GBR		—			●

**b) Major stages of the Earth's evolutionary history (previously criterion (i) now criterion (viii)):**  
outstanding examples representing the major stages of the Earth's evolutionary history

	Area	Condition and trend				Confidence
		Very good	Good	Poor	Very poor	
<b>Overview:</b> The Region remains a globally outstanding example of an ecosystem that has evolved over millennia, and almost all geomorphological evolutionary processes remain intact. Examples of all stages of reef development remain, although the overall health of reefs, especially in the southern two-thirds, has declined significantly.	GBR	↔				●

Excerpt from statement	Comment	Area	Condition and trend				Confidence
			Very good	Good	Poor	Very poor	
<i>Globally outstanding example of an ecosystem that has evolved over millennia</i>	The Reef remains an outstanding example of evolutionary history. Recent research has identified deepwater reefs that extend for hundreds of kilometres along the outer shelf at between 40 and 70 metres depth.	GBR	↔				◐
<i>Area has been exposed and flooded by at least four glacial and interglacial cycles, and over the past 18,000 years reefs have grown on the continental shelf</i>	The deepwater reefs are providing valuable records of past coral reef responses to climate and sea level change.	GBR	↔				●
<i>Today, the Great Barrier Reef forms the world's largest coral reef ecosystem... Including examples of all stages of reef development</i>	The Great Barrier Reef remains the world's largest coral reef ecosystem and, while its condition has deteriorated, it remains one of the world's most healthy reef systems, including examples of all stages of reef development.	GBR		↓			●



Excerpt from statement	Comment	Area	Condition and trend				Confidence
			Very good	Good	Poor	Very poor	
<i>Processes of geological and geomorphological evolution are well represented, linking continental islands, coral cays and reefs</i>	Geomorphological features and processes are well represented. Most remain in good condition but some processes are declining, especially in the inshore southern two-thirds.	GBR					
<i>The varied seascapes and landscapes that occur today have been moulded by changing climates and sea levels, and the erosive power of wind and water, over long time periods</i>	The impacts of modern climate change are beginning to have effects on seascapes; for example, through reduced reef building.	GBR					
<i>One-third of the Great Barrier Reef lies beyond the seaward edge of the shallower reefs (and) comprises continental slope and deep oceanic waters and abyssal plains</i>	Evidence of cold water coral communities has been found on deepwater knolls along the edge of the Great Barrier Reef at depths of more than 1000 metres, but these deep areas are hardly known.	GBR					

- c) **Ecological and biological processes (previously criterion (ii) now criterion (ix)):** outstanding examples representing significant ongoing geological processes, biological evolution and man's interaction with his natural environment

	Area	Condition and trend			
		Very good	Good	Poor	Very poor
<b>Overview:</b> Many ecosystem processes remain in good condition; however some, such as recruitment and reef building, are declining. Any processes associated with species groups that are in decline (for example, corals and seagrasses) have likely also declined. In the inshore southern two-thirds, there are particular concerns about some processes such as connectivity, nutrient cycling and sedimentation, principally associated with land-based activities in the catchment. Traditional Owners maintain their cultural practices and customs; however, Indigenous heritage values are under pressure, especially in the southern two-thirds of the Region.	GBR				

Excerpt from statement	Comment	Area	Condition and trend				Confidence
			Very good	Good	Poor	Very poor	
<i>Significant diversity of reef and island morphologies reflects ongoing geomorphic, oceanographic and environmental processes</i>	There remains a diverse range of reef and island morphologies. Most geomorphic, oceanographic and environmental processes remain in good condition, but some are declining, especially in the inshore southern two-thirds.	GBR					
<i>Complex cross-shelf, longshore and vertical connectivity is influenced by dynamic oceanic currents</i>	Most marine species and habitats are thought to remain well connected. There is increasing evidence of intensified flow and accelerated warming in the East Australian Current.	GBR					

Excerpt from statement	Comment	Area	Condition and trend				Confidence
			Very good	Good	Poor	Very poor	
<i>Ongoing ecological processes such as upwellings, larval dispersal and migration</i>	Ecological processes remain in good condition in northern areas. Some processes are in poor condition inshore in the southern two-thirds of the Region and are deteriorating.	GBR					
<i>Ongoing erosion and accretion of coral reefs, sand banks and coral cays combine with similar processes along the coast and around continental islands</i>	Reef building is likely to be in good condition for much of the Region, especially in the north, but has been affected by cyclones and reduced coral cover, especially in the southern two-thirds of the Region.	GBR					
<i>Extensive beds of Halimeda algae represent active calcification and accretion over thousands of years</i>	<i>Halimeda</i> beds are poorly studied, but are likely to be in very good condition given their isolation from land-based impacts and level of protection from trawling.	GBR					
<i>Biologically, the unique diversity of the Great Barrier Reef reflects the maturity of an ecosystem that has evolved over millennia; evidence exists for the evolution of hard corals and other fauna</i>	The diversity of species remains high, but some species are in poor condition, especially inshore in the southern two-thirds of the Region.	GBR					
<i>Vegetation on the cays and continental islands exemplifies the important role of birds...in seed dispersal and plant colonisation</i>	Many islands are national parks or protected within the Marine Park. There are introduced plants on most islands.	GBR					
<i>Human interaction with the natural environment is illustrated by strong ongoing links between Aboriginal and Torres Strait Islanders and their sea country, and includes numerous shell deposits (middens) and fish traps, plus the application of story places and marine totems</i>	Traditional Owners with connections to the Great Barrier Reef maintain their cultural practices and customs. Indigenous heritage is under pressure, especially in the southern two-thirds of the Region.	GBR					

**d) Habitats for conservation of biodiversity (previously criterion (iv) now criterion (x)):** habitats where populations of rare or endangered species of plants and animals still survive

	Area	Condition and trend			
		Very good	Good	Poor	Very poor
<p><b>Overview:</b> There are significant concerns about some key habitats, particularly coral reefs and seagrass meadows, the latter of which is significant for species such as dugong and turtles. Many of the key habitats and species, for which the Great Barrier Reef was inscribed on the World Heritage List (coral reefs, seagrasses, islands, open waters, dugongs, turtles, whales, dolphins and seabirds) are in poor and declining condition, especially in the southern two-thirds of the property. The concerns are not as great in far northern areas, which remain relatively intact. Populations of humpback whales, estuarine crocodiles, loggerhead turtles and green turtles (southern stock) are recovering from historical declines. There have been no records of species extinction, although there is concern that spartooth shark has not been recorded in or near the Region since 1982.</p>	GBR				

Excerpt from statement	Comment	Area	Condition and trend				Confidence
			Very good	Good	Poor	Very poor	
<i>One of the richest and most complex natural ecosystems on Earth, and one of the most significant for biodiversity conservation</i>	The Great Barrier Reef remains a complex ecosystem, rich in biodiversity. Some key habitats are under pressure, especially in southern inshore areas.	GBR					
<i>Tens of thousands of marine and terrestrial species, many of which are of global conservation significance</i>	Some populations (dugong, sharks, seabirds and marine turtles) are known to have declined. Others such as humpback whales, loggerhead turtles and estuarine crocodiles are increasing.	GBR					
<i>The world's most complex expanse of coral reefs...Contain some 400 species of corals in 60 genera</i>	Although there is no published evidence of loss of species associated with coral reefs, there has been a serious decline in hard coral cover and deterioration of coral reef habitats in the southern two-thirds of the Region.	GBR					
<i>Large ecologically important interreefal areas. The shallower marine areas support half the world's diversity of mangroves</i>	The Region's mangrove forests remain very diverse with at least 39 mangrove species and hybrids recorded.	GBR					
<i>Large ecologically important interreefal areas. The shallower marine areas support...many seagrass species</i>	Seagrass diversity remains; however, there have been recent severe declines in abundance and community composition in the inshore southern two-thirds of the Region.	GBR					
<i>Waters also provide major feeding grounds for one of the world's largest populations of the threatened dugong</i>	The dugong population in northern areas remains robust. The population in the southern two-thirds of the Region was very low at the time of listing, and remains so. Declines in the condition of seagrass meadows have had profound effects on dugongs in recent years.	GBR					
<i>At least 30 species of whales and dolphins occur here</i>	Little is known about the populations of most whale species. Two inshore dolphin species are known to be at risk.	GBR					
<i>A significant area for humpback whale calving</i>	The humpback whale population is recovering strongly after being decimated by whaling. The calving habitats are well protected.	GBR					
<i>Six of the world's seven species of marine turtle occur in the Great Barrier Reef. As well as the world's largest green turtle breeding site at Raine Island, the Great Barrier Reef also includes many regionally important marine turtle rookeries</i>	Of the habitats that support marine turtles, the condition of seagrass meadows and coral reefs have declined significantly. While nesting habitats are generally in good condition, sea level rise, increasing air temperature and extreme weather events are affecting their condition.	GBR					

Excerpt from statement	Comment	Area	Condition and trend				Confidence
			Very good	Good	Poor	Very poor	
<i>Some 242 species of birds have been recorded in the Great Barrier Reef. Twenty-two seabird species breed on cays and some continental islands, and some of these breeding sites are globally significant</i>	While the nesting habitats for seabirds remain in generally good condition, declines of up to 70 per cent in some nesting populations have been recorded. There is evidence this may relate to reduced availability of pelagic prey.	GBR					
<i>The continental islands support thousands of plant species, while the coral cays also have their own distinct flora and fauna</i>	Plant diversity is generally well protected, with about one-third of the islands contained within national parks.	GBR					

**Table 7.12 Benchmarking the integrity of the Great Barrier Reef World Heritage Area**

Based on the extent to which the property meets the criteria set out in the *World Heritage Convention Operational Guidelines*.<sup>348</sup>

	Area	Condition and trend			
		Very good	Good	Poor	Very poor
<b>Overview:</b> The Great Barrier Reef is the world's third largest World Heritage Area and encompasses all but the most northerly part of the Great Barrier Reef ecosystem. Except for small exclusions, it is all within a marine protected area and is, therefore, afforded a high level of direct protection and management. External pressures such as climate change, catchment run-off and coastal development are affecting its overall integrity.	GBR				

	Area	Condition and trend				Confidence	
		Very good	Good	Poor	Very poor	Condition	Trend
<b>Includes all elements necessary to express its outstanding universal value:</b> The Great Barrier Reef meets all four natural criteria. While some ecosystems, habitats and populations are under pressure, the elements remain largely intact, particularly in the northern third of the Region.	GBR						
<b>Is of adequate size to ensure the complete representation of the features and processes which convey the property's significance:</b> The Great Barrier Reef Region is vast, covering 14 degrees of latitude and extending 80 to 250 kilometres from the coast. Except for some small exclusions and about 600 of the 1050 islands, almost all of the World Heritage Area is within marine or national parks and is, therefore, afforded a high level of protection and management. While it is of adequate size to ensure complete representation of features and processes, increasing pressures from outside the Region are affecting them.	GBR						
<b>Is protected from the adverse effects of development and/or neglect:</b> While activities within the property are comprehensively managed and use is generally sustainable, the remoteness of some of the property poses challenges for managing agencies. This, and previous assessments, have demonstrated that the most significant impacts on the property's values arise from external pressures such as climate change, catchment run-off and coastal development. In the southern two-thirds of the Region, where there are greater levels of development, the condition and trend of some values are in decline.	GBR						



## 7.6.2 Great Barrier Reef Marine Park

Habitats in the northern third of the Great Barrier Reef are believed to remain in very good condition and are able to support dependent species. Southern habitats, especially those inshore, have deteriorated, particularly seagrass meadows and coral reefs. Terrestrial habitats that support the Reef are generally in very good condition in the northern catchment. However, these habitats have been substantially modified in the southern two-thirds of the catchment, especially saltmarshes, wetlands, woodlands and forests. The functioning of waterbodies connecting the catchment to the Region has deteriorated.

Of those species and species groups for which there is information, there have been significant declines in many, especially those in the inshore southern two-thirds of the Region, and some iconic and cultural keystone species such as dugongs and some marine turtles. Serious declines have been recorded in most hard corals and seagrasses, some fishes and sharks, plus some seabird and shorebird populations. There are four examples of species showing good recovery after past serious declines: humpback whales, estuarine crocodiles, loggerhead turtles and green turtles (southern stock).

Glacio-hydroisostatic processes have resulted in several metres of sea level variation over the past 6500 years. However, geomorphological features close to the populated coast are likely to have been impacted more due to increased human activity and development in this area. The effects of future climate change impacts on geological features are difficult to predict, but the geomorphology of reef islands is predicted to be vulnerable to climate change scenarios.

Traditional Owners with connections to the Great Barrier Reef Marine Park maintain their cultural practices and customs. Places of Indigenous heritage values have not been systematically identified and many have deteriorated, especially around development areas and on islands. Stories, languages and songlines are being affected by activities in the Region. Most places of historic significance are poorly recorded and their condition is not well understood. Heritage values are being maintained or restored at some lightstations.

The Marine Park's environment provides a range of benefits to catchment communities and the nation. It is a source of national pride for many Australians. Many catchment residents understand and appreciate the Reef and have a strong personal attachment to it. Reef-dependent industries contribute strongly to the Australian economy. In recent years, some community benefits have been adversely impacted by the direct and indirect effects of extreme weather.

## 7.6.3 National heritage places

As the Great Barrier Reef was listed as a national heritage place because of its status as a world heritage property, the summary of condition of its outstanding universal value (Section 7.6.1) is also relevant to this matter of national environmental significance. In addition, five of the criteria for national heritage places were identified at the time of listing as being relevant to the Great Barrier Reef. Table 7.13 presents a summary of condition and trend for each.

**Table 7.13 Current condition and trend of the Great Barrier Reef national heritage place**

The Great Barrier Reef was listed as a national heritage place because it is a world heritage property. Five of the listing criteria for national heritage places have been identified as relevant to the Great Barrier Reef.

Relevant national heritage place criteria	Summary of current condition and trend
(a) the place has outstanding heritage value to the nation because of the place's importance in the course, or pattern, of Australia's natural or cultural history	The place retains its importance in the course and pattern of Australia's natural and cultural history.
(b) the place has outstanding heritage value to the nation because of the place's possession of uncommon, rare or endangered aspects of Australia's natural or cultural history	The Region has many uncommon, rare and endangered aspects of Australia's natural and cultural history (for example, marine turtles, seabirds, shorebirds, shell middens and fish traps). There are few examples of recovering populations of endangered species. The Region supports globally significant populations of dugong (an iconic and cultural keystone species). There have been significant declines in dugong and some marine turtles. While Traditional Owner connections remain strong, places of Indigenous heritage value have not been systematically identified and many have deteriorated, especially near developed areas and on islands.
(c) the place has outstanding heritage value to the nation because of the place's potential to yield information that will contribute to an understanding of Australia's natural or cultural history	The Great Barrier Reef is renowned for its potential to yield information that will contribute to an understanding of Australia's natural or cultural history. Natural history is well studied, and the place has vast potential to yield cultural history information.

Relevant national heritage place criteria	Summary of current condition and trend
(d) the place has outstanding heritage value to the nation because of the place's importance in demonstrating the principal characteristics of: <ul style="list-style-type: none"> <li>i. a class of Australia's natural or cultural places; or</li> <li>ii. a class of Australia's natural or cultural environments</li> </ul>	Some of the natural and cultural places and environments are currently in poor condition and declining (for example, seagrass meadows, coral reefs, and some Indigenous heritage values). Natural and cultural places and environments that are more remote (for example north of about Port Douglas and further offshore such as outer barrier reefs) are in good to very good condition.
(e) the place has outstanding heritage value to the nation because of the place's importance in exhibiting particular aesthetic characteristics valued by a community or cultural group.	A recent review of the aesthetic value of the Great Barrier Reef found it had a significant profile in the minds of international visitors and Australians as an iconic, international attraction. It was closely associated with the range of natural environments and experiences it offers. <sup>251</sup> A general decline in the condition (and trend) of a number of habitats, species, processes and values in the southern two-thirds of the Region (for example, coral cover, Indigenous heritage values and charismatic fauna such as dugong and turtles) means it is likely that this criterion has been affected.

#### 7.6.4 Commonwealth marine area

The assessment summary for the Great Barrier Reef Marine Park provided above is also relevant to the Commonwealth marine area.

#### 7.6.5 Listed migratory and listed threatened species

Listed migratory and listed threatened species are assessed as being in generally poor condition, with the exception of humpback whales, estuarine crocodiles and green turtles (southern stock). Threatened inshore and estuarine species or migratory species that rely on habitats in these areas (for example, seabirds and shorebirds) are the most affected by cumulative impacts, both within and beyond the Region. A summary of condition and trend for each group of listed migratory or threatened species is provided below.

Each of the six species of **marine turtle** (listed migratory and threatened) has different population status and current trends. Although the nesting populations of some species are thought to be increasing (for example, loggerhead and green turtle—southern stock), other nesting stocks are declining (for example, green turtle — northern stock and hawksbill turtle). Little is known about the foraging components of most populations, but there are concerns for some species particularly due to pressures outside the Region. Declines in seagrass meadows following the 2011 floods and cyclones have placed additional stress on southern green turtle stocks.

**Estuarine crocodiles** (listed migratory) are in good condition throughout the Region and populations are increasing.

Little is known about the population dynamics of most listed **whale** species. The blue, fin, sei, Antarctic minke, Bryde's, killer and sperm whales are believed to be seasonal migrants to the Region and are rarely sighted. The humpback whale population is in good condition and population numbers are recovering.

**Australian snubfin and Indo-Pacific humpback dolphins** (listed migratory) are considered to be at risk and in decline in the southern inshore two-thirds of the Region due to a number of factors, including habitat loss from declining water quality, climate change and coastal development (for example, all three of the known snubfin dolphin aggregations are in the vicinity of either a planned port expansion or a proposal for a new development).

The northern population of **dugong** (listed migratory) remains healthy with a stable population trend. There has been a substantial decline in dugong in waters south of Cooktown since the 1960s, but the population was thought to have stabilised in 2009. However, a series of extreme weather events from 2009 to 2011 affected seagrass meadows and, therefore, dugong numbers. Populations in the southern inshore area are now assessed as being in very poor condition.

There are significant concerns about a small number of **sharks** which are listed migratory or threatened species, including the speartooth shark which may now be extinct in the Region. The life histories and inshore habitat preferences of many listed shark and ray species make them vulnerable to impacts.

Twenty-four species of **seabirds** are listed as either migratory or threatened. There are also 30 species of listed migratory **shorebirds**. Monitoring of listed seabirds and shorebirds that breed in the Region is currently insufficient to reliably determine condition and trend. Although all seabirds are vulnerable to impacts of climate change, it is the offshore and pelagic foraging seabirds which are most at risk. Some catastrophic nesting failure has already been recorded in wedge-tailed shearwaters in the south. Australia-wide declines in shorebirds of between 70 and 80 per cent have been recorded in the past 24 years.

### 7.6.6 Wetlands of international importance

The Shoalwater and Corio Bay wetland encompasses coastal and subcoastal ecosystems which are relatively undisturbed and in very good condition.<sup>349</sup> This can be attributed mainly to its restricted access as a defence training area. Military use of the site is strictly controlled, managed and monitored and has not caused any known changes to the ecological character of the site. This relies in part upon the fact that almost all training activities and associated infrastructure are outside the boundaries of the site. The Defence Department has pest animal management programs for the site, and a regional oil spill response plan is in place.<sup>350</sup>

A high level of habitat diversity is present, varying from open sandy and rocky shorelines, estuarine embayments and inlets, to a number of remarkable groundwater-dependent freshwater wetlands. These wetlands support a broad range of natural values, including threatened wetland flora and fauna species, and a significant diversity and abundance of waterbirds. The area was thought to provide a refuge for dugong and turtle after the extreme weather events of 2011, as much of its seagrass habitat was in good condition. It continues to be valued for its Indigenous heritage, with Traditional Owners continuing to access the area with the support of the Australian Defence Force.

## 7.7 Key information gaps

The following is a summary of the key information gaps in relation to the condition and trend of values underpinning matters of national environmental significance.

### 7.7.1 Biodiversity — habitats and species

Monitoring programs and research have provided a good understanding of status and trends of many habitats and species underpinning matters of national environmental significance, but there are still gaps for some important elements.

- A small percentage of **islands** are monitored annually for pest species but information is lacking on the status and trends of island habitats, including about 700 islands in the World Heritage Area that are not part of the protected area estate (many of which have important values such as seabird rookeries).
- While the spatial extent of **mangrove forests** is relatively well mapped, trends for species composition remain a gap.
- Many intertidal and limited subtidal **seagrass meadows** are monitored for health and species composition; but the majority of subtidal and deeper meadows are not monitored.
- **Macroalgae** on reefs are monitored as part of coral reef monitoring, but elsewhere in the Region very little is known about this group.
- Emergent mid-shelf and offshore **coral reefs** are well studied and monitored except in the far northern area. Trends in inshore fringing reefs are not as well known, and knowledge of their status is a key gap. **Deeper reefs** (including coral dominated shoals) and their corals are only beginning to be discovered and are not as well understood. These deepwater reefs may play an important role in the future resilience of the Great Barrier Reef.
- **Lagoon floor** habitats (including *Halimeda* banks and non-coral dominated shoals) have been mapped as a one-off project, but there is no information on trends.
- Knowledge of **continental slope** habitats and species remains a gap. Recent surveys are providing insights into what is there; however, there is little information on habitat and species and no trend information. Information on the upper continental slope in the south-east of the Region is required to improve future ecological risk assessments.<sup>238</sup>
- **Invertebrates**, both marine and terrestrial (other than corals and some commercially important species like prawns), are not well understood.
- Information on **phytoplankton** populations comes mostly from chlorophyll analysis of water samples and remote sensing as well as through ad hoc bloom observations. There is very little knowledge of phytoplankton species diversity and condition and trend, while zooplankton are very poorly understood.
- Although knowledge of **microbes** is improving through studies of diseases of coral and other species, understanding of this area remains relatively poor.
- **Bony fish** are included in surveys of coral reefs, but outside of this habitat there is little information except through fishery catch data. More information on targeted fish stocks and bycatch, particularly species identified as being 'at risk', is required to improve fisheries management.
- Recent studies have improved knowledge on movement and behaviour for some **shark and ray** species, but generally there is a poor understanding of their distribution, status and trends. A clear understanding is needed of the biology, ecology, species status and harvest of the lesser known bycatch species (for example, the shark-like batoids).
- Most information on **sea snakes** comes from studying bycatch from the trawl fishery. Sea snake populations and trends are poorly understood.
- Trends in **marine turtle** populations are modelled and inferred from long-term nesting and foraging ground data from a limited number of sites. However, there is a general lack of long-term data for any

species in the far north, and further information on juvenile recruitment and foraging animals in inshore waters and benthic habitats of the Region is required. As migratory species, a better understanding of impacts on populations outside the Region is required. Little is known about olive ridley turtles or foraging flatback turtles.

- Monitoring of trends in **seabirds** that breed in the Region is currently insufficient to reliably determine condition and trend. There is a lack of biological and ecological information on seabirds to help inform a robust management response.
- There are no population estimates for **shorebirds** in the Region.
- There is improving information on inshore **dolphins** but there remains very poor information on population trends and dynamics, as well as key habitats and dietary requirements for these species.
- Ongoing five-yearly aerial surveys of **dugongs** continue to be required, coupled with a better understanding of movement patterns and diving behaviour to inform the scales at which dugongs should be managed.
- The **spatial distributions** of most species and habitats are poorly collated, synthesised and mapped. Intertidal habitats are relatively well mapped and a few studies have employed existing field datasets to predictively model distribution and condition of other biodiversity values.

### 7.7.2 Geomorphological features

An initial study on the important features and processes that make up the geomorphological aspects of the World Heritage Area identified that further work is required on these attributes, including:

- a more detailed assessment of impacts where sensitivity and/or risk is identified as medium to high
- identification of geological features and processes that are susceptible to cumulative impacts.

### 7.7.3 Indigenous and historic heritage values

There is very little information available on Indigenous or historic heritage. Many places of significance are not well documented and only some are listed on a heritage register (under the EPBC Act). It is important to also recognise that both Indigenous and historic heritage values will continue to evolve to represent the flow of history and changing community perceptions.

#### Indigenous heritage

Indigenous heritage is a unique, irreplaceable part of Australia's national cultural heritage which requires improved recognition, protection and understanding. With regard to the Indigenous heritage of the Region, the Authority recognises:

- existing information is limited and not well collated to provide a **baseline of knowledge**
- there is no appropriate **consultation process** with Traditional Owners to re-establish the permissions relating to access and use of information (for example, intellectual property)
- there is no cultural heritage **information management** system.

The Authority also recognises gaps in its knowledge of the following specific aspects of Indigenous heritage values and their appropriate management:

- **places of significance** for Traditional Owners, including sacred sites, spiritual sites, burial sites, songlines and ceremonial sites — some of which may not be at all obvious
- **tangible places of importance** for Indigenous people, including middens, fish traps, scarred trees, camp sites and rock art sites
- **archaeological sites or Indigenous places** recognised as being of national significance (for example, the axe quarry on South Molle Island and the 'contact' rock art of Flinders Island)
- intangible **story places and songlines** and their connections to biodiversity values and ecological processes
- **places or totems** and the reasons they are of contemporary value to Indigenous people
- Indigenous **place names and language** relevant to the Region. Some have been recorded (for example, Bandjin Reefs or Woopaburra place names) but many more need to be before this knowledge is lost.

#### Historic heritage

Specific aspects of the Region's historic heritage have been documented; for example, the locations of historic lighthouses and some 470 historic shipwrecks are recorded. There is also information about the heritage values of many islands. However, there remain gaps in knowledge about many historic places or events. Knowledge gaps of particular significance include:

- the location and condition of **European exploration sites**, such as places visited by early French explorers; many of the places located by Captain Cook in his voyage of discovery; and those relevant to Captain Bligh's voyage after the wreck of the *Pandora*
- key **World War II sites**, including training areas, vessel and aircraft wrecks, structures, and sites for testing and dumping of ordnance and chemicals



- places that trace the history of **extractive industries**, from the early days of guano mining, through to dugong and turtle factories, the mining of limestone and granite, oil exploratory leases, and the evolution of fishing and collecting.

#### 7.7.4 Community benefits of the environment

Some aspects of social, cultural and economic benefits, such as employment and income, have been assessed intermittently, but there has been little systematic monitoring of community benefits derived from the Great Barrier Reef environment. For example, there is very little quantitative information on recreational activity, including where recreational users go and what they do, or about the benefits derived from the environment by Indigenous communities. The following aspects are pivotal to a better understanding about the ways people use and derive benefit from the Region's environment:

- **peoples' activities** in the Region, including:
  - where they go
  - what they do
  - how they access these locations
  - their levels of satisfaction, enjoyment, understanding and appreciation
  - their motivations for visiting
- **economic contributions** of Reef-dependent and non-Reef dependent activities undertaken in the Region and its catchment; plus Reef-dependent industries outside of the catchment
- **stakeholder and community perceptions** of matters such as crowdedness, threats to the Reef, incompatible use, pollution, biophysical changes and aesthetic qualities
- **personal attachment** (including cultural, spiritual and traditional ties) to the Reef by Traditional Owners, stakeholders and visitors
- the extent of **knowledge** that different groups of people have about the Great Barrier Reef and the factors that affect its condition
- the uptake of **new and emerging technologies** and how these influence use, appreciation, enjoyment and understanding of the Region
- different types and components of **media and social networks**, including the different types of Region-related information that is gathered and communicated
- number, types and scale of **health benefits** derived from the Great Barrier Reef and from living adjacent to the Reef
- the range and importance of **ecosystem services** provided by the Great Barrier Reef
- social and cultural limits around **acceptable use** of the Reef
- the **aesthetic and superlative natural phenomena** that make up the world heritage property. While an initial study was completed in early 2013, further work is required to understand the management implications of aesthetic values, including experiential attributes.

#### 7.7.5 Environmental processes

Knowledge of physical and chemical processes is improving but ecological processes are still very poorly understood. It is these processes that connect the elements of the ecosystem into a whole, and understanding this interconnectivity remains an important gap.

- A quantitative understanding of the **interconnectedness** of the ecosystem is largely lacking. There is a critical need and increased emphasis on developing qualitative, quantitative and spatial models to inform system-level understanding and management.
- There is very little information available on **microbial processes** apart from some disease-causing species.
- There is some information on pelagic **primary production**, particularly through remote sensing, while benthic primary production is inferred through surveys of primary producers. There is no system-level understanding of primary production.
- There is moderate information on herbivore and predator populations, but quantitative information about the processes of **herbivory** and **predation** is poor.
- **Symbiotic relationships** are well known for many organisms, including corals, but there is little information available on the spatial distribution of key organisms and the condition of the relationships, except through coral bleaching studies.
- There is reasonable information on the ecosystem components that contribute to **reef building**, but the interplay between accretion and erosion is not well monitored.
- There is some information on **competition**, for instance between coral and algae on reefs; however' this needs to be better understood in the face of climate change. Information on other forms of competition is generally lacking.
- There is some knowledge of fish and coral larval **connectivity** as well as the way some fish populations use different habitats through their life cycle (for example, mangroves or rivers to coral reefs). There is a critical need for an ongoing effort to better understand connectivity.

- While there is a good understanding of nutrient cycling in general, a better understanding of nutrient and carbon cycling, especially in relation to factors such as crown-of-thorns starfish outbreaks and small and large scale effects of ocean acidification, remains pertinent.

### 7.7.6 Processes to address information gaps

The information gaps identified above will be addressed using the same approaches outlined in Chapter 6, Section 6.11, including both scientific and community knowledge.

## 7.8 Summary of conclusions

- Information on condition and trend of **habitats and species** is highly variable with some well-known (for example, shallow coral reefs) and others poorly-known, particularly habitats in remote areas or deep waters.
- Marine habitats **north of Port Douglas** are believed to remain in good or very good condition and are able to support dependent species.
- For **inshore areas south of about Port Douglas**, the condition of species such as dugongs and inshore dolphins and critical habitats such as coral reefs and seagrass meadows is assessed as being poor to very poor. Other species assessed as being in poor condition include marine turtles, sharks and rays including sawfish, sea snakes, seabirds, shorebirds, blue and threadfin salmon and snapper — many of which are listed migratory or listed threatened species.
- **Terrestrial habitats** that support the Great Barrier Reef ecosystem are generally in very good condition north of Port Douglas. Further south, in the bulk of the Region's catchment, all habitats that support the Great Barrier Reef have been substantially modified. This has affected connectivity and the capacity for these habitats to support marine habitats and species.
- Information is lacking for many **environmental processes**, particularly ecological processes. They are critical to the functioning of the Great Barrier Reef. The assessment found individual environmental processes were in good or very good condition in the northern third of the Region, and in good or poor condition further south.
- **Geomorphological features** of the Region are interconnected with its ecological systems. They are largely unchanged over many millennia and are in good or very good condition. Some features are likely to have been affected in the southern two-thirds of the Region.
- **Indigenous heritage** values including sites, cultural keystone species, structures and stories, are being affected by coastal development, climate change and extreme weather, as well as activities taking place within the Region. These values are assessed as being in poor condition overall.
- Some **historic heritage** values such as shipwrecks and structures have been comprehensively identified and recorded. They are assessed as being in good condition, though many will deteriorate with the passage of time.
- **Community benefits** are dependent on the condition of biophysical values of the Great Barrier Reef. Economic contributions have been relatively stable at \$5.7 billion (2012) and employment at 69,000. Impacts such as extreme weather have affected some values (for example, access to reef resources and enjoyment) but community benefits are generally assessed as being in good to very good condition.
- The **natural beauty** of most of the Region remains intact, especially for offshore coral reefs and aerial vistas, as well as for neighbouring islands. The significant loss of coral cover, especially in the southern two-thirds of the Region, has reduced underwater aesthetic value.
- For the Great Barrier Reef World Heritage Area, three of the four relevant criteria of **outstanding universal value** plus integrity are assessed as being in good to very good condition when benchmarked against their condition in 1981. The attributes that make up the habitats for conservation of biodiversity (previously criterion (iv) now criterion (x)) are assessed as being in poor condition overall. Of particular concern, is that more than half of the attributes for all criteria show a deteriorating trend since inscription of the area.

A summary of the current condition of all key values is presented in Table 7.14.

**Table 7.14 Current condition of key values**

The table is a summary of the assessment of condition presented above. Where the condition of a value is different for different areas of the Region, the poorest grade is shown.

Condition	Key values			
	Biodiversity	Geomorphological features	Indigenous and historic heritage values	Community benefits
Very good	<ul style="list-style-type: none"> <li>• Heath and shrublands</li> <li>• <i>Halimeda</i> banks</li> </ul>	<ul style="list-style-type: none"> <li>• Channels and canyons</li> <li>• <i>Halimeda</i> banks</li> </ul>		<ul style="list-style-type: none"> <li>• Enjoyment</li> <li>• Personal connection</li> <li>• Health benefits</li> </ul>
Good	<ul style="list-style-type: none"> <li>• Islands</li> <li>• Beaches and coastlines</li> <li>• Mangrove forests and mangroves</li> <li>• Deeper reefs (&gt;30 m)</li> <li>• Lagoon floor</li> <li>• Shoals</li> <li>• Continental slope</li> <li>• Rainforests</li> <li>• Estuarine crocodiles</li> <li>• Whales</li> <li>• Macroalgae</li> <li>• Benthic microalgae</li> <li>• Other invertebrates</li> <li>• Plankton and microbes</li> <li>• Bony fish</li> </ul>	<ul style="list-style-type: none"> <li>• Coral reefs</li> <li>• Islands and shorelines</li> <li>• River deltas</li> </ul>	<ul style="list-style-type: none"> <li>• Indigenous cultural practices, observances, customs and lore</li> <li>• Places of historic significance — World War II features and sites</li> <li>• Places of historic significance — lightstations</li> <li>• Places of historic significance — other</li> <li>• Places of scientific significance</li> <li>• Places of social significance — iconic sites</li> </ul>	<ul style="list-style-type: none"> <li>• Income</li> <li>• Employment</li> <li>• Understanding and appreciation</li> <li>• Access to Reef resources</li> <li>• Aesthetics</li> </ul>
Poor	<ul style="list-style-type: none"> <li>• Freshwater wetlands</li> <li>• Sea snakes</li> <li>• Sharks and rays</li> <li>• Marine turtles</li> <li>• Open waters</li> <li>• Seabirds</li> <li>• Shorebirds</li> <li>• Forested floodplain</li> <li>• Saltmarshes</li> <li>• Dolphins</li> </ul>	<ul style="list-style-type: none"> <li>• Seagrass meadows</li> </ul>	<ul style="list-style-type: none"> <li>• Indigenous sacred sites, sites of particular significance, places important for cultural tradition</li> <li>• Indigenous stories, songlines, totems and languages</li> <li>• Indigenous structures, technology, tools and archaeology</li> <li>• Places of historic significance — historic shipwrecks</li> </ul>	
Very poor	<ul style="list-style-type: none"> <li>• Connecting waterbodies</li> <li>• Coral reefs (&lt;30 m) and corals</li> <li>• Dugongs</li> <li>• Grass and sedgeland</li> <li>• Seagrass meadows and seagrasses</li> <li>• Woodlands</li> <li>• Forests</li> </ul>			

## References

1. Knowlton, N. and Jackson, J.B.C. 2008, Shifting baselines, local impacts, and global change on coral reefs, *PLoS Biology* 6(2): e54.
2. Pauly, D. 1995, Anecdotes and the shifting baseline syndrome of fisheries, *Trends in Ecology and Evolution* 10(10): 430.
3. Bellwood, D.R., Hughes, T.P., Folke, C. and Nyström, M. 2004, Confronting the coral reef crisis, *Nature* 429: 827-833.
4. De'ath, G., Fabricius, K.E., Sweatman, H. and Puotinen, M. 2012, The 27-year decline of coral cover on the Great Barrier Reef and its causes, *Proceedings of the National Academy of Sciences* 109(44): 17995-17999.
5. Fernbach, M. and Nairn, K. 2007, *Reef recollections: An oral history of the Great Barrier Reef*, Great Barrier Reef Marine Park Authority, Townsville.
6. Wachenfeld, D. 1997, Long-term trends in the status of coral reef-flat benthos: the use of historical photographs, in *Proceedings of the State of the Great Barrier Reef World Heritage Area Workshop: 27-29 November 1995, Townsville, Queensland*, eds. D. Wachenfeld, J. Oliver and K. Davis, Great Barrier Reef Marine Park Authority, Townsville, pp.134-148.

7. McCook, L.J. and Price, I.R. 1997, Macroalgal distributions on the Great Barrier Reef: A review of patterns and causes, in *Proceedings of the National Great Barrier Reef: Science, Use and Management Conference, 25-29 November 1996, James Cook University, Townsville*, eds. J. Campbell and C. Dalliston, Great Barrier Reef Marine Park Authority, Townsville, pp.37-46.
8. McCook, L.J., Ayling, T., Cappel, M., Choat, J.H., Evans, R.D., DeFreitas, D.M., Heupel, M., Hughes, T.P., Jones, G.P., Mapstone, B., Marsh, H., Mills, M., Molloy, F.J., Pitcher, C.R., Pressey, R.L., Russ, G.R., Sutton, S., Sweatman, H., Tobin, R., Wachenfeld, D. and Williamson, D. 2010, Adaptive management of the Great Barrier Reef: A globally significant demonstration of the benefits of networks of marine reserves, *Proceedings of the National Academy of Sciences* 107(43): 18278-18285.
9. Roff, G., Clark, T.R., Raymond, C.E., Zhao, J., Feng, Y., McCook, L.J., Done, T.J. and Pandolfi, J.M. 2013, Palaeoecological evidence of a historical collapse of corals at Pelorus Island, inshore Great Barrier Reef, following European settlement, *Proceedings of the Royal Society B: Biological Sciences* 280(1750): 2012-2100.
10. Turner, M. and Batianoff, G.N. 2007, Vulnerability of island flora and fauna of the Great Barrier Reef to climate change, in *Climate change and the Great Barrier Reef: a vulnerability assessment*, eds J.E. Johnson and P.A. Marshall, Great Barrier Reef Marine Park Authority and Australian Greenhouse Office, Townsville, pp. 621-666.
11. Walker, T.A. 1991, Pisonia islands of the Great Barrier Reef: Part I. Distribution, abundance and dispersal by seabirds of *Pisonia grandis*, *Atoll Research Bulletin* 350: 1-23.
12. Congdon, B.C. 2008, Seabirds, in *The Great Barrier Reef: biology, environment and management*, eds P.A. Hutchings, M.J. Kingsford and O. Hoegh-Guldberg, CSIRO, Collingwood, pp. 359-368.
13. Smithers, S.G., Harvey, N., Hopley, D. and Woodroffe, C.D. 2007, Vulnerability of geomorphological features in the Great Barrier Reef to climate change, in *Climate change and the Great Barrier Reef: a vulnerability assessment*, eds J.E. Johnson and P.A. Marshall, Great Barrier Reef Marine Park Authority and Australian Greenhouse Office, Townsville, pp. 667-716.
14. Hopley, D., Smithers, S. and Parnell, K.E. 2007, *The geomorphology of the Great Barrier Reef: development, diversity and change*, Cambridge University Press, Cambridge, UK.
15. Heatwole, H., Done, T. and Cameron, E. 1981, *Community ecology of a coral cay: A study of One-Tree Island, Great Barrier Reef, Australia*, W. Junk, The Hague.
16. Galloway, R.W., Story, R., Cooper, R. and Yapp, G.A. 1984, *Coastal lands of Australia*, CSIRO Division of Water and Land Resources, Canberra.
17. Great Barrier Reef Marine Park Authority 2012a, *Informing the outlook for Great Barrier Reef coastal ecosystems*, GBRMPA, Townsville.
18. Clemens, R.S., Haslem, A., Oldland, J., Shelley, L., Weston, M.A. and Abdullah Abu Diyan, M. 2008, *Identification of significant shorebird areas in Australia: mapping, thresholds and criteria*, Birds Australia, Carlton.
19. Hamann, M., Grech, A., Wolanski, E. and Lambrechts, J. 2011, Modelling the fate of marine turtle hatchlings, *Ecological Modelling* 222: 1515-1521.
20. Duke, N.C., Bella, A.M., Pederson, D.K., Roelfsema, C.M. and Nash, C.B. 2005, Herbicides implicated as the cause of severe mangrove dieback in the Mackay region, NE Australia: consequences for marine plant habitats of the GBR World Heritage Area, *Marine Pollution Bulletin* 51: 308-324.
21. Duke, N. 1997, Mangroves in the Great Barrier Reef World Heritage Area: Current status, long-term trends, management implications and research, in *Proceedings of the State of the Great Barrier Reef World Heritage Area Workshop: 27-29 November 1995, Townsville, Queensland*, eds. D. Wachenfeld, J. K. Oliver, K. Davis, J. K. Oliver and K. Davis, Great Barrier Reef Marine Park Authority, Townsville, pp.288-299.
22. Goudkamp, K. and Chin, A. 2006, Mangroves and saltmarshes, in *The state of the Great Barrier Reef*, ed. A. Chin, Great Barrier Reef Marine Park Authority, Townsville.
23. Duke, N.C. and Larkum, A.W.D. 2008, Mangroves and seagrasses, in *The Great Barrier Reef: biology, environment and management*, eds P. Hutchings, M. Kingsford and O. Hoegh-Guldberg, CSIRO Publishing, Collingwood, pp. 156-170.
24. Duke, N.C., Lawn, T., Roelfsema, C.M., Phinn, S., Zahmel, K.N., Pedersen, D.K., Harris, C., Steggles, N. and Tack, C. 2003, *Assessing historical change in coastal environments: Port Curtis, Fitzroy Estuary and Moreton Bay regions: final report to the CRC for coastal zone and Estuary Waterway Management*, Marine Botany Group, Centre for Marine Studies, University of Queensland, Brisbane.
25. Brodie, J.E. and Mitchell, A.W. 2005, Nutrients in Australian tropical rivers: Changes with agricultural development and implications for receiving environments, *Marine and Freshwater Research* 56(3): 279-302.
26. Coles, R., McKenzie, L.J., Rasheed, M., Mellors, J., Taylor, H., Dew, K., McKenna, S., Sankey, T., Carter, A. and Grech, A. 2007, *Status and trends of seagrass habitats in the Great Barrier Reef World Heritage Area: Report to the Marine and Tropical Sciences Research Facility*, Reef and Rainforest Research Centre, Cairns.
27. McKenzie, L.J., Collier, C. and Waycott, M. 2012a, *Reef Rescue Marine Monitoring Program: Inshore seagrass, annual report for the sampling period 1st July 2010 – 31st May 2011*, Fisheries Queensland, Cairns.
28. Fairweather, C., McKenna, S. and Rasheed, M. 2011a, *Long-term seagrass monitoring in the Port of Mourilyan: November 2010*, Fisheries Queensland, Cairns.
29. Fairweather, C.L., McKenna, S.A. and Rasheed, M.A. 2011b, *Long term seagrass monitoring in Cairns Harbour and Trinity Inlet: December 2009 and 2010*, Fisheries Queensland, Cairns.
30. Taylor, H.A. and Rasheed, M.A. 2010, *Port of Townsville long-term seagrass monitoring: October 2010*, Department of Employment, Economic Development and Innovation, Cairns.
31. Sankey, T.L. and Rasheed, M.A. 2011, *Gladstone permanent transects seagrass monitoring sites - February and March 2011 update*, Fisheries Queensland, Cairns.
32. McKenzie, L.J., Yoshida, R., Grech, A. and Coles, R. 2010, *Queensland seagrasses: Status 2010 - Torres Strait and East Coast*, Department of Employment, Economic Development and Innovation, Brisbane.
33. McKenna, S.A., Rasheed, M.A., Unsworth, R.K.F., Taylor, H.A., Chartrand, K.M. and Sankey, T.L. 2009, *Long term seagrass monitoring in Cairns Harbour and Trinity Inlet – October/December 2008*, Department of Employment, Economic Development and Innovation, Northern Fisheries Centre, Cairns.
34. McKenzie, L.J., Unsworth, R. and Waycott, M. 2010, *Reef Rescue Marine Monitoring Program: Intertidal seagrass annual report for sampling period 1 September 2009 - 31 May 2010*, Fisheries Queensland, Cairns.
35. Fabricius, K.E., Okaji, K. and De'ath, G. 2010, Three lines of evidence to link outbreaks of the crown-of-thorns seastar *Acanthaster planci* to the release of larval food limitation, *Coral Reefs* 29: 593-605.
36. Hughes, T.P., Bellwood, D.R., Baird, A.H., Brodie, J., Bruno, J.F. and Pandolfi, J.M. 2011, Shifting base-lines, declining coral cover, and the erosion of reef resilience: Comment on Sweatman et al (2011), *Coral Reefs* 30(3): 653-660.
37. Sweatman, H. and Syms, C. 2011, Assessing loss of coral cover on the Great Barrier Reef: A response to Hughes et al. (2011), *Coral Reefs* 30: 661-664.
38. Sweatman, H., Cheal, A.J., Coleman, G.J., Emslie, M.J., Johns, K., Jonker, M., Miller, I.R. and Osborne, K. 2008, *Long-term monitoring of the Great Barrier Reef: Status Report 8*, Australian Institute of Marine Science, Townsville.
39. Hoegh-Guldberg, O., Anthony, K., Berkelmans, R., Davis, S., Fabricius, K., Lough, J., Marshall, P., van Oppen, M.J.H., Negri, A.P. and Willis, B. 2007, Vulnerability of reef-building corals on the Great Barrier Reef to climate change, in *Climate change and the Great Barrier Reef: A Vulnerability Assessment*, eds J.E. Johnson and P.A. Marshall, Great Barrier Reef Marine Park Authority and Australian Greenhouse Office, Townsville, pp. 271-307.



40. Done, T., Whetton, P., Jones, R.J., Berkelmans, R., Lough, J.M., Skirving, W. and Wooldridge, S. 2003, *Global climate change and coral bleaching on the Great Barrier Reef*, Department of Natural Resources and Mines, Brisbane.
41. Berkelmans, R., De'ath, G., Kininmonth, S. and Skirving, W.J. 2004, A comparison of the 1998 and 2002 coral bleaching events on the Great Barrier Reef: Spatial correlation, patterns and predictions, *Coral Reefs* 23(1): 74-83.
42. Brodie, J.E., Kroon, F.J., Schaffelke, B., Wolanski, E.C., Lewis, S.E., Devlin, M.J., Bohnet, I.C., Bainbridge, Z.T., Waterhouse, J. and Davis, A.M. 2012, Terrestrial pollutant runoff to the Great Barrier Reef: An update of issues, priorities and management responses, *Marine Pollution Bulletin* 65: 81-100.
43. Great Barrier Reef Marine Park Authority 2009, *Great Barrier Reef Outlook Report 2009*, GBRMPA, Townsville.
44. Osborne, K., Dolman, A.M., Burgess, S.C. and Johns, K.A. 2011, Disturbance and the dynamics of coral cover on the Great Barrier Reef (1995–2009), *PLoS ONE* 6(3): e17516.
45. Webster, J.M., Beaman, R.J., Bridge, T., Davies, P.J., Byrne, M., Williams, S., Manning, P., Pizarro, O., Thornborough, K., Woolsey, E., Thomas, A. and Tudhope, S. 2008, From corals to canyons: The Great Barrier Reef margin, *EOS, Transactions, American Geophysical Union* 89(24): 217-218.
46. Bridge, T.C.L., Done, T.J., Friedman, A., Beaman, R.J., Williams, S.B., Pizarro, O. and Webster, J.M. 2011, Variability in mesophotic coral reef communities along the Great Barrier Reef, Australia, *Marine Ecology Progress Series* 428: 63-75.
47. Bridge, T.C.L., Fabricius, K.E., Bongaerts, P., Wallace, C.C., Muir, P., Done, T.J. and Webster, J.M. 2012, Diversity of scleractinian and octocorallia in the mesophotic zone of the Great Barrier Reef, Australia, *Coral Reefs* 31(1): 179-189.
48. van Oppen, M.J.H., Bongaerts, P., Underwood, J.N., Peplow, L.M. and Cooper, T.F. 2011a, The role of deep reefs in shallow reef recovery: An assessment of vertical connectivity in a brooding coral from west and east Australia, *Molecular Ecology* 20(8): 1647-1660.
49. Pears, R.J., Morison, A.K., Jebreen, E.J., Dunning, M.C., Pitcher, C.R., Courtney, A.J., Houlden, B. and Jacobsen, I.P. 2012a, *Ecological risk assessment of the East Coast Otter Trawl Fishery in the Great Barrier Reef Marine Park: summary report*, Great Barrier Reef Marine Park Authority, Townsville.
50. Pitcher, C.R., Burridge, C.Y., Wassenberg, T.J., Hill, B.J. and Poiner, I.R. 2009, A large scale BACI experiment to test the effects of prawn trawling on seabed biota in a closed area of the Great Barrier Reef Marine Park, Australia, *Fisheries Research* 99(3): 168-183.
51. Pitcher, C.R., Doherty, P., Arnold, P., Hooper, J., Gribble, N.A., Bartlett, C., Browne, M., Campbell, N., Cannard, T., Cappel, M., Carini, G., Chalmeres, S., Cheers, S., Chetwynd, D., Colegax, A., Coles, R., Cook, S., Davie, P., De'ath, G., Devereux, D., Done, B., Donovan, T., Ehrke, B., Ellis, N., Ericson, G., Fellegara, I., Forcey, K., Furey, M., Gledhill, D., Good, N., Gordon, S., Haywood, M., Hendricks, M., Jacobsen, I., Johnson, J., Jones, M., Kininmonth, S., Kistler, S., Last, P., Leite, A., Amrks, S., McLeod, I., Oczkowicz, S., Robinson, M., Rose, C., Seabright, D., Sheils, J., Sherlock, M., Skelton, P., Smith, D., Smith, G., Speare, P., Stowar, M., Strickland, C., Van der Geest, C., Venables, W., Walsh, C., Wassenberg, T.J., Welna, A. and Yearsley, G. 2007a, *Seabed biodiversity on the continental shelf of the Great Barrier Reef World Heritage Area*. AIMS/CSIRO/QM/QDPI CRC Reef Research Task final report, CSIRO Marine and Atmospheric Research, Hobart.
52. Pitcher, C.R., Venables, B., Browne, M., Doherty, P. and De'ath, G. 2007b, *Indicators of protection levels for seabed habitats, species and assemblages on the continental shelf of the Great Barrier Reef World Heritage Area*, Reef and Rainforest Research Centre, Cairns.
53. Stieglitz, T.C. 2012, Habitat engineering by decadal-scale bioturbation around shipwrecks on the Great Barrier Reef mid-shelf, *Marine Ecology Progress Series* 477: 29-40.
54. Cappel, M., MacNeil, M.A., Stowar, M.J. and Doherty, P.J. 2009a, *The influence of zoning (closure to fishing) on fish communities of the deep shoals and reef bases of the southern Great Barrier Reef. Part 1: Baited video surveys of the Pompey, Swain and Capricorn-Bunker groups of reefs off Mackay and Gladstone. Report to the Marine and Tropical Sciences Research Facility*, Reef and Rainforest Research Centre, Cairns and Australian Institute of Marine Science, Townsville.
55. Cappel, M., De'ath, G., Stowar, M.J., Johannson, C. and Doherty, P.J. 2009b, *The influence of zoning (closure to fishing) on fish communities of the deep shoals and reef bases of the southern Great Barrier Reef Marine Park. Part 2: development of protocols to improve accuracy in baited video techniques used to detect effects of zoning*, Reef and Rainforest Research Centre, Cairns.
56. Stieglitz, T.C. 2011, The Yongala's Halo of Holes - Systematic Bioturbation Close to a Shipwreck, in *Seafloor Geomorphology as Benthic Habitat: GeoHab Atlas of seafloor geomorphic features and benthic habitats*, eds P.T. Harris and E.K. Baker, Elsevier, London, pp. 277-287.
57. Haywood, M., Pitcher, C., Ellis, N., Wassenberg, T., Smith, G., Forcey, K., McLeod, I., Carter, A., Strickland, C. and Coles, R. 2008, Mapping and characterisation of the inter-reefal benthic assemblages of the Torres Strait, *Continental Shelf Research* 28(16): 2304-2316.
58. Orme, G.R. and Salama, M.S. 1988, Form and seismic stratigraphy of *Halimeda* banks in part of the northern Great Barrier Reef Province, *Coral Reefs* 6(3-4): 131-137.
59. Rees, S.A., Opdyke, B.N., Wilson, P.A. and Henstock, T.J. 2007, Significance of *Halimeda* bioherms to the global carbonate budget based on a geological sediment budget for the Northern Great Barrier Reef, Australia, *Coral Reefs* 26(1): 177-188.
60. Sinutok, S., Hill, R., Doblin, M.A., Wuhner, R. and Ralph, P.J. 2011, Warmer more acidic conditions cause decreased productivity and calcification in subtropical coral reef sediment-dwelling calcifiers, *Limnology and Oceanography* 56(4): 1200-1212.
61. Beaman, R.J., Webster, J.M. and Wust, R.A.J. 2008, New evidence for drowned shelf edge reefs in the Great Barrier Reef, Australia, *Marine Geology* 247(1-2): 17-34.
62. Puga-Bernabéu, Á., Webster, J.M., Beaman, R.J. and Guilbaud, V. 2011, Morphology and controls on the evolution of a mixed carbonate-siliciclastic submarine canyon system, Great Barrier Reef margin, north-eastern Australia, *Marine Geology* 289(1-4): 100-116.
63. Brodie, J.E., Devlin, M., Haynes, D. and Waterhouse, J. 2011, Assessment of the eutrophication status of the Great Barrier Reef lagoon (Australia), *Biogeochemistry* 106(2): 281-302.
64. De'ath, G., Lough, J.M. and Fabricius, K.E. 2009, Declining coral calcification on the Great Barrier Reef, *Science* 323(5910): 116-119.
65. Guinotte, J.M. and Fabry, V.J. 2008, Ocean acidification and its potential effects on marine ecosystems, *Annals of the New York Academy of Sciences* 1134(1): 320-342.
66. Devlin, M., Waterhouse, J., McKinna, L. and Lewis, S. 2010, *Terrestrial runoff in the Great Barrier Reef: Marine Monitoring Program (3.7.2b) Tully and Burdekin case studies*, Australian Centre for Tropical and Freshwater Research, James Cook University, Townsville.
67. Devlin, M., McKinna, L. and Harkness, P. 2010, *Reef Rescue Marine Monitoring Program: terrestrial runoff in the Great Barrier Reef (3.7.2b): Flood plume monitoring for 2009/10 annual report*, Australian Centre for Tropical and Freshwater Research, Townsville.
68. Devlin, M., Wenger, A., Waterhouse, J., Alvarez-Romero, J., Abbott, B. and Teixeira da Silva, E. 2011, *Reef Rescue Marine Monitoring Program: flood plume monitoring annual report 2010-11, Incorporating results from the Extreme Weather Response Program flood plume monitoring, Report for the Great Barrier Reef Marine Park Authority*, Australian Centre for Tropical Freshwater Research, James Cook University, Townsville.
69. Lovelock, C. and Clarke, S. 1993, *Field guide to the mangroves of Queensland*, Australian Institute of Marine Science, Townsville.
70. Wolanski, E., Mazda, Y., Furukawa, K., Ridd, P., Kithaka, J., Spagnol, S. and Stieglitz, T. 2001, Water circulation in mangroves and its implications for biodiversity, in *Oceanographic processes of coral reefs: physical and biological*

links in the Great Barrier Reef, ed. E. Wolanski, CRC Press, Boca Raton, Florida, USA, pp. 53-76.

71. Duke, N.C. 2006, *Australia's mangroves: the authoritative guide to Australia's mangrove plants*, University of Queensland, Brisbane.

72. Waycott, M., Collier, C., McMahon, K., Ralph, P., McKenzie, L.J., Udy, J. and Grech, A. 2007, Vulnerability of seagrasses in the Great Barrier Reef to climate change, in *Climate Change and the Great Barrier Reef: a vulnerability assessment*, eds J.E. Johnson and P.A. Marshall, Great Barrier Reef Marine Park Authority and Australian Greenhouse Office, Townsville, pp. 193-235.

73. Diaz-Pulido, G., Gouezo, M., Tilbrook, B., Dove, S. and Anthony, K.R.N. 2011, High CO<sub>2</sub> enhances the competitive strength of seaweeds over corals, *Ecology Letters* 14: 156-162.

74. Diaz-Pulido, G., Harri, S., McCook, L.J. and Hoegh-Guldberg, O. 2010, The impact of benthic algae on the settlement of a reef-building coral, *Coral Reefs* 29(1): 203-208.

75. Diaz-Pulido, G. 2008, Macroalgae, in *The Great Barrier Reef: biology, environment and management*, eds P. Hutchings, M. Kingsford and O. Hoegh-Guldberg, CSIRO Publishing, Collingwood, pp. 145-155.

76. Diaz-Pulido, G. & McCook, L.J. 2008, *State of the Great Barrier Reef Report: Environmental status: macroalgae (seaweeds)*, Great Barrier Reef Marine Park Authority, Townsville, <[http://www.gbrmpa.gov.au/\\_data/assets/pdf\\_file/0019/3970/SO\\_RR\\_Macroalgae.pdf](http://www.gbrmpa.gov.au/_data/assets/pdf_file/0019/3970/SO_RR_Macroalgae.pdf)>.

77. Cheshire, A.C., Collings, G.J., Edyvane, K.S. and Westphalen, G. 2000, *Overview of the conservation status of Australian marine macroalgae: A report to Environment Australia*, University of Adelaide, Adelaide.

78. Anthony, K.R.N., Maynard, J.A., Diaz-Pulido, G., Mumby, P.J., Marshall, P.A., Cao, L. and Hoegh-Guldberg, O. 2011, Ocean acidification and warming will lower coral reef resilience, *Global Change Biology* 17: 1798-1808.

79. Hughes, T.P., Linares, C., Dakos, C., van de Leemput, I.A. and van Nes, E.H. 2012, Living dangerously on borrowed time during slow, unrecognized regime shifts, *Trends in Ecology and Evolution* 28(3): 149-155.

80. Hughes, T.P., Rodrigues, M.J., Bellwood, D.R., Ceccarelli, D., Hoegh-Guldberg, O., McCook, L.J., Moltschniowskyj, N.A., Pratchett, M.S., Steneck, R.S. and Willis, B. 2007a, Phase shifts, herbivory, and the resilience of coral reefs to climate change, *Current Biology* 17(4): 360-365.

81. Roelfsema, C.M., Phinn, S.R. and Dennison, W.C. 2002, Spatial distribution of benthic microalgae on coral reefs determined by remote sensing, *Coral Reefs* 21: 264-274.

82. Heil, C.A., Chaston, K., Jones, A., Bird, P., Longstaff, B., Costanzo, S. and Dennison, W.C. 2004, Benthic microalgae in coral reef sediments of the southern Great Barrier Reef, Australia, *Coral Reefs* 23: 336-343.

83. Sweatman, H., Delean, S. and Syms, C. 2011, Assessing loss of coral cover on Australia's Great Barrier Reef over two decades, with implications for longer-term trends, *Coral Reefs* 30: 521-531.

84. De'ath, G. and Fabricius, K.E. 2010, Water quality as a regional driver of coral biodiversity and macroalgae on the Great Barrier Reef, *Ecological Applications* 20(3): 840-850.

85. Alderslade, P. and Fabricius, K.E. 2008, Octocorals, in *The Great Barrier Reef: biology, environment and management*, eds P. Hutchings, M. Kingsford and O. Hoegh-Guldberg, CSIRO Publishing, Collingwood, pp. 222-245.

86. Veron, J. 1986, *Corals of Australia and the Indo-Pacific*, Angus and Robertson, North Ryde, Australia.

87. Veron, J.E.N., DeVantier, L.M., Turak, E., Green, A.L., Kininmonth, S., Stafford-Smith, M. and Peterson, N. 2009, Delineating the Coral Triangle, *Galaxea* 11(2): 91-100.

88. Pickerell, J. 2009, *Drowned reef: world's largest fossil*. *Cosmos Online*, Cosmos Media, <<http://www.cosmosmagazine.com/features/drowned-reef/>>.

89. Queensland Department of Employment, Economic Development and Innovation 2011a, *Stock status of Queensland's fisheries resources 2011*, DEEDI, Brisbane.

90. Queensland Department of Employment, Economic Development and Innovation 2010a, *Stock status of Queensland's fisheries resources 2009-10*, DEEDI, Brisbane.

91. Richardson, A.J., McKinnon, A.D. and Swadling, K.M. 2012, Zooplankton. in *Marine Climate Change Impacts and Adaptation Report Card for Australia 2012*, eds E.S. Poloczanska, A.J. Hobday and A.J. Richardson, CSIRO, Canberra.

92. Webster, N.S. and Bourne, D.G. 2012, Microbes, in *A marine climate change impacts and adaptation report card for Australia 2012*, eds E.S. Poloczanska, A.J. Hobday and A.J. Richardson, CSIRO, Canberra.

93. Harrison, H.B., Williamson, D.H., Evans, R.D., Almany, G.R., Thorrold, S.R., Russ, G.R., Feldheim, K.A., van Herwerden, L., Planes, S., Srinivasan, M., Berumen, M.L. and Jones, G.P. 2012, Larval export from marine reserves and the recruitment benefit for fish and fisheries, *Current Biology* 22(11): 1023-1028.

94. Queensland Department of Employment, Economic Development and Innovation 2010b, *Annual status report Rocky Reef Fin Fish Fishery 2010*, DEEDI, Brisbane.

95. Welch, D.J., Ballagh, A., Newman, S.J., Lester, R.J.G., Moore, B.R., vanHerwerden, L., Horne, J.B., Allsop, Q., Saunders, T., Stapley, J. and Gribble, N.A. 2010, *Defining the stock structure of northern Australia's threadfin salmon species, Final report to the Fisheries Research and Development Corporation, Project 2007/032*, Fisheries Research Development Corporation, Canberra.

96. Welch, D., Gribble, N.A. and Garrett, R. 2002, *Assessment of the threadfin salmon fishery in Queensland, 2002*, Department of Primary Industries and Fisheries, Brisbane.

97. Welch, D.J., Hoyle, S.D., McPherson, G.R. and Gribble, N.A. 2002, *Preliminary assessment of the Queensland east coast spanish mackerel fishery, Information Series Q102110*, Department of Primary Industries, Brisbane.

98. Craik, W. 1979, *Amateur Fishing on the Great Barrier Reef. Technical Memorandum No. 4*, Great Barrier Reef Marine Park Authority, Townsville.

99. Goeden, G.B. 1979, Is the Great Barrier Reef being overfished? *Australian Fisheries* (9): 18-20.

100. Craik, G.J.S. 1981, Recreational fishing on the Great Barrier Reef, in *Proceedings of the 4th International Coral Reef Symposium*, eds E. Gomez D. and C. E. Birkeland, Marine Sciences Center, University of the Philippines, Manila, Philippines, pp.47-52.

101. Goeden, G.B. 1982, Intensive fishing and a 'keystone' predator species: ingredients for community instability, *Biological Conservation* 22(4): 273-281.

102. Evans, R.D. and Russ, G.R. 2004, Larger biomass of targeted reef fish in no-take marine reserves on the Great Barrier Reef, Australia, *Aquatic Conservation: Marine and Freshwater Ecosystems* 14(5): 505-519.

103. Williamson, D.H., Russ, G.R. and Ayling, A.M. 2004, No-take marine reserves increase abundance and biomass of reef fish on inshore fringing reefs of the Great Barrier Reef, *Environmental Conservation* 31(2): 149-159.

104. Frisch, A. and Van Herwerden, L. 2006, Field and experimental studies of hybridization between coral trouts, *Plectropomus leopardus* and *Plectropomus maculatus* (Serranidae), on the Great Barrier Reef, Australia, *Journal of Fish Biology* 68(4): 1013-1025.

105. Pogonoski, J.J., Pollard, D.A. and Paxton, J.R. 2002, *Conservation overview and action plan for Australian threatened and potentially threatened marine and estuarine fishes*, Environment Australia, Canberra.

106. Queensland Department of Employment, Economic Development and Innovation 2011b, *Annual status report 2010 Coral Reef Fin Fish Fishery*, DEEDI, Brisbane.

107. Cornish, A. and Kiwi, L.K. 2004, *Plectropomus leopardus*. In: *IUCN 2011, IUCN Red List of Threatened Species, Version 2013.1*, IUCN the International Union for Conservation of Nature, Cambridge, UK, <<http://www.iucnredlist.org/details/44684/0>>.

108. Queensland Department of Employment, Economic Development and Innovation 2011c, *Framework for defining stock status*, DEEDI, Brisbane.



109. Campbell, A.B., O'Neill, M.F., Staunton-Smith, J., Atfield, J.C. and Kirkwood, J. 2012, *Stock assessment of the Australian East Coast Spanish mackerel (Scomberomorus commerson) fishery*, Department of Agriculture, Fisheries and Forestry, Brisbane.
110. Queensland Department of Employment, Economic Development and Innovation 2010c, *Barramundi results update 2007-09*, DEEDI, Brisbane.
111. Queensland Department of Employment, Economic Development and Innovation 2011d, *Annual status report 2010: East Coast Inshore Fin Fish Fishery*, DEEDI, Brisbane.
112. Chin, A., Kyne, P.M., Walker, T.I. and McAuley, R.B. 2010, An integrated risk assessment for climate change: analysing the vulnerability of sharks and rays on Australia's Great Barrier Reef, *Global Change Biology* 16(7): 1936-1953.
113. Chin, A. and Kyne, P.M. 2007, Vulnerability of chondrichthyan fishes of the Great Barrier Reef to climate change, in *Climate change and the Great Barrier Reef: a vulnerability assessment*, eds J.E. Johnson and P.A. Marshall, Great Barrier Reef Marine Park Authority and Australian Greenhouse Office, Townsville, pp. 393-425.
114. Kyne, P.M. 2008, *Chondrichthyans and the Queensland East Coast Trawl Fishery: Bycatch reduction, biology, conservation status and sustainability*. PhD thesis, University of Queensland, Brisbane.
115. Welch, D.J., Ovenden, J., Simpfendorfer, C.A., Tobin, A.J., Morgan, J.A.T., Street, R., White, J., Harry, A.V., Schroeder, R. and Macbeth, W.G. 2011, *Stock structure of exploited shark species in north-eastern Australia: report to the Fisheries Research and Development Corporation project 2007/35*, Fishing and Fisheries Research Centre, Townsville.
116. Heupel, M.R., Simpfendorfer, C.A. and Fitzpatrick, R. 2010, Large-Scale Movement and Reef Fidelity of Grey Reef Sharks, *PLoS ONE* 5(3): 9650.
117. Knip, D.M., Heupel, M.R. and Simpfendorfer, C.A. 2010, Sharks in nearshore environments: models, importance, and consequences, *Marine Ecology Progress Series* 402: 1-11.
118. Lynch, A.-J., Sutton, S.G. and Simpfendorfer, C.A. 2009, Implications of recreational fishing for elasmobranch conservation in the Great Barrier Reef Marine Park, *Aquatic Conservation: Marine and Freshwater Ecosystems* 20(3): 312-320.
119. Heupel, M.R., Williams, A.J., Welch, D.J., Ballagh, A., Mapstone, B.D., Carlos, G., Davies, C. and Simpfendorfer, C.A. 2009, Effects of fishing on tropical reef associated shark populations on the Great Barrier Reef, *Fisheries Research* 95(2-3): 350-361.
120. Ayling, A.M. and Choat, J.H. 2008, *Abundance patterns of reef sharks and predatory fishes on differently zoned reefs in the offshore Townsville region*, Great Barrier Reef Marine Park Authority, Townsville.
121. Stevens, J.D., Pillans, R.D. and Salini, J.P. 2005, *Conservation assessment of Glyptis sp A (spear-tooth shark), Glyptis sp. C (northern river shark), Pristis microdon (freshwater sawfish) and Pristis zijsron (green sawfish): final report to the Department of the Environment and Heritage*, CSIRO Marine Research, Hobart.
122. Robbins, W.D., Hisano, M., Connolly, S.R. and Choat, J.H. 2006, Ongoing collapse of coral reef shark populations, *Current Biology* 16: 2314-2319.
123. Grech, A. and Coles, R. 2011, Interactions between a trawl fishery and spatial closures for biodiversity conservation in the Great Barrier Reef World Heritage Area, Australia, *PLoS ONE* 6(6): e21094.
124. Courtney, A.J., Schemel, B.I., Wallace, R.M., Campbell, M.J. and Mayer, D.J. 2010, *Reducing the impact of Queensland's trawl fisheries on protected sea snakes*, Fisheries Research and Development Corporation and Department of Employment, Economic Development and Innovation, Brisbane.
125. Courtney, A.J., Schemel, B.I., Wallace, R.M., Campbell, M.J. and Mayer, D.J. 2008, *Quantifying and reducing the incidental catch and fishing mortality of sea snakes in the Queensland (Australia) trawl fishery, FRDC Project No. 2005/053*, Department of Employment, Economic Development and Innovations, Brisbane.
126. Lukoschek, V., Heatwole, H., Grech, A., Burns, G. and Marsh, H. 2007, Distribution of two species of sea snakes, *Aipysurus laevis* and *Emydocephalus annulatus*, in the southern Great Barrier Reef: Metapopulation dynamics, marine protected areas and conservation. *Coral Reefs* 26(2): 291-307.
127. Limpus, C.J. 2008a, *A biological review of Australian marine turtle species, 1 Loggerhead turtle, Caretta caretta (Linnaeus)*, Environmental Protection Agency, Brisbane.
128. Limpus, C.J. 2008b, *A biological review of Australian marine turtle species, 2 Green turtle, Chelonia mydas (Linnaeus)*, Environmental Protection Agency, Brisbane.
129. Limpus, C.J. 2009a, *A biological review of Australian marine turtle species, 3 Hawksbill turtle, Eretmochelys imbricata*, Environmental Protection Agency, Brisbane.
130. Limpus, C.J. 2008c, *A biological review of Australian marine turtle species, 4 Olive ridley turtle Lepidochelys olivacea (Eschscholtz)*, Environmental Protection Agency, Brisbane.
131. Limpus, C.J. 2007, *A biological review of Australian marine turtle species, 5 Flatback turtle, Natator depressus (Garman)*, Environmental Protection Agency, Brisbane.
132. Limpus, C.J. 2009b, *A biological review of Australian marine turtle species, 6 Leatherback turtle, Dermochelys coriacea (Vandelli)*, Environmental Protection Agency, Brisbane.
133. Limpus, C.J., Miller, J.D., Parmenter, C.J. and Limpus, D.J. 2003, The green turtle, *Chelonia mydas*, population of Raine Island and the northern Great Barrier Reef: 1843-2001, *Memoirs of the Queensland Museum* 49(1): 349-440.
134. Limpus, C.J. 2010, Personal communication, *East Australian population trend and nesting census for the loggerhead turtle 2009 - 2010*.
135. Chaloupka, M.Y., Bjørndal, K.A., Balazs, G.H., Bolten, A.B., Ehrhart, L.M., Limpus, C.J., Suganuma, H., Tröeng, S. and Yamaguchi, M. 2008, Encouraging outlook for recovery of a once severely exploited marine megaherbivore, *Global Ecology and Biogeography* 17(2): 297-304.
136. Fuentes, M.M.P.B., Limpus, C.J., Hamann, M. and Dawson, J. 2010, Potential impacts of projected sea-level rise on sea turtle rookeries, *Aquatic Conservation: Marine and Freshwater Ecosystems* 20(2): 132-139.
137. Fuentes, M.M.P.B., Hamann, M. and Limpus, C.J. 2010, Past, current and future thermal profiles of green turtle nesting grounds: Implications from climate change, *Journal of Experimental Marine Biology and Ecology* 383(1): 56-64.
138. Fuentes, M.M.P.B., Limpus, C.J. and Hamann, M. 2011, Vulnerability of sea turtle nesting grounds to climate change, *Global Change Biology* 17(1): 140-153.
139. Fuentes, M.M.P.B. and Abbs, D. 2010, Effects of projected changes in tropical cyclone frequency on sea turtles, *Marine Ecology Progress Series* 412: 283-292.
140. Read, M.A., Grigg, G.C., Irwin, S.R., Shanahan, D., Engle, K., Thomson, S., Irwin, S., Jackson, R., Warriors, W. and One, W. 2007, Satellite tracking reveals long distance coastal travel and homing by translocated estuarine crocodiles, *Crocodylus porosus*, *PLoS ONE* 2(9): e949.
141. Campbell, H.A., Watts, M.E., Sullivan, S., Read, M.A., Choukroun, S., Irwin, S.R. and Franklin, C.E. 2010, Estuarine crocodiles ride surface currents to facilitate long-distance travel, *Journal of Animal Ecology* 79(5): 955-964.
142. Read, M.A., Miller, J.D., Bell, I.P. and Felton, A. 2004, The distribution and abundance of the estuarine crocodile, *Crocodylus porosus*, in Queensland, *Wildlife Research* 31(5): 527-534.
143. Read, M.A. 2012, Estuarine crocodile, in *Queensland's Threatened Animals*, eds L.K. Curtis, A.J. Dennis, K.R. McDonald, P.M. Kyne and S.J.S. Debus, CSIRO Publishing, Collingwood, pp. 185-186.
144. Sullivan, S., Holden, J. and Williams, C. 2010, *Report on the distribution and abundance of the estuarine crocodile, Crocodylus porosus, in Queensland. Waterways of the populated east coast area - research conducted September 2009 to February 2010*, Department of Environment and Resource Management, Brisbane.

145. Congdon, B.C., Erwin, C.A., Peck, D.R., Baker, B.G., Double, M.C. and O'Neill, P. 2007, Vulnerability of seabirds on the Great Barrier Reef to climate change, in *Climate change and the Great Barrier Reef: a vulnerability assessment*, eds J.E. Johnson and P.A. Marshall, Great Barrier Reef Marine Park Authority and Australian Greenhouse Office, Townsville, pp. 427-463.
146. Hulsman, K., O'Neill, P.O. and Stokes, T. 1997, *Current status and trends of seabirds on the Great Barrier Reef*, Great Barrier Reef Marine Park Authority, Townsville.
147. Hulsman, K., O'Neill, P., Stokes, T. and Warnett, M. 1997, Threats, status, trends and management of seabirds on the Great Barrier Reef, Vol 1, in *The Great Barrier Reef, science, use and management*, eds J. Campbell and C. Dalliston, Great Barrier Reef Marine Park Authority, Townsville, pp. 164-177.
148. King, B.R. 1993, The status of Queensland seabirds, *Corella* 17(3): 65-92.
149. Batianoff, G.N. and Cornelius, N.J. 2005, Birds of Raine Island: population trends, breeding behaviour and nesting habitats, *Proceedings of the Royal Society of Queensland* 112: 1-29.
150. Dyer, P.K. 2003, A decline in the number of wedge-tailed shearwaters breeding on Raine Island, *Corella* 27(1): 13-17.
151. Dyer, P.K., O'Neill, P. and Hulsman, K. 2005, Breeding numbers and population trends of wedge-tailed shearwater (*Puffinus pacificus*) and black noddy (*Anous minutus*) in the Capricorn Cays, southern Great Barrier Reef, *Emu* 105: 249-257.
152. Devney, C.A., Short, M. and Congdon, B.C. 2009a, Sensitivity of tropical seabirds to El Niño precursors, *Ecology* 90(5): 1175-1183.
153. Devney, C.A., Short, M. and Congdon, B.C. 2009b, Cyclonic and anthropogenic influences on tern populations, *Wildlife Research* 36: 368-378.
154. De'ath, G. 2001, 'Population changes from 1984-2001 in a seabird colony at Michaelmas Cay, Queensland, unpublished report to the Great Barrier Reef Marine Park Authority'.
155. Peck, D.R., Smithers, B.V., Krockenberger, A.K. and Congdon, B.C. 2004, Sea surface temperature constrains wedge-tailed shearwater foraging success within breeding seasons, *Marine Ecology Progress Series* 281: 259-266.
156. Smithers, B.V., Peck, D.R., Krockenberger, A.K. and Congdon, B.C. 2003, Elevated sea-surface temperature, reduced provisioning and reproductive failure of wedge-tailed shearwaters (*Puffinus pacificus*) in the southern Great Barrier Reef, Australia, *Marine and Freshwater Research* 54(8): 973-977.
157. Fisk, G., Hodge, J., Hulse, L., Steven, A., Turner, L., Waldron, J. and Zeller, B. 2003, The coastal zone, in *State of the environment Queensland 2003*, ed. Environmental Protection Agency, Environmental Protection Agency, Brisbane, pp. 644-673.
158. Heatwole, H., O'Neill, P., Jones, M. and Preker, M. 1996, *Long-term population trends of seabirds on the Swain Reefs, Great Barrier Reef*, CRC Reef Research Centre, Townsville.
159. Fuller, R.A. and Dhanjal-Adams, K.L. 2012, 'Monitoring seabirds in the Great Barrier Reef: a power analysis. Report to Queensland Department of Environment and Resource Management'.
160. Batianoff, G.N., Naylor, G.C., Olds, J. and Neldner, V.J. 2009, Distribution patterns, weed incursions and origins of terrestrial flora at the Capricorn-Bunker Islands, Great Barrier Reef, Australia, *Cunninghamia* 11(1): 107-121.
161. Jaquemet, S., LeCorre, M. and Weimerskirch, H. 2004, Seabird community structure in a coastal tropical environment: importance of associations with sub-surface predators and fish aggregating devices, *Marine Ecology Progress Series* 268: 281-292.
162. Olsen, P. 2008, *The state of Australia's birds 2008*, Birds Australia, Carlton.
163. Nebel, S., Porter, J.L. and Kingsford, R.T. 2008, Long-term trends of shorebird populations in eastern Australia and impacts of freshwater extraction, *Biological Conservation* 141: 971-980.
164. Geering, A., Agnew, L. and Harding, S. 2007, *Shorebirds of Australia*, CSIRO Publishing, Collingwood.
165. Noad, M.J., Dunlop, R.A., Paton, D. and Kniest, H. 2011, *Abundance estimates of the east Australian humpback whale population: 2010 survey and update*, University of Queensland, Brisbane.
166. Noad, M.J., Dunlop, R.A., Cato, D. and Paton, D. 2008a, *Abundance estimates of the east Australian humpback whale population: Final report for the Australian Department of the Environment, Water, Heritage and the Arts*, University of Queensland, Brisbane.
167. Noad, M.J., Dunlop, R.A., Paton, D. and Cato, D.H. 2008b, *Unpublished report, An update of the east Australian humpback whale population (E1) rate of increase*, International Whaling Commission Scientific Committee, Santiago, Chile.
168. Lawler, I.R., Parra, G.J. and Noad, M.J. 2007, Vulnerability of marine mammals in the Great Barrier Reef to climate change, in *Climate change and the Great Barrier Reef: a vulnerability assessment*, eds J.E. Johnson and P.A. Marshall, Great Barrier Reef Marine Park Authority and Australian Greenhouse Office, Townsville, pp. 497-513.
169. Noad, M.J., Paton, D. and Cato, D.H. 2006, Absolute and relative abundance estimates of Australian east coast humpback whales (*Megaptera novaeangliae*), *Journal of Cetacean Research and Management* (Special Issue 3): 243-252.
170. Parra, G.J. 2005, *Behavioural ecology of Irrawaddy, Orcaella brevirostris* (Owen in Gray, 1866), and Indo-Pacific humpback dolphins, *Sousa chinensis* (Osbeck, 1765), in northeast Queensland, Australia: A comparative study, PhD thesis, James Cook University, Cairns.
171. Parra, G.J., Azuma, C., Preen, A.R., Corkeron, P.J. and Marsh, H. 2002, Distribution of Irrawaddy dolphins, *Orcaella brevirostris*, in Australian waters, *Raffles Bulletin of Zoology* 10(Suppl.): 141-154.
172. Cagnazzi, D., Harrison, P.L., Ross, G.J.B. and Lynch, P. 2011, Abundance and site fidelity of Indo-Pacific humpback dolphins in the Great Sandy Strait, Queensland, Australia, *Marine Mammal Science* 27(2): 255-281.
173. Corkeron, P.J., Morissette, N.M., Porter, L. and Marsh, H. 1997, Distribution and status of hump-backed dolphins, *Sousa chinensis*, in Australian waters, *Asian Marine Biology* 14: 49-59.
174. Corkeron, P.J. 1997, The status of cetaceans in the Great Barrier Reef Marine Park, in *Proceedings of the State of the Great Barrier Reef World Heritage Area workshop: 27-29 November 1995, Townsville, Queensland*, eds D. Wachenfeld, J. Oliver and K. Davis, Great Barrier Reef Marine Park Authority, Townsville, pp.28-287.
175. Karczmarski, L., Cockcroft, V.G. and McLachlan, A. 1999, Group size and seasonal pattern of occurrence of humpback dolphins *Sousa chinensis* in Algoa Bay, South Africa, *South African Journal of Marine Science* 21: 89-97.
176. Parra, G.J., Corkeron, P.J. and Marsh, H. 2006, Population sizes, site fidelity and residence patterns of Australian snubfin and Indo-Pacific humpback dolphins: Implications for conservation, *Biological Conservation* 129(2): 167-180.
177. Parra, G.J., Corkeron, P.J. and Marsh, H. 2004, The Indo-Pacific humpback dolphin, *Sousa chinensis* (Osbeck 1765) in Australian waters: a summary of current knowledge, *Aquatic Mammals* 30(1): 197-206.
178. Greenland, J.A. and Limpus, C.J. 2008, *Marine wildlife stranding and mortality database annual report 2007, II Cetacean and Pinniped*, Environmental Protection Agency, Brisbane.
179. Greenland, J.A. and Limpus, C.J. 2007, *Marine wildlife stranding and mortality database annual report 2006, II Cetacean and pinniped*, Environmental Protection Agency, Brisbane.
180. Greenland, J.A. and Limpus, C.J. 2006, *Marine wildlife stranding and mortality database annual report 2005, II Cetacean and pinniped*, Environmental Protection Agency, Brisbane.
181. Greenland, J.A., Limpus, C.J. and Brieze, I. 2005, *Marine wildlife stranding and mortality database annual report*



2004, *II Cetacean and pinniped*, Environmental Protection Agency, Brisbane.

182. Cockcroft, V.G., DeKock, A.C., Lord, D.A. and Ross, G.J.B. 1989, Organochlorines in bottlenose dolphins *Tursiops truncatus* from the east coast of South Africa, *South African Journal of Marine Science* 8(1): 207-217.

183. Bowater, R.O., Norton, J., Johnson, S., Hill, B., O'Donoghue, P. and Prior, H. 2003, Toxoplasmosis in Indo-Pacific humpbacked dolphins (*Sousa chinensis*), from Queensland, *Australian Veterinary Journal* 81(10): 627-632.

184. Jefferson, T.A., Hung, S.K. and Lam, P.K.S. 2006, Strandings, mortality and morbidity of Indo-Pacific humpback dolphins in Hong Kong, with emphasis on the role of organochlorine compounds, *Journal of Cetacean Research and Management* 8(2): 181-193.

185. Ross, G.J.B. 2006, *Review of the conservation status of Australia's smaller whales and dolphins*, Australian Government, Canberra.

186. Fury, C.A. and Harrison, P.L. 2011, Impact of flood events on dolphin occupancy patterns, *Marine Mammal Science* 27(3): E185-E205.

187. Murdoch, E.M., Reif, J.S., Mazzoil, M., McCulloch, S.D., Fair, P.A. and Bossart, G.D. 2008, Lobomycosis in bottlenose dolphins (*Tursiops truncatus*) from the Indian River Lagoon, Florida: estimation of prevalence, temporal trends and spatial distribution, *EcoHealth* 5(5): 289-297.

188. VanParijs, S.M. and Corkeron, P.J. 2001, Boat traffic affects the acoustic behaviour of Pacific humpback dolphins *Sousa chinensis*, *Journal of the Marine Biological Association of the United Kingdom* 81: 533-538.

189. Lukoschek, V. and Chilvers, B.L. 2008, A robust baseline for bottlenose dolphin abundance in coastal Moreton Bay: a large carnivore living in a region of escalating anthropogenic impacts, *Wildlife Research* 35(7): 593-605.

190. Sobotzick, S., Hagihara, R., Grech, A. and Marsh, H. 2012, *Aerial survey of the urban coast of Queensland to evaluate the response of the dugong population to the widespread effects of the extreme weather events of the summer of 2010-11, Final report to the Australian Marine Mammal Centre and the National Environmental Research Program*, James Cook University, Townsville.

191. Marsh, H., Hodgson, A., Lawler, I.R., Grech, A. and Delean, S. 2007, *Condition, status and trends and projected futures of the dugong in the northern Great Barrier Reef and Torres Strait: Including identification and evaluation of the key threats and evaluation of available management options to improve its status*, Report to the Marine and Tropical Sciences Research Facility, Reef and Rainforest Research Centre, Cairns.

192. Pollock, K.H., Marsh, H., Lawler, I.R. and Alldredge, M.W. 2006, Estimating animal abundance in heterogeneous environments: an application to aerial surveys for dugongs, *Journal of Wildlife Management* 70(1): 255-262.

193. Marsh, H. and Lawler, I.R. 2001, *Dugong distribution and abundance in the southern Great Barrier Reef Marine Park and Hervey Bay: Results of an aerial survey in October-December 1999*, Great Barrier Reef Marine Park Authority, Townsville.

194. Marsh, H. and Lawler, I.R. 2006, *Dugong distribution and abundance on the urban coast of Queensland: A basis for management: Final report to Marine and Tropical Research Sciences Facility interim projects 2005-6*, James Cook University, Townsville.

195. Marsh, H. and Lawler, I.R. 2002, *Dugong distribution and abundance in the northern Great Barrier Reef Marine Park - November 2000*, Great Barrier Reef Marine Park Authority, Townsville.

196. Marsh, H., De'ath, G., Gribble, N.A. and Lane, B. 2001, *Shark control records hindcast serious decline in dugong numbers off the urban coast of Queensland*, Great Barrier Reef Marine Park Authority, Townsville.

197. Hodgson, A.J. and Marsh, H. 2007, Response of dugongs to boat traffic: The risk of disturbance and displacement, *Journal of Experimental Marine Biology and Ecology* 340(1): 50-61.

198. Preen, T. 2001, *Dugongs, boats, dolphins and turtles in the Townsville-Cardwell region and recommendations for a boat traffic management plan for the Hinchinbrook Dugong Protection Area*, Great Barrier Reef Marine Park Authority, Townsville.

199. Erftemeijer, P.L. and Lewis, R.R. 2006, Environmental impacts of dredging on seagrasses: A review, *Marine Pollution Bulletin* 52(12): 1553-1572.

200. Biddle, T.M., Boyle, M. and Limpus, C.J. 2011, *Marine wildlife stranding and mortality database annual report 2009 and 2010: Dugong*, Department of Environment and Resource Management, Brisbane.

201. Haynes, D.B., Brodie, J., Waterhouse, J., Bainbridge, Z., Bass, D. and Hart, B. 2007, Assessment of the water quality and ecosystem health of the Great Barrier Reef (Australia): conceptual models, *Environmental Management* 40(6): 993-1003.

202. Haynes, D., Carter, S., Gaus, C., Muller, J. and Dennison, W. 2005, Organochlorine and heavy metal concentrations in blubber and liver tissue collected from Queensland (Australia) dugong (*Dugong dugon*), *Marine Pollution Bulletin* 51(1-4): 361-369.

203. Hutchings, P.A., Haynes, D., Goudkamp, K. and McCook, L.J. 2005, Catchment to reef: water quality issues in the Great Barrier Reef Region—an overview of papers, *Marine Pollution Bulletin* 51(1-4): 3-8.

204. Haynes, D. 2001, *Pesticide and heavy metal concentrations in Great Barrier Reef sediment, seagrass and dugongs (Dugong dugon)*, PhD, University of Queensland, St. Lucia, Brisbane.

205. Lough, J. 2007, Climate and climate change on the Great Barrier Reef, in *Climate change and the Great Barrier Reef: A Vulnerability Assessment*, eds J.E. Johnson and P.A. Marshall, Great Barrier Reef Marine Park Authority and Australian Greenhouse Office, Townsville, pp. 15-50.

206. Great Barrier Reef Marine Park Authority 2013a, *Great Barrier Reef Biodiversity Conservation Strategy 2013*, GBRMPA, Townsville.

207. McCook, L.J., Folke, C., Hughes, T., Nyström, M., Obura, D. and Salm, R. 2007, Ecological resilience, climate change and the Great Barrier Reef, in *Climate Change and the Great Barrier Reef: A vulnerability assessment*, eds J.E. Johnson and P.A. Marshall, Great Barrier Reef Marine Park Authority and Australian Greenhouse Office, Townsville, pp. 75-96.

208. Brodie, J., Fabricius, K.E., De'ath, G. and Okaji, K. 2005, Are increased nutrient inputs responsible for more outbreaks of crown-of-thorns starfish? An appraisal of the evidence, *Marine Pollution Bulletin* 51(1-4): 266-278.

209. Sweatman, H. 2008, No-take reserves protect coral reefs from predatory starfish, *Current Biology* 18(14): R598-R599.

210. Thompson, A., Costello, P., Davidson, J., Schaffelke, B., Uthicke, S. and Liddy, M. 2013, *Reef Rescue Marine Monitoring Program. Report of AIMS activities: inshore coral reef monitoring 2012. Report for Great Barrier Reef Marine Park Authority*, Australian Institute of Marine Science, Townsville.

211. Walker, B. and Salt, D. 2006, *Resilience thinking: sustaining ecosystems and people in a changing world*, Island Press, Washington DC.

212. Fabricius, K.E., De'ath, G., McCook, L.J., Turak, E. and Williams, D.M. 2005, Changes in algal, coral and fish assemblages along water quality gradients on the inshore Great Barrier Reef, *Marine Pollution Bulletin* 51(1-4): 384-398.

213. McKenzie, L.J., Collier, C. and Waycott, M. 2012b, *Reef Rescue Marine Monitoring Program: nearshore seagrass, annual report for the sampling period 1st July 2010 - 31st May 2011*, Fisheries Queensland, Cairns.

214. Great Barrier Reef Marine Park Authority 2012b, *A vulnerability assessment for the Great Barrier Reef: Seagrass*, GBRMPA, Townsville.

215. Great Barrier Reef Marine Park Authority Unpublished report, 'A vulnerability assessment for the Great Barrier Reef: Dugongs'.

216. Department of Environment and Heritage Protection 2012a, *Marine wildlife strandings*, DEHP, Brisbane,

<[http://www.ehp.qld.gov.au/wildlife/caring-for-wildlife/marine\\_strandings.html](http://www.ehp.qld.gov.au/wildlife/caring-for-wildlife/marine_strandings.html)>.

217. Department of Environment and Heritage Protection 2012b, *Back on Track Actions for Biodiversity documents*, DEHP, Brisbane, <<http://www.ehp.qld.gov.au/wildlife/prioritisation-framework/bot-biodiversity-documents.html>>.

218. Great Barrier Reef Marine Park Authority 2012c, *A vulnerability assessment for the Great Barrier Reef: Indo-Pacific humpback and Australian snubfin dolphins*, GBRMPA, <[http://www.gbrmpa.gov.au/\\_data/assets/pdf\\_file/0020/21746/gbrmpa-VA-Snub-IPhumpback-11-7-12.pdf](http://www.gbrmpa.gov.au/_data/assets/pdf_file/0020/21746/gbrmpa-VA-Snub-IPhumpback-11-7-12.pdf)>.

219. Cagnazzi, D., Parra, G.J., Westley, S. and Harrison, P.L. 2013, At the heart of the industrial boom: Australian snubfin dolphins in the Capricorn Coast, Queensland, need urgent conservation action, *PLoS ONE* 8(2): e56729.

220. Parra, G.J., Schick, R. and Corkeron, P.J. 2006, Spatial distribution and environmental correlates of Australian snubfin and Indo-Pacific humpback dolphins, *Ecography* 29(3): 396-406.

221. Cagnazzi, D. 2010, *Conservation status of Australian snubfin dolphin, Orcaella heinsohni, and Indo-Pacific humpback dolphin, Sousa chinensis, in the Capricorn Coast, Central Queensland, Australia*, PhD, Southern Cross University, NSW.

222. Department of Sustainability, Environment, Water, Population and Communities 2009, *Death or injury to marine species following capture in beach meshing (nets) and drum lines used in Shark Control Programs*, DSEWPaC, Canberra, <<http://www.environment.gov.au/biodiversity/threatened/kt/shark-control-programs.html#attachment>>.

223. Fury, C.A. and Reif, J.S. 2012, Incidence of pox-virus like lesions in two estuarine dolphin populations in Australia: links to flood events, *Science of The Total Environment* 416: 536-540.

224. Macdonald, D. and Rushton, S. 2003, Modelling space use and dispersal of mammals in real landscapes: a tool for conservation, *Journal of Biogeography* 30(4): 607-620.

225. Mitchell, M. and Powell, R. 2004, A mechanistic home range model for optimal use of spatially distributed resources. *Ecological Modelling*, 177: 209-232.

226. Berg Soto, A. 2012, *Evaluation of different management approaches to reduce the bycatch of the Indo-Pacific humpback dolphin (Sousa chinensis) and the Australian snubfin dolphin (Orcaella heinsohni) in Queensland, Australia*, PhD thesis, James Cook University, Townsville.

227. Grech, A., Sheppard, J.K. and Marsh, H. 2011, Informing species conservation at multiple scales using data collected for marine mammal stock assessments, *PLoS ONE* 6(3): e17993.

228. Marsh, H., De'ath, G., Gribble, N.A. and Lane, B. 2005, Historical marine population estimates: triggers or targets for conservation? The dugong case study, *Ecological Applications* 15(2): 481-492.

229. Great Barrier Reef Marine Park Authority 2011a, *A vulnerability assessment for the Great Barrier Reef: threadfin salmon*, GBRMPA, Townsville.

230. Great Barrier Reef Marine Park Authority 2011b, *A vulnerability assessment for the Great Barrier Reef: grey mackerel*, GBRMPA, Townsville.

231. Evans, R.D., Williamson, D.H., Sweatman, H., Russ, G.R., Emslie, M.J., Cheal, A.J. and Miller, I.R. 2006, *Surveys of the effects of rezoning of the GBR Marine Park in 2004 on some fish species: preliminary findings*, Marine and Tropical Sciences Research Facility, Townsville.

232. Department of Agriculture Fisheries and Forestry 2013, *Stock status of Queensland's fisheries resources 2012 summary*, DAFF, Brisbane.

233. Great Barrier Reef Marine Park Authority 2012d, *A vulnerability assessment for the Great Barrier Reef: Sharks and rays*, GBRMPA, Townsville.

234. Department of Primary Industries and Fisheries 2006, *A report on the Queensland Shark Safety Program*, DPIF, Brisbane.

235. Chin, A., Tobin, A.J., Simpfendorfer, C. and Heupel, M. 2012, Reef sharks and inshore habitats: patterns of occurrence and implications for vulnerability, *Marine Ecology Progress Series* 460: 115-125.

236. Harry, A.V., Tobin, A.J., Simpfendorfer, C.A., Welch, D.J., Mapleston, A., White, J., Williams, A.J. and Stapley, J. 2011, Evaluating catch and mitigating risk in a multispecies, tropical, inshore shark fishery within the Great Barrier Reef World Heritage Area, *Marine and Freshwater Research* 62(6): 710-721.

237. Tobin, A.J., Simpfendorfer, C.A., Mapleston, A., Currey, L., Harry, A.V., Welch, D.J., Ballagh, A.C., Chin, A., Szczecinski, N., Schlaff, A., White, J. and Moore, B. 2010, *A quantitative ecological risk assessment of sharks and finfish of the Great Barrier Reef World Heritage Area inshore waters: a tool for fisheries and marine park managers. Identifying species at risk and potential mitigation strategies*, Marine and Tropical Sciences Research Facility, Cairns.

238. Pears, R.J., Morison, A.K., Jebreen, E.J., Dunning, M.C., Pitcher, C.R., Courtney, A.J., Houlden, B. and Jacobsen, I.P. 2012b, *Ecological risk assessment of the East Coast Otter Trawl Fishery in the Great Barrier Reef Marine Park: Technical report*, Great Barrier Reef Marine Park Authority, Townsville.

239. White, J., Heupel, M.R., Simpfendorfer, C.A. and Tobin, A.J. 2013, Shark-like batoids in Pacific fisheries: prevalence and conservation concerns, *Endangered Species Research* 19: 277-284.

240. Whiteway, T., Smithers, S., Potter, A. and Brooke, B. 2013, *Geological and geomorphological features of Outstanding Universal Value in the Great Barrier Reef World Heritage Area: Technical report prepared for the Department of Sustainability, Environment, Water, Population and Communities*, Geoscience Australia and James Cook University, Townsville.

241. Department of the Environment 2013, *Australian National Shipwreck Database*, Department of the Environment, Canberra, <<http://www.environment.gov.au/heritage/shipwrecks/database.html>>.

242. Barry, G. 2012a, *Transcript of an interview with a Traditional Owner from Mamu Country*.

243. Garibaldi, A. and Turner, N.J. 2004, Cultural keystone species: Implications for ecological conservation and restoration, *Ecology and Society* 9(3): Part 1.

244. Barry, G. 2012b, *Transcript of an interview with Woppaburra Traditional Owners*.

245. Great Barrier Reef Marine Park Authority 2008, *Story place - information on traditional connections to sea*, GBRMPA, Brisbane, <[http://www.gbrmpa.gov.au/corp\\_site/info\\_services/library/resources/story\\_place](http://www.gbrmpa.gov.au/corp_site/info_services/library/resources/story_place)>.

246. Great Barrier Reef Marine Park Authority 2012e, *Recreation management strategy for the Great Barrier Reef Marine Park*, GBRMPA, Townsville.

247. Deloitte Access Economics 2013, *Economic contribution of the Great Barrier Reef*, Great Barrier Reef Marine Park Authority, Townsville.

248. Office of Economic and Statistical Research 2007, *Queensland Household Survey Summary Report: Prepared for Great Barrier Reef Marine Park Authority*, Queensland Treasury, Brisbane.

249. Rolfe, J., Gregg, D., and Tucker, G., 2011, *Valuing local recreation in the Great Barrier Reef, Australia, Research Report No. 102 August 2011*, Crawford School of Economics and Government, Australian National University, Canberra.

250. Office of Economic and Statistical Research 2012, *Estimated population by local government area*, Queensland Treasury and Trade, Brisbane, <<http://www.oesr.qld.gov.au/subjects/demography/population-estimates/tables/erp-lga-qld/index.php>>.

251. Context Pty Ltd 2013, *Defining the aesthetic values of the Great Barrier Reef*, Department of Sustainability, Environment, Water, Population and Communities, Canberra.

252. Lawrence, K., van Putten, I. and Fernbach, M. 2010, *Profiles of recreational use of the Great Barrier Reef Marine Park*, Great Barrier Reef Marine Park Authority, Townsville.

253. Queensland Government (Unpublished data), 'Vessel Registration Data by LGA for December 2000, 2006 and 2011. Vessel Registration Data by Postcode for 2007, 2009, 2011. Data retrieved from Department of Transport and Main Roads (DTMR)



Transport Registration and Integrated Licensing System (TRAILS)'.

254. Stoeckl, N., Brodie, J., Larson, S., and Prideaux B. 2012, *Project 10.2: Socio economic systems and reef resilience, milestone report*, Reef and Rainforest Research Centre on behalf of the NERP Tropical Ecosystems Hub, Townsville.
255. Great Barrier Reef Marine Park Authority 2012f, *Environmental management charge*, GBRMPA, Townsville, <<http://www.gbrmpa.gov.au/zoning-permits-and-plans/environmental-management-charge>>.
256. Great Barrier Reef Marine Park Authority 2012g, 'Visitor satisfaction survey - ReefHQ Aquarium. Unpublished data'.
257. Murphy, L., Moscardo, G. and Schurmann, A. 2012, *Reef Guardian Campaign Evaluation Report, Stage 2*, Great Barrier Reef Marine Park Authority, Townsville.
258. Great Barrier Reef Marine Park Authority 2013b, *Great Barrier Reef Region strategic assessment stakeholder and community engagement report*, Unpublished, Townsville.
259. Marshall, N.A. and Tobin, R. 2012, "More than meets the eye": The social and economic impacts of recent natural disasters on marine resource dependent industries of the Great Barrier Reef Region, Great Barrier Reef Marine Park Authority, Townsville.
260. Parks Forum 2008, *Value of parks: inspire, refresh, conserve, protect, play*, Parks Forum, Fitzroy, Vic.
261. Ridgway, K. and Hill, K. 2012, East Australian Current, in *A marine climate change impacts and adaptation report card for Australia 2012*, eds E.S. Poloczanska, A.J. Hobday and A.J. Richardson, CSIRO, Canberra, pp. 47-60.
262. Steinberg, C. 2007, Impacts of climate change on the physical oceanography of the Great Barrier Reef, in *Climate change and the Great Barrier Reef: a vulnerability assessment*, eds J.E. Johnson and P.A. Marshall, Great Barrier Reef Marine Park Authority and Australian Greenhouse Office, Townsville, pp. 51-74.
263. Williams, K.J., Dunlop, M., Bustamante, R.H., Murphy, H.T., Ferrier, S., Wise, R.M., Liedloff, A., Skewes, T.D., Harwood, T.D., Kroon, F., Williams, R.J., Joehnk, K., Crimp, S., Smith, M.S., James, C. and Booth, T. 2012, *Queensland's biodiversity under climate change: Impacts and adaptation – synthesis report*, CSIRO Climate Adaptation Flagship, Canberra.
264. Abbs, D. 2012, *The impact of climate change on the climatology of tropical cyclones in the Australia region*, Working Paper No. 11, CSIRO Climate Adaptation Flagship.
265. Delean, S. and De'ath, G. 2008, *Spatial and temporal patterns of indicators of reef health on the Great Barrier Reef. Report to the Marine and Tropical Sciences Research Facility*, Reef and Rainforest Research Centre, Cairns.
266. Fuentes, M.M.P.B., Bateman, B.L. and Hamann, M. 2011, Relationship between tropical cyclones and the distribution of sea turtle nesting grounds, *Journal of Biogeography* 38: 1886-1896.
267. Great Barrier Reef Marine Park Authority 2011c, *Extreme weather and the Great Barrier Reef*, GBRMPA, Townsville.
268. Callaghan, J. and Power, S. 2011, Variability and decline in the number of severe tropical cyclones making land-fall over eastern Australia since the late nineteenth century, *Climate Dynamics* 37: 647-662.
269. Knutson, T.R., McBride, J.L., Chan, J., Emanuel, K., Holland, G., Landsea, C., Held, I., Kossin, J.P., Srivastava, A.K. and Sugi, M. 2010, Tropical cyclones and climate change, *Nature Geoscience* 3: 157-163.
270. Kuleshov, Y., Fawcett, R., Qi, L., Trewin, B., Jones, D., McBride, J. and Ramsay, H. 2010, Trends in tropical cyclones in the South Indian Ocean and the South Pacific Ocean, *Journal of Geophysical Research* 115: doi:10.1029/2009JD012372.
271. Williams, K.J. and Crimp, S. 2012, *Queensland's biodiversity under climate change: An overview of climate change in Queensland, Climate Adaptation Flagship Working Paper No 12A*, CSIRO Climate Adaptation Flagship, Canberra.
272. Intergovernmental Panel on Climate Change 2012, Summary for policymakers, in *Managing the risks of extreme events and disasters to advance climate change adaptation. A special report of Working Groups I and II of the Intergovernmental Panel on Climate Change*, eds C.B. Field, V. Barros, T.F. Stocker, D. Qin, D.J. Dokken, K.L. Ebi, M.D. Mastrandrea, K.J. Mach, G. Plattner, S.K. Allen, M. Tignor and P.M. Midgley, Cambridge University Press, Cambridge, UK and New York, NY, USA, pp. 3-21.
273. Troccoli, A., Muller, K., Coppin, P., Davy, R., Russell, C. and Hirsch, A. 2011, Long-term wind speed trends over Australia, *Journal of Climate* 25(1): 170-183.
274. Reef Water Quality Protection Plan Secretariat 2011, *Great Barrier Reef First Report Card: 2009 baseline, Reef Water Quality Protection Plan*, Reef Water Quality Protection Plan Secretariat, Brisbane.
275. Department of the Premier and Cabinet 2011, *Interim activity report and performance outlook - Reef Water Quality Protection Plan*, Reef Water Quality Protection Plan Secretariat, Brisbane.
276. Department of the Premier and Cabinet 2013a, *Great Barrier Reef Report Card 2011, Reef Water Quality Protection Plan*, Reef Water Quality Protection Plan Secretariat, Brisbane.
277. Church, J.A., White, N.J., Hunter, J.R. and McInnes, K.L. 2012, Sea level, in *A Marine Climate Change Impacts and Adaptation Report Card for Australia 2012*, eds E.S. Poloczanska, A.J. Hobday and A.J. Richardson, CSIRO, Canberra, pp. 27-46.
278. Hobday, A.J. and Lough, J.M. 2011, Projected climate change in Australian marine and freshwater environments, *Marine and Freshwater Research* 62(9): 1000-1014.
279. Department of Climate Change and Energy Efficiency 2010, *Climate change risks to Australia's coast: A first pass national assessment*, DCCEE, Canberra.
280. Department of Climate Change and Energy Efficiency 2011, *Climate change risks to coastal buildings and infrastructure: A supplement to the First Pass national assessment*, DCCEE, Canberra.
281. Lough, J., Gupta, A.S. and Hobday, A.J. 2012, Temperature, in *A marine climate change: impacts and adaptation report card for Australia 2012*, eds E.S. Poloczanska, A.J. Hobday and A.J. Richardson, CSIRO, Canberra.
282. Hale, R., Calosi, P., McNeill, L., Mieszkowska, N. and Widdicombe, S. 2011, Predicted levels of future ocean acidification and temperature rise could alter community structure and biodiversity in marine benthic communities, *Oikos* 120: 661-674.
283. Steffen, W. and Hughes, L. 2013, *The critical decade 2013: Climate change science, risks and responses*, Climate Commission Secretariat, Canberra.
284. Kamrowski, R.L., Limpus, C.J., Moloney, J. and Hamann, M. 2012, Coastal light pollution and marine turtles: Assessing the magnitude of the problem, *Endangered Species Research* 19: 85-98.
285. Reef Water Quality Protection Plan Secretariat 2013, *Great Barrier Reef Report Card 2011: Reef Water Quality Protection Plan*, Reef Water Quality Protection Plan Secretariat, Brisbane.
286. Department of Environment and Heritage Protection 2013, *Reefwise Farming*, DEHP, Brisbane, <<http://www.reefwisefarming.qld.gov.au/>>.
287. Department of the Premier and Cabinet 2013b, *2013 Scientific Consensus Statement - Land use impacts on Great Barrier Reef water quality and ecosystem condition*, DPC, Brisbane.
288. Doropoulos, C., Ward, S., Diaz-Pulido, G., Hoegh-Guldberg, O. and Mumby, P.J. 2012, Ocean acidification reduces coral recruitment by disrupting intimate larval-algal settlement interactions, *Ecology Letters* 15(4): 338-346.
289. Howard, W.R., Nash, M., Anthony, K., Schmutter, K., Bostock, H., Bromhead, D., Byrne, M., Currie, K., Diaz-Pulido, G., Eggs, S., Ellwood, M., Eyre, B., Haese, R., Hallegraeff, G., Hill, K., Hurd, C., Law, C., Lenton, A., Matear, R., McNeil, B., McCulloch, M., Muller, M.N., Munday, P., Opdyke, B., Pandolfi, J.M., Richards, R., Roberts, D., Russell, B.D., Smith, A.M., Tilbrook, B., Waite, A. and Williamson, J. 2012, Ocean acidification, in *A Marine Climate Change Impacts and Adaptation Report Card for Australia 2012*, eds E.S.

Poloczanska, A.J. Hobday and A.J. Richardson, CSIRO, Canberra.

290. Pandolfi, J.M., Connolly, S.R., Marshall, D.J. and Cohen, A.L. 2011, Projecting coral reef futures under global warming and ocean acidification, *Science* 333(6041): 418-422.

291. Schaffelke, B., Carleton, J., Costello, P., Davidson, J., Doyle, J., Furnas, M., Gunn, K., Skuza, M., Wright, M. and Zagorskis, I. 2012, *Reef Rescue Marine Monitoring Program. Final report of AIMS activities 2011/12: Inshore water quality monitoring. Produced for the Great Barrier Reef Marine Park Authority*, Australian Institute of Marine Science, Townsville.

292. Danovaro, R., Corinaldesi, C., Dell'Anno, A., Fuhrman, J.A., Middelburg, J.J., Noble, R.T. and Suttle, C.A. 2011, Marine viruses and global climate change, *FEMS Microbiology Reviews* 35(6): 993-1034.

293. Harvell, C.D., Jordan-Dahlgren, E., Merkel, S., Rosenberg, E., Raymundo, L., Smith, G., Weil, G. and Willis, B. 2007, Coral disease, environmental drivers and the balance between coral and microbial associates, *Oceanography* 20(1): 36-59.

294. Harvell, C.D., Mitchell, C.E., Ward, J.R., Altizer, S., Dobson, A.P., Ostfeld, R.S. and Samuel, M.D. 2002, Climate warming and disease risks for terrestrial and marine biota, *Science* 296(5576): 2158-2162.

295. Haapkylä, J., Unsworth, R.K.F., Flavell, M., Bourne, D.G., Schaffelke, B. and Willis, B.L. 2011, Seasonal rainfall and runoff promote coral disease on an inshore reef, *PLoS ONE* 6(2): e16893.

296. Haapkylä, J., Melbourne-Thomas, J., Flavell, M. and Willis, B.L. 2010, Spatiotemporal patterns of coral disease prevalence on Heron Island, Great Barrier Reef, Australia, *Coral Reefs* 29(4): 1035-1045.

297. Uthicke, S. 2004, Overfishing of holothurians: Lessons from the Great Barrier Reef, in *Advances in sea cucumber aquaculture and management*, eds A. Lovatelli, C. Conand, S. Purcell, S. Uthicke, J.F. Hamel and A. Mercier, Food and Agriculture Organization of the United Nations, Rome, Italy, pp. 163-171.

298. Uthicke, S., Welch, D. and Benzie, J.A.H. 2004, Slow growth and lack of recovery in overfished holothurians on the Great Barrier Reef: evidence from DNA fingerprints and repeated large-scale surveys, *Conservation Biology* 18(5): 1395-1404.

299. Anthony, K. 2000, Enhanced particle-feeding capacity of corals on turbid reefs (Great Barrier Reef, Australia), *Coral Reefs* 19: 59-67.

300. Broadhurst, M.K., Millar, R.B., Brand, C.P. and Uhlmann, S.S. 2009, Modified sorting technique to mitigate the collateral mortality of trawled school prawns (*Metapenaeus macleayi*), *Fishery Bulletin* 107(3): 286-297.

301. Waterhouse, J. 2010, *Optimising water quality and impact monitoring, evaluation and reporting programs, Synthesis report prepared for the Marine and Tropical Sciences Research Facility with contributions from K. Fabricius, R. Pearson, A. Arthington, J. Wallace, M. Sheaves, M. Waycott, L. McKenzie, P. Kuhnert, J. Brodie, S. Lewis, Z. Bainbridge, M. van Grieken, C. Robinson, T. Lynam and their research teams*, Reef and Rainforest Research Centre, Cairns.

302. Brodie, J., De'ath, G., Devlin, M., Furnas, M. and Wright, M. 2007, Spatial and temporal patterns of near-surface chlorophyll a in the Great Barrier Reef lagoon, *Marine and Freshwater Research* 58(4): 342-353.

303. Brodie, J., Schroeder, T., Rohde, K., Faithful, J., Masters, B., Dekker, A., Brando, V. and Maughan, M. 2010, Dispersal of suspended sediments and nutrients in the Great Barrier Reef lagoon during river-discharge events: conclusions from satellite remote sensing and concurrent flood-plume sampling, *Marine and Freshwater Research* 61(6): 651-664.

304. Cheal, A.J., MacNeil, M.A., Cripps, E., Emslie, M.J., Jonker, M., Schaffelke, B. and Sweatman, H. 2010, Coral-macroalgal phase shifts or reef resilience: links with diversity and functional roles of herbivorous fishes on the Great Barrier Reef, *Coral Reefs* 29(4): 1005-1015.

305. Kirkman, H. 1997, *Seagrasses of Australia (Australia: State of the environment technical paper series (Estuaries and the Sea))*, Department of the Environment, Canberra.

306. Collier, C. and Waycott, M. 2009, *Drivers of change to seagrass distributions and communities on the Great Barrier Reef, literature review and gaps analysis. Report to the Marine and Tropical Sciences Research Facility*, Reef and Rainforest Research Centre, Cairns.

307. Hughes, T.P., Bellwood, D.R., Folke, C.S., McCook, L.J. and Pandolfi, J.M. 2007b, No-take areas, herbivory and coral reef resilience, *Trends in Ecology and Evolution* 22: 1-3.

308. Choat, J.H., Robbins, W.D. and Clements, K.D. 2004, The trophic status of herbivorous fishes on coral reefs, *Marine Biology* 145(3): 445-454.

309. Jompa, J. and McCook, L.J. 2002a, The effects of nutrients and herbivory on competition between a hard coral (*Porites cylindrica*) and a brown alga (*Lobophora variegata*), *Limnology and Oceanography* 47: 527-534.

310. Jompa, J. and McCook, L.J. 2002b, Effects of competition and herbivory on interactions between a hard coral and a brown alga, *Journal of Experimental Marine Biology and Ecology* 271: 25-39.

311. McCook, L.J. 1999, Macroalgae, nutrients and phase shifts on coral reefs: scientific issues and management consequences for the Great Barrier Reef, *Coral Reefs* 18: 357-367.

312. McCook, L.J. 1996, Effects of herbivores and water quality on *Sargassum* distribution on the central Great Barrier Reef: cross-shelf transplants, *Marine Ecology Progress Series* 139: 179-192.

313. McCook, L.J. 1997, Effects of herbivory on zonation of *Sargassum* spp. within fringing reefs of the central Great Barrier Reef, *Marine Biology* 129: 713-722.

314. Ceccarelli, D. and Ayling, T. 2010, *Role, importance and vulnerability of top predators on the Great Barrier Reef: a review*, Great Barrier Reef Marine Park Authority, Townsville.

315. Russ, G.R., Cheal, A.J., Dolman, A.M., Emslie, M.J., Evans, R.D., Miller, I.R., Sweatman, H. and Williamson, D.H. 2008, Rapid increase in fish numbers follows creation of world's largest marine reserve network, *Current Biology* 18(12): R514-R515.

316. Hohman, T.C., McNeil, P.L. and Muscatine, L. 1982, Phagosome-lysosome fusion inhibited by algal symbionts of *Hydra viridis*, *Journal of Cell Biology* 94: 56-63.

317. Oliver, T.A. and Palumbi, S.R. 2011, Many corals host thermally resistant symbionts in high-temperature habitat, *Coral Reefs* 30(1): 241-250.

318. Császár, N.B.M., Ralph, P.J., Frankham, R., Berkelmans, R. and van Oppen, M.J.H. 2010, Estimating the potential for adaptation of corals to climate warming, *PLoS ONE* 5(3): e9751.

319. Mieog, J.C. 2009, *Flexibility of the coral-algal symbiosis in the face of climate change: Investigating the adaptive bleaching hypothesis*, PhD thesis, University of Groningen, Groningen, Netherlands.

320. Mieog, J.C., Olsen, J.L., Berkelmans, R., Bleuler-Martinez, S., Willis, B.L. and van Oppen, M.J.H. 2009, The roles and interactions of symbiont, host and environment in defining coral fitness, *PLoS ONE* 4(7): e6364.

321. Stat, M., Loh, W.K., LaJeunesse, T.C., Hoegh-Guldberg, O. and Carter, D.A. 2009, Stability of coral-endosymbiont associations during and after a thermal stress event in the southern Great Barrier Reef, *Coral Reefs* 28: 709-710-713.

322. van Oppen, M.J.H., Baker, A.C., Coffroth, M.A. and Willis, B.L. 2009, Bleaching resistance and the role of algal endosymbionts, in *Coral bleaching: patterns, processes, causes and consequences*, eds M.J.H. van Oppen and J.M. Lough, Springer, Berlin, pp. 83-102.

323. Mao-Jones, J., Ritchie, K.B., Jones, L.E. and Ellner, S.P. 2010, "How microbial community composition regulates coral disease development", *PLoS Biology*, [Online], vol. 8, no. 3, pp. e1000345.

324. Bourne, D.G., Garren, M., Work, T.M., Rosenberg, E., Smith, G.W. and Harvell, C.D. 2009, Microbial disease and the coral holobiont, *Trends in Microbiology* 17(12): 554-562.

325. Littman, R.A., Willis, B.L. and Bourne, D.G. 2009, Bacterial communities of juvenile corals infected with different



*Symbiodinium* (dinoflagellate) clades, *Marine Ecology Progress Series* 389: 45-59.

326. van Oppen, M.J.H., Leong, J. and Gates, R.D. 2009, Coral-virus interactions: A double-edged sword? *Symbiosis* 47: 1-8.

327. Jones, A.M., Berkelmans, R., van Oppen, M.J.H., Mieog, J.C. and Sinclair, W. 2008, A community change in the algal endosymbionts of a scleractinian coral following a natural bleaching event: Field evidence of acclimatization, *Proceedings of the Royal Society B-Biological Sciences* 275(1641): 1359-1365.

328. van Oppen, M.J.H., Souter, P., Howells, E.J., Heyward, A. and Berkelmans, R. 2011b, Novel genetic diversity through somatic mutations: fuel for adaptation of reef corals? *Diversity* 3(3): 405-423.

329. Lewis, J., Hoole, D. and Chappell, L.H. 2003, Parasitism and environmental pollution: parasites and hosts as indicators of water quality, *Parasitology* 126(7): S1-S3.

330. Fearon, R.J. and Cameron, A.M. 1997, Preliminary Evidence Supporting the Ability of Hermatypic Corals to Affect Adversely Larvae and Early Settlement Stages of Hard Coral Competitors, *Journal of chemical ecology* 23(7): 1769-1780.

331. McCook, L.J., Jompa, J. and Diaz-Pulido, G. 2001, Competition between corals and algae on coral reefs: a review of evidence and mechanisms, *Coral Reefs* 19(4): 400-417.

332. McCook, L.J. 2001, Competition between corals and algal turfs along a gradient of terrestrial influence in the nearshore central Great Barrier Reef, *Coral Reefs* 19: 419-425.

333. Norström, A.V., Nyström, M., Lokrantz, J. and Folke, C. 2009, Alternative states on coral reefs: Beyond coral-macroalgal phase shifts, *Marine Ecology Progress Series* 376: 295-306.

334. Jones, G., Almany, G.R., Russ, G., Sale, P., Steneck, R., van Oppen, M.J.H. and Willis, B.L. 2009, Larval retention and connectivity among populations of corals and reef fishes: history, advances and challenges, *Coral Reefs* 28(2): 307-325.

335. Howells, E., van Oppen, M.J.H. and Willis, B.L. 2009, High genetic differentiation and cross-shelf patterns of genetic diversity among Great Barrier Reef populations of *Symbiodinium*, *Coral Reefs* 28(1): 215-225.

336. Souter, P., Willis, B.L., Bay, L.K., Caley, M.J., Muirhead, A. and van Oppen, M.J.H. 2010, Location and disturbance affect population genetic structure in four coral species of the genus *Acropora* on the Great Barrier Reef, *Marine Ecology Progress Series* 416: 35-45.

337. Devlin, M.J. and Waterhouse, J. 2010, *Improved understanding of biophysical and socio-economic connections between catchment and reef ecosystems: wet and dry tropics case studies*, Reef and Rainforest Research Centre, Cairns.

338. Almany, G.R., Connolly, S.R., Heath, D.D., Hogan, J.D., Jones, G.P., McCook, L.J., Mills, M., Pressey, R.L. and Williamson, D.H. 2009, Connectivity, biodiversity conservation, and the design of marine reserve networks for coral reefs, *Coral Reefs* 28: 339-351.

339. Waycott, M. and McKenzie, L.J. 2010, *Final report project 1.1.3 to the Marine and Tropical Sciences Research Facility: Condition, trend and risk in coastal habitats: Seagrass indicators, distribution and thresholds of potential concern*, Reef and Rainforest Research Centre, Cairns.

340. Thompson, A., Costello, P., Davidson, J., Logan, M., Schaffelke, B., Uthicke, S. and Takahashi, M. 2011, *Reef Rescue Marine Monitoring Program, Report of AIMS Activities – Inshore coral reef monitoring 2011, Report for Great Barrier Reef Marine Park Authority*, Australian Institute of Marine Science, Townsville.

341. Munday, P.L., Dixon, D.L., McCormick, M.I., Meekan, M., Ferrari, M.C.O. and Chivers, D.P. 2010, Replenishment of fish populations is threatened by ocean acidification, *Proceedings of the National Academy of Sciences* 107(29): 12930.

342. Anthony, K., Kleypas, J.A. and Gattuso, J.P. 2011, Coral reefs modify their seawater carbon chemistry—implications for impacts of ocean acidification, *Global Change Biology* 17(12): 3655-3666.

343. Fabricius, K.E., Langdon, C., Uthicke, S., Humphrey, C., Noonan, S., De'ath, G., Okazaki, R., Muehlehner, N., Glas,

M. and Lough, J.M. 2011, Losers and winners in coral reefs acclimatized to elevated carbon dioxide concentrations, *Nature Climate Change* 1: 165-169.

344. Pennisi, E. 2009, Calcification rates drop in Australian reefs, *Science* 323(5910): 27.

345. Cooper, T.F., De'ath, G., Fabricius, K.E. and Lough, J.M. 2008, Declining coral calcification in massive *Porites* in two nearshore regions of the northern Great Barrier Reef, *Global Change Biology* 14(3): 529-538.


346. Hutchings, P. and Hoegh-Guldberg, O. 2008, Calcification, erosion and the establishment of the framework of coral reefs, in *The Great Barrier Reef: biology, environment and management*, eds P. Hutchings, O. Hoegh-Guldberg and M. Kingsford, CSIRO Publishing, Collingwood, pp. 74-84.

347. Lough, J. 2008, Coral calcification from skeletal records revisited, *Marine Ecology Progress Series* 373: 257-264.

348. UNESCO Intergovernmental Committee for the Protection of the World Cultural and Natural Heritage 2012, *Operational guidelines for the implementation of the World Heritage Convention*, UNESCO, <<http://whc.unesco.org/archive/opguide12-en.pdf>>.

349. Department of Defence, Environmental Protection Agency, Great Barrier Reef Marine Park Authority and Environment Australia 1999, *Information sheet on Ramsar Wetlands (RIS)*, Environment and Heritage Section, Department of Defence, Canberra.

350. O'Neill, P. 2009, *State of the Environment Report for Shoalwater Bay Training Area 2008*, Department of Defence, Canberra.






## Chapter 8

---

### **Management effectiveness — an independent assessment**



<b>8.6</b>	<b>Effectiveness of specific measures to address impacts .....</b>	<b>8–29</b>
8.6.1	Understanding impacts and considering issues.....	8–30
8.6.2	Managing impacts .....	8–31
8.6.3	Providing certainty.....	8–32
8.6.4	Halt and reverse declines.....	8–44
8.6.5	Adapting to climate change .....	8–44
8.6.6	Integrating with government programs .....	8–45
8.6.7	Meeting Australia’s international obligations .....	8–46
8.6.8	Monitoring, evaluating and reporting .....	8–47
<b>8.7</b>	<b>Effectiveness of specific measures to protect values .....</b>	<b>8–48</b>
8.7.1	Identifying values and determining their condition and trend.....	8–48
8.7.2	Monitoring, evaluating and reporting of values.....	8–49
<b>8.8</b>	<b>Effectiveness of specific measures to protect matters of national environmental significance .....</b>	<b>8–50</b>
<b>8.9</b>	<b>Summary of conclusions .....</b>	<b>8–55</b>
	<b>References.....</b>	<b>8–56</b>

## Table of contents

<b>8.1</b>	<b>Background .....</b>	<b>8–3</b>
<b>8.2</b>	<b>Assessment method .....</b>	<b>8–4</b>
8.2.1	Impacts and values assessed .....	8–6
8.2.2	Calculation of grades .....	8–7
8.2.3	Traditional Owner and stakeholder input.....	8–9
<b>8.3</b>	<b>Protecting values .....</b>	<b>8–10</b>
8.3.1	Biodiversity protection.....	8–10
8.3.2	Indigenous and historic heritage values .....	8–11
8.3.3	Community benefits .....	8–14
<b>8.4</b>	<b>Assessing external impacts .....</b>	<b>8–15</b>
8.4.1	Climate change and extreme weather .....	8–15
8.4.2	Water quality protection (catchment run-off) .....	8–17
8.4.3	Coastal development (protection of coastal ecosystems) .....	8–18
<b>8.5</b>	<b>Managing direct use.....</b>	<b>8–19</b>
8.5.1	Tourism.....	8–19
8.5.2	Fishing – commercial.....	8–21
8.5.3	Fishing – recreational .....	8–22
8.5.4	Recreation .....	8–23
8.5.5	Port activities .....	8–24
8.5.6	Shipping.....	8–26
8.5.7	Defence activities.....	8–27
8.5.8	Research activities.....	8–28

**Cover page image:** Scott’s fairy wrasse (*Cirrhitilabrus scottorum*).



## **Extract from Great Barrier Reef Region Strategic Assessment terms of reference**

### **4.1 Current Program**

4.1.1 Describe and assess the effectiveness of the Program to:

- a. *identify the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area, and determine their current condition and trend, including spatial and non-spatial approaches*
- b. *identify and analyse direct, indirect, consequential and cumulative impacts, including the methods used to determine these types of impacts*
- c. *consider environmental, social, cultural and economic issues*
- d. *avoid, mitigate, offset and adaptively manage impacts*
- e. *address uncertainty and risk*
- f. *provide certainty regarding where uses may occur, the type of activities allowed, conditions under which activities may proceed and circumstances where impacts are likely to be unacceptable*
- g. *halt and reverse any declines and enhance the condition of the relevant matters of national environmental significance, including mechanisms to deliver a 'net benefit' to the condition of the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area*
- h. *adapt to reasonable climate change scenarios*
- i. *integrate with related local, Queensland and Australian government programs to protect and manage the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area*
- j. *meet Australia's international responsibilities in relation to the environment and protection of world heritage*
- k. *monitor, evaluate and report on the:*
  - i. *condition and trends of the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area*
  - ii. *impacts of activities, including the setting of targets to benchmark management performance.*

*Note: In this report 'current Program' is referred to as 'current management arrangements'.*

## **8.1 Background**

The current management arrangements of the Great Barrier Reef Marine Park Authority (the Authority) are described in Chapter 3.

This chapter provides a summary of key findings from the independent assessment of management effectiveness undertaken by Professor Marc Hockings (UniQuest Pty Ltd), Dr Andrea Leverington (UniQuest Pty Ltd) and Brian Gilligan. It describes:

- assessment method
- the effectiveness of the Authority's management arrangements to:
  - protect values relevant to the matters of national environmental significance (biodiversity, Indigenous and historic heritage values and community benefits)
  - address external impacts (climate change, water quality and coastal development)
  - manage direct use
  - meet the criteria set out in item 4.1.1 of the terms of reference for the strategic assessment (see Appendix 2)
  - protect each of the matters of national environmental significance.

The terms of reference for the independent review are contained in Appendix 4, and its full report is available on the Authority's website at [www.gbrmpa.gov.au](http://www.gbrmpa.gov.au). In this chapter, direct excerpts from the independent assessment report are shown in italics. The independent assessment report was prepared in early 2013, based on information available at that time. Notes about significant changes to management since then have been provided.

Outcomes from the independent assessment inform recommendations for management improvements (see Chapter 12) and the Program Report.

## 8.2 Assessment method

Management effectiveness evaluation is defined as the assessment of how well protected areas are being managed — primarily the extent to which they are protecting values and achieving goals and objectives.

The independent assessment used the International Union for Conservation of Nature (IUCN) World Commission on Protected Areas framework for evaluating management effectiveness<sup>1</sup>, which has been widely applied around the world to assess management effectiveness in a range of circumstances. The framework (Figure 8.1) focuses on six management elements and the links between them. Within the framework, effective management:

- begins with understanding the **context** of the protected area, including its values, the threats it faces and opportunities available, its stakeholders, and the management and political environment
- progresses through **planning**: establishing vision, goals, objectives and strategies to conserve values and reduce threats
- allocates **inputs** (resources) of staff, money and equipment to work towards the objectives
- implements management actions according to accepted **processes**
- eventually produces **outputs** (goods and services) which should usually be outlined in management plans and work plans
- results in impacts or **outcomes**, hopefully achieving defined goals and objectives.




Figure 8.1 Framework for assessing management effectiveness of protected areas<sup>1</sup>

The assessment system was adapted from that used in the *Great Barrier Reef Outlook Report 2009*<sup>2</sup>. The key difference is that this evaluation focused on the management activities within the Authority's jurisdiction and any joint management arrangements with the Queensland Government (for example, joint permitting arrangements and the Field Management Program), whereas the assessment conducted for the Outlook Report 2009 considered all relevant management activities across the Australian and Queensland governments.

In accordance with items 4 and 8 of the terms of reference for the independent assessment, it also:

- took more explicit account of matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area; direct, indirect, consequential and cumulative impacts; and condition and trend of the values underpinning matters of national environmental significance
- considered whether the Authority's management arrangements provide certainty regarding where uses may occur, the types of activities allowed, conditions under which activities may proceed and circumstances where impacts are likely to be unacceptable
- considered jurisdictional consistency of management arrangements to achieve outcomes
- assessed the effectiveness of the range of management tools (Table 8.1) used to protect and manage the values.

**Table 8.1 Management tools considered in assessing management effectiveness**

Management tool	Purpose
<b>Great Barrier Reef Marine Park Act and Regulations</b>	The <i>Great Barrier Reef Marine Park Act 1975</i> and Regulations govern the protection and management of the Great Barrier Reef Marine Park. They provide for the Zoning Plan and plans of management and govern permitting decisions.
<b>Zoning Plan</b>	Provides control of use (predominantly extractive activities) and, to a lesser extent, access within the Great Barrier Reef Marine Park. Establishes the need for permits for some uses in the Marine Park, such as tourism, infrastructure and research. There are complementary arrangements in adjacent areas under Queensland jurisdiction.
<b>Plans of management</b>	Set out specific arrangements for areas, species, ecological communities or activities (for example, Cairns Area and Whitsundays Plans of Management). They complement zoning and permitting arrangements. Some components are legally binding.
<b>Permits</b>	Facilitate opportunities for use of the Great Barrier Reef. Permits are issued mainly for marine tourism, research, harvest fisheries, dredging, dumping and infrastructure (for example, jetties and marinas) and include detailed environmental impact assessments. Matched in adjacent areas of Queensland jurisdiction, generally providing a joint permit. Fisheries licences are issued and managed by the Queensland Government.
<b>Fees and charges</b>	Three main fees and charges apply in the Marine Park: <ul style="list-style-type: none"> <li>• The cost of assessing an application for a permit for commercial activities is partly recovered through payment of a permit application assessment fee.</li> <li>• The environmental management charge applies to some commercial activities operating under a permit issued by the Authority. The revenue is applied to Marine Park management.</li> <li>• Bonds (usually as a bank guarantee) may be held by the Authority to cover the risks associated with a proposed activity.</li> </ul>
<b>Traditional Owner agreements</b>	Formal Traditional Use of Marine Resources Agreements describe how Traditional Owner groups work with Australian and Queensland governments to manage traditional use activities in sea country. Indigenous Land Use Agreements are between one or more native title groups and other people or parties, about the use and management of land and waters.
<b>Compliance</b>	Activities that encourage adherence with legal requirements, both through education and enforcement. Includes formal (for example, the Field Management Program jointly undertaken with the Queensland Government) and informal (for example, the Eyes and Ears Incident Reporting program) activities.
<b>Site infrastructure</b>	On-ground infrastructure installed to better protect the values of individual sites (for example, reef protection markers, public moorings and signs). Implemented and maintained by the Authority and Queensland Parks and Wildlife Service through the Field Management Program.
<b>Policy</b>	Specific arrangements that guide decision makers and the public. Types of policy documents include: strategies, policies, site management arrangements, position statements and guidelines.
<b>Partnerships</b>	Formal arrangements, often executed through a memorandum of understanding or an agreement, to enable a partnership approach to management of the Marine Park (for example, intergovernmental agreement with the Queensland Government, Reef Advisory Committees, Local Marine Advisory Committees, memorandum of understanding with a government authority and partnership with Ecotourism Australia).
<b>Stewardship and best practice</b>	Voluntary arrangements with stakeholders that provide the opportunity for contributions to protection and management (for example, Reef Guardian programs, Pro-vision Reef Stewardship Action Plan, best environmental practices).
<b>Education and community awareness</b>	Programs to inform and motivate members of the community about the Great Barrier Reef and its protection and management, including ways they can contribute (for example, Reef HQ Aquarium, the Authority's website, information sheets, zoning maps).
<b>Research and monitoring</b>	Undertaken, commissioned or partnered by the Authority to better inform decisions on protection and management of the Great Barrier Reef (for example, reef health and impact surveys, Eye on the Reef monitoring and climate change research programs).

### 8.2.1 Impacts and values assessed

While all of the management topics assessed in the Outlook Report 2009 are considered in this assessment, additional topics were included to address the requirements of a strategic assessment conducted under the *Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act). The following 15 management topics were assessed:

#### Values

- biodiversity protection
- heritage (both Indigenous and historic)
- community benefits

#### External impacts on values

- climate change and extreme weather
- water quality protection (catchment run-off)
- coastal development (protection of coastal ecosystems)

#### Direct uses

- tourism (marine-based)
- fishing – commercial
- fishing – recreational
- recreation (non-extractive)
- port activities
- shipping
- defence activities
- research activities.

The scale of the management topics range from localised issues that affect only a small proportion of the total area (for example, defence activities), to others which have implications across all or most of the Region (for example, climate change and extreme weather, recreation, commercial fishing). The management topics were not weighted or graded to account for differences in scale and complexity.

Each management topic was assessed independent of the others to ensure the grades were based on that value, impact or use alone. Consequently, the grades given need to be interpreted in relation to the scale and complexity of the management topic (Table 8.2).



Table 8.2 A comparison of the scale and complexity of management topics assessed

Management topic	Scale	Complexity		
		Social	Bio-physical	Jurisdictional
Values				
Biodiversity protection	Region-wide	minor	major	moderate
Indigenous heritage	Region-wide but variable in intensity	major	moderate	moderate
Historic heritage	Region-wide	moderate	minor	moderate
Community benefits	Region-wide	major	moderate	minor
External impacts on values				
Climate change and extreme weather	Region-wide	major	major	major
Water quality protection	Great Barrier Reef catchment and mainly inshore waters	major	major	major
Coastal development	Coastal areas and mainly inshore waters	major	major	major
Direct uses				
Tourism	Region-wide but variable in intensity	major	moderate	moderate
Fishing – commercial	Region-wide but variable in intensity	moderate	major	moderate
Fishing – recreational	Region-wide but variable in intensity	moderate	major	moderate
Recreation	Region-wide but variable in intensity	major	moderate	moderate
Port activities	Concentrated around ports	moderate	major	major
Shipping	Concentrated around shipping areas	moderate	moderate	moderate
Defence activities	Limited in area and duration	minor	minor	minor
Research activities	Region-wide but limited in intensity	minor	moderate	minor

### 8.2.2 Calculation of grades

Assessment criteria were developed for each of the six management elements of the evaluation framework, with a total of 45 criteria across the elements (Table 8.3). Criteria varied slightly from the Outlook Report 2009 — they were refined to take account of lessons from that assessment and other subsequent assessments. Additional indicators were incorporated to align with the requirements of the strategic assessment's terms of reference. In addition, some criteria were deleted as they were considered not relevant to the strategic assessment or had proved difficult to reliably assess during the Outlook Report process.

Each management topic was assessed against all 45 criteria to determine a grade of management effectiveness. The independent assessors provided a justification for the rating and documented the main evidence they considered in reaching their judgement. These individual ratings were added and then scaled to produce an overall rating for each management element of the framework. A four-point rating scale commonly used in management effectiveness evaluation systems, and adopted in the Outlook Report 2009, was used in this assessment.

**Table 8.3 Assessment criteria to assess effectiveness of management**

A total of 45 criteria guided the calculation of the grades for management effectiveness.

Criteria Number	Criteria Description
<b>Context</b>	
CO1	The values that underpin matters of national environmental significance in the Great Barrier Reef (including outstanding universal value of the Great Barrier Reef World Heritage Area) relevant to.....are understood by managers.
CO2	Direct and indirect impacts associated with.....are understood by managers.
CO3	Consequential and cumulative impacts associated with.....are understood by managers.
CO4	The current condition and trend of matters of national environmental significance (spatial and non-spatial) relevant to.....are known by managers.
CO5	The stakeholders relevant to.....are well known by managers.
<b>Planning</b>	
PL1	There is a planning system in place that effectively addresses.....
PL2	The planning system for.....addresses the major pressures and drivers impacting on the Great Barrier Reef's values.
PL3	Actions for implementation regarding.....are clearly identified within the plan.
PL4	Clear, measurable and appropriate objectives for management of.....have been documented.
PL5	The main stakeholders and/or the local community are effectively engaged in planning to address.....
PL6	Sufficient policy currently exists to effectively address.....
PL7	There is consistency across jurisdictions when planning for.....
PL8	Plans provide certainty regarding where uses may occur, the type of activities allowed, conditions under which activities may proceed and circumstances where impacts are likely to be acceptable.
<b>Inputs</b>	
IN1	Current financial resources are adequate and prioritised to meet management objectives to address.....
IN2	Current human resources within the managing organisations are adequate to meet specific management objectives to address.....
IN3	The right skill sets and expertise are currently available to the managing organisations to address.....
IN4	The necessary biophysical information is currently available to address.....
IN5	The necessary socio-economic information is currently available to address.....
IN6	The necessary traditional (Indigenous) knowledge is currently available to address.....
IN7	There are additional sources of non-government input (e.g. volunteers) contributing to address.....
<b>Processes</b>	
PR1	The main stakeholders and/or industry(ies) are effectively engaged in the ongoing management of.....
PR2	The local community is effectively engaged in the ongoing management of.....
PR3	There is a sound governance system in place to address.....
PR4	There is effective performance monitoring to gauge progress towards the objective(s)
PR5	Appropriate training is available to the managing agencies to address.....
PR6	Management of.....is consistently implemented across the relevant jurisdictions.
PR7	There are effective processes applied to resolve differing views/conflicts regarding.....
PR8	Direct and indirect impacts of activities associated with.....are appropriately considered.
PR9	Consequential and cumulative impacts of activities associated with.....are appropriately considered.

Criteria Number	Criteria Description
PR10	The best available biophysical research and/or monitoring information is applied appropriately to make relevant management decisions regarding.....
PR11	The best available socio-economic research and/or monitoring information is applied appropriately to make relevant management decisions regarding.....
PR12	The best available traditional (Indigenous) knowledge is applied appropriately to make relevant management decisions regarding.....
PR13	Relevant standards are identified and being met regarding.....
PR14	Targets have been established to benchmark management performance.
<b>Outputs</b>	
OP1	To date, the actual management program has (or activities have) progressed in accordance with the planned work program for.....
OP2	Implementation of management documents and/or programs relevant to.....have progressed in accordance with timeframes specified in those documents.
OP3	The results (refer OP1) have achieved their stated management objectives.
OP4	To date, products or services have been produced in accordance with the stated management objectives for.....
<b>Outcomes</b>	
OC1	The relevant managing agencies are to date effectively addressing.....and moving towards the attainment of the desired outcomes.
OC2	The outputs relating to.....are on track to ensure the values of the Great Barrier Reef are protected (refer CO1).
OC3	The outputs (refer OP1 and OP3) for.....are reducing the major risks and the threats to the Great Barrier Reef.
OC4	Use of the Great Barrier Reef relating to.....is demonstrably environmentally sustainable.
OC5	Use of the Great Barrier Reef relating to.....is demonstrably economically sustainable.
OC6	Use of the Great Barrier Reef relating to.....has demonstrably enhanced community understanding and/or enjoyment.
OC7	The relevant managing agencies have developed effective partnerships with local communities and/or stakeholders to address.....

The rating scale was:

- ineffective – zero to 20 per cent of optimal condition
- partially effective – 21 to 50 per cent of optimal condition
- mostly effective – 51 to 80 per cent of optimal condition
- effective – 81 to 100 per cent of optimal condition.

For the outcomes element, two grades were assigned — one for outcomes related to biodiversity, the other for all outcomes, taking into account biodiversity outcomes and those relating to social, economic and management objectives.

### 8.2.3 Traditional Owner and stakeholder input

Members of Local Marine Advisory Committees, Reef Advisory Committees and participants in the regional strategic assessment workshops were asked to provide their views on the strengths and weaknesses of the identified management tools (Table 8.1). The assessment allowed respondents to choose the management topics they wished to address (they could respond to as many or as few as they wished). They were asked to indicate strengths and weaknesses in the Authority's management of the topic and to rate the effectiveness the management tools employed on a four-point scale (very effective, mostly effective, partially effective, not effective). They could also indicate if they had no opinion or believed the tool was not applicable to the topic. Respondents were only asked to assess tools the Authority had indicated were applicable to management of a particular issue (for example, permits are not relevant to recreational fishing). Participants were able to provide this information electronically using a web-based survey.

Ninety-five respondents provided 222 assessments of the management topics. The majority of respondents (69 per cent) were members of Local Marine Advisory Committees, while 16 per cent were members of Reef Advisory Committees. A third of the respondents attended one of the strategic assessment workshops. A small number of respondents were Traditional Owners. The average length of time that respondents had been involved in the Great Barrier Reef was 18.5 years, but the length of experience was highly variable ranging from none to 60 years. Only three respondents lived outside Queensland, while a further 11 lived in Queensland outside the Great Barrier Reef catchment.

The method for undertaking the assessment of management effectiveness was presented at the Local Marine Advisory Committee chairs' workshop in Townsville in October 2012, and at three stakeholder workshops coordinated by the Authority in Townsville, Cairns and Rockhampton in December 2012. The independent reviewers presented the preliminary assessment of management effectiveness and the assessment results concerning the management tools at the stakeholder workshops. Participants were provided an opportunity to discuss outcomes for particular topics.

Feedback from stakeholders has been included in a summary statement for each of the management topics. Full results are available on the Authority's website at [www.gbrmpa.gov.au](http://www.gbrmpa.gov.au).

## 8.3 Protecting values

This section provides a summary of the effectiveness of the Authority's current management arrangements to protect the values of the Region. Direct excerpts from the report are in italics, whereas non-italicised text has been added by the Authority for contextual purposes. A grading table is provided for each management topic (Table 8.4 to Table 8.7).

### 8.3.1 Biodiversity protection

Table 8.4 Management effectiveness scores for biodiversity protection					
Biodiversity protection		Effective	Mostly effective	Partially effective	Ineffective
Context					
Planning					
Inputs					
Processes					
Outputs					
Outcomes	Overall				
	Biodiversity				

*The Authority is the lead agency for managing biodiversity protection in the Region and uses a number of regulatory and non-regulatory tools.*

*Protection of the biodiversity of the Great Barrier Reef Region is the primary objective for much of the management action undertaken in the Great Barrier Reef and its catchment and forms part of the primary objective under the Great Barrier Reef Marine Park Act 1975 (the Act). Management of biodiversity is undertaken using an array of tools, principally the Zoning Plan, but also management plans, permits, policy documents, site management, stewardship, education and best practices. A number of Queensland and Commonwealth agencies also have responsibility for protection of biodiversity in the Region. This potentially complex management regime has been simplified through intergovernmental coordination and cooperation.*

*Evidence has shown a declining trend in coral cover over time,<sup>3</sup> with an estimated 50 per cent decline in coral cover on the Great Barrier Reef over the past 27 years,<sup>4</sup> which is cause for considerable concern. This highlights the importance of considering cumulative and consequential impacts which are currently less well understood by managers, although these*

*issues are now receiving greater attention in assessments and other considerations of the Authority.*

*Threat abatement plans, recovery plans and wildlife conservation plans under the EPBC Act, as well as specific on-the-water actions by the Authority (for example, reef protection markers and special management areas), are in place to address individual biodiversity issues for a small number of species and habitats that occur within the Region. Some of those plans, such as the recovery plan for marine turtles, have not been updated in many years. With regard to iconic and threatened species, such as dugong and some marine turtles, these plans and actions have had some effect in stabilising populations. Status and trend assessments prepared by the Authority indicate that more species are continuing to decline than have stabilised or are increasing resulting in the partially effective biodiversity outcomes. However, status and trend are uncertain for many groups because of lack of data.*

*The information base for biodiversity protection continues to improve through scientific research and the compilation and assessment of information by Authority staff (such as vulnerability assessments and assessment of status of habitats and species). Gaps in knowledge — for example, status and trend for some specific plant and animal groups, habitats and ecosystems — are well recognised. Traditional Owners are increasingly involved in biodiversity management, though their knowledge and understanding is often not available or accessible for decision making.*

*Planning for biodiversity protection has been significantly improved since the Outlook Report 2009 through preparation of the draft Biodiversity Conservation Strategy 2012 and the Informing the outlook for the Great*



Barrier Reef Coastal Ecosystems technical report. However, targets in the draft Biodiversity Conservation Strategy tend to be process and output-focused and should be complemented by additional outcome-focused targets. On a Reef-wide scale, the Zoning Plan and the complementary plans for the adjacent Great Barrier Reef Coast Marine Park have made the most significant contribution to biodiversity protection.

These have provided a robust framework and are already demonstrating positive results.<sup>5</sup> However, the zoning provisions only address biodiversity protection from direct extractive uses, particularly fishing. Major threats to biodiversity, such as climate change, coastal development (with the exception of Commonwealth managed islands) and catchment run-off are not addressed by either the zoning provisions or individual biodiversity protection measures, although programs such as the Reef Plan which are designed to improve water quality will have a positive impact on biodiversity. Many of the actions to address biodiversity are long term and will take time to manifest an improvement in biodiversity outcomes.

Major risks and threats to biodiversity protection are well documented, and risk assessments and management procedures are in place for the major threats. Vulnerability assessments are being prepared for key species and habitats. These provide comprehensive documentation of risks to biodiversity values and mitigation measures within the Region. However, there remains little capacity to track the Authority's resource allocations that specifically target biodiversity objectives or to assess which outputs and outcomes result from management actions. Secure long-term funding for monitoring will be needed to provide data on changes to biodiversity outcomes over time.

Key stakeholders in biodiversity protection have been identified and are generally well known to managers especially through Reef Advisory Committees, Local Marine Advisory Committees and other consultative mechanisms.

#### **Effectiveness of management tools as assessed by stakeholders**

Surveyed stakeholders considered the Zoning Plan to be the most effective of the Authority's management tools used to manage biodiversity conservation in the Marine Park. Zoning, the Great Barrier Reef Marine Park Act and Regulations, education and community awareness, plans of management, and stewardship and best practice were considered to be effective or mostly effective by more than 50 per cent of the respondents. Permits and compliance were seen as the weakest tools in biodiversity protection.

Zoning, along with education and community awareness, were singled out as management strengths. Problems raised by jurisdictional limitations on the Authority to manage activities in the coastal zone and commercial fisheries were commonly cited as weaknesses. Recommendations for improvement included increased field presence and compliance action with greater penalties for breaches.

Workshop participants acknowledged that the Authority undertook an extensive biodiversity protection program; however, they expressed concern that the work being undertaken was not sufficient to protect the values of the reef.

### **8.3.2 Indigenous and historic heritage values**

#### **Indigenous heritage values**

The Authority has a lead role in the protection of Indigenous heritage in the Region.

Aboriginal and Torres Strait Islander peoples are the Traditional Owners of the Great Barrier Reef Region. There are more than 70 Aboriginal and Torres Strait Islander Traditional Owner clan groups that maintain heritage values for their land and sea country. These values may be cultural, spiritual, economic, social or physical, and demonstrate continuing connections with the Reef and its natural resources.

Traditional Owners have recognised inherent rights and interests over their sea country in which the Authority has a management responsibility. Key management measures for Indigenous heritage include partnerships, education and community awareness, stewardship and best practice, legislation and Traditional Use of Marine Resources Agreements and Indigenous Land Use Agreements.

The Authority acknowledges the Convention on Biological Diversity that states in Article 10<sup>6</sup> Sustainable Use of Components of Biological Diversity that each contracting party shall, as far as possible and as appropriate (among other things):

Indigenous heritage		Effective	Mostly effective	Partially effective	Ineffective
Context					
Planning					
Inputs					
Processes					
Outputs					
Outcomes	Overall				
	Biodiversity				

*“Protect and encourage customary use of biological resources in accordance with traditional cultural practices that are compatible with conservation or sustainable use requirements”.*

*The Authority's knowledge of Indigenous values is improving with the Reef Rescue Land and Sea Country Indigenous Partnerships Program<sup>7</sup> and the development of Traditional Use of Marine Resources Agreements. However, direct and indirect impacts, cumulative impacts associated with traditional use of marine resources, and the less tangible Indigenous heritage such as traditional knowledge and maintenance of cultural practice are not widely available and therefore not well understood nor considered across the Authority. The importance of the Reef in Indigenous economies is also not well understood and not incorporated fully into management.*

*Planning for Indigenous heritage was considered in the Authority's Great Barrier Reef Marine Park Heritage Strategy (Heritage Strategy).<sup>8</sup> Actions are included in this document, although there are no targets or timeframes associated with the actions. The Reef Rescue Land and Sea Country Indigenous Partnerships Program articulates a set of objectives and targets to ensure "the continued use, support and reinvigoration of traditional ecological knowledge to underpin biodiversity conservation". This program will be evaluated at its completion, however to date there has been 100 per cent completion against milestones. The Authority's Strategic Plan<sup>9</sup> also includes specific objectives concerning working with Aboriginal and Torres Strait Islanders to take into account traditional affiliations, culture, heritage values and rights of management in the Marine Park.*

*The Australian Government, under the Caring for our Country initiative, committed \$10 million over five years towards the Reef Rescue Land and Sea Country Indigenous Partnerships Program in 2008. This funding expires in June 2013. [Note: An extension of the Reef Rescue programs was announced after preparation of the independent assessment report.]*

*Identifying Traditional Owners who can speak for country can be difficult. The Authority has established an Indigenous Reef Advisory Committee to bring together expertise and experience in Indigenous partnership initiatives and sea country management from within the Great Barrier Reef and other parts of Australia to provide issues-based advice that informs the operations of the Authority. One of the benefits of the Traditional Use of Marine Resources Agreement program is the relationship between sea country and a Traditional Owner group that is identified and agreed upon as part of the Traditional Use of Marine Resources Agreement development process.*

*Sea Country Plans, such as the Kuku Yalanji example (near Mossman), have been developed by Traditional Owners for their own country. This plan identifies values, planning needs and management outcomes and is funded by the Authority and the Department of Sustainability, Environment, Water, Population and Communities through a small grants program. The Authority's environment impact management assessment processes need to consider these plans to ensure permitted activities are consistent with community plans. The Field Management Program also includes planning, development, engagement of Traditional Owners, training, mentoring, resourcing, monitoring, compliance, planning and reporting.*

*The Authority's framework for Traditional Use of Marine Resources Agreements complements existing community-based measures developed by some Traditional Owner groups to manage their use of some of these resources and recognises entitlements enshrined in the Native Title Act 1993. These agreements promote the sustainable use of threatened species such as dugong and turtle, and iconic species such as barramundi cod and giant clams within the Marine Park. There are currently five Traditional Use of Marine Resources Agreements and one Indigenous Land Use Agreement covering more than 20 per cent of the Great Barrier Reef, engaging 14 Traditional Owner clan groups (jointly signed by the Authority and Queensland Government). [Note: An additional Traditional Use of Marine Resources Agreement was accredited after preparation of the independent assessment report.]*

*While progress has been made by the Authority in engaging with key Indigenous stakeholders in the Great Barrier Reef, further work is needed to develop a mutually agreed and culturally appropriate process for joint planning. An Indigenous cultural heritage strategy would enable a shared vision to be developed with Traditional Owner groups with actions and timeframes for implementation. The strategy should include protocols for managing culturally sensitive information within the Authority and externally.*

#### **Effectiveness of management tools as assessed by stakeholders**

*Traditional Use of Marine Resources Agreements were the only management tool considered to be effective or mostly effective by more than 50 per cent of respondents. Policy and permits, as well as the Great Barrier Reef Marine Park Act and Regulations, were seen as ineffective in addressing Indigenous heritage issues.*

*Respondents generally recognised an improving engagement of the Authority with Traditional Owners, with a better policy and program platform as well as positive results from the development of Traditional Use of Marine Resources Agreements. At the same time, respondents — especially those who are Traditional Owners — recognise there is more to be done, particularly in relation to better informing and involving Traditional Owners in planning and management.*

*Traditional Owners who participated in the stakeholder workshops expressed wide-ranging views on this topic, including the desire for the development of stronger partnerships and greater involvement in decision making and management. The need for education and communication with the wider community was also discussed. The need for acknowledgement of native title rights in the management of the Region was also raised.*

## Historic heritage values

The Authority has a regulatory role in the protection of historic heritage aspects of the environment (however, it has no jurisdiction over Queensland island heritage). The Authority works with other agencies to protect these values.

For the purposes of this report, historic heritage encompasses historic shipwrecks, World War II features and sites, lighthouses and associated structures, historic landscapes, and historic places and structures that embody a specific cultural, historic or scientific value. Natural heritage is considered under biodiversity protection.

There is generally a reasonable understanding by the Authority of the Region's values relevant to historic issues. However, while the Authority has a good understanding of the impacts on natural heritage, the consequential and cumulative impacts associated with historic heritage are less well understood. Information on condition and trend has improved through the 2010–2011 audit of Commonwealth islands' historic heritage. However, little is known about the condition and trend of shipwrecks, World War II sites, or heritage places such as Endeavour Reef.

The Great Barrier Reef Marine Park Act and Regulations provide the legislative power for the protection of historic heritage values and consideration of potential impacts to heritage through the permitting process. There are also strong drivers for heritage management under the Commonwealth and world heritage obligations. These obligations also drive consistency of implementation of historic heritage management across jurisdictions.

The Authority's Heritage Strategy<sup>8</sup>, endorsed in 2005, provides guidance for protection of the Marine Park's heritage values through a range of planning instruments and policies. Historic shipwrecks are protected through specific legislation and entry controls. The Heritage Strategy identifies actions but does not set a timetable for implementation or indicate relative priorities. Risks and threats have been more explicitly addressed in new periodic reporting processes for world heritage sites. The Authority's Heritage Strategy requires updating to effectively address major pressures and drivers impacting on historic heritage.

An assessment of the Heritage Strategy showed reasonable progress was being made, with 65 per cent of the actions completed or good or satisfactory progress having been made towards completion. Thirty-five per cent of actions had not commenced or been considered. Only one heritage management plan is currently in place, although two others are underway. Policy documents to address issues associated with historic heritage require updating and implementation.

Resourcing of the management of historic heritage is poor, with less than one full-time equivalent position dedicated to working in this area. The frameworks for engagement with stakeholders, industry and the community is good, but limited due to staff resources, with expert advice and consultation provided through the Reef Advisory Committees, the Australian Heritage Council, the tourism industry, Local Marine Advisory Committees, the World Heritage Committee and the Queensland Museum.

The Authority is currently developing a heritage register that will capture all values relevant to historic heritage. The register will assist managers and the community in knowing the location of and an understanding of heritage values.

## Effectiveness of management tools as assessed by stakeholders

Insufficient responses were received to evaluate the effectiveness of management tools for historic heritage. Workshop participants raised the issue of confusing federal and state jurisdictions as a barrier to effective management.

**Table 8.6 Management effectiveness scores for historic heritage values**

Historic heritage		Effective	Mostly effective	Partially effective	Ineffective
Context					
Planning					
Inputs					
Processes					
Outputs					
Outcomes	Overall				
	Biodiversity				

### 8.3.3 Community benefits

Table 8.7 Management effectiveness scores for community benefits				
Community benefits	Effective	Mostly effective	Partially effective	Ineffective
Context				
Planning				
Inputs				
Processes				
Outputs				
Outcomes	Overall			
	Biodiversity			

The Authority has a regulatory role in the protection of the environment including the social and economic aspects of the environment. The Authority, together with other agencies, works to adopt an integrated approach to the management of the social, economic and environmental aspects of the Region.

Community benefits of the Reef encompass socio-economic aspects such as employment and income, in addition to less tangible attributes such as understanding, appreciation, enjoyment, personal connection, health benefits and access to the Reef. Many of these attributes are values based.

Management by the Authority is undertaken using a range of measures, including stewardship and best practice, legislation, zoning plans, permits and site infrastructure. While information and understanding of the value of the Reef from an economic perspective is assessed and monitored, an understanding of the broader community benefits that the Region supports is not well understood nor quantified. This has been recognised and the Authority, as part of the strategic assessment, is beginning to seek community views about threats, values and pressures in relation to community benefits as part of the strategic assessment.

There is likely to be increasing conflict between economic benefits of the Reef — associated with port and coastal development — and the personal, recreational and value based benefits. The consequential and cumulative impacts on community benefits [from these developments] have not been well documented or assessed thoroughly.

Many of the pressures associated with community benefits, such as population change, coastal development, economic growth and climate change, are outside the Authority's direct jurisdiction. However, the Authority's Recreation Management Strategy<sup>10</sup> made some progress towards recognising the enjoyment and personal attachment to the Reef. Social, cultural and heritage values are considered when assessing permits applications, but this application is limited by the Authority's knowledge of these aspects in specific locations.

In particular, the Reef Guardian program, Reef HQ Aquarium, and the Recreation Management Strategy have objectives for management associated with community benefits. The Field Management Program, jointly managed with the Queensland Government, also has a strong commitment to ensuring public access to the Reef and islands.

Many of the community benefit issues are considered under programs and policies developed for other purposes such as access to resources, conservation and multiple use. Currently there are no guidelines or benchmarks for social or economic impact assessments for the World Heritage Area. An overarching strategy that outlines the objectives for community benefits, while showing the linkages across the programs, would clarify the Authority's roles and responsibilities. It would also provide an improved framework to assess the management effectiveness with greater accuracy.

Stakeholder engagement through the Reef Advisory Committees, Local Marine Advisory Committees and stakeholders brought together for the strategic assessment enable managers to gain a better understanding of community values and issues of concern. Volunteer programs, such as components of the marine monitoring program (for example, Seagrass Watch) and Eye on the Reef, also provide avenues for community involvement in protecting the Great Barrier Reef.

#### Effectiveness of management tools by stakeholders

The Zoning Plan, the Great Barrier Reef Marine Park Act and Regulations were seen as most effective tools used by the Authority in promoting community benefits, primarily by making the Marine Park accessible to the community. The permits system, stewardship programs such as Reef Guardians, and research and monitoring were seen as very or mostly effective by more than half of the respondents.

Community partnership programs such as Reef Guardians and educational materials were viewed as strengths of the Authority's management. A lack of adequate resources for management was identified as a weakness.

Stakeholders at the workshops commented that the Authority did not prioritise managing community benefits and that with growing coastal populations there was potential for greater conflict between user groups, which was not being addressed.



## 8.4 Assessing external impacts

This section provides a summary of the effectiveness of the Authority's current management arrangements to address external impacts. Direct excerpts from the report are in italics, whereas non-italicised text has been added by the Authority for contextual purposes. A grading table is provided for each management topic (Table 8.8 to Table 8.10).

### 8.4.1 Climate change and extreme weather

*The Authority has had a lead role in facilitating awareness of the impacts from climate change and extreme weather and in building resilience in the Marine Park. It has an advisory role to other agencies in relation to mitigation and adaptation to climate change and extreme weather in the Region.*

*The Authority has limited jurisdictional responsibility for addressing climate change in the broad sense. However, the Authority contributes significantly to the development of international best practice for managing responses to climate change and extreme weather issues as they relate to the Reef's marine ecosystems. This is chiefly done through research and monitoring, partnerships with research institutions, government agencies and stakeholder groups and education, community awareness and stakeholder engagement programs.*

*The Great Barrier Reef Climate Change Adaptation Strategy and Action Plan 2012–2017 (the Action Plan)<sup>11</sup> acknowledges the important role the Authority plays in informing national and international climate policy and providing knowledge to support effective management of inshore areas. However, the direct management role of the Authority is statutorily limited, requiring a focus on ecosystem-based adaptation to increase the resilience of the Reef by reducing impacts from other sources such as land-based sediments and nutrients. Assessing the effectiveness of the Authority in this role is challenging, as so much depends on the actions of others.*

*The Authority's planning initiatives are focused on enhancing the capacity of ecological, socio-economic and management systems to adapt to change in ecological and social variables. Dedicated staff positions have been established and specialist expertise exists within the Authority to coordinate efforts in this area, as well to facilitate national, regional and international collaboration which is necessary to address the complex issues involved. Climate change considerations have now been incorporated into many business areas across the Authority, rather than being confined to one work unit.*

*Work completed from 2007 to 2012 under the Great Barrier Reef Climate Change Action Plan 2007–2012<sup>12</sup> has included raising awareness of the implications of climate change for the Great Barrier Reef and the ecosystem services it provides. Effort has also been focused on building interest in adaptation planning as a way to reduce future risks from climate change and minimise its impact on the environment. This has included work with reef-dependent industries such as tourism and fishing, as well as communities, and considers planning and adaptation actions at all levels from individuals and businesses through to managers and government.<sup>13</sup>*

*Vulnerability assessments<sup>14,15</sup> for specific functions and activities that have the potential to impact on the Region's values continue to provide good contextual information for management of climate change implications. The assessment for the Outlook Report 2009 noted that managers and key community stakeholders recognised that while climate change drivers and influences are largely global in nature, regional and local activities give rise to specific vulnerabilities that require action.*

*Climate change plans and strategies were in place in 2009 and the focus has now moved to implementing the Action Plan. This plan aims to translate objectives into specific policies and measurable actions for on-ground management by the Authority to increase the resilience of the Reef ecosystem. Specific threats such as increasing sea temperatures, ocean acidification, increased storm events and associated changes to freshwater inputs, currents and connectivity have been identified as matters to be addressed by adaptation initiatives that seek to minimise the impacts of these events.*

*A number of the Authority's management initiatives contribute to offsetting the impacts of climate change by reducing risks from other stressors. The Authority's climate change work provides adaptation resources to help minimise the vulnerability of coastal communities and Reef-dependent industries while also increasing their capacity to collaborate in building ecosystem resilience to climate change. From a review of the supporting evidence, implementation and evaluation of the Action Plan appears to be on schedule.*

*While direct and indirect impacts of activities related to climate change and extreme weather are understood by managers — and risk-based approaches have been applied in adaptation work — they are not yet appropriately*

**Table 8.8 Management effectiveness scores for climate change and extreme weather**

Climate change and extreme weather		Effective	Mostly effective	Partially effective	Ineffective
Context		Effective	Mostly effective	Partially effective	Ineffective
Planning		Effective	Mostly effective	Partially effective	Ineffective
Inputs		Effective	Mostly effective	Partially effective	Ineffective
Processes		Effective	Mostly effective	Partially effective	Ineffective
Outputs		Effective	Mostly effective	Partially effective	Ineffective
Outcomes	Overall	Effective	Mostly effective	Partially effective	Ineffective
	Biodiversity	Effective	Mostly effective	Partially effective	Ineffective

considered in all aspects of Reef management. The amendment of policy documents and procedures, along with the development of practical guidelines for relevant permit-holders and applicants, has been identified as priorities for attention.

Work continues on identifying the gaps in available biophysical information. The Authority recognises the particular need to clarify socio-economic factors such as the resource dependency<sup>16</sup> of different stakeholder groups on various marine and coastal natural resources that are likely to be altered under a changing climate. Some of these information gaps are now being addressed. Evaluation of resilience indicators and completion of a reef resilience atlas were identified in 2009 as particularly important to address cumulative impacts, both biophysical and socio-economic.

By 2012, the Authority's climate change work was instrumental in improving understanding and consideration of consequential and cumulative impacts, but this remains a challenge. Critical elements of current condition and trend cannot be confidently determined and monitored by managers. Efforts continue, including work with Traditional Owners, to apply available traditional knowledge to consider climate change implications, particularly through Traditional Use of Marine Resources Agreements.

Community engagement relating to climate change and extreme weather continues through the Authority's initiatives such as Eye on the Reef and the Reef Guardian program which includes schools, councils, fishers and farmers. Work under the Action Plan contributed to substantially strengthened partnerships with key sectors of the commercial fishing industry and the marine tourism industry. A Reef Health incident response system was developed in 2011,<sup>17</sup> in consultation with stakeholders, and includes response plans for:

- coral bleaching
- cyclone
- coral disease
- crown-of-thorns starfish (in preparation)

Case studies of actions that can enhance the Reef's resilience to climate change and extreme weather are being documented by the Authority.

Despite significant progress in building the ecological resilience of the Reef, and the social and economic resilience of Reef industries, the work of the Authority cannot make the Reef and its industries invulnerable to the impacts of extreme weather. A series of extreme weather events such as cyclones, floods and heatwaves between 2002 and 2011 caused significant impacts to corals, seagrasses, dugongs and green turtles, as well as the tourism and fishing industries. This decade of extreme weather provided an indication of what is to come if the planet's climate continues to change at the current rate. The future success of the Authority's work in adapting to climate change and extreme weather is dependent on successful global efforts to mitigate climate change.

The planned and systematic approach being applied gives confidence that the Authority is achieving all that might be reasonably expected to reduce the threats posed to matters of national environmental significance from climate change and extreme weather in the Great Barrier Reef Region. However, in spite of good systems and processes, the long-term trend for Great Barrier Reef ecosystems is still poor, and the extent to which specific initiatives can effectively address particular problems will only become clear over time. This situation highlights the importance of robust performance monitoring and adaptive management by the Authority.

#### **Effectiveness of management tools as assessed by stakeholders**

Research and monitoring was considered to be the most effective of the management tools used by the Authority to manage climate change and extreme weather in the Great Barrier Reef Region. The Great Barrier Reef Marine Park Act and Regulations were considered to be only partially effective, while education and community awareness was mostly effective. Surprisingly, given the effort the Authority has expended in these areas, stewardship and best practice and partnerships were not regarded as particularly effective by a majority of respondents in this survey. However, feedback from evaluation processes conducted in relation to projects with partners in the fishing industry provides positive evidence that the Authority's partnership and stewardship approach is valued and has been effective in progressing climate change and extreme weather related activities and management. This suggests there may be a need to promote awareness about the success of these partnerships. Workshop participants recognised the work that the Authority was undertaking, but felt the Authority lacked jurisdictional responsibility in this area.

### 8.4.2 Water quality protection (catchment run-off)

The Authority has a lead role for the management activities that impact water quality within the Marine Park, as well as an advisory or partnership role with other agencies in relation to activities that occur outside the Region that may impact on the water quality in the Marine Park.

The Authority's management of water quality is through legislation and permits for point source discharges into the Marine Park, as well as undertaking the marine monitoring functions of [the Reef Water Quality Protection Plan] Reef Plan<sup>18</sup>. However, the most significant contribution to water quality decline in the Great Barrier Reef is from activities outside the Marine Park associated with agricultural practices. This limits the Authority's capacity to take direct action with respect to water quality decline. The non-point source discharges, such as run-off from agriculture, is managed through partnerships with the Queensland Government, the Department of Sustainability, Environment, Water, Population and Communities, land holders and industry groups, and through education and community awareness, stewardship and best practice.

The assessment for management effectiveness undertaken for the Outlook Report 2009 concluded there was slow progress being made towards the attainment of the desired outcomes and to reducing the risks and threats to the Reef's values. The Outlook Report 2009 recognised the positive work that was being done through the Reef Plan, but highlighted the little evidence of change through a lack of monitoring, and the lack of delivering with the planning framework.

The values that underpin the matters relevant to water quality are well understood by the Authority. While many of the direct and indirect impacts of poor water quality are well known, knowledge is not as comprehensive concerning the consequential and cumulative impacts of water quality. Current condition and trend are known for specific species such as corals and seagrass; however, despite early evidence of a reduction in loads of pollutants, there is still little documented evidence concerning improvements in water quality in the Region.

Since the Outlook Report 2009, a review of Reef Plan has provided better focus and direction for managers including the Authority, including targets for water quality and land management improvement. The revised Reef Plan reduces the list of actions from 65 to 11 to provide a more strategic and adaptive plan. It is focused on outcomes and takes into account new policy documents and regulatory frameworks. Measureable targets, improved accountability, and coordinated monitoring, evaluation and reporting underpin it. Targets are focused on short and medium-term outcomes for water quality and land management. In addition, Water Quality Guidelines for the Great Barrier Reef Marine Park<sup>19</sup> and the development of a Coastal Ecosystems Assessment Framework<sup>20</sup> by the Authority set limits for water entering the Marine Park and provide a framework for assessment of ecosystem services within the basins located in the catchment with a focus on improving the health and resilience of the Reef. In terms of assessing performance, a Reef Plan monitoring, evaluation and reporting strategy has been developed and the first Reef Plan report card has been published. (Note: since preparation of the independent report, a second and third Reef Plan report card have been published.)

The Authority engages with key stakeholders and has expanded the Reef Guardians program (which commenced with schools and local councils), to include farmers, graziers and fishers. However, as the management of water quality leaving the Great Barrier Reef catchment is significantly within the Queensland Government's jurisdiction, the Authority can only work with the community and stakeholders to encourage best practice land management.

The Authority has its greatest direct influence on water quality through the Great Barrier Reef Marine Park Act and Regulations (Section 66(2)e of the Act) under which it is illegal to discharge waste into the Marine Park except for some permissible actions. Permitting the discharge of wastewater into the Reef is assessed on a case by case basis, against guideline trigger levels. (Note: trigger levels in the Regulations are for aquaculture only.)

The Authority has allocated significant resources into understanding the water quality issues from a biophysical aspect, though information is still limited with respect to the socio-economic impact of loss of ecosystem services from poor water quality. The Authority monitors the long-term health of key marine ecosystems and the condition of water quality in the inshore lagoon. The Australian Government has committed more than \$200 million to the improvement of water quality in the Reef, including water quality grants, Reef Partnerships, Land and Sea Country Indigenous Partnerships, research and development, and monitoring and reporting (including the publication of an annual Great Barrier Reef water quality report card). (Note: the Queensland Government spent \$175 million between 2009 and 2013 and has committed to an additional \$35 million per year through to 2020 under Reef Plan.)

**Table 8.9 Management effectiveness scores for water quality protection (catchment run-off)**

Water quality protection		Effective	Mostly effective	Partially effective	Ineffective
Context					
Planning					
Inputs					
Processes					
Outputs					
Outcomes	Overall				
	Biodiversity				



### Effectiveness of management tools as assessed by stakeholders

Education and community awareness, followed by research and monitoring, were considered to be the most effective of the management tools used to manage water quality in the Great Barrier Reef Marine Park. Compliance and the Great Barrier Reef Marine Park Act and Regulations were seen as the weakest tools in managing water quality.

Reef Plan and the Reef Guardian program are regarded by many respondents as strengths of management. Water quality is seen as an area where effective partnerships are crucial for effective management. A lack of jurisdictional responsibilities, the sheer scale of the issue, and lack of corresponding resources and political will are seen as the major weaknesses and impediments. A number of respondents called for greater attention to compliance with existing Regulations, stiffer penalties for breaches and an extension of the Authority's powers to control activities outside the Marine Park that impact on Great Barrier Reef values.

These findings were supported at the workshops where discussions focused on management issues outside the Authority's jurisdiction, and where participants supported the positive changes achieved by stewardship programs.

### 8.4.3 Coastal development (protection of coastal ecosystems)

Table 8.10 Management effectiveness scores for coastal development (protection of coastal ecosystems)					
Coastal development (protection of coastal ecosystems)		Effective	Mostly effective	Partially effective	Ineffective
Context					
Planning					
Inputs					
Processes					
Outputs					
Outcomes	Overall				
	Biodiversity				

The Authority has an advisory role in the management of coastal development and the protection of coastal ecosystems.

Fourteen broad categories of coastal ecosystems are important to the functioning of the Great Barrier Reef Region: coral reefs, lagoon floors, islands, open water, seagrasses, coastline, estuaries, freshwater wetlands, forested floodplains, heath and shrublands, grass and sedgeland, woodlands, forests and rainforests. These ecosystems provide important links between land, freshwater and marine environments, as well as feeding and breeding grounds for many terrestrial, wetland and marine species. Changes to, or loss of these coastal ecosystems, can lead to a variety of adverse environmental, social and cultural impacts, with serious implications for matters of national environmental significance and for industries and communities dependent on the Great Barrier Reef Region.

Critical habitats in the Great Barrier Reef catchment provide for marine ecosystem functions directly through ecosystem connectivity. They also support healthy marine ecosystems by preventing excess sediments, nutrients and pollutants from being transported to the Reef lagoon. Reef coastal ecosystems support a variety of ecosystem functions, such as nursery areas for marine species, floodwater buffers and sediment and nutrient sinks.

As a result of the Outlook Report 2009, the Authority has developed a comprehensive document, *Informing the outlook for the Great Barrier Reef Coastal Ecosystems*<sup>21</sup> which assessed the pressures affecting coastal ecosystems that have the potential to influence the Reef's health and resilience. This document provides an effective context for management as it describes the functioning, as well as the threats, pressures, risks and trends of the Reef's coastal ecosystems.

The Authority's regulatory role in managing coastal ecosystems is restricted to addressing the impacts of coastal development works such as jetties, marinas and dredging which occur below mean low water (within the Marine Park) or on Authority-managed Commonwealth islands. The Authority has an advisory role to the Department of Sustainability, Environment, Water, Population and Communities on the EPBC Act referrals, for actions likely to have a significant impact on the Great Barrier Reef World Heritage Area and a joint assessment role for activities impacting on matters within the Marine Park. Such actions may be located within or outside the Region. The Authority also works in partnership with Queensland and local governments to provide technical advice and facilitate improved understanding of the values of the World Heritage Area and the critical role coastal ecosystems play in the functioning of the Reef. Additionally, the Authority's stewardship programs, including its Reef Guardians program, focus on working with industry sectors and the broader community to improve understanding of the connectivity between coastal and marine ecosystems and to facilitate improved management practices.

The Queensland Government has jurisdiction over most development and management in the coastal zone (above the mean low water mark). The former Queensland Coastal Protection State Planning Policy has been replaced with the new draft Coastal Protection State Planning Regulatory Provision. This potentially has significant implications for the Great Barrier Reef Region because it removes many of the specific requirements placed on local government and potential developers to undertake best practice and to minimise environmental harm.

It is evident the Authority has a reasonable understanding of the direct and indirect impacts associated with coastal ecosystems, though there has been little quantification of these impacts. The consequential and cumulative impacts require better understanding and monitoring. Urgent action (by the Authority) is also needed



to improve the health of coastal ecosystems to boost the health and resilience of the Reef. Vulnerability assessments for each of the at-risk coastal ecosystems have also been undertaken.

Targets and performance measures for coastal ecosystems included in the Authority's draft Biodiversity Conservation Strategy,<sup>22</sup> lack outcome specific targets. Reef Plan, which focuses on non-point source water quality, also contains a number of performance measures relating to coastal ecosystems. The Authority's stakeholder engagement remains effective.

The Authority's stakeholder engagement on coastal ecosystem management is increasing. The Australian Government's Reef Rescue grants to improve land management practices laid the groundwork for building relationships with farmers. Partnerships continue to grow through the Authority's Reef Guardian Farmers stewardship program. Coastal ecosystems management is also the focus of coastal Reef Guardian Councils and Schools, and is regularly discussed at Local Marine Advisory Committees and Reef Advisory Committees.

The work done in the Authority's draft Biodiversity Conservation Strategy and Informing the outlook for Great Barrier Reef coastal ecosystems suggests the long-term condition and trend for coastal ecosystems is very poor if joint management action is not taken soon to halt and reverse the decline in inshore and coastal ecosystems, particularly south of Port Douglas.

Given the number of agencies involved in managing coastal ecosystems, an integrated framework for planning, assessment and approvals is needed to ensure the values of the Great Barrier Reef Region are maintained and where possible improved.

### Effectiveness of management tools as assessed by stakeholders

Most respondents rated the Authority's management tools for coastal ecosystems as either not effective, or only partially effective. The only tools to be rated as effective or very effective by more than 40 per cent of the respondents were research and monitoring along with education and community awareness.

The work of the Authority's Reef Guardians program with local farmers and other partnership programs were regarded as a strength by respondents, while a lack of influence (by the Authority) on coastal planning and development decisions (and by some, a perceived political will to control development) are regarded as the most significant weaknesses. Respondents see a need to extend the Authority's jurisdictional control over activities that impact on the values of the Marine Park, to improve compliance action (especially in relation to permit conditions and compliance action by state authorities), and an increased focus on partnerships between the Authority, local government and local industry.

Stakeholder comments at workshops reflected the survey results, consistently raising concerns over the Authority's limited influence over the management of coastal ecosystems and the consequential impacts on Reef health and resilience.

## 8.5 Managing direct use

This section provides a summary of the effectiveness of the Authority's current management arrangements for each of the direct use categories. Direct excerpts from the report are in italics, whereas non-italicised text has been added by the Authority for contextual purposes. A grading table is provided for each management topic (Table 8.11 to Table 8.18).

### 8.5.1 Tourism

*The Authority is the lead agency for management of marine tourism in the Region.*

*Management of tourism in the Great Barrier Reef employs a mix of management tools, with emphasis on plans of management for intensively used areas and partnerships with accredited tour operators, as well as permits — often with specific conditions.*

*Tourism within the Reef is recognised by managers as one of the most significant uses of the Region and the presentation of its values. Tourism is acknowledged to be a major driver for economic growth and employment for coastal Queensland. Tourism, commercial fishing, recreation and scientific research contribute \$5.7 billion per annum — of this, \$5.2 billion is attributed to tourism.*<sup>23</sup>

*The Outlook Report 2009 recognised high levels of visitor satisfaction and significant economic benefits to local communities but noted that longer-term environmental performance would depend on clearer understanding of the implications of latent capacity in the permit system and improved documentation of the levels of specific activities undertaken in each location over time. This has been partially*

		Effective	Mostly effective	Partially effective	Ineffective
Tourism	Context				
	Planning				
	Inputs				
	Processes				
	Outputs				
Outcomes	Overall				
	Biodiversity				

improved with the implementation of the Authority's environmental management charge online system and the development of the Reef Management System (a permitting system).

With the launch of the Great Barrier Reef as a National Landscape in March 2012, the Authority sought to reinforce the mutual understanding between tourism operators and managers on values underpinning matters of national environmental significance in the Great Barrier Reef for tourism. The Authority chaired the regional Steering Committee for the Great Barrier Reef National Landscape which has members from Tourism Queensland, Queensland Parks and Wildlife Service, key tourism industry representatives, and the conservation sector.

Plans of management covering less than 10 per cent of the total Great Barrier Reef area are in place for the areas that receive more than 80 per cent of tourist visits but enhanced planning capability within the Authority and a schedule of regular reviews would improve consistency. The Outlook Report 2009 assessment noted some tourism activities occur over a wide area and that the planning system did not include an overarching strategy to guide tourism activities. An overarching strategy, although under development, is still outstanding and an expansion of site planning has not occurred to areas where growth is increasing such as the southern Great Barrier Reef. If site planning is not pre-emptive, protection of the Region's values in these expanding areas will not be as effective.

Policies covering many aspects of tourism are now outdated and work is still needed to address shortcomings identified in 2009. Implementation of the Action Plan, marine tourism contingency planning and development of a co-ordination framework to assist with consistency across jurisdictions are priorities in working towards an overarching strategy for managing tourism in the Region.

Sound governance, industry partnerships and management processes are in place to address tourism issues; the Authority is widely recognised as a world leader in this area. Joint permitting and assessment processes support consistency in the approach across jurisdictions. Most tourists are carried by a small number of highly accredited tour operators. The Authority's permit conditions seek to limit cumulative impacts. The potential problems arising from significant latent capacity within the permit system have been recognised and, at least, partially addressed through the Authority's plans of management, capping permits and a booking system for sensitive sites. While the permitting system manages tourism well, a tourism permit can be more than 16 pages long and contain more than 50 conditions. Whether these permits are effective at informing and educating tourism operators about what they can and can't do — while delivering required outcomes for the environment, social, cultural and heritage values — requires evaluation. A key issue is the complexity of the current management arrangements for tourism. Knowing where you can go and what you can do relies on knowledge of zoning plans, plans of management, complex permissions and best practices. Simplification and alignment of current arrangements through regular ongoing reviews is required.

Since 2012, environmental management charge records can be recorded online ... improving efficiencies for the operators and the government agencies involved. Permit bookings are displayed on the Authority's website, providing a new level of public transparency.

The tourism industry has experienced difficulty in retaining and attracting experienced staff due to industry downturns and competition from the mining sector. This, alongside a reduction in educational opportunities (the reef discovery training program is no longer operational) has generally led to a decline in the delivery of education on World Heritage Area values and the Authority's obligations regarding presentation of these values. However, within the Authority there continues to be high levels of skills related to marine tourism management and impact assessment. Expert advice also continues to be sought through the Tourism and Recreation Reef Advisory Committee and also the Indigenous Reef Advisory Committee. Representatives from the tourism industry are also members of Local Marine Advisory Committees.

The Authority endeavours to build on best available biophysical and monitoring research information to make relevant management decisions, including reef health monitoring information provided by tourism operators through the Eye on the Reef program. This program collects information through various methods including the Sightings Network, the Eyes and Ears Incident Reporting Network, Tourism Weekly Monitoring, Rapid Monitoring and Reef Health and Impact Survey sub-programs. Eye on the Reef has been enhanced and now integrates data and reporting across programs, while having a user-friendly data portal and the provision for online training.

#### **Effectiveness of management tools as assessed by stakeholders**

The Zoning Plan, followed by plans of management, was seen as the Authority's most effective tools in managing tourism. All tools, with the exception of site infrastructure (for example, reef protection markers, public moorings and signage), were seen as very effective by some respondents. More than half of the respondents understood compliance to be only partially effective.

Good industry relationships and strong partnerships with groups such as the Association of Marine Park Tourism Operators, together with the zoning and permit systems and plans of management, are seen as strengths. Nevertheless, respondents identified a need to improve standards in some Reef tourism operations with more experienced and trained interpretive staff. They saw the Authority could play a role in improving standards through the permit and compliance monitoring systems. More regular reviews of plans of management and making permits for commercial tourism operators easier to understand were suggested improvements.

Other areas for improvement identified include:

- more effective engagement at the management level with tourism operators
- reduced bureaucracy and enhanced flexibility in permit processing
- more human resources allocated to speed up permit processing
- streamlining processes for reviewing and amending plans of management
- more compliance activity and 'sticks' to deal with the poor performing operators who are 'damaging the brand'
- innovative policy development rather than continuing rounds of 'document to exhaustion'.

The desirability of the Authority in taking a proactive leadership role in defining the iconic tourism experience, rather than simply responding to proposals by tour operators, was also raised, as was the need for site infrastructure to follow destination development planning rather than being reactive. Workshop participants raised the inflexibility of plans of management to respond to changes in the tourism industry.

## 8.5.2 Fishing — commercial

The Authority has a regulatory role in the management of commercial fishing through its Zoning Plan, Regulations and permits for a limited number of Queensland managed commercial and developmental fisheries. The Authority has an advisory role to other agencies in relation to the management of commercial fishing in the Region.

Fishing is the principal extractive use of the Great Barrier Reef. Commercial fishing targets a range of species including fish, sharks, crabs, lobsters and prawns. Viable commercial fishing industries depend on a healthy ecosystem. The total revenue from commercial fishing in the World Heritage Area in 2010–11 was estimated at \$192 million.<sup>23</sup>

Management of fishing and its dependent aquatic environment is shared between the Australian and Queensland governments. While the Authority has a direct role in ensuring that fishing is ecologically sustainable in the Marine Park it has an advisory role on the management of commercial fisheries, and works with other agencies to improve fisheries management.

The Authority's primary management tools with respect to commercial fishing are the Zoning Plan, associated Regulations, stewardship (Reef Guardian Fishers) and permits for a small number of harvest fisheries and developmental fisheries. Special management areas are in place to regulate fishing practices in areas important to dugongs. These tools complement the Queensland legislation that licences commercial fishing. Compliance, undertaken by the joint Australian and Queensland governments' Field Management Program, also plays a significant role in managing the impacts of commercial fishing.

As identified in the Outlook Report 2009, the Authority has a good understanding of commercial catch information. Cumulative impacts associated with commercial fishing are reasonably well known. The Authority's recent Ecological risk assessment of the East Coast Otter Trawl Fishery in the Great Barrier Reef found the overall ecological risks from trawling are relatively low, but there are some high ecological risks for deep water skates, several rays, and sea-snakes.<sup>24</sup> A reduction in trawl fishing effort of more than 40 per cent between 2005 and 2009, driven by prevailing economic conditions rather than management intervention, has significantly reduced the ecological risk from trawling. However, higher effort levels are still allowable under existing management arrangements. Further reductions in trawl bycatch and other efforts to reduce risks for species of conservation concern are important for the sustainability of the fishery.

Stewardship among commercial fishers is promoted through the Authority's Reef Guardian Fishers program which currently involves nine fishing operations and up to 50 fishing vessels. These commercial fishers demonstrate leadership in their sector by going well beyond Regulations to maximise the ecological sustainability of their operations. A significant example of stewardship by commercial fishers is the self-imposed moratorium on commercial coral collection in a large part of the Keppel Islands following flooding impacts.

The Authority has completed vulnerability assessments for threadfin salmon and grey mackerel. These assessments show the minimum legal size for both species are smaller than reported sizes at first maturity, exposing these species to fishing pressure before fish can breed. ...The Authority is also actively involved with a range of fishery related projects, such as assessment of coral trout stocks. However, the status and trend of many other fish stocks are unknown. Protection of dugongs from netting in the Townsville region has improved through rule changes for commercial fishing within part of the Bowling Green Bay Species Conservation (Dugong Protection) Special Management Area in 2011. However, illegal fishing is considered one of the greatest risks to the environmental sustainability of commercial fishing.

Fishing — commercial		Effective	Mostly effective	Partially effective	Ineffective
Context	Context				
	Planning				
	Inputs				
	Processes				
	Outputs				
	Overall				
Outcomes	Biodiversity				



Commercial fisheries arrangements in the Reef are accredited against national sustainability guidelines. The Authority is involved in this role in an informal advisory capacity only. Engagement with stakeholders is critical for the Authority to positively influence the management of fishing. The Authority partners with the Queensland Government, the fishing industry and other stakeholders such as the CSIRO and James Cook University to better understand the impacts of fishing on the Reef and to help ensure use of the Reef is ecologically sustainable. The Authority has also worked with commercial fishers to develop a Reef Guardian Fishers stewardship program to recognise and promote sustainable fishing practices. The sector is also represented on most Reef Advisory Committees and Local Marine Advisory Committees.

The Authority's Field Management Program resources are focused on compliance, partnerships and engagement, with further work undertaken through the Climate Change Adaptation Strategy and Action Plan. However, funding for the Field Management Program has been static since 2008 and declining in real terms. Funding has not kept pace with an increase in the protected area.

#### Effectiveness of management tools as assessed by stakeholders

Zoning was considered to be the most effective of the Authority's management tools used to manage commercial fishing in the Great Barrier Reef Marine Park although the compliance effort and political will to enforce restrictions is seen as weak. Getting the balance right between the interests of commercial and recreational fishers is seen as an area requiring attention by some respondents. Vessel monitoring systems are regarded as effective but need to be extended. Research and monitoring, along with the Great Barrier Reef Marine Park Act and Regulations, were considered to be very or mostly effective by more than 50 per cent of the survey respondents. The Authority's partnerships, compliance, plans, permits and policy were seen as ineffective or only partially effective by more than 60 per cent of respondents.

### 8.5.3 Fishing — recreational

Table 8.13 Management effectiveness scores for fishing — recreational					
Recreational fishing		Effective	Mostly effective	Partially effective	Ineffective
Context					
Planning					
Inputs					
Processes					
Outputs					
Outcomes	Overall				
	Biodiversity				

The Authority has a limited regulatory role in relation to recreational fishing and collecting (primarily through the Zoning Plan and Regulations). The Authority has an advisory role to the Queensland Government in relation to the management of recreational fishing in the Region.

Recreational fishing is one of the most significant recreational activities undertaken on the Reef. The main target species are coral trout and other cod, emperor, tropical snapper, barramundi, bream, mackerel, whiting, crabs, lobster and bait fish. Recreational fishers report that they release nearly half of all fish caught<sup>25</sup> but the survival of released fish is highly variable and dependent on a number of factors including the experience of the fisher.

Management of the ecological sustainability of recreational fishing by the Authority is primarily through the Zoning Plan and Regulations, through public appreciation areas that limit spearfishing (and some commercial fishing activities), in addition to education and awareness programs. As is the case with commercial fishing, compliance plays a role in the management of recreational fishing. Recreational fishing is predominately managed by the Queensland Department of Agriculture, Fisheries and Forestry through size, take and

possession limits; however, there is no total allowable catch for this fishery. Fishing gear constraints and limitations of the take of certain species contained in the Zoning Plan and the Great Barrier Reef Marine Park Regulations also contribute to the management of recreational fishing and collecting.

The Authority's Recreation Management Strategy<sup>10</sup> identified the key threats to the Reef from recreational fishing but did not consider this activity in detail. Ecosystem effects and cumulative impacts of fishing are poorly understood, but are likely to be concentrated in inshore areas close to major population centres. Local depletion, particularly of some inshore species, is of concern in some areas.<sup>5</sup> Economic data estimated that more than \$57 million per annum was spent on recreational fishing, with more than 3.4 million fishing trips undertaken in 2012.<sup>23</sup>

While anchor damage in the Whitsundays led to the establishment of no-anchoring areas, the Recreation Management Strategy identified that recreational fishing practices are unlikely to result in significant direct or indirect impacts on the habitats of the Region. However, an increase in the coastal population and the fly-in fly-out community could lead to a decline in the Region's values if not spatially monitored and managed.

The Authority has no specific policy document on recreational fishing, and there are no explicit objectives, actions or targets articulated for recreational fishing. Responsible reef practices<sup>26</sup> are detailed on the Authority's website, providing best practice guidance for a number of activities associated with recreational fishing, such as boating, anchoring and mooring. Consequential or cumulative impacts are less well understood by the Authority and will require long-term trend and habitat monitoring to give confidence that potential impacts can be effectively addressed. Compliance and wider Field Management Program financial resources, associated with the surveillance of recreational fishing, appears inadequate and on the decline. For the Reef to be environmentally



sustainable, illegal fishing needs to be identified and ceased. The current Field Management Program is operating at capacities set in the 1990s when Reef usage and pressures were lower, and prior to the seven-fold expansion of the area of highly protected zones and the emergence of climate change.

Stakeholder engagement is good with respect to user groups, with representatives on Local Marine Advisory Committees. However, many individuals do not belong to groups or clubs. The Recreation Management Strategy identifies that engagement and collaboration with recreational users, Traditional Owners and the community more broadly are likely to play a key role in further reducing risks associated with recreational fishing. As identified in the Outlook Report 2009, effective collaboration in management efforts and getting a better indication of the total recreational fishing effort are particularly challenging in the diffuse recreational fishing sector.

#### **Effectiveness of management tools as assessed by stakeholders**

The Zoning Plan was the only tool considered to be very or mostly effective in managing recreational fishing by more than 50 per cent of respondents. The Zoning Plan, associated educational maps and materials, and a broad ecosystem-based approach to management were seen as strengths of the Authority's management of recreational fishing.

Policy, along with stewardship and best practice, were seen as being the least effective. A lack of capacity to manage recreational fishing effort or take and a lack of rangers in the field were regarded as weaknesses. A lack of effective compliance of the Zoning Plan, the Great Barrier Reef Marine Park Act and Regulations was also highlighted by many respondents. Some respondents proposed additional Marine National Park Zones (green zones), especially along the coastline, to improve fish recruitment and the sustainability of recreational fishing.

Workshop participants consistently raised the need for more baseline information about recreational fishing, a greater compliance presence and the cross-jurisdictional issues with state fisheries management. The introduction of a recreational fishing licence to improve user compliance and raise additional revenue for management of the Marine Park was also proposed. Improved information based on better research and monitoring was seen as necessary to improve management and the understanding of users.

### **8.5.4 Recreation**

The Authority has a regulatory role (predominantly through the Zoning Plan) and a strong partnership role to manage recreation within the Region.

The Authority manages recreation through a variety of management tools including legislation, the Zoning Plan, plans of management, site management, partnerships, education and community awareness. Permits are not required for low impact activities, including recreational activities; however, permits may be required for group activities where the impact is likely to be significant. Specific policies for interactions with whales and dolphins and on seabird breeding islands are also in place. However, the diffuse nature of recreation and lack of permit control means that plans of management and site planning arrangements do not provide certainty regarding where uses may occur and where impacts are likely to be acceptable.

The Authority considers tourism as those travelling or carried by a commercial tourism operation, while recreation includes locals recreating and a portion of traditional tourists under international definitions (for example, free and independent tourists).

The assessment of management effectiveness for recreation undertaken for the Outlook Report 2009 concluded that despite the lack of an overarching planning framework and targeted management objectives, quality recreation products and services have been provided, and that good governance and management processes are in place to address recreation issues. In response, the Authority prepared a Recreation Management Strategy. The strategy is designed to provide an overarching framework for the management of recreation in a coordinated manner and to inform the public of the management approach. The strategy also identifies which management tools are in place for each recreation activity. The values that attract large numbers of visitors are well documented, and threats and risks to those values are clearly articulated in the strategy. The risk analysis determined there were no threats that posed a very high or high risk to the values. Cumulative impacts are recognised as an issue in the strategy, but are not specifically addressed. The condition and trend of recreation is referenced, but the strategy does not address the impacts of recreation on condition and trend as a whole.

The implementation of the Recreation Management Strategy is identified in the Authority's Strategic Plan, and resources are allocated in line with the objectives. Funding is concentrated mainly on describing what and where activities are undertaken and on stewardship and education programs, and less on site planning, site infrastructure and recreation-specific policy document development.

**Table 8.14 Management effectiveness scores for recreation**

Recreation		Effective	Mostly effective	Partially effective	Ineffective
Context					
Planning					
Inputs					
Processes					
Outputs					
Outcomes	Overall				
	Biodiversity				

Stakeholder engagement remains strong with the Tourism and Recreation Reef Advisory Committee consulted regularly as part of recreation planning. Recreation representatives are also included in the 12 Local Marine Advisory Committees, many of which have produced regional brochures specifically for recreational users. Authority staff in the Cairns, Mackay and Rockhampton regional offices interact with recreational users particularly through Community Access Points. However, the diversity and informality inherent in the sector presents an ongoing challenge to engage with the majority of recreational users and to document their values and activities spatially.

Coordination between relevant agencies (the Authority, Queensland Parks and Wildlife Service, Queensland Boating and Fisheries Patrol, Maritime Safety Queensland, Australian Water Police) to enforce Marine Park Acts, Regulations, Zoning Plans and plans of management is high. In addition, some products and services such as maps and brochures are jointly prepared and presented.

The Recreation Management Strategy provides excellent data concerning the risk-based approach adopted by the Authority, while identifying the major risks and threats associated with recreation and avenues to reduce those risks. However, timeframes and targets to meet the objectives will assist in assessing the performance on the strategy in achieving the desired outcomes.

#### Effectiveness of management tools as assessed by stakeholders

All of the Authority's management tools, except compliance and policy, were seen as very or mostly effective by more than 40 per cent of respondents, with the Zoning Plan and the Great Barrier Reef Marine Park Act and Regulations being seen as the most effective tools overall. The Zoning Plan, Reef Guardian programs and educational programs are seen as strengths in recreation management, while a lack of compliance enforcement and availability of field staff are the main identified weaknesses.

Workshop participants raised the need for better understanding of the changing recreational sector, in particular the growing number of new recreational users as a result of the mining boom. Suggested future management actions included more reef protection markers and moorings, as well as increased education.

### 8.5.5 Port activities

Table 8.15 Management effectiveness scores for ports Including dredging and dredge material disposal.					
Ports		Effective	Mostly effective	Partially effective	Ineffective
Context					
Planning					
Inputs					
Processes					
Outputs					
Outcomes	Overall				
	Biodiversity				

The Authority has an advisory role in the management of ports, due to most ports being located outside of the Marine Park (within port exclusion areas) (see Table 5.2, Chapter 5).

The planning and development responsibilities for ports lie with the Queensland Government and the Department of Sustainability, Environment, Water, Population and Communities. [Note: The role of the Department is generally confined to the assessment of referred actions that will or are likely to have a significant impact on matters of national environmental significance. The department has no broader planning, development.] The role of the Authority is therefore restricted to the consequential impacts from ports. However, actions may affect matters of environmental significance and trigger a referral requirement under the EPBC Act, requiring assessment from the Department of Sustainability, Environment, Water, Population and Communities and potentially the Authority if it affects the Marine Park or World Heritage Area. Outside the Marine Park, the Authority's role is advisory. This means that while the impact of the port development may be assessed, the Authority has limited responsibilities over the planning of port locations and port activities that occur within the exclusion areas. For example, dredging often occurs outside the Marine Park or within the excluded area. Dredging and disposal may also occur within

the Marine Park, but in these instances the Authority has some accountability and will assess the activity under its permit process.

The increase in the number of proposals for new and expanded ports along the Great Barrier Reef coast is of particular concern. There is also an increase in requests for dredging to extend outside of port exclusion areas and into the Marine Park — for example, dredged entrance channels to allow for larger draft ships to access ports such as Townsville and Hay Point. The Authority has identified risks from the proposed port developments to the inshore area of the Region, which have the potential to cause significant negative impacts on species and habitats critical to the healthy functioning of the Reef's ecosystem.

While dredging associated with ports usually occurs within port limits, dredge material disposal often occurs within the Marine Park. There is generally a good understanding by staff of the Authority of the direct and indirect impacts of dredging on the values of the Reef and policies for the disposal of dredge material and environmental impact management require direct and indirect impacts to be considered. However, this is often considered on an application-by-application basis and the consequential and cumulative impacts of dredging and spoil disposal are

less well understood. The Authority has little knowledge about the condition and trend of many of the ecosystems and species at risk from port development.

The Authority has a role to play in the implementation of offsets associated with some port approvals under the EPBC Act (for example, Curtis Island and Abbot Point). This is a new role for the Authority; it is currently developing guidelines and procedures for managing the funds, while ensuring the offsets meet the intended effects and expenditure is not duplicated with other offsets (for example, those required under Queensland Government approvals).

There is no overarching strategy for port development in Queensland although a consultation paper was recently released, and there are multiple port expansions and new port development proposals. This has a significant impact on the Authority's ability to protect the Region's values, as each port proposal is assessed individually on its merits. Where there are activities within the Authority's jurisdiction, this case-by-case approach results in increased workloads and does not enable issues, such as cumulative impacts, to be addressed at a strategic level. The complex jurisdictional environment within which ports operate — combined with the lack of Authority policy documents for ports — continues to negatively impact on the Authority's capacity to avoid, mitigate, offset and adaptively manage the impacts from ports on the Region's values. Greater certainty and coordinated planning and approval processes are required between the Commonwealth and Queensland jurisdictions to inform where and how ports can operate within and adjacent to the Region.

Where the Authority has jurisdiction over impacts of port development (for example, dredge material disposal within the Marine Park), objectives, actions for implementation, guidelines and standards have been developed. Guidelines for hydrodynamic numerical modelling for dredging projects<sup>27</sup> provide guidance and standards for all new applications. The draft Biodiversity Conservation Strategy is committed to "reducing threats to protected and at risk habitats, species and groups of species" are also in draft form. The environmental management charge currently does not extend to dredge material disposal. Given the charge is a contribution towards management of the Marine Park, extension of this charge to high risk activities such as ports and dredging should be considered.

Resourcing for assessing and managing the impacts of port proposals or developments has increased but is not sufficient to keep pace. While staff have been committed to developing policy positions, assessments have also increased. Stakeholder engagement by the Authority is sound, with regular meetings held for key stakeholders such as Local Marine Advisory Committees and Reef Advisory Committees, as well as regular meetings with the Queensland Ports Association. Community input is limited to occasions when a proposal triggers the environmental impact assessment process under the EPBC Act or the Great Barrier Reef Marine Park Act because it may have impacts on matters of national environmental significance or the proposal occurs within the Marine Park. Documentation associated with environmental impact assessment processes are made available for public comment.

There is little evidence of outcomes to support ports being environmentally sustainable. However, ports are seen to be economically sustainable and local communities are reliant on ports for goods and services, as well as employment. This is balanced against the loss of amenity and 'enjoyment' and potential economic losses of other industries (for example, fishing) near port developments and potentially further afield given the potential for dredge material to resuspend and disperse further than the dredge disposal areas.

#### **Effectiveness of management tools as assessed by stakeholders**

None of the management tools available to the Authority were seen to be particularly effective in managing ports. Permits (including impact assessment), the Great Barrier Reef Zoning Plan 2003 (the Zoning Plan), research and monitoring, and the Great Barrier Reef Marine Park Act and Regulations were considered to be the most effective of the management tools. Policy was seen as the weakest tool in managing ports with more than 90 per cent of respondents rating it as ineffective or only partially effective. Two contrasting views of the Authority's involvement in planning and management of ports were evident in the responses. One reflected a view that the Authority does not fully understand port development and their environmental management systems and is "driven by a culture that regards port activity as inherently harmful". The other view sees ports as a major threat to the Reef but that the Authority lacks the jurisdictional authority, capacity or willingness to address the impacts posed by port development and management.

The engagement of the Authority with port authorities was seen positively, although there was generally little strength in management identified by respondents. Jurisdictional limitations and a corresponding lack of influence of port development and management decisions, weak compliance enforcement, a lack of understanding by the Authority of the ports industry and a lack of consistent messages from the Authority concerning port development were identified as weaknesses.

A number of respondents saw an extension of the jurisdictional reach of the Authority to have a greater say over port development as a potential solution, while a number called for more explicit policy from the Authority on port planning and management.

Extensive discussions about ports occurred at the stakeholder workshops. The main management issues raised were: the need to improve assessment processes, overlapping state and federal responsibilities, concerns about dredge disposal, improved governance arrangements for ports, and the Authority's lack of influence over decision making and management.



## 8.5.6 Shipping

Table 8.16 Management effectiveness scores for shipping					
Shipping		Effective	Mostly effective	Partially effective	Ineffective
Context					
Planning					
Inputs					
Processes					
Outputs					
Outcomes	Overall				
	Biodiversity				

The Authority has a limited regulatory role and a strong partnership role with other agencies to manage shipping within the Region. There are stringent management requirements for commercial shipping in the waters of the Great Barrier Reef which was designated a Particularly Sensitive Sea Area by the International Maritime Organisation in 1990. Shipping in the Great Barrier Reef is managed by several government agencies including the Australian Maritime Safety Authority, Maritime Safety Queensland, and the Authority.

The main management tools used by the Authority to manage shipping include legislation and the Zoning Plan. Shipping traffic is confined to designated shipping areas in the Great Barrier Reef Region, defined in the Zoning Plan 2003, unless a specific permit is issued. Permits, compliance, policy documents and education are used to a lesser extent.

The assessment of management effectiveness undertaken for the Outlook Report 2009 concluded there were relatively few incidents threatening the Reef's values relative to the large number of shipping movements in and through the area, and that invasive species were the highest risk with respect to shipping.

Statistics show there is a general upward trend in the number of pollution and vessel incidents reported.<sup>28</sup> [Note: While there is an upward trend in reported vessel incidents, the number of shipping incidents (i.e. for vessels greater than 50 metres in length) has fallen since the introduction of the Reef vessel tracking service.<sup>29</sup>] The most concerning issues with regards to shipping are the threat of oil, chemical and cargo releases, groundings and the introduction of invasive marine species. In addition, a greater range of hazardous substances (such as chemicals and liquefied natural gas) is now being shipped through the Reef.

Since the release of the Outlook Report 2009 and in response to mining industry growth, there has been a significant increase in development proposals to expand all of Queensland's major trading ports. There are also several new port development proposals. Mining industry growth will result in the construction of new shipping berths, infrastructure development, dredging and a rise in shipping traffic through Reef waters. The extent to which this increase in ports and shipping activities will emerge as a threat to the health of the Reef will undoubtedly depend on how well these activities are managed by all stakeholders, industry and regulators.

Potential direct and indirect environmental impacts associated with shipping activities are understood by the Authority, although the cumulative impacts are less well known. A reasonable planning and legislative framework for managing shipping is in place. However, there are exceptions for threats posed by bio-fouling, ballast water discharges, containerised chemicals, noise, turbidity and light, antifouling biocide leaching, physical impacts from vessel groundings, response and rehabilitation of environment following grounding events, and increasing anchorage concerns (where the Authority has fewer measures in place).

The Great Barrier Reef Marine Park is one of the world's most regulated shipping areas. However, experience from the Shen Neng 1 grounding in 2010<sup>30</sup> highlighted significant gaps in the Authority's ability to take action and rehabilitate following this incident. Agencies that manage shipping, including the Authority, need to address a major gap in the capacity to respond to non-oil spills with rapid clean up (for example, access to emergency funds sufficient to clean up in a timely and cost-effective manner).

There is concern (from the independent assessors) about resourcing levels given the increase in shipping movements. With the decline in funding for the Field Management Program, there is significant risk for appropriate preparedness and response times to a shipping incident.

There is good stakeholder engagement with regular discussions about shipping issues at Local Marine Advisory Committee and Reef Advisory Committee meetings, with representatives from the shipping sector members on some of the committees. Regular meetings are held with the Queensland Ports Association. The Australian Maritime Safety Authority, the Authority and Maritime Safety Queensland work closely together to protect the marine environment from adverse consequences of shipping operations and provide for the safety of life of ship crews, passengers and other users of the Reef.

The Authority is working with other agencies to assess the effectiveness of current shipping safety measures to inform any additional measures needed to protect the value of the Reef. The Authority's plans for oil pollution response are well documented and managed. However, there continues to be an absence of clear measurable objectives and targets with respect to shipping, while progress on addressing other contaminants and marine pests remains slow.



### Effectiveness of management tools as assessed by stakeholders

The Zoning Plan and compliance were seen as very or mostly effective by more than 50 per cent of respondents. The Great Barrier Reef Marine Park Act and Regulations, along with education and community awareness, were seen as least effective with 60 per cent of respondents regarding them as ineffective or only partially effective.

Collaboration with other authorities and arrangements for compulsory pilotage and designated shipping lanes are regarded as strengths of the Authority in management of shipping, while the capacity to address likely increases in shipping and potential impacts from marine pests were cited as weaknesses.

Workshop participants consistently raised management issues which are outside the Authority's jurisdiction, such as the need for improved pilotage, bilge and ballast water management, marine debris and other impacts from 'parked' ships, exclusion of other users in high shipping areas, and management response to groundings.

### 8.5.7 Defence activities

The Authority has a limited regulatory role in the management of defence activities within the Region. Defence activities and movements by the navy, air force and army occur throughout the Region. Three defence training areas are within the Great Barrier Reef: Shoalwater Bay, Cowley Beach and Halifax Bay. The Shoalwater Bay, Tully, High Range and Mt Stuart training areas also include significant catchment areas which drain into the Reef.

The Defence Department is responsible for the conduct of training activities by the defence forces, including visiting overseas defence force members. This responsibility includes the management of the environmental impacts of those activities, subject to the application of zoning plans. The management of defence operations within the Great Barrier Reef Region therefore relies substantially on the environmental management standards of the Defence Department and the strength of the partnership agreement between this body and the Authority.<sup>31</sup> The assessment for the Outlook Report 2009 concluded the limited area of operations and the high level of performance by the Defence Department in managing the environmental impacts means that defence activities pose minimal threats to Great Barrier Reef values.

The Department of Defence has an environmental management strategy that includes objectives to implement best practice environmental management in its operations. In addition, a strategic environmental assessment for defence activities in the Great Barrier Reef World Heritage Area (the Strategic Environmental Assessment)<sup>32</sup> was developed by the Defence Department and the Authority, covering defence activities in the Great Barrier Reef World Heritage Area. This document clearly outlines the values of the Great Barrier Reef and adjacent areas, including matters of national environmental significance and the potential risks to these values. The assessment was developed in 2006 and is currently being updated.

As noted in the Outlook Report 2009, defence activities are allowed under the Zoning Plan. Strategic documents, policies and regular meetings facilitate implementation of the management agreement and ensure a consistent approach with other management agencies. Specific defence resources are devoted to environmental management, while staff exchanges, secondments, patrols and systematic application of the Defence Heritage Toolkit help boost the department's expertise to manage defence activities and their potential impacts on Reef values.

Adequate biophysical information within defence training areas continues to be available for the Authority's management decisions. Navy hydrographic surveys improve knowledge of benthic habitats, ocean and weather conditions relevant to management. Surveys and studies about a range of ecological communities and species are undertaken. Local communities continue to be engaged in planning for specific exercises and routinely through Defence Environmental Advisory Committees.

Training exercises are thoroughly planned and include good performance monitoring, debriefs and post exercise monitoring. The identification of clear performance indicators, particularly those related to addressing cumulative impact issues, remains a challenge. The other challenge relates to the high level of reliance by the Authority on planning, assessment and performance monitoring work undertaken by the Defence Department. While Authority staff have an appropriate mix of skills to fulfil their statutory responsibilities for defence activities, the liaison and monitoring work is undertaken as a relatively low priority by a small number of Authority staff in competition with other core business. The skills required to assess defence activities are limited to a few core staff. An increase in training and sharing of corporate knowledge is essential to ensure this activity is effectively managed into the future.

**Table 8.17 Management effectiveness scores for defence activities**

Defence activities		Effective	Mostly effective	Partially effective	Ineffective
Context					
Planning					
Inputs					
Processes					
Outputs					
Outcomes	Overall				
	Biodiversity				

A systematic approach generally ensures that statutory and planning timeframes are routinely met and results are reported by the Defence Department in a timely manner. The knowledge base for confident management of defence activities in the Great Barrier Reef, both within the Authority and in the wider community, continues to increase as a result of consultative meetings and reports documenting efforts to minimise environmental impacts, along with a lack of evidence of death or injury to species such as dugong and marine turtles.

#### Effectiveness of management tool as assessed by stakeholders

Insufficient responses were received to assess the effectiveness of management tools for defence activities.

### 8.5.8 Research activities

Table 8.18 Management effectiveness scores for research activities					
Research activities		Effective	Mostly effective	Partially effective	Ineffective
Context					
Planning					
Inputs					
Processes					
Outputs					
Outcomes	Overall				
	Biodiversity				

The Authority has a lead role in relation to the management of research activities (that is, access to the Marine Park to undertake research) within the Region. The Great Barrier Reef is known internationally as a premier site in which to conduct scientific studies. The majority of research occurs at the four major research stations at Lizard Island, Orpheus Island, One Tree Island and Heron Island. Diffuse research is conducted at other locations.

The Authority manages the impacts of research activities on Reef values, including extractive and observational research use, through a mix of management tools. As provided for in the Great Barrier Reef Marine Park Act and Regulations, permits for specific research projects and accreditation of partner research institutions are the principal means of managing the potential impacts from this activity. These mechanisms cross reference to the Zoning Plan and plans of management for specific geographic areas, as well as to specific policy documents on managing research and provisions for compliance and enforcement of permit conditions.

The Authority's Scientific information needs for the management of the Great Barrier Reef Marine Park report<sup>33</sup>

identifies key information needed to better inform management of the area. This document, along with partnership agreements with key research institutions, provides the basis for research alignments which should deliver valuable outcomes for improved management. The Authority is updating this document as a key input to the development of an integrated monitoring framework for the World Heritage Area.

The assessment of management effectiveness undertaken for the Outlook Report 2009 concluded that management of research in the Reef was moving towards desired outcomes, reducing risks and threats to Reef values. While research activities were generally considered to be environmentally sustainable, it was recognised there was a need to more effectively manage research through the implementation of a relational database designed to capture and manage permitted use. To date, this system (now renamed the Reef Management System) is still under development and unable to inform intended and actual research take through queries and reports. This limits the ability to manage cumulative use. This has been identified as an area requiring improvement and will be particularly valuable for the scientific research zones that surround research stations where research is heavily concentrated.

Further opportunity to improve the management of cumulative impacts has been identified through the development and implementation of environmental management plans for high use scientific research zones. In 2009, the development of these plans was judged to be slow, with only one plan completed and another two in draft form. While little progress has been made since then, other management arrangements are in place to assist with the effective management of these zones. For example, permit conditions require consultation and negotiation between researchers and research station staff regarding the location and timing of permitted research activities. Researchers are also required to submit reports detailing their collection throughout the permit duration.

Historically, research has been focused on biophysical systems but there has been more recent interest in socio-economic drivers. While more biophysical information is available upon which to address management of scientific research, there is limited traditional knowledge and socio-economic information is only slowly becoming available. Research proposals with the potential to impact Indigenous values are referred to the relevant Indigenous liaison staff for guidance. However, Traditional Owner participation in research within their sea country, and dissemination of research results to them, is limited (in contrast to their high level of interest in this area). Direct risks and threats to Reef values from specific research activities are recognised but potential cumulative impacts on the Region's values are unclear and warrant more focused attention. Ironically, the Reef's reputation as one of the best managed reef systems in the world is likely to make it a preferred site for more research which could significantly add to the pressure on research sites.

The Authority has very successful and effective relationships with the research community in relation to how scientific research is managed in the Great Barrier Reef. This is shown through the accreditation process and

strong partnerships with research institutions. Co-accreditation arrangements are in place to ensure consistency between jurisdictions, while permits require stakeholder and local community engagement as necessary. Management staff have appropriate research skills and expertise to effectively manage research activities; however, resources available for the management of research activities are limited. Consequently there is limited capacity to contribute to the formulation of Regulations and policies associated with research or to develop environmental management plans for scientific research zones. The 2004 Policy on Managing Scientific Research in the Great Barrier Reef Marine Park<sup>34</sup> needs to be reviewed and updated to reflect this focus on cumulative impacts. In addition, the limited impact and collection limits in the Regulations need to be revised to ensure the take limits are ecologically sustainable (for example, a researcher could collect five tiger sharks from one location under the current accreditation scheme).

Uncertainties resulting from the limited understanding of potential cumulative impacts and limited attention to compliance with permit conditions make it impossible to state categorically that outcomes for management of research activities are being met and objectives are as clear and measurable as they need to be.

Research activities on the Reef are well regarded internationally and there are robust management processes and governance arrangements in place to manage research on the Reef. In general, research is not seen to have a large and detrimental impact on the Reef ecosystem. However, confidence that research is environmentally sustainable is reduced by the limited knowledge of cumulative impacts of collection undertaken for research. It will be important for an updated policy, coupled with information needs documentation, to set clear targets to guide the management of research of the next 25 years. Particular attention should be paid to addressing potential cumulative impacts in intensively researched areas.

#### **Effectiveness of management tools as assessed by stakeholders**

The Zoning Plan, partnerships, and education and community awareness were seen as the Authority's most effective management tools for scientific research. Compliance, policy, stewardship and permits were seen as ineffective or only partially effective.

The existing partnerships with research institutions are identified as a major strength. However, the limited research capability within the Authority itself is noted as a weakness and is attributed to the lack of long-term secure funding to support strategic research. The possibility of establishing a specialist board to guide research was raised by some respondents. The limited amount of compliance monitoring, especially on the cumulative impacts of research, is identified as a weakness, along with the need for better housekeeping and management of research sites.

## **8.6 Effectiveness of specific measures to address impacts**

This section contains an assessment of the effectiveness of the Authority's current management arrangements to address impacts. It is structured around the relevant criteria in item 4.1.1 of the terms of reference for the strategic assessment (see Appendix 2) which were also part of the assessment criteria for the independent review. The relevant criteria are:

- a. [not applicable]
- b. identify and analyse direct, indirect, consequential and cumulative impacts, including the methods used to determine these types of impacts
- c. consider environmental, social, cultural and economic issues
- d. avoid, mitigate, offset and adaptively manage impacts
- e. address uncertainty and risk
- f. provide certainty regarding where uses may occur, the type of activities allowed, conditions under which activities may proceed and circumstances where impacts are likely to be unacceptable
- g. halt and reverse any declines and enhance the condition of the relevant matters of national environmental significance, including mechanisms to deliver a 'net benefit' to the condition of the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area
- h. adapt to reasonable climate change scenarios
- i. integrate with related local, Queensland and Australian government programs to protect and manage the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area
- j. meet Australia's international responsibilities in relation to the environment and protection of world heritage
- k. monitor, evaluate and report on the:
  - i. [not applicable]
  - ii. impacts of activities, including the setting of targets, to benchmark management performance.

The grades were calculated by averaging the scores for each criterion across all management topics. A summary of the individual scores is provided in Appendix 1 to the full report of the independent assessment, available on the Authority's website. In order to understand the relative influence of impacts on values, a contextual

understanding of the condition and trend of values is needed. The assessment considered the Authority's understanding of the current condition and trend of matters of national environmental significance (items (a) and (k)(i) above) for each management topic. This is discussed in Section 8.8. The summary below was reviewed and endorsed by the independent assessors.

### 8.6.1 Understanding impacts and considering issues

The independent assessment found the Authority has generally a good understanding of direct and indirect impacts associated with impacts of highest concern. Impacts are better understood where the Authority has greater jurisdictional responsibility; for example, tourism, defence activities, and research activities. An understanding of the direct and indirect impacts of water quality, dredging and coastal development on the Region's values is improving within the Authority. An understanding of, and plans and processes to address cumulative use and impacts on outstanding universal value, was assessed as lacking across the Authority's entire management scheme.

The assessment outcomes in relation to understanding impacts are summarised in Table 8.19.

**Table 8.19 Effectiveness of the Authority's management arrangements to understand impacts**

Indicator	Grade				Management tools
	Effective	Mostly effective	Partially effective	Ineffective	
4.1.1(b) identify and analyse direct, indirect, consequential and cumulative impacts on relevant matters of national environmental significance, including the methods used to determine these types of impacts					
Direct and indirect impacts are understood by managers					<p>Strengths:</p> <ul style="list-style-type: none"><li>• Zoning Plan (for extractive activities)</li><li>• Plans of management (for tourism and recreation)</li><li>• Reef Plan</li><li>• Policy documents (for example <i>Biodiversity Conservation Strategy</i>)</li><li>• Research and monitoring (for example Outlook Report)</li></ul> <p>Weaknesses:</p> <ul style="list-style-type: none"><li>• Permits (case-by-case assessments and limited understanding of cumulative use)</li><li>• Policy documents (for example, limits on acceptable use, cumulative impact assessment policy)</li><li>• Research and monitoring (of cumulative impacts)</li></ul>
Consequential and cumulative impacts are understood by managers					

The Great Barrier Reef Marine Park Regulations explicitly require consideration of environmental, social, cultural and heritage values as part of every permit assessment for activities which occur within the Great Barrier Reef Marine Park. However, the effectiveness of these permit assessments relies on the Authority's knowledge of these aspects in specific locations. The independent assessment found limitations in the Authority's understanding of Indigenous heritage, historic heritage and community benefits. It identified that further work is needed to maintain relationships with Traditional Owners, support co-management and develop a mutually agreed and culturally appropriate process for the integration of Indigenous knowledge into management. For a number of historic heritage matters, the assessors found there is insufficient understanding of their location, condition and trend, and suggested that a database with spatial capacity would improve consideration of the potential impacts from proposed activities on heritage values in the Region. The consideration of community benefits was found to vary according to the understanding and perspective of the community. The independent assessors noted there is likely to be increasing conflict between economic benefits associated with port and coastal development, and the aesthetic and recreational benefits. Currently, the Authority has no guidelines or benchmarks for assessing or managing community benefits derived from the Region or policies to guide the assessment of cumulative impacts.

Stakeholder engagement and application of biophysical information were identified as the strongest aspects of management across all management topics. Respondents (from stakeholder surveys) saw a need to extend the Authority's jurisdictional control over activities that impact on the values of the Marine Park, to improve compliance action and to increase the focus on partnerships between the Authority, local government and local industry. The assessment outcomes in relation to considering issues are summarised in Table 8.20.



**Table 8.20 Effectiveness of the Authority's management arrangements to consider issues**

Indicator	Grade				Management tools
	Effective	Mostly effective	Partially effective	Ineffective	
4.1.1(c) consider environmental, social, cultural and economic issues in its decision-making processes					
Engagement with stakeholders and local community					<p>Strengths:</p> <ul style="list-style-type: none"><li>• Act and Regulations (Regulations 88Q and 88R)</li><li>• Partnerships (for example, Reef Advisory Committees, Local Marine Advisory Committees)</li></ul> <p>Weaknesses:</p> <ul style="list-style-type: none"><li>• Policy documents (for example, lack of integrated strategies and processes to address Indigenous heritage, lack of minimum requirements for public consultation)</li><li>• Research and monitoring (traditional and social information)</li></ul>
Availability of information to enable consideration of issues:					
Biophysical					
Indigenous					
Social					
Local community and stakeholders effectively engaged in ongoing management					

## 8.6.2 Managing impacts

Assessors found that where the Authority has a high level of control over activities, its effectiveness at avoiding, mitigating and adaptively managing impacts was effective or mostly effective (for example, tourism, research activities, defence activities, recreation). Where the Authority has 'diffuse' or limited jurisdictional responsibility, its ability to avoid, mitigate and adaptively manage impacts was found to be only partially effective. The partially effective grade was found to be pronounced in areas of planning, process and outcomes and for activities occurring outside the Marine Park (for example, port activities, coastal development) and those occurring in the Marine Park, but not under the Authority's direct jurisdictional control (for example, commercial and recreational fishing).

An assessment of the effectiveness of the Authority's management in the south of the Great Barrier Reef, around the Keppel Islands, found a number of examples of planning that support the protection of the Keppel Bay area. This includes zoning plans, specific threat abatement plans with no-anchorage areas, declared fish habitat, and Reef Plan. However, it was identified that further work is required to develop a plan of management in this area to address increased use (Table 8.21).

On a Reef-wide scale, assessors found the Authority's Zoning Plan and the complementary plans for the adjacent Great Barrier Reef Coast Marine Park have made the most significant contribution to biodiversity protection. The Representative Areas Program (1999–2004)<sup>35</sup> involved a systematic planning and consultative program to develop new zoning for the Marine Park. The outcomes resulted in the protection of at least 20 per cent of every bioregion, leading to the protection of 'representative' examples of all different habitat types, reducing risks to the Marine Park. This process also maximised the benefits and minimised the negative impacts of the rezoning on the existing users of the Marine Park, addressing uncertainty for extractive uses in particular. The Zoning Plan has been in operation since 1 July 2004. However, the zoning provisions only address biodiversity protection from direct extractive uses, particularly fishing, and do not address uncertainty for a range of other activities (for example, where some development can and cannot occur).

















Across the Authority's entire management scheme an understanding of, and plans and processes to address cumulative use and impacts on *outstanding universal value* was assessed as 'lacking'.

The assessors noted that recent changes by the Queensland Government to the state's coastal planning provisions potentially has significant implications for the Great Barrier Reef Region because it removes many of the specific requirements placed on local government and potential developers to undertake best practice and to minimise environmental harm. These changes potentially reduce the capacity of management agencies, like the Authority, to impose requirements to avoid, minimise or mitigate impacts from coastal development activities on the Reef, its ecosystems and species.

In addition, the complex jurisdictional environment within which ports operate, combined with the lack of Authority policy documents for ports, was found to reduce the capacity to effectively avoid, mitigate, offset and adaptively manage the impacts from port activities on the Region's values. Given the positive results of Reef Plan, the independent assessors suggested an extension of this arrangement may be needed for other activities.

The assessment outcomes in relation to managing impacts are summarised in Table 8.21.

**Table 8.21 Effectiveness of the Authority's management arrangements to manage impacts**

Indicator	Grade				Management tools
	Effective	Mostly effective	Partially effective	Ineffective	
4.1.1(d) avoid, mitigate, offset and adaptively manage impacts from activities					
Direct and indirect impacts are appropriately considered					<p>Strengths:</p> <ul style="list-style-type: none"><li>Climate Change Adaptation Strategy and Action Plan (both 2007–2012 and 2012–2017)</li><li>Permits (for example, for tourism and research activities)</li><li>Partnerships (defence activities, High Standard Tourism program)</li></ul> <p>Weaknesses:</p> <ul style="list-style-type: none"><li>Plans of management (for example, consistency across jurisdictions and need for development of new plans in growth areas)</li><li>Policy documents (for example, cumulative impact management, net environmental benefits and offsets policies)</li></ul>
Consequential and cumulative impacts are appropriately considered					
Management is effectively addressing impacts and moving toward attainment of desired outcomes (NOTE: an ineffective rating was given with regard to climate change)					
Use of the Great Barrier Reef is demonstrably environmentally sustainable (NOTE: an ineffective rating was given with regard to climate change)					

### 8.6.3 Providing certainty













One of the key desired outcomes of the strategic assessment process is to guide management of the Great Barrier Reef by providing greater certainty on where sustainable uses can occur, the type of activities that will be allowed and the conditions under which activities may proceed. The assessors found that, in accordance with the Act, the Authority continues to place primary emphasis on the long-term protection and conservation of the environment, biodiversity and heritage values of the Great Barrier Reef Region. The Authority allows ecologically sustainable use, so far as this is consistent with the Region's protection and conservation.

The independent assessment found zoning provides a high level of certainty for extractive activities. However, zoning does not provide explicit guidance on where many activities not directly regulated by the Zoning Plan can be carried out. The assessors considered that greater certainty and coordinated planning and approval processes are required by Commonwealth and Queensland jurisdictions to inform where and how ports can operate within and adjacent to the Region. In addition, they considered the diffuse nature of recreation and lack of permit control mean plans of management and site planning arrangements do not currently provide a high level of certainty regarding where uses may occur and where impacts are likely to be acceptable.

The Authority's plans of management were assessed as effective tools to manage activities in specific areas. The current plans are predominantly designed to address tourism and recreation in areas that were, and continue to be, high use areas. The assessors noted these plans have not been updated for some time, and do not account for the full range of direct uses. In addition, emerging high use areas (for example, offshore Townsville and Rockhampton) have no plans of management in place to address uncertainty and increasing multiple uses.

The assessment outcomes in relation to providing certainty are summarised in Table 8.22.


**Table 8.22 Effectiveness of the Authority's management arrangements to provide certainty**

Indicator	Grade				Management tools
	Effective	Mostly effective	Partially effective	Ineffective	
4.1.1(e) address uncertainty and risk					
There is a planning system in place that effectively addresses the management topic					<p>Strengths:</p> <ul style="list-style-type: none"><li>• Zoning Plan</li><li>• Plans of management (for tourism and recreation)</li><li>• Policy documents (for example, environmental impact management policy)</li></ul> <p>Weaknesses:</p> <ul style="list-style-type: none"><li>• Plans of management (require updating and new plans needed in growth areas)</li><li>• Policy documents (for example, limits on acceptable use, cumulative impact assessment policy)</li></ul>
The outputs are reducing the major risks and the threats to the Great Barrier Reef (NOTE: an ineffective rating was given with regard to climate change)					
4.1.1(f) provide certainty regarding where uses may occur, the type of activities allowed, conditions under which activities may proceed and circumstances where impacts are likely to be unacceptable					
Plans provide certainty regarding where uses may occur, the type of activities allowed, conditions under which activities may proceed and circumstances where impacts are likely to be acceptable					<p>Strengths:</p> <ul style="list-style-type: none"><li>• Zoning Plan</li><li>• Plans of management</li></ul> <p>Weaknesses:</p> <ul style="list-style-type: none"><li>• Plans of management (require updating and new plans needed in growth areas)</li><li>• Policy documents (for example, limits on acceptable use)</li></ul>

The level of certainty provided to activities in the Region through the current management arrangements varies according to the activity. The levels of certainty provided for the major activities are summarised in Table 8.23 and detailed in Figure 8.2 to Figure 8.10. There is a high degree of certainty for many activities in many areas, including tourism, fishing, research activities, shipping, recreation and defence activities. Using tourism as an example, certainty is provided through a range of management tools, including the Zoning Plan, plans of management and permits (Figure 8.2).

**Table 8.23 Summary of management certainty provided for activities that occur in the Region**

Activity	Level of certainty
Traditional use	Medium
Tourism	High
Fishing and collecting	High
Recreation	Medium
Ports	Low
Shipping	High
Defence activities	High
Research activities	High



**Figure 8.2 Certainty provided for tourism activities by the Authority's management arrangements**


Plans of management and site management, in combination with other management arrangements, provide certainty about how tourism can be undertaken, especially in the most frequently used areas.



For shipping, a Designated Shipping Area (defined in the Zoning Plan), compulsory pilotage areas and designated ship anchorages (adjacent to some ports) provide spatial certainty for shipping activities in the Region. Additional management tools, including permits for barging of bulk fuel, sea dumping permits and the Vessel Traffic Service (managed by the Australian Maritime Safety Authority), set requirements for ship operations and reporting.

The Authority's current management arrangements for the operation of ports provide a low level of certainty (Figure 8.10), principally because many are located outside its jurisdiction. The Marine Park boundary provides a degree of certainty — 10 port areas are outside the Marine Park and two minor ports are within it. Where ports occur in the Marine Park, the provision of clear environmental permit conditions provides certainty for the implementation of port developments. An improved understanding of cumulative impacts and development of guidelines to assess the effects of multiple impacts will increase certainty for the ecological system, as well as increasing certainty for adjacent and overlapping uses.

The independent assessment recognised that certainty for particular activities can never be absolute when operating in a changing natural environment about which understanding is continually improving and society's expectations are evolving. The spatial scale over which different activities operate is not uniform, and neither are the multiple uses in any one location. Providing certainty will always depend to some extent on relevant considerations at the time. For Reef-dependent activities, certainty is strongly linked to the status of the environment; for example, the connection between a healthy Reef and a healthy tourism industry. In addition, the assessors noted management of the Region needs to take into consideration the potential for actions arising from one activity affecting other activities in an area, either by affecting the activity directly or by altering the state of the ecological system on which it depends. They also recognised that industries and other users require flexibility in management arrangements to allow scope for adaptation to new technologies, changing market demand and changing environmental conditions.



**Figure 8.3 Certainty provided for fishing and collecting activities by the Authority's management arrangements**

Zoning and Special Management Areas, in combination with other management arrangements, provide certainty about how fishing and collecting can be undertaken. The activities allowed in each zone, are set out in the activities guide (Figure 8.4).

## ACTIVITIES GUIDE

(see relevant *Zoning Plans and Regulations* for details)

	General Use Zone	Habitat Protection Zone	Conservation Park Zone	Buffer Zone	Scientific Research Zone <sup>2</sup>	Marine National Park Zone	Preservation Zone
Aquaculture	Permit	Permit	Permit <sup>1</sup>	×	×	×	×
Bait netting	✓	✓	✓	×	×	×	×
Boating, diving, photography	✓	✓	✓	✓	✓ <sup>2</sup>	✓	×
Crabbing (trapping)	✓	✓	✓ <sup>3</sup>	×	×	×	×
Harvest fishing for aquarium fish, coral and beachworm	Permit	Permit	Permit <sup>1</sup>	×	×	×	×
Harvest fishing for sea cucumber, trochus, tropical rock lobster	Permit	Permit	×	×	×	×	×
Limited collecting	✓ <sup>4</sup>	✓ <sup>4</sup>	✓ <sup>4</sup>	×	×	×	×
Limited spearfishing (snorkel only)	✓	✓	✓ <sup>1</sup>	×	×	×	×
Line fishing	✓ <sup>5</sup>	✓ <sup>5</sup>	✓ <sup>6</sup>	×	×	×	×
Netting (other than bait netting)	✓	✓	×	×	×	×	×
Research (other than limited impact research)	Permit	Permit	Permit	Permit	Permit	Permit	Permit
Shipping (other than in a designated shipping area)	✓	Permit	Permit	Permit	Permit	Permit	×
Tourism programme	Permit	Permit	Permit	Permit	Permit	Permit	×
Traditional use of marine resources	✓ <sup>7</sup>	✓ <sup>7</sup>	✓ <sup>7</sup>	✓ <sup>7</sup>	✓ <sup>7</sup>	✓ <sup>7</sup>	×
Trawling	✓	×	×	×	×	×	×
Trolling	✓ <sup>5</sup>	✓ <sup>5</sup>	✓ <sup>5</sup>	✓ <sup>5,8</sup>	×	×	×

**PLEASE NOTE:** This guide provides an introduction to Zoning in the Great Barrier Reef Marine Parks.

1. Restrictions apply to aquaculture, spearfishing and harvest fishing for aquarium fish, beachworm and coral in the Conservation Park Zone.
2. Except for One Tree Island Reef (SR-23-2010) and Australian Institute of Marine Science (SR-19-2008) which are closed to public access and shown as orange, all other Scientific Research Zones are shown as green with an orange outline.
3. Limited to 4 catch apparatus per person (eg. crab pots, collapsible traps or dillies).
4. By hand or hand-held implement and generally no more than 5 of a species.
5. Maximum of 3 lines/rods per person with a combined total of 6 hooks per person.
6. Limited to 1 line/rod per person and 1 hook per line. Only 1 dory detached from a commercial fishing vessel.
7. Apart from traditional use of marine resources in accordance with s.211 of the *Native Title Act 1993*, an accredited Traditional Use of Marine Resources Agreement or permit is required.
8. Pelagic species only. Seasonal Closures apply to some Buffer Zones.

Detailed information is contained in the *Great Barrier Reef Marine Park Zoning Plan 2003* and *Regulations* and the *Marine Parks (Great Barrier Reef Coast) Zoning Plan 2004*.


- Permits are required for most other activities not listed above.
- Commonwealth owned islands in the Great Barrier Reef Marine Park are zoned "Commonwealth Islands Zone" - shown as cream.
- All Commonwealth Islands may not be shown.
- Special Management Areas may provide additional restrictions at some locations.
- The Zoning Plan does not affect the operation of s.211 of the *Native Title Act 1993*.

**ACCESS TO ALL ZONES IS PERMITTED IN AN EMERGENCY.**

**Figure 8.4 Activities guide — an indication of what can be done in each zone**

The activities guide appears on all zoning maps and provides introductory guidance as to what activities can occur in each zone and under what conditions (for example, maximum of three lines or rods per person with a combined total of six hooks per person when trolling in General Use and Habitat Protection zones).






**Figure 8.5 Certainty provided for research activities by the Authority's management arrangements**


Zoning, accreditation and permits, in combination with other management arrangements, provide certainty about how research activities can be undertaken.






**Figure 8.6 Certainty provided for recreation activities by the Authority's management arrangements**

Plans of management, public moorings and no-anchoring areas, in combination with other management arrangements, provide certainty about how recreation activities other than fishing can be undertaken.



**Figure 8.7 Certainty provided for traditional use by the Authority's management arrangements**


Traditional Use of Marine Resources Agreements and Indigenous Land Use Agreements, in combination with other management arrangements, provide certainty for Traditional Owners and opportunities for co-management of sea country.



**Figure 8.8 Certainty provided for defence activities by the Authority's management arrangements**

Designated defence training areas, in combination with other management arrangements, provide greater certainty about the conduct of defence activities in the Region.






**Figure 8.9 Certainty provided for shipping activities by the Authority's management arrangements**

Shipping traffic is confined to Designated Shipping Areas in the Region. Compulsory pilotage and other measures increase navigational safety and reduce the risk of ship groundings. Ship anchorages are designated by Maritime Safety Queensland.





**Figure 8.10 Certainty provided for commercial port operations by the Authority's management arrangements**

Only two ports are located within the Marine Park. A Remote Natural Area is designated in the north of the Marine Park to ensure the area is recognised and managed for its natural and undeveloped character, free from structures, and to restrict activities such as dredging.

### 8.6.4 Halt and reverse declines

In order to halt and reverse declines and enhance the condition of relevant matters of national environmental significance, the assessors considered that greater traction in threat reduction is needed. Improved threat reduction is, in turn, dependent on the adoption of significant changes to current policies regarding coastal development, resource use, control of other activities and sufficient resourcing to implement threat reduction programs in the field (for example, compliance and wider field management programs). The Authority's strengths in relation to halting and reversing declines was ensuring the Zoning Plan was complied with, and existing plans of management were fully implemented. For the Reef to be environmentally sustainable, illegal activities need to be identified and effectively controlled. The assessors highlighted that the Authority has additional work to do in updating existing and implementing new plans in areas that require them and developing policies that would lead to net conservation benefits.

An ineffective rating for the achievement of biodiversity outcomes was assigned to the Authority's management of climate change and extreme weather. The assessors indicated that despite significant progress in building the ecological resilience of the Reef, and the social and economic resilience of Reef industries, the Authority's work alone is currently ineffective at reducing the impacts of climate change and extreme weather. The future effectiveness of the Authority's work is dependent on successful global efforts to mitigate climate change. Recent information suggests that, so far, these are not sufficient to maintain conditions compatible with coral-dominated reef habitats in the Region.

The assessment outcomes in relation to halting and reversing declines are summarised in Table 8.24.

**Table 8.24 Effectiveness of the Authority's management arrangements to halt and reverse declines**

















Indicator	Grade				Management tools
	Effective	Mostly effective	Partially effective	Ineffective	
4.1.1(g) halt and reverse any declines and enhance the condition of the relevant matters of national environmental significance, including mechanisms to deliver a 'net benefit' to the condition of the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area					
Management is effectively addressing impacts and moving toward attainment of desired outcomes	<div></div>	<div></div>	<div></div>	<div></div>	Strengths: <ul style="list-style-type: none"><li>Zoning Plan</li><li>Plans of management</li></ul>
The outputs are on track to ensure the values of the Great Barrier Reef are protected (NOTE: an ineffective rating was given with regard to climate change)	<div></div>	<div></div>	<div></div>	<div></div>	Weaknesses: <ul style="list-style-type: none"><li>Plans of management (require updating and new plans in areas that need them)</li><li>Policy documents (to deliver net environmental benefits)</li></ul>

### 8.6.5 Adapting to climate change

Despite significant progress in building the ecological resilience of the Reef, and the social and economic resilience of Reef industries, the assessors reported that the Authority's work cannot make the Region and its industries invulnerable to the impacts of climate change and extreme weather. The outlook for the Reef is poor, and addressing climate change is a local, national and global challenge. The integration of climate change throughout the Authority's core business was found to be largely occurring by building the capacity of external groups to respond to the implications of climate change. They highlighted that through the action plan, the concepts of risk and resilience have been cemented and have strongly influenced the Authority's Strategic Plan.

The assessment outcomes in relation to adapting to climate change are summarised in Table 8.25.

Table 8.25 Effectiveness of the Authority's management arrangements to adapt to climate change













Indicator	Grade				Management tools
	Effective	Mostly effective	Partially effective	Ineffective	
4.1.1(h) adapt to reasonable climate change scenarios					
The planning system for climate change and extreme weather addresses the major pressures and drivers impacting on the Great Barrier Reef's values					<p>Strengths:</p> <ul style="list-style-type: none"><li>Policy documents (for example, <i>Climate Change Adaptation Strategy and Action Plan 2012–2017</i>)</li><li>Partnerships (Reef Rescue Land and Sea Country Indigenous Partnerships Program, <i>Climate Change Action Strategy</i> (tourism))</li></ul> <p>Weaknesses:</p> <ul style="list-style-type: none"><li>Policy documents (a number require updates to more explicitly integrate climate change and net conservation benefits policy)</li></ul>
Actions for implementation regarding climate change and extreme weather are clearly identified within the plan					
Clear, measurable and appropriate objectives for management of climate change and extreme weather have been documented					
The relevant managing agencies have developed effective partnerships with local communities and/or stakeholders to address climate change and extreme weather					

### 8.6.6 Integrating with government programs

The assessors noted that under the Great Barrier Reef Intergovernmental Agreement, the Authority and the Queensland Parks and Wildlife Service have effective joint permit arrangements which streamline management, improve certainty and provide consistency. They noted that this integration works extremely well. However, they highlighted management effectiveness challenges for those broadscale issues which are complex socially, biophysically and jurisdictionally. Some inconsistencies in planning systems remain on management of wastewater generated outside the Region (for example, in creeks or from the mainland). This has downstream effects on the Region. For agricultural activities, the assessors considered that Reef Plan, and the Authority's role in that plan, provides an effective, integrated approach to water quality improvement. However, the impact of changes to Queensland's Coastal Plan on downstream effects from land-based water quality is not known at this stage. Assessors further noted that changes to the Queensland coastal planning policies '*have significant implications for the Great Barrier Reef Region because it removes many of the specific requirements placed on local government and potential developers to undertake best practice and to minimise environmental harm*'.

The assessment outcomes in relation to integrating with government programs are summarised in Table 8.26.

Table 8.26 Effectiveness of the Authority's management arrangements to integrate with government programs

Indicator	Grade				Management tools
	Effective	Mostly effective	Partially effective	Ineffective	
4.1.1(i) integrate with related local, Queensland and Australian government programs to protect and manage the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area					
There is consistency across jurisdictions when planning for relevant management topics					<p>Strengths:</p> <ul style="list-style-type: none"><li>Partnerships (for example, Great Barrier Reef Intergovernmental Agreement</li><li>Compliance (joint Field Management Program)</li><li>Permits (joint permits with the Queensland Government)</li></ul> <p>Weaknesses:</p> <ul style="list-style-type: none"><li>Management plans</li><li>Act and Regulations</li><li>Partnerships</li></ul>
There is a sound governance system in place to address protection and management of management topics					
Management of management topics are consistently implemented across the relevant jurisdictions					

### 8.6.7 Meeting Australia's international obligations





















The Great Barrier Reef Marine Park Act and associated Regulations provide some legislative power to contribute to the protection of the world heritage property. The Authority has an advisory role to the Commonwealth Department of Sustainability, Environment, Water, Population and Communities, which has a statutory responsibility to protect the Great Barrier Reef World Heritage Area. Risks and threats to the World Heritage Area have been more explicitly addressed in new periodic reporting processes for world heritage sites to which the Authority contributes.

The Authority's partnerships and planning across jurisdictions are not yet fully integrated to deliver desired outcomes for the World Heritage Area. In particular, the attainment of desired outcomes is weak for climate change and coastal development. Some inconsistencies across the planning systems of different jurisdictions remain in relation to management of wastewater generated outside the Region (for example, in creeks or from the mainland). This has downstream effects on the Region. The Authority's permit process is mostly effective at considering potential impacts to heritage. However, a reduction in educational opportunities (such as stopping the Authority's reef discovery training program) has led to a decline in the delivery of education on world heritage values and the Authority's obligations regarding the presentation of these values.

The assessment outcomes in relation to meeting Australia's international obligations are summarised in Table 8.27.



**Table 8.27 Effectiveness of the Authority's management arrangements to meet Australia's international obligations**













Indicator	Grade				Management tools
	Effective	Mostly effective	Partially effective	Ineffective	
4.1.1(j) meet Australia's international responsibilities in relation to the environment and protection of world heritage					
Relevant standards are identified and being met					<p>Strengths:</p> <ul style="list-style-type: none"><li>• Act and Regulations</li><li>• Zoning Plan</li><li>• Permits</li><li>• Compliance</li><li>• Partnerships (for example, Intergovernmental Agreement)</li><li>• Research and monitoring (for example, Outlook Report, world heritage periodic reporting)</li></ul> <p>Weaknesses:</p> <ul style="list-style-type: none"><li>• Management plans (integration and alignment across jurisdictions)</li><li>• Partnerships (integration and alignment across jurisdictions)</li><li>• Education and community awareness (world heritage and outstanding universal value)</li></ul>
Management is effectively addressing impacts and moving toward attainment of desired outcomes					
The outputs are on track to ensure the values of the Great Barrier Reef are protected					
The outputs are reducing the major risks and threats					
Use of the Great Barrier Reef is demonstrably environmentally sustainable					

### 8.6.8 Monitoring, evaluating and reporting

For a number of values (Section 8.3), the independent assessors considered that the Authority has an insufficient understanding of their location, condition and trend. This reflects the vast area of the Region, its complexity and diversity. It affects the Authority's ability to ascertain the impacts of activities on these values, and to measure whether desired targets have been achieved. Since the Outlook Report 2009, a review of Reef Plan has provided better focus and direction for managers, including setting targets for water quality and land management improvement. The Authority's ability to address consequential and cumulative impacts, apply socio-economic and Indigenous knowledge, and set targets to benchmark performance are problematic for most management topics assessed. Targets and performance measures for coastal ecosystems are included in the Authority's *Biodiversity Conservation Strategy*,<sup>22</sup> but currently lack outcome-specific targets.

The assessment outcomes in relation to monitoring, evaluating and reporting are summarised in Table 8.28.

**Table 8.28 Effectiveness of the Authority's management arrangements to monitor, evaluate and report**

Indicator	Grade				Management tools
	Effective	Mostly effective	Partially effective	Ineffective	
4.1.1(k) monitor, evaluate and report on the: ii. impacts of activities, including the setting of targets to benchmark management					
There is effective performance monitoring to gauge progress towards the objective(s)					<p>Strengths:</p> <ul style="list-style-type: none"><li>Partnerships (for example, Reef Plan)</li><li>Research and monitoring (for example, Outlook Report, <i>Informing the outlook for Great Barrier Reef coastal ecosystems report</i>)</li></ul> <p>Weaknesses:</p> <ul style="list-style-type: none"><li>Policy documents (setting targets, addressing consequential and cumulative impacts, applying socio-economic and Indigenous knowledge)</li></ul>
Relevant standards are identified and being met					
Targets have been established to benchmark management performance					

## 8.7 Effectiveness of specific measures to protect values

This section contains an assessment of the effectiveness of the Authority's current management arrangements to protect the Region's values. It is structured around the relevant criteria in item 4.1.1 of the terms of reference for the strategic assessment, which were also part of the assessment criteria for the independent review. The relevant criteria are:

- a. identify the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area, and determine their current condition and trend, including spatial and non-spatial approaches
- k. monitor, evaluate and report on the:
  - i. condition and trends of the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area.

The grades were calculated by averaging the scores for each criterion across all management topics. A summary of the individual scores is provided in the full report of the assessment of management effectiveness, available on the Authority's website. The summary was reviewed and endorsed by the independent assessors.

### 8.7.1 Identifying values and determining their condition and trend

The Authority's understanding of the Region's values is strong for most management topics assessed. While knowledge of Indigenous values is improving within the Authority, the less tangible Indigenous heritage values, such as traditional knowledge and maintenance of cultural practice, are not well understood. For a number of historic heritage matters, the Authority has insufficient understanding of their location, condition and trend.

The Authority's five-yearly Outlook Report's assessment of condition and trend of values provides a consistent benchmark for management. In addition, the Authority's development of comprehensive vulnerability assessments and *Informing the outlook for the Great Barrier Reef coastal ecosystems*<sup>21</sup> report have substantially contributed to an understanding of condition and trend. The Authority's *Scientific Information Needs for the Management of the Great Barrier Reef Marine Park* report<sup>33</sup> identifies key information needed to better inform management of the area. This document, along with partnership agreements with key research institutions, provides an effective basis for research alignment which the assessors consider should deliver valuable outcomes for improved management.

The size, diversity and complexity of the Region means that the Authority's spatial understanding of the Region's values is limited in some places (for example, inshore ecosystems) and for some attributes (for example, Indigenous heritage, recreational fishing and community benefits). In particular, spatial approaches to understand the extent and importance of elements of matters of national environmental significance, including the outstanding universal value, and essential processes on land and sea that maintain these values are not well developed. The Authority's knowledge about the condition and trend of aesthetic value, geological and geomorphological features of outstanding universal value remains a gap. Independent reports (completed in April 2013)<sup>36,37</sup> have assisted in developing a framework to enable assessment of these features; however, information on the condition and trend of these attributes is poor.

The assessment outcomes in relation to identifying matters of national environmental significance and determining their condition and trend are summarised in Table 8.29.

**Table 8.29 Effectiveness of the Authority's management arrangements to identify values and determine their condition and trend**













Indicator	Grade				Management tools
	Effective	Mostly effective	Partially effective	Ineffective	
4.1.1(a) identify the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area, and determine their current condition and trend, including spatial and non-spatial approaches					
The values that underpin matters of national environmental significance in the Great Barrier Reef (including outstanding universal value of the World Heritage Area) are understood by managers	<div></div>	<div></div>	<div></div>	<div></div>	Strengths: <ul style="list-style-type: none"> <li>Partnerships (Reef Rescue Land and Sea Country Indigenous Partnerships Program)</li> <li>Traditional Owner agreements</li> <li>Research and monitoring (for example, Outlook Report, condition and trend for well-known and iconic species)</li> </ul>
The current condition and trend of matters of national environmental significance (spatial and non-spatial) are known by managers	<div></div>	<div></div>	<div></div>	<div></div>	Weaknesses: <ul style="list-style-type: none"> <li>Policy documents (for example, out-of-date recovery plans, policy)</li> <li>Research and monitoring (for values associated with Indigenous heritage, historic heritage, community benefits and recreational fishing)</li> <li>Spatial understanding of values relevant to matters of national environmental significance</li> <li>Understanding of cumulative impacts on outstanding universal value</li> </ul>

## 8.7.2 Monitoring, evaluating and reporting of values

Where the Authority has a good understanding of values relevant to matters of national environmental significance (for example, some aspects of biodiversity), monitoring, evaluation and reporting is partially effective (Table 8.29). Vulnerability assessments<sup>14,15</sup> and condition and trend assessments (through the Eye on the Reef program, Field Management Program and climate change activities) provide useful information on biophysical aspects of the environment, enabling benchmarking. With regard to community benefits, issues are considered under programs and policies developed by the Authority for other purposes. Currently, there are no guidelines or benchmarks for social or economic impact assessments for the World Heritage Area. The assessors advised that an overarching strategy that outlines the objectives for community benefits, while showing the linkages across the programs, would clarify the Authority's roles and responsibilities. It would also provide an improved framework to assess management effectiveness with greater accuracy.

The assessment outcomes in relation to monitoring, evaluating and reporting are summarised in Table 8.30.

**Table 8.30 Effectiveness of the Authority's management arrangements to monitor, evaluate and report on condition and trend**

Terms of reference criteria	Indicator and grade				Management tools
	Effective	Mostly effective	Partially effective	Ineffective	
4.1.1(k) monitor, evaluate and report on the: (ii) condition and trends of the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area					
There is effective performance monitoring to gauge progress towards the objective(s)					<p>Strengths:</p> <ul style="list-style-type: none"><li>Partnerships (for example, Reef Plan)</li><li>Research and monitoring (for example, Outlook Report, <i>Informing the outlook for Great Barrier Reef coastal ecosystems</i>)</li></ul> <p>Weaknesses:</p> <ul style="list-style-type: none"><li>Policy documents (setting targets)</li></ul>
Relevant standards are identified and being met					
Targets have been established to benchmark management performance					

## 8.8 Effectiveness of specific measures to protect matters of national environmental significance

A summary of the effectiveness of the Authority's management arrangements to protect each of the matters of national environmental significance is presented in Table 8.31. The results are drawn from the independent assessment of management effectiveness. The summary was reviewed and endorsed by the independent assessors.



**Table 8.31 Effectiveness of the Authority's management arrangements to protect each matter of national environmental significance**

The overall outcomes of the independent assessment of management effectiveness are presented in the left-hand column. The summary for each matter of national environmental significance was reviewed and endorsed by the independent assessors.

Management effectiveness finding for each element	World heritage properties	Great Barrier Reef Marine Park	National heritage places	Commonwealth marine areas	Listed migratory species	Listed threatened species	Wetlands of international importance
<p><b>Context:</b> <i>Understanding of values, direct and indirect threats and stakeholders is strong for most management topics assessed — with the exception of Indigenous heritage, historic heritage, recreational fishing and community benefits. In particular, defence activities, tourism, research activities and water quality protection are well understood. This reflects a solid information and research base and a very mature understanding of the key values of the Great Barrier Reef in a national and international context, along with the direct and indirect threats to those values. Understanding of cumulative and consequential impacts, as well as condition and trend, is more variable.</i></p> <p><i>An understanding of stakeholders is consistently strong across all topics (in fact, it showed the strongest performance across the entire range of assessment criteria).</i></p>					<p>The species involved are generally well understood; however, there is a limited understanding of condition and trend, especially for listed migratory shorebirds. In particular, an understanding of consequential and cumulative impacts on listed species is limited.</p>	<p>Listed threatened species are well understood and their status, condition and trend are considered in the Authority's <i>Biodiversity Conservation Strategy</i> and vulnerability assessments. However, a greater understanding of the condition and trend of inshore species is needed (for example sharks and rays and shorebirds). Consequential and cumulative impacts are not well known.</p>	<p>Partnerships with the Defence Department have provided some condition and trend information (for example, the Strategic Environmental Assessment for Defence Activities). However, a greater understanding of consequential and cumulative impacts is needed (for example, underwater noise).</p>
<p><b>Planning:</b> <i>Significant efforts have been made in planning for a number of issues such as biodiversity protection, coastal development and recreation. Some plans are still in draft form, considerably outdated or are yet to be fully implemented, although planning effectiveness is improving. In the case of coastal development, the Authority's development of the Coastal Ecosystems Assessment Framework is very positive, however the fractured nature of the planning regime is problematic and recent changes to coastal planning in Queensland have raised concerns.</i></p> <p><i>In the case of ports, the project specific nature of planning means that a Great Barrier Reef-wide strategic view to direct development and maintain and enhance values is</i></p>					<p>A lack of consistency across jurisdictions is currently the weakest aspect of planning. While threat abatement plans, recovery plans and wildlife conservation plans are in place, some are considerably outdated. The <i>Biodiversity Conservation Strategy</i> and vulnerability assessments are effective planning documents for listed migratory and threatened species.</p>	<p>A lack of consistency across jurisdictions is currently the weakest aspect of planning. While threat abatement plans, recovery plans and wildlife conservation plans are in place, some are considerably outdated. Status and trend assessments prepared by the Authority indicate that many species are continuing to decline. The <i>Biodiversity Conservation Strategy</i> and vulnerability assessments are</p>	<p>The location of the Shoalwater and Corio Bays wetland predominantly within a defence training area, fortuitously affords this area effective protection from many high risk impacts. The Strategic Environmental Assessment outlines responsibilities for the planning and implementation of environmental protection measures during the planning and conduct of defence</p>

Management effectiveness finding for each element	World heritage properties	Great Barrier Reef Marine Park	National heritage places	Commonwealth marine areas	Listed migratory species	Listed threatened species	Wetlands of international importance
<i>generally not available. Recent plans for expanding the number and size of ports in response to mining industry growth in Queensland are of concern. Lack of resourcing means an updating of the Heritage Strategy and the preparation of heritage management plans has not been satisfactorily progressed. Lack of consistency across jurisdictions is the weakest aspect of planning.</i>	Shipping Management Plan is a strategic step forward to plan for and integrate actions across the Commonwealth marine area.					effective planning documents for listed migratory and threatened species.	exercises. Comprehensive environmental management plans are in place for the Shoalwater Bay defence training area. Zoning in the Shoalwater and Corio Bays Area (outside the defence area) is the primary planning tool.
<b>Inputs:</b> <i>Adequacy of inputs is variable across management issues, being least effective for community benefits, coastal development, ports, shipping, commercial and recreational fishing, and historic heritage management. The adequacy of socio-economic and Indigenous knowledge is a problem for most issues and is among the worst performing criteria across the whole assessment. Secure resourcing is a significant ongoing issue for many management areas and a better understanding and documentation of management resource requirements is needed.</i>	Security and adequacy of resources are lacking in many areas and are insufficient to address the scale of a number of issues (i.e. consequential or cumulative impacts, heritage and community benefits). Funding for the Field Management Program has been static since 2008 and declining in real terms. Funding has not kept pace with an increase in the protected area.				Adequacy of resources (see Great Barrier Reef World Heritage Area). A large majority of biophysical information on listed species is sourced from the research community and volunteer programs.	Adequacy of resources (see Great Barrier Reef World Heritage Area). A large majority of biophysical information on listed species is sourced from the research community and volunteer programs. For listed threatened species there is a better knowledge base and more targeted programs; however, resources for these programs are not secure in the long-term (for example, Strandings database, Reef HQ turtle hospital).	Adequate biophysical information within defence training areas continues to be available for the Authority's management decisions (for example, hydrographic surveys improve knowledge of benthic habitats). The skills required to assess defence activities are limited to a few core staff. An increase in training and sharing of corporate knowledge is essential, to ensure management of the wetland is effective.
<b>Process:</b> <i>Management processes are particularly strong for defence activities. They are weakest for coastal development, community benefits, commercial and recreational fishing, ports and Indigenous heritage. Addressing consequential and cumulative impacts, application of socio-economic and Indigenous knowledge, and setting of targets to benchmark performance are problematic for most issues. As was found in the Outlook Report 2009 assessment, the extent to which consequential and cumulative impacts are being addressed and the</i>	Processes are strong where the Authority has lead management over a topic or where there is strong integration and aligned vision between jurisdictions (for example, Reef Plan). A framework to address consequential and cumulative impacts is lacking. Stakeholder engagement continues to be one				Significant recent progress has been made by completion of status, trend and vulnerability assessments for many key species and habitats. However, comprehensive targets need to be established and measured. The Marine Wildlife Stranding program remains an effective tool to respond to and monitor listed species incidents.		Comprehensive arrangements are in place for the protection of this wetland with the Department of Defence (i.e. Strategic Environmental Assessment). A framework to address consequential and cumulative impacts is still required.

Management effectiveness finding for each element	World heritage properties	Great Barrier Reef Marine Park	National heritage places	Commonwealth marine areas	Listed migratory species	Listed threatened species	Wetlands of international importance
<i>application of Indigenous knowledge remain the weakest indicators across the entire management effectiveness assessment. Stakeholder engagement and application of biophysical information are the strongest aspects of management across all issues. Note that none of the topics was scored 'ineffective'.</i>	of the strongest aspects of management.						
<b>Outputs:</b> <i>Delivery of desired outputs has been weakest for coastal ecosystems, commercial and recreational fishing and ports. They are strongest in relation to climate change and extreme weather, defence, commercial marine tourism and research activities, with a noticeable improvement in the delivery of outputs relating to water quality management compared to the Outlook Report 2009. The knowledge base of the management agencies and community has consistently improved. While the majority of management programs are progressing satisfactorily, time frames frequently slip and it is not yet clear that the programs are achieving all their desired objectives. Note that none of the topics was scored 'ineffective'.</i>	While planning documents are in place for a number of matters, it is too early to judge the success of some of these (e.g. <i>Biodiversity Conservation Strategy</i> ). Delivery of outputs and programs are strong for climate change and extreme weather and tourism. Review and update of management plans has been delayed. The pace of development is affecting the capacity to adaptively manage and achieve outputs.				While planning documents are in place for a number of matters, it is too early to judge the success of some of these (e.g. <i>Biodiversity Conservation Strategy</i> ). Review and update of plans related to threatened species has been limited.		While planning documents are in place for a number of matters, it is too early to judge the success of some of these (e.g. <i>Biodiversity Conservation Strategy</i> ). Environmental plans related to the Shoalwater and Corio Bays area are updated regularly, in partnership with the Authority.
<b>Biodiversity outcomes:</b> <i>When environmental outcomes are separated out from the broader outcome context, the biodiversity protection outcomes score drops to a partially effective rating, while the climate change and extreme weather outcomes score drops to an ineffective rating.</i>	Achievement of biodiversity outcomes is limited where drivers and activities that occur outside the Marine Park affect matters of national environmental significance. While all other elements of the management cycle may be effective, the achievements of desired outcomes for biodiversity protection are not being achieved. Despite significant progress in building the ecological resilience				Data indicates improvements in humpback whale numbers. However, five of the six species of turtles in the Great Barrier Reef have declined. Dugongs in the southern inshore area have also declined. While all other elements of the management cycle may be effective, the achievement of desired outcomes for biodiversity protection are not being achieved.	Five of the six species of turtles in the Great Barrier Reef have declined. Dugongs in the southern inshore area have also declined. While all other elements of the management cycle may be effective, the achievement of desired outcomes for biodiversity protection are not being achieved. Despite significant progress in building the ecological resilience of the Reef, and the	Achievement of biodiversity outcomes is limited where drivers and activities that occur outside the Marine Park affect matters of national environmental significance. While all other elements of the management cycle may be effective, the achievement of desired outcomes for biodiversity protection are not being achieved. Despite significant progress in building the ecological

Management effectiveness finding for each element	World heritage properties	Great Barrier Reef Marine Park	National heritage places	Commonwealth marine areas	Listed migratory species	Listed threatened species	Wetlands of international importance
	of the Reef, and the social and economic resilience of Reef industries, the current capacity of the Authority to influence outcomes is limited. The projected outlook in relation to climate change is poor.				Despite significant progress in building the ecological resilience of the Reef, and the social and economic resilience of Reef industries, the current capacity of the Authority to influence outcomes is limited. The projected outlook in relation to climate change is poor.	social and economic resilience of Reef industries, the current capacity of the Authority to influence outcomes is limited. The projected outlook in relation to climate change is poor.	resilience of the Reef, and the social and economic resilience of Reef industries, the current capacity of the Authority to influence outcomes is limited. The projected outlook in relation to climate change is poor.
<b>Overall outcomes:</b> <i>Achievement of desired outcomes (values protected, threats reduced, long-term environmental and economic sustainability) is highly variable across issues. Objectives in relation to community understanding of issues and development of effective partnerships are being achieved. Overall, the greatest concern in relation to achievement of desired outcomes relates to climate change, coastal ecosystems, ports, commercial and recreational fishing, and water quality, where management is led by other agencies and outside the Authority's jurisdiction.</i>		In spite of good systems and processes, the long-term trend for Great Barrier Reef ecosystems is still poor, and the extent to which specific initiatives can effectively address particular problems will only become clear over time. This situation highlights the importance of robust performance monitoring and adaptive management.					



## 8.9 Summary of conclusions

- The effectiveness of the Authority's current management arrangements was independently assessed using the International Union for Conservation of Nature (IUCN) World Commission on Protected Areas evaluation framework. This framework has been widely applied around the world.
- The independent assessment found the Authority is working towards effective management in all areas, and there have been considerable improvements in a number of areas since the *Outlook Report 2009*.
- The assessment recognised the difficulties for the Authority in achieving positive outcomes on the ground, given the spatial and temporal scales of the threats facing matters of national environmental significance and the diminishing resource base to implement actions.
- The assessment highlighted that the Authority required greater traction in threat reduction for an improvement in outcomes. This is dependent on the adoption of significant changes to current policies regarding coastal development, resource use and control of other human impacts, as well as sufficient resourcing to implement threat-reduction programs in the field.
- The assessment found the Authority's management of activities within the Region, for which it has direct responsibility, is effective. Its significant management attention on tourism, which operates across much of the Region, and is moderately complex, was identified as being effective, primarily because of its direct jurisdictional control.
- The independent assessment highlighted that management effectiveness challenges for the Authority were evident for those issues which are broad in scale and complex socially, biophysically and jurisdictionally (that is, port activities, shipping, climate change and extreme weather, coastal development, water quality protection, commercial and recreational fishing and Indigenous heritage) or those that are poorly resourced (for example, historic heritage).
- For the management topics of climate change and extreme weather, coastal development and water quality protection, the assessment identified particular management challenges in consistency of approaches across jurisdictions which impacts on planning.
- For commercial and recreational fishing, the assessment identified particular challenges for the Authority in the areas of monitoring and compliance, especially as they relate to addressing potential cumulative impacts.
- For Indigenous heritage, the management challenges faced by the Authority are particularly in areas of understanding the context and processes to better incorporate Indigenous heritage across the Authority's management arrangements.
- Inputs, process and outcomes were the elements of the management cycle where the Authority's current management arrangements were least likely to meet the endorsement criteria. Elements of context and planning in relation to biodiversity protection and water quality performed strongly; however, the trend of this grade depended on the continuation of key programs.
- The independent assessment found the Authority generally has a good understanding of direct and indirect impacts and, where the Authority has a high level of control over activities, its effectiveness at avoiding, mitigating and adaptively managing impacts was effective or mostly effective. Its effectiveness at halting and reversing declines and enhancing the condition of relevant matters of national environmental significance was rated as less effective, especially for topics that originate beyond the Region and are jurisdictionally complex.
- The Authority's ability to address consequential and cumulative impacts, apply socio-economic and Indigenous knowledge, and set targets to benchmark performance was assessed as problematic for most management topics.
- The overall assessment results are summarised in Table 8.32. The independent assessor's full report and recommendations are available on the Authority's website at [www.gbrmpa.gov.au](http://www.gbrmpa.gov.au).

**Table 8.32 Summary of management effectiveness results for management topics**

The overall rating for outcomes relates to the Authority's effectiveness for all seven outcome criteria (that is, outcomes relating to social, economic, biodiversity and managerial aspects). The second outcome rating relates only to the effectiveness of achieving biodiversity outcomes separate from the influence of scores for other outcomes.

Understanding the table							
Effective: 81 to 100 per cent of optimal condition	Mostly effective: 51 to 80 per cent of optimal condition	Partially effective: 21 to 50 per cent of optimal condition			Ineffective: zero to 20 per cent of optimal condition		
Management topic	Effectiveness of existing measures to protect and manage						
	Context	Planning	Inputs	Processes	Outputs	Outcomes	
						Overall	Biodiversity
Values							
Biodiversity							
Indigenous heritage							
Historic heritage							
Community benefits							
External impacts on values							
Climate change and extreme weather							
Water quality protection							
Coastal development							
Direct uses							
Tourism							
Fishing – commercial							
Fishing – recreational							
Recreation							
Port activities							
Shipping							
Defence activities							
Research activities							

## References

1. Hockings, M., Stolton, S., Leverington, F., Dudley, N. and Courrau, J. 2006, *Evaluating effectiveness: a framework for assessing management effectiveness of protected areas*, 2nd edn, IUCN, Gland, Switzerland.
2. Great Barrier Reef Marine Park Authority 2009a, *Great Barrier Reef Outlook Report 2009*, GBRMPA, Townsville.
3. Bellwood, D.R., Hughes, T.P., Folke, C. and Nyström, M. 2004, Confronting the coral reef crisis, *Nature* 429: 827-833.

4. De'ath, G., Fabricius, K.E., Sweatman, H. and Puotinen, M. 2012, The 27-year decline of coral cover on the Great Barrier Reef and its causes, *Proceedings of the National Academy of Sciences* 109(44): 17995-17999.
5. McCook, L.J., Ayling, T., Cappel, M., Choat, J.H., Evans, R.D., DeFreitas, D.M., Heupel, M., Hughes, T.P., Jones, G.P., Mapstone, B., Marsh, H., Mills, M., Molloy, F.J., Pitcher, C.R., Pressey, R.L., Russ, G.R., Sutton, S., Sweatman, H., Tobin, R., Wachenfeld, D. and Williamson, D. 2010, Adaptive management

of the Great Barrier Reef: A globally significant demonstration of the benefits of networks of marine reserves, *Proceedings of the National Academy of Sciences* 107(43): 18278-18285.

6. Convention on Biological Diversity. *Article 10 Sustainable Use of Components of Biological Diversity*, CBD, Montreal, Canada, <<http://www.cbd.int/convention/articles/default.shtml?a=cbd-10>>.

7. Great Barrier Reef Marine Park Authority 2011a, *Reef Rescue Land and Sea Country Indigenous Partnerships Program*, GBRMPA, Townsville, <<http://www.gbrmpa.gov.au/our-partners/traditional-owners/reef-rescue>>.

8. Great Barrier Reef Marine Park Authority 2005, *Great Barrier Reef Marine Park Heritage Strategy*, GBRMPA, Townsville.

9. Great Barrier Reef Marine Park Authority 2012a, *Great Barrier Reef Marine Park Strategic Plan 2012-2016*, GBRMPA, Townsville, <[http://www.gbrmpa.gov.au/\\_data/assets/pdf\\_file/0014/3281/gbrmpa-Strategic-Plan-2012.pdf](http://www.gbrmpa.gov.au/_data/assets/pdf_file/0014/3281/gbrmpa-Strategic-Plan-2012.pdf)>.

10. Great Barrier Reef Marine Park Authority 2012b, *Recreation management strategy for the Great Barrier Reef Marine Park*, GBRMPA, Townsville.

11. Great Barrier Reef Marine Park Authority 2012c, *Great Barrier Reef Climate Change Adaptation Strategy and Action Plan 2012 to 2017*, GBRMPA, Townsville.

12. Great Barrier Reef Marine Park Authority 2007, *Great Barrier Reef climate change action plan (2007-2012)*, Great Barrier Reef Marine Park Authority, Townsville.

13. Great Barrier Reef Marine Park Authority 2012d, *Climate change adaptation: Outcomes from the Great Barrier Reef Climate Change Action Plan 2007-2012*, GBRMPA, Townsville.

14. Great Barrier Reef Marine Park Authority 2012e, *Vulnerability assessments*, GBRMPA, Townsville, <<http://www.gbrmpa.gov.au/about-the-reef/biodiversity/draft-biodiversity-conservation-strategy/vulnerability-assessments>>.

15. Marshall, P.A. and Johnson, J.E. 2007, The Great Barrier Reef and climate change: vulnerability and management implications, in *Climate change and the Great Barrier Reef: a vulnerability assessment*, eds J.E. Johnson and P.A. Marshall, Great Barrier Reef Marine Park Authority and Australian Greenhouse Office, Townsville, pp. 773-801.

16. Marshall, N. 2011, Assessing Resource Dependency on the Rangelands as a Measure of Climate Sensitivity, *Society and Natural Resources* 24(10): 1-11.

17. Great Barrier Reef Marine Park Authority 2011b, *Reef health incident response system*, GBRMPA, Townsville.

18. Department of the Premier and Cabinet 2009, *Reef Water Quality Protection Plan 2009 for the Great Barrier Reef World Heritage Area and adjacent catchments*, Reef Water Quality Protection Plan Secretariat, Brisbane.

19. Great Barrier Reef Marine Park Authority 2009b, *Water quality guidelines for the Great Barrier Reef Marine Park*, GBRMPA, Townsville.

20. Great Barrier Reef Marine Park Authority 2012f, *Great Barrier Reef Coastal Ecosystems Assessment Framework*, GBRMPA, Townsville.

21. Great Barrier Reef Marine Park Authority 2012g, *Informing the outlook for Great Barrier Reef coastal ecosystems*, GBRMPA, Townsville.

22. Great Barrier Reef Marine Park Authority 2013, *Great Barrier Reef Biodiversity Conservation Strategy*, GBRMPA, Townsville.

23. Deloitte Access Economics 2013, *Economic Contribution of the Great Barrier Reef 2011-2012*, Great Barrier Reef Marine Park Authority, Townsville.

24. Pears, R.J., Morison, A.K., Jebreen, E.J., Dunning, M.C., Pitcher, C.R., Courtney, A.J., Houlden, B. and Jacobsen, I.P. 2012, *Ecological risk assessment of the East Coast Otter Trawl Fishery in the Great Barrier Reef Marine Park: technical report*, Great Barrier Reef Marine Park Authority, Townsville.

25. Taylor, S., Webley, J. and McInnes, K. 2012, *2010 State wide recreational fishing survey*, Department of Agriculture, Fisheries and Forestry, Brisbane.

26. Great Barrier Reef Marine Park Authority 2010, *Responsible reef practices for tourism operators in the Great Barrier Reef Marine Park*, Revised edition, GBRMPA, Townsville.

27. Great Barrier Reef Marine Park Authority 2012h, *Guidelines for the use of hydrodynamic numerical modelling for dredging projects in the Great Barrier Reef Marine Park*, GBRMPA, Townsville, <[http://www.gbrmpa.gov.au/\\_data/assets/pdf\\_file/0018/26532/Guidelines-on-Hydrodynamics-Modelling-15-Aug-2012.pdf](http://www.gbrmpa.gov.au/_data/assets/pdf_file/0018/26532/Guidelines-on-Hydrodynamics-Modelling-15-Aug-2012.pdf)>.

28. Turner, M., Beattie, S. and Edison, K. 2011, *Great Barrier Reef Marine Park Authority Environment and Science Coordinators Report 2011-2012*, Great Barrier Reef Marine Park Authority, Townsville.

29. Polglaze, Griffin, Miller and Associates 2012, *Great Barrier Reef Shipping: Review of environmental implications*, PGM Environment, Safety Bay, Western Australia.

30. Great Barrier Reef Marine Park Authority 2011c, *Grounding of the Shen Neng 1 on Douglas Shoal, April 2010: Impact Assessment Report*, GBRMPA, Townsville.

31. Department of Defence and Great Barrier Reef Marine Park Authority 2008, *Management agreement between Department of Defence and the Great Barrier Reef Marine Park Authority on implementation of the strategic environmental assessment of defence activities in the Great Barrier Reef Marine Park*, GBRMPA, Townsville.

32. URS Australia Pty Ltd 2006, *Strategic Environmental Assessment of Defence Activities in the Great Barrier Reef World Heritage Area*, Directorate of Environmental Stewardship, Department of Defence, Canberra.

33. Great Barrier Reef Marine Park Authority 2009c, *Scientific information needs for the management of the Great Barrier Reef Marine Park 2009-2014*, GBRMPA, Townsville.

34. Great Barrier Reef Marine Park Authority 2004, *Policy on managing scientific research in the Great Barrier Reef Marine Park*, GBRMPA, Townsville, <[http://www.gbrmpa.gov.au/\\_data/assets/pdf\\_file/0017/3860/gbrmpa\\_ManagingScientificResearchPolicy\\_2004.pdf](http://www.gbrmpa.gov.au/_data/assets/pdf_file/0017/3860/gbrmpa_ManagingScientificResearchPolicy_2004.pdf)>.

35. Day, J., Fernandes, L., Lewis, A., De'ath, G., Slegers, S., Barnett, B., Kerrigan, B., Breen, D., Innes, J., Oliver, J.K., Ward, T. and Lowe, D. 2000, The Representative Areas Program - protecting the biodiversity of the Great Barrier Reef World Heritage Area, in *Proceedings 9th International Coral Reef Symposium, 23-27 October, Bali*, eds. Anonymous, Ministry of Environment, Indonesian Institute of Sciences, International Society for Reef Studies, Indonesia, pp. 687-696.

36. Context Pty Ltd 2013, *Defining the aesthetic values of the Great Barrier Reef*, Department of Sustainability, Environment, Water, Population and Communities, Canberra.

37. Whiteway, T., Smithers, S., Potter, A. and Brooke, B. 2013, *Geological and geomorphological features of Outstanding Universal Value in the Great Barrier Reef World Heritage Area: Technical report prepared for the Department of Sustainability, Environment, Water, Population and Communities*, Geoscience Australia and James Cook University, Townsville.







## Chapter 9

---

### Demonstration case studies



## Table of contents

<b>9.1</b>	<b>Case studies selected.....</b>	<b>9-3</b>
9.1.1	Topics chosen.....	9-3
9.1.2	Meeting the selection criteria.....	9-4
<b>9.2</b>	<b>Assessing management effectiveness.....</b>	<b>9-5</b>
<b>9.3</b>	<b>Dugongs.....</b>	<b>9-6</b>
9.3.1	Significance .....	9-6
9.3.2	Key issues .....	9-6
9.3.3	Current management and its effectiveness.....	9-8
9.3.4	Outcomes and potential improvements .....	9-9
<b>9.4</b>	<b>Corals .....</b>	<b>9-10</b>
9.4.1	Significance .....	9-10
9.4.2	Key issues .....	9-10
9.4.3	Current management and its effectiveness.....	9-11
9.4.4	Outcomes and potential improvements .....	9-12
<b>9.5</b>	<b>Islands .....</b>	<b>9-13</b>
9.5.1	Significance .....	9-13
9.5.2	Responsibilities for islands .....	9-13
9.5.3	Key issues .....	9-14
9.5.4	Current management and its effectiveness.....	9-15
9.5.5	Outcomes and potential improvements .....	9-17
<b>9.6</b>	<b>Princess Charlotte Bay .....</b>	<b>9-18</b>
9.6.1	Significance .....	9-18

9.6.2	Key issues .....	9-20
9.6.3	Current management and its effectiveness .....	9-20
9.6.4	Outcomes and potential improvements.....	9-21
<b>9.7</b>	<b>Cairns Planning Area.....</b>	<b>9-21</b>
9.7.1	Significance .....	9-21
9.7.2	Key issues .....	9-22
9.7.3	Current management and its effectiveness .....	9-22
9.7.4	Outcomes and potential improvements.....	9-24
<b>9.8</b>	<b>Mackay Whitsunday — water quality improvement.....</b>	<b>9-25</b>
9.8.1	Significance .....	9-25
9.8.2	Key issues .....	9-25
9.8.3	Current management and its effectiveness .....	9-27
9.8.4	Outcomes and potential improvements.....	9-29
<b>9.9</b>	<b>Keppel Bay .....</b>	<b>9-29</b>
9.9.1	Significance .....	9-29
9.9.2	Key issues .....	9-30
9.9.3	Current management and its effectiveness .....	9-31
9.9.4	Outcomes and potential improvements.....	9-32
<b>9.10</b>	<b>East Coast Trawl Fishery .....</b>	<b>9-32</b>
9.10.1	Significance .....	9-32
9.10.2	Key issues .....	9-33
9.10.3	Current management and its effectiveness .....	9-36
9.10.4	Outcomes and potential improvements.....	9-37
<b>9.11</b>	<b>Summary of conclusions and guidance for future management .....</b>	<b>9-37</b>
<b>References.....</b>		<b>9-38</b>

Cover page image: A dugong (*Dugong dugon*)

## **Extract from Great Barrier Reef Region Strategic Assessment terms of reference**

### **4.1 Demonstration cases**

4.2.1 *Develop demonstration cases to assess in finer detail the effectiveness of the Program to protect and manage the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area, and to guide improvements to the Program.*

4.2.2 *Demonstration cases are to be chosen by the Australian and Queensland governments. Criteria that will be used to guide this selection process include, but are not limited to:*

- a) *where multiple impacts are acting or predicted to act upon a region, locality or value*
- b) *to examine a specific management approach or method to identify a set of values/attributes or to assess a range of impacts/pressures*
- c) *to demonstrate connectivity across coastal and marine systems*
- d) *to demonstrate the integration of environmental, social, cultural and economic benefits in decision making*
- e) *to improve understanding of factors affecting Great Barrier Reef ecosystem resilience*
- f) *where lessons or outcomes could transfer to other areas*
- g) *opportunities to build capacity for future management*
- h) *to examine the effectiveness of management across local, Queensland and Australian government jurisdictions.*

The terms of reference for the strategic assessment provide for demonstration case studies to be undertaken. The purpose of the case studies is to assess in finer detail the effectiveness of current management arrangements to protect and manage the relevant matters of national environmental significance, including outstanding universal value, and to guide improvements to management arrangements. Demonstration cases also provide an opportunity to examine cumulative impacts on values at a local or regional scale and their wider implications for management by government agencies and the community.

This chapter sets out the case study topics and their alignment with the terms of reference. It presents the case studies, including their outcomes and potential improvements for future management. The technical reports that support each case study are available on the Authority's website at [www.gbrmpa.gov.au](http://www.gbrmpa.gov.au). The *Ecological risk assessment of the East Coast Otter Trawl Fishery in the Great Barrier Reef Marine Park: technical report*<sup>1</sup> supports the East Coast Trawl Fishery case study.

## **9.1 Case studies selected**

### **9.1.1 Topics chosen**

The eight demonstration case studies are:

- dugongs
- corals
- islands
- Princess Charlotte Bay
- Cairns Planning Area
- Mackay Whitsunday — water quality improvement
- Keppel Bay
- East Coast Trawl Fishery.

The case studies examine individual values underpinning matters of national environmental significance (corals, dugongs and islands); major uses of the Region (tourism and fishing); a major management initiative (water quality improvement in the Mackay Whitsundays); and the combined effect of multiple uses on values in specific locations (Keppel Bay and Princess Charlotte Bay). The demonstration case studies on dugong, islands and Mackay Whitsunday water quality improvement have been prepared collaboratively with the Queensland Government for development of both strategic assessment reports.

A joint demonstration case study on shipping was also proposed, however this has been superseded by development of the North-East Shipping Management Plan. This assesses the effectiveness of current risk control measures and identifies new and enhanced control measures that will be required, commensurate with the projected shipping growth in the Region (see Section 5.4.6, in Chapter 5).



Recognising the extensive separate investigations on ports and their management being undertaken in parallel to the strategic assessment, a demonstration case study on ports was not carried out. As they have become available, the preliminary outcomes of these investigations have been taken into account in the strategic assessment. The investigations include:

- *Independent review of the Port of Gladstone*: This review was commissioned in response to the June 2012 decision of the World Heritage Committee. A panel of experts is examining and reporting on environmental management and governance arrangements for the port and may make recommendations to the Environment Minister relevant to the review's terms of reference. This may include recommendations relating to optimisation of port development and operations. At the time of writing this report, the Review was yet to conclude.
- *Improved dredge material management for the Great Barrier Reef Region*<sup>2</sup>: This has provided proposed strategies for improved material management for the major ports and one boat harbour/marina (Rosslyn Bay) in the Great Barrier Reef World Heritage Area. Independent consultants undertook the project under the Sustainable Regional Development Program, funded by the Australian Government.
- *Identification of impacts and proposed management strategies associated with offshore ship anchorages in the Great Barrier Reef World Heritage Area*<sup>3</sup>: This has identified environmental impacts and risks from anchorages for each port and potential management options to avoid and mitigate environmental, social and heritage impacts from existing and future ship anchorages, given likely increases in future shipping. This is also a Sustainable Regional Development Program project, undertaken by independent consultants.
- *Environmental best practice port development: an analysis of international approaches*<sup>4</sup>: This has identified international benchmarks in the management of environmental impacts of ports and their potential application in an Australian context.

### 9.1.2 Meeting the selection criteria

Item 4.2.2 of the terms of reference for the strategic assessment sets out criteria to guide selection of demonstration case studies. The way in which each of the case studies meets these criteria is outlined in Table 9.1. Furthermore, the selection of case studies considered matters consistently raised by stakeholders, key risks identified in the *Great Barrier Reef Outlook Report 2009*<sup>5</sup>, a geographic spread of areas and the availability of sufficient information for a detailed analysis.

**Table 9.1 Meeting the selection criteria for demonstration case studies**

Terms of reference criteria	Demonstration case study
a) Where multiple impacts are acting or predicted to act upon a region, locality or value	Place-based case studies such as Princess Charlotte Bay, Cairns Planning Area and Keppel Bay provide insight into areas where there are multiple impacts on a locality. The studies on dugongs, corals and islands examine multiple impacts on values.
b) To examine a specific management approach or method to identify a set of values/attributes or to assess a range of impacts/pressures	Each of the case studies examines a number of management approaches: from research and monitoring to policies, planning and compliance. The Cairns Planning Area case study examines the effectiveness of the plan of management for that area. The East Coast Trawl Fishery study examines the effectiveness of all management approaches for that fishery. The coral and dugong demonstration cases have incorporated qualitative impact assessment methods.
c) To demonstrate connectivity across coastal and marine systems	Connectivity across coastal and marine systems is addressed in all demonstration cases.
d) To demonstrate the integration of environmental, social, cultural and economic benefits in decision making	Each of the place-based case studies examines how community benefits have been integrated into decision making, especially the Cairns Planning Area study. The trawl fishery case study considers the integration of social and economic considerations.
e) To improve understanding of factors affecting Great Barrier Reef ecosystem resilience	The case studies on dugongs and corals are particularly relevant to an examination of resilience.
f) Where lessons or outcomes could transfer to other areas	All case studies provide lessons and outcomes that can be transferred to other areas.
g) Opportunities to build capacity for future management	The outcomes of all the case studies will inform capacity building for future management.
h) To examine the effectiveness of management across local, Queensland and Australian government jurisdictions	All the case studies address effectiveness of management.



## 9.2 Assessing management effectiveness

An assessment of the effectiveness of management has been undertaken by the Authority for each demonstration case (except the Mackay Whitsunday — water quality improvement and East Coast Trawl Fishery). These assessments were reviewed by the same consultants who undertook the independent review of management effectiveness<sup>6</sup> (see Chapter 8). Direct excerpts from their report are shown in this chapter in italics. For the Mackay Whitsunday — water quality improvement and East Coast Trawl Fishery, direct excerpts are reproduced from Chapter 8.

The assessment approach followed the framework for evaluating management effectiveness described in Chapter 8. Current management activities relevant to each stage of the management cycle (context, planning, inputs, process, outputs and outcomes) were considered. The effectiveness of the management arrangements to achieve desired results for each of the six elements were qualitatively assessed on a four point rating scale (81 to 100 per cent; 51 to 80 per cent; 21 to 50 per cent; one to 20 per cent) and graded (effective; mostly effective; partially effective; ineffective).

The scope of this assessment included management activities within the Authority's jurisdiction and any joint management arrangements with the Queensland Government (for example, joint Marine Park permitting, and joint compliance and enforcement through the Field Management Program).

The relationship between the demonstration cases and the management topics and tools assessed are summarised in Table 9.2 and Table 9.3.

**Table 9.2 Matching demonstration case studies to management topics**

Management topic	Dugongs	Corals	Islands	Princess Charlotte Bay	Cairns Planning Area	Mackay Whitsunday — water quality improvement	Keppel Bay	East Coast Trawl Fishery
Biodiversity protection	•	•	•	•	•	•	•	•
Indigenous heritage	•		•	•	•		•	
Historic heritage			•					
Community benefits	•	•	•	•	•	•	•	•
Climate change and extreme weather	•	•	•		•		•	•
Water quality protection	•	•			•	•	•	
Coastal development	•	•		•	•	•	•	•
Tourism		•	•		•	•	•	
Fishing — commercial	•			•	•		•	•
Fishing — recreational	•			•	•		•	
Recreation		•	•	•		•	•	
Port activities	•	•			•			
Shipping				•	•	•	•	
Defence activities	•							
Research activities	•	•	•				•	

**Table 9.3 Matching demonstration case studies to management tools**

Management tool	Dugongs	Corals	Islands	Princess Charlotte Bay	Cairns Planning Area	Mackay/Whitsunday — water quality improvement	Keppel Bay	East Coast Trawl Fishery
Act and Regulations	●	●	●	●	●	●		
Zoning Plan	●	●	●	●	●	●	●	●
Plans of management	●	●	●		●	●		
Permits		●	●	●	●	●	●	
Traditional Owner agreements	●		●	●			●	
Compliance	●	●	●	●	●	●	●	●
Site infrastructure		●	●		●	●	●	
Fees and charges					●			
Policy	●				●	●		
Partnerships	●	●	●	●	●	●		●
Stewardship and best practice	●	●	●	●	●	●	●	●
Education and community awareness	●	●	●		●	●	●	
Research and monitoring	●	●	●		●	●	●	●
Reporting	●	●	●			●		

## 9.3 Dugongs

### 9.3.1 Significance

The Great Barrier Reef World Heritage Area is home to a globally significant population of dugongs, which was one of the reasons why it was inscribed on the World Heritage List in 1981.<sup>7</sup> Dugongs, together with the seagrass habitats upon which they depend, contribute to the recognition of the Reef's outstanding universal value, including its significant ongoing ecological and biological processes, and significant natural habitats for the conservation of biological diversity.

Dugongs are of great cultural, spiritual and social importance, especially to the Reef's Traditional Owners. These large marine mammals are a matter of national environmental significance as a listed migratory species under the *Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act), and are listed as a marine species under that Act. Dugongs are listed as vulnerable under the *Nature Conservation Act 1992* (Qld) and as a protected species under the Great Barrier Reef Marine Park Regulations 1983.

Dugongs are found in the coastal waters of some 40 countries and territories, but their status is either declining, locally extinct, or unknown in most of their range.<sup>8</sup> Australia's northern waters (from Moreton Bay in southern Queensland to Shark Bay in Western Australia) contain approximately one quarter of the world's dugong habitats. Australia is the only developed country with a significant dugong population, and with extensive areas of coastline at relatively low risk from coastal development.<sup>9,10</sup>

### 9.3.2 Key issues

Dugongs feed almost exclusively on seagrass and they must consume large quantities each day to fulfil their energy requirements.<sup>11</sup> The dugong's preference for inhabiting inshore areas exposes them to a range of human-related marine and land-based activities.<sup>10</sup>

As long-lived, slow-breeding animals, dugongs have slow rates of recovery from population declines. Survival of adult dugongs must be more than 90 per cent each year to maintain a population.<sup>12</sup> Population simulations based on adult survival estimates from manatees (the most closely related group of species) and empirical reproductive data from dugongs in the Torres Strait estimate the maximum population growth rate to be up to six per cent annually; in most cases the population growth rate will be much less than this.<sup>13,14</sup>


Dugong life history suggests that the impact of mortality will typically be more significant than habitat loss. However these two categories of threat are related; habitat loss can lead to starvation and mortality. The greatest impacts on dugong populations in the World Heritage Area vary with location. Along the urban coast the most

serious impacts are habitat loss and degradation including impacts from: cyclone activity and extreme weather; nutrients, pesticides and sediment from catchment run-off; clearing or modifying of coastal habitats; coastal reclamation; direct impacts of dredging; dumping and resuspension of dredge material.<sup>10</sup> Dugongs are also affected directly by disease, their incidental capture in nets (death of discarded species from the commercial net fishery and the Queensland Shark Control Program), marine debris, boat strike, illegal fishing and poaching, and hunting for traditional use to varying degrees.<sup>10</sup> Along the coast of Cape York, mortality from Indigenous hunting and incidental drowning in commercial gill nets are the main impacts, however, hunting is restricted to inshore waters around settlements and a very high percentage of dugong habitat in the region is not subject to hunting.

Commercial harvesting, which began in Queensland in 1847, took a significant toll on the species, particularly south of Cairns, where virtually all of the operations were based apart from some operations at Indigenous communities to supply oil for medicinal use. While commercial dugong harvesting was a cottage industry, there were large harvests of the animals, especially in the late 1800s to 1930s. In some years more than 100 dugongs were harvested from Moreton Bay (south of the Region).<sup>15</sup> Harvesting was banned in Queensland in 1969, however, government-sponsored operations based in a few Indigenous communities continued until the mid-1970s.

The introduction and widespread use of monofilament fishing nets from about the 1960s also resulted in a significant incidental capture of dugong, prompting the progressive introduction of Dugong Protection Areas, coastal zoning and changes to net designs and gear over the past 20 years.

Despite management actions to protect them, dugong population numbers in the World Heritage Area south of Cooktown have not recovered. In fact, the numbers of dugongs continued to decline between the early 1960s and 1990s. By the 1990s, it was suggested dugong numbers had declined by more than 90 per cent along the coast of Queensland south of Cooktown.<sup>10,16,17</sup> Evidence of decline was derived retrospectively from their incidental catch in the Queensland Shark Control Program between 1962 and 1999, during which time at least 837 dugongs were killed.<sup>18</sup> The trends in the average number of dugongs caught in nets at 47 beaches showed a decline of 8.7 per cent per year between 1962 and 1999 (Figure 9.1), which was assumed to reflect the rate of population decline from all causes.<sup>17,19</sup> Initiatives began in 1992 to reduce this impact.<sup>18</sup> Baited hooks on drumlines have replaced shark nets in many localities. Drumlines still catch the targeted shark species, but catch few dugongs or cetaceans.



**Figure 9.1 Dugongs caught in the Queensland Shark Control Program, 1962–1999**

Profile of the annual estimated mean numbers of dugongs (log-linear model and 95 per cent confidence intervals) caught in the Queensland Shark Control Program from six shark netting contract areas at 47 beaches (green dots) between Cairns and the Gold Coast for the period 1962 to 1999.<sup>16</sup> The estimated rate of decline averages 8.7 per cent per year. There is no statistical evidence that the changes to the program introduced after the 1992 review of practices<sup>18</sup> changed the pattern of declining catches up to 1999, however the power to detect any change would be weak.

Aerial surveys suggested there were about 2000 dugongs in the World Heritage Area south of Cooktown in 2005 and that the decline had halted.<sup>20</sup> This estimate is a standardised relative index of dugong abundance and is less than the actual abundance. Although this was only a small fraction of pre-European levels, the population was thought to have stabilised.<sup>5,20,21,22</sup> However, widespread destruction of seagrass meadows (from heavy flooding and cyclone Yasi in 2010–11, coupled with two preceding years of higher than average rainfall and the chronic impact of poor water quality),<sup>23,24</sup> led to a further decline as dugongs died<sup>25</sup> or emigrated from the affected areas. The standardised index of dugong relative abundance gained from aerial surveys in 2011 showed a decline to approximately 600 animals in waters between the Daintree River and the southern limit of the World Heritage Area.<sup>26</sup> The condition of the southern population is assessed as poor and declining (with good confidence in both assessments).<sup>5,20,21,22</sup> In contrast, dugong populations south of the World Heritage Area in Hervey Bay and Moreton Bay have not declined since the previous aerial surveys in 2005.<sup>26</sup>

North of Cooktown, the current status of dugongs is considered to be good with a stable population trend (with good confidence in both assessments).<sup>17,19,27,28,29</sup> In 2006, standardised index of relative abundance of dugongs between Cooktown and Cape York was approximately 8800.<sup>19</sup> Outside the World Heritage Area to the north, the relative abundance of the population in the Torres Strait is estimated to be greater than 14,000 individuals.<sup>19</sup>

### 9.3.3 Current management and its effectiveness

Dugongs have been a focus of management agencies for many decades, with high level leadership from the Great Barrier Reef ministerial forum. In the 1980s, the Authority's zoning plans protected some important dugong habitats in Marine National Park (green) zones and Preservation (pink) zones. The level of protection was increased significantly in 1997 in response to the long-term decline in the dugong population south of Cooktown and the ongoing and unsustainable levels of interactions between dugongs and fishing nets and shark control nets.<sup>30</sup> These measures established 16 Dugong Protection Areas under Queensland fisheries legislation which imposed spatial and fishing gear restrictions and prohibited the use of some types of fishing nets.<sup>31</sup> Additionally, changing shark control nets to drumlines has reduced the number of interactions with dugongs.

Soon after, measures to protect seagrass habitats from trawling were introduced, as well as netting restrictions and net attendance rules to reduce the incidental capture of dugong within the East Coast Inshore Fin Fish Fishery. The use of nets in the Queensland Shark Control Program was also reviewed<sup>18</sup> and modified.<sup>32</sup> Other management arrangements were introduced by the Department of Defence for activities in the Shoalwater Bay Defence Training Area.<sup>33</sup> While there is limited traditional hunting of dugong as a native title right, some Traditional Owners have voluntarily adopted reduced levels of hunting under Traditional Use of Marine Resources Agreements, with some clans choosing not to hunt this species at all.

Dugong habitats were explicitly taken into consideration when the Great Barrier Reef Marine Park was rezoned in 2003.<sup>34</sup> As a result, approximately 96 per cent of high conservation value dugong habitats are highly protected from extractive industries,<sup>35,36</sup> and 24 per cent of known shallow water seagrass meadows are included in highly protected green zones which prohibit unpermitted extractive activities.<sup>37</sup> While it had been recognised for some time that shallow water seagrass meadows within Dugong Protection Areas are important to dugong and are at risk from declining water quality,<sup>38</sup> it is now also recognised that deeper water seagrass meadows are used by dugongs as a refuge and when preferred shallow water meadows are affected by floods or cyclones.<sup>26</sup> While a decline in shallow inshore seagrass meadows south of Cooktown is recorded, there are major gaps in our understanding of the condition, abundance and distribution of deeper water seagrass meadows, which are likely to be more important to dugongs than previously supposed. Major efforts to improve water quality through the *Reef Water Quality Protection Plan* (Reef Plan) are expected to have positive outcomes for seagrass in the medium term.<sup>39,40</sup>

The independent review of the Authority's assessment of management effectiveness concluded:

- **Context** is effective. *Evidence presented on legal status and reasons for designation of dugong as a matter of national environmental significance, status and trend in dugong population numbers and distribution; nature, extent and impact of threatening processes; stakeholder interests all indicate that the context of dugong conservation and management is well understood by the Authority.*
- **Planning** is mostly effective. *The demonstration case provides evidence of multiple levels of planning for dugong conservation over a 40-year period starting with the cessation of a commercial dugong fishery in 1969 and with a turtle and dugong conservation strategy first prepared in 1994. Dugong Protection Areas date back to 1997 and recovery plans [Qld] to 1999. Other plans, policies, regulations and strategies have been developed and implemented by Commonwealth and state agencies to address the major anthropogenic impacts on dugong. Protection and enhancement of dugong habitat has been addressed through the rezoning and water quality improvement plans. Management of impacts of coastal activities on the extent and condition of seagrass beds remains as an area where improved planning and regulation could improve prospects for the species.*
- **Inputs** are only partially effective. *Resources for key management actions (for example, mitigating impacts of coastal land management), field management (surveillance and compliance, emergency response) and monitoring of dugong populations and habitat condition are either currently inadequate or not secure in the long term.*
- **Process** is mostly effective. *Plans and strategies developed for dugong management are generally being implemented. An effective partnership with Indigenous people is being developed and the number of Traditional Use of Marine Resources Agreements is being extended to cover the main areas of relevance. The ways in which the plans and strategies are being implemented is appropriate. The methods used in surveillance and enforcement are of a high standard, but field management capacity to ensure a high level of compliance is lacking.*
- The delivery of **outputs** is mostly effective; however *implementation of relevant plans is being hampered by a shortage of resources.*
- Achieving desired **outcomes** overall is partially effective. *Concerns relating to the cumulative impacts of climate change and extreme weather and increased development in coastal ecosystems limit the effectiveness of planning and management measures introduced to conserve dugong populations in the Region. Biodiversity outcomes for dugong management are partially effective. Consideration of the southern population of dugong in the Great Barrier Reef Region alone would lead to a conclusion that*



*management had been ineffective in conserving the species. However, the fact that the larger northern population is considered to be stable and in good condition means that the overall biodiversity assessment should be rated as partially effective.*

### 9.3.4 Outcomes and potential improvements

Dugong numbers today are the result of cumulative impacts and actions over a long time.<sup>10</sup> Temporal changes in local and regional dugong population estimates are compounded by large-scale dugong movements. This is a challenge for management because trends in numbers can take decades to detect with a high degree of statistical confidence, and it may take decades to gauge the effectiveness of management strategies. Indeed it may be impossible to detect the effect of management because of the confounding effects of extreme weather events.

Ongoing and effective management action in the World Heritage Area is required to protect dugong populations and, most critically, to halt and reverse the decline in the southern populations. Effective management will depend on minimising not only short-term or acute impacts, but also cumulative or chronic impacts that may seem insignificant in isolation. The influence of climate change and proposed urban and industrial expansion throughout the Region means greater engagement of all levels of government and the community will be needed to achieve recovery of dugongs. Because dugong movements can be substantial, habitat connectivity and integrity are critical, and management efforts must be coordinated across local, state, national and international levels to ensure dugongs are protected throughout their range.

Impacts of greatest significance to dugongs south of Cooktown are habitat loss and degradation from cyclone activity and extreme weather; sediment, nutrient and pesticides from catchment run-off; modification of supporting terrestrial habitats; coastal reclamation; dredging and dumping and resuspension of dredge material; disease; vessel strike and vessel disturbance; death from incidental capture in nets (commercial net fishery and shark control program); and marine debris. North of Cooktown, seagrass is exposed to fewer water quality-related impacts. There, impacts to dugongs include incidental capture in nets (commercial net fishery), illegal fishing and poaching, and hunting for traditional use, habitat loss and degradation from cyclone activity and extreme weather and disease.

Mapping<sup>35,36</sup> and qualitative models<sup>41</sup> have been used to aid understanding of cumulative impacts on dugongs. The condition for dugong populations is projected to decline to very poor in the southern World Heritage Area, and to remain good in the north in coming decades. Model outputs based on a possible increase in the intensity of storms and ocean warming predict a likely downward trend in dugong populations, even with a possible reduction in nutrients and sediments (based on Reef Plan targets), unless direct human-induced dugong mortality from all sources is reduced at the same time. To enable population recovery in the World Heritage Area south of Cooktown, modelling<sup>42</sup> suggests an annual human-induced mortality limit of less than 10, with a target of zero.<sup>17,20</sup> This target will be very difficult to achieve — for example, five dugong deaths from human activities were reported south of Cooktown in 2011.<sup>43</sup> Sustainable limits for human-induced mortality have been estimated to be between 56 and 112 for dugongs in the World Heritage Area north of Cooktown (with recommendations to strive for the lower limit).<sup>19</sup>

#### Enhancing protection and restoration

A critical management action for dugongs across their range in the Great Barrier Reef is to maintain, enhance and restore the health of seagrass meadows, which they rely on for food. Risks to the main seagrass meadows in Dugong Protection Areas from poor water quality were identified in 2001, based on the best knowledge at that time. This issue is being addressed by actions under Reef Plan.<sup>39</sup> The importance of landscape connectivity between terrestrial and marine systems for maintaining healthy Great Barrier Reef inshore biodiversity has been recognised<sup>44</sup> with increased investment by the Australian Government under Reef Rescue 2013–2018.<sup>45</sup>

#### Minimising impacts

It is also critical that impacts on dugongs and their habitat are reduced and minimised. Strengthening a number of ongoing management actions is important and the wider community can play a more prominent role in formulation and implementation of management arrangements to recover dugong populations. For example, continuously improving net fishing techniques, with the cooperation of fishers, technology and practices (such as, codes of practice) is important to reduce drowning in fishing nets.

Community action and compliance activities at a local and regional scale are needed to help reduce and minimise mortality or ill health of dugongs (for example, from vessel strike, marine debris, disease, noise and interference). Traditional Use of Marine Resources Agreements will continue to act as a valuable tool to facilitate sustainable traditional hunting and support the role of Indigenous communities and rangers in compliance and enforcement activities.

In addition, supporting the objectives of the Queensland Government's current net buyback program to reduce fishing effort in the East Coast Inshore Fin Fish Fishery is important, given remaining concerns about set mesh netting, the potential for the mortality of vulnerable species in nets, and some failures to report these interactions in the Lockhart River and Princess Charlotte Bay areas. Another potential initiative could include a revision of apparatus used in the Queensland Shark Control Program to further reduce risks to dugongs, while meeting the public safety imperative.

## Improving adaptive management

Adaptive management requires an ongoing commitment to regular monitoring, evaluation and review of the abundance and distribution of dugong and seagrass to understand trends. A greater understanding of cumulative impacts, including impacts of extreme weather, should lead to an improvement in the adequacy of measures to avoid or mitigate impacts in important dugong areas. This could even include investigating ways of restoring and rehabilitating seagrass meadows.

Continuous improvement of best practice standards for all activities likely to affect dugong (for example, in relation to vessel operation, vessel waste management, recreational use, tertiary sewage treatment and re-cycling, urban storm water management, and erosion and sediment control) will help ensure activities are undertaken to the highest standard, based on the most up-to-date information. Part of this continuous improvement is the implementation of new technologies to reduce illegal fishing and increase compliance with fisheries management arrangements, as well as the synthesis of information on trends and regional and local differences. Traditional Use of Marine Resources Agreements and other Indigenous management tools such as community management plans will continue to be a valuable tool, providing for the exchange and incorporation of traditional ecological knowledge into management.

Continued commitment to scientific and community-based monitoring, reporting and stewardship programs will allow management agencies to understand the effectiveness of their actions. Such programs may include those with a focus on understanding movements of dugongs and condition of seagrass — providing real-time information to managers, industry and the public; long-term, structured, large-scale aerial surveys — providing information on population abundance, distribution and trends; and a better understanding of existing and emerging impacts on dugongs (for example, noise).

## 9.4 Corals

### 9.4.1 Significance

Corals contribute to the recognition of the Reef's outstanding universal value for all four of the natural criteria for World Heritage listing: exceptional natural beauty, significant geomorphic features, significant ongoing ecological and biological processes, and significant natural habitats for the conservation of biological diversity.

The Great Barrier Reef is the largest coral reef ecosystem in the world, stretching 2600 kilometres along the north-eastern coast of Australia. Coral reefs themselves cover an area of about 26,000 square kilometres and make up seven per cent of the Great Barrier Reef Marine Park. The Reef is also one of the world's most diverse ecosystems. A network of more than 2900 separate reefs forms 30 distinct reef bioregions (see Figure 4.4 in Chapter 4), comprising a total of 10 per cent of all the world's coral reefs.<sup>46</sup> The Reef's habitat complexity is founded on about 410 species of hard coral, which support a rich diversity of reef-associated species.<sup>5</sup>

The coral reef environment supports a range of Reef-dependent industries and uses, and directly underpins the community's cultural, social and economic wellbeing. The economic contribution to the Australian economy generated by tourism, recreation, commercial fishing and scientific research in the Reef catchment and the World Heritage Area in 2012 was \$5.7 billion. It also supported about 69,000 full-time equivalent jobs.<sup>47</sup>

Globally, coral reefs are under serious pressure — 75 per cent of the world's coral reefs are currently threatened by local and global pressures, more than 90 per cent of the world's reefs will be threatened by 2030 and nearly all of them could be at risk in less than 40 years.<sup>48</sup>

### 9.4.2 Key issues

Despite the Great Barrier Reef being recognised as one of the best managed coral reef ecosystems in the world,<sup>49</sup> research and anecdotal information indicates there have been substantial changes to coral cover and species composition on inshore fringing reefs dating from at least the 1920s<sup>50</sup> (see Section 7.1.2 for an example and discussion of the decline of inshore reefs) and a 40 per cent decline in the past 27 years.<sup>51,52,53</sup> While there is debate over the exact extent, cause and severity of this loss,<sup>51</sup> there is no doubt that coral cover overall and specifically in some regions is declining.


Long-term monitoring by the Australian Institute of Marine Science since 1986 shows average hard coral cover across the Great Barrier Reef has declined from 28 per cent to 13.8 per cent — two-thirds of that loss has occurred since 1998.<sup>54</sup> A majority of the loss has been recorded since 2006 in central and southern regions of the Great Barrier Reef. North of Cooktown, coral reefs are in good condition, with hard coral cover remaining relatively stable over the past 27 years. The status of coral reefs in the Region reflects the pattern of extreme weather events, coastal development and water quality. There is no long-term data on the condition and trend of deep water (more than 30 metres) reefs.<sup>55</sup>

The major drivers of loss of coral cover on a Reef-wide scale are:

- Outbreaks of the crown-of-thorns starfish (*Acanthaster planci*).<sup>56</sup> Large-scale crown-of-thorns starfish outbreaks were first recorded in the 1960s.<sup>57</sup> Since then, there have been three major outbreaks (1962 to 1976, 1978 to 1990, 1993 to 2005<sup>58</sup>).
- Direct impacts of cyclones.<sup>5,56</sup> For example, cyclone Hamish was a major factor in reducing the average coral cover on southern offshore reefs in the Swains region of the World Heritage Area from 35 per cent in 2006 to 8 per cent.<sup>54</sup>

- Declining water quality in the Great Barrier Reef lagoon — including an increase in nutrients, pollutants, sediment and freshwater — from catchment run-off, urban and industrial discharge and dredge material disposal.<sup>52,53,59</sup> Coral disease has also emerged as a chronic problem for many southern inshore reefs after flood events.<sup>60</sup>
- Increasing sea surface temperatures, which have induced coral bleaching.<sup>61,62</sup>

Many of these drivers act in synergy and are highly interconnected. For example, climate change predictions include increased intensity of cyclones<sup>63</sup>, increased nutrients is likely to be linked to crown-of-thorns starfish outbreaks<sup>64,65,66</sup>, and thermal stress tolerance in corals is reduced with degraded water quality<sup>67</sup>, both of which are linked to increased disease susceptibility<sup>68,69,70</sup>. Hence it is critical to improve the management of direct anthropogenic impacts to reduce the overall stress on coral reefs in the Region, noting that once coral reefs reach a tipping point towards non coral-dominated ecosystems, it is unlikely they will ever recover<sup>71</sup>.



Crown-of-thorns starfish (*Acanthaster planci*) on a coral reef

### 9.4.3 Current management and its effectiveness

Average coral cover has declined throughout the Region, this is despite high community interest and concern for corals, comprehensive partnership programs with industry stakeholders, and increased efforts by managers to enhance coral reef resilience (particularly through improving water quality).

In the Great Barrier Reef Marine Park, the Authority has managed localised impacts on coral reef habitats from dredging, dredge material disposal, coral collection and anchoring through the Zoning Plan, permit conditions, plans of management, best practices and site management. In addition, the Authority is working in partnership with the Australian and Queensland governments through Reef Plan to halt and reverse the decline in water quality. This partnership focuses specifically on agricultural pollutants which account for about 90 per cent of the pollutant loads. The Australian Government recently recommitted a further \$200 million to its Reef Rescue 2013–2018 initiative to enhance the Reef's resilience to climate change and catchment run-off. The Queensland Government has also announced it would continue its \$35 million per year investment in Reef Plan.<sup>39</sup>

Other pressures on corals such as those derived from coastal development (for example local effects of urban and industrial discharge), port operations (for example through sedimentation) and shipping (such as impacts of spills and antifoulants) are predominantly outside the direct regulatory control of the Authority.

With regard to coral, the Authority's overall management effectiveness and effectiveness for biodiversity outcomes was endorsed by the reviewers as partially effective. The independent review of the Authority's assessment concluded:

- Context** is effective. *The values and impacts of corals reefs is well understood as demonstrated by the large amount of scientific literature published about the corals in the Great Barrier Reef. Australia is recognised as a worldwide leader in coral research, much of which is focused on the Great Barrier Reef. Qualitative models are used to consider the potential for recovery or decline of the reefs under various scenarios, enabling guidance on potential management interventions.*
- Planning** is mostly effective. *A range of planning tools including the Zoning Plan and Reef Plan are reasonably effective in protecting corals. The Region has been mapped into 30 Reef bioregions, providing good scientific-based evidence to support planning for biodiversity through the zoning plans. Planning for improving water quality, one of the key drivers impacting on corals, is through the Reef Plan.*
- Inputs** are mostly effective. *There have been significant inputs into the protection of shallow water corals through Reef Plan, scientific research projects and community monitoring programs. However, the joint management program that is critical to the ongoing protection of the corals requires increased long-term funding that is adequate to undertake the required tasks.*

- **Process** is mostly effective. *Sound governance is evident through legislation, the intergovernmental agreement and effective stakeholder partnerships. Plans and strategies are being implemented appropriately. Monitoring of coral reef condition is undertaken in a systematic manner, with over 9000 reef health and impact surveys undertaken since 2009. Targets have been set for recovering coral cover.*
- **Outputs** are mostly effective. *Outputs to protect the values of corals including the representative protection of corals through the Zoning Plan, the reef protection markers, and improved water quality through the Reef Plan have been implemented.*
- **Outcomes** are partially effective. *Irrespective of the resources and outputs for protecting corals that have been implemented, coral cover is declining and this is predicted to continue. Many of the drivers that negatively impact on corals such as water quality are long-term issues and the improvements in water quality will take time to be reflected in improved coral cover. The reviewers considered **Biodiversity outcomes** to be partially effective. Although coral reefs south of Cooktown have experienced declines over the past [27] years, those north of Cooktown have remained relatively stable. One of the key drivers of coral loss in the southern region has been from the impact of cyclones (an example is the Swains Reef where modelled decline was from 38 to 8 per cent since 2006). However, there is evidence that the corals are recovering from the impact, with large numbers of coral recruits observed in recent surveys. Commitment to long-term improvement in water quality is necessary.*

#### 9.4.4 Outcomes and potential improvements

Mapping and qualitative models<sup>41</sup> have helped managers understand cumulative impacts on corals. They have also provided guidance on management interventions that may be most effective at avoiding or mitigating impacts to build coral resilience and aid recovery.

Global climate change, poor water quality, and increasing prevalence and severity of extreme weather are continuing sources of environmental stress that cumulatively impede the recovery and resilience of coral reefs on the Great Barrier Reef. Consequently, with current management, the condition for corals is projected to decline to 'very poor' in the southern two-thirds of the Region, and to 'poor' in the northern third in coming decades.

##### Enhancing protection and restoration

Reducing impacts related to climate change (for example, increased sea temperature and cyclone severity) by ensuring the concentration of carbon dioxide in the atmosphere does not approach the threshold value of 430 parts per million is central to the long-term health and resilience of coral reefs.

So too is a whole of government approach to reversing declines in water quality. It is critical that water quality objectives are met, and that land clearing, dredging and disposal of dredge material do not adversely affect coral cover on inshore, mid-shelf, outer shelf and deep sea reefs.

Improving the consistency of and strengthening the application of water quality guidelines across the Great Barrier Reef World Heritage Area, including through regulatory and partnership approaches, would likely strengthen the adoption of best practices in activities like dredging and dredge material disposal.

Proactive site planning for high growth areas for recreation and other uses would provide certainty and targeted on-ground actions.

Restoring coral reefs following incidents such as vessel groundings or cyclones would help increase the resilience of the ecosystem as a whole. Such restoration could be guided by a policy that reviews current best practice and international developments in technology, methods and capacities.

##### Minimising impacts

Recognising that crown-of-thorns starfish outbreaks are the major cause of coral cover decline, returning outbreaks to their natural cycles is a major component of improving the condition of coral reefs. This is likely to require a reduction of nutrients in Reef waters, direct control of crown-of-thorns starfish at the source of outbreaks and compliance with the zoning and spatial management arrangements.

Increasing investments in site infrastructure and compliance activities Reef-wide would assist with the prevention of direct damage of coral (for example, public moorings, reef protection markers, enforceable no-anchoring areas). As the adjacent coastal population continues to increase, catering for expanded use of the Region is critical.

##### Improving adaptive management

Integrated long-term monitoring of the condition and trend of coral reefs and understanding the adaptive capacities of coral reefs under a changing climate are required to ensure management actions are effectively reducing the impacts of highest risk to coral reefs. Another aspect to long-term monitoring is improving managing agencies' understanding of the social, cultural and economic benefits associated with coral reef habitats.

Maintaining managing agencies' close collaborations with recreational and commercial users of the Region (for example, tourism operators, commercial fisheries) to collect relevant data (for example through the Eye on the Reef monitoring program) will continue to allow early warnings of impacts such as crown-of-thorns starfish outbreaks or thermal bleaching events.



## 9.5 Islands


### 9.5.1 Significance

Islands contribute to all four of the natural criteria for which the World Heritage Area was listed in 1981: exceptional natural beauty, significant geomorphic features, significant ongoing ecological and biological processes, and significant natural habitats for the conservation of biological diversity.

There are about 1050 islands in the World Heritage Area — these are exceptionally diverse in terms of their geography, geomorphology and ecology. Islands are a unique component of the World Heritage Area and critical to its integrity. Interconnected reef and island ecosystems support some of the richest biodiversity on the planet. For example, continental islands and cays in the World Heritage Area support more than 200 bird species, many of which are in breeding colonies, while providing globally important nesting sites for marine turtles.

Islands function as important refugia for plants and animals, protecting them from impacts prevalent on the mainland, and some habitats are found only on islands of the World Heritage Area (such as *Pisonia* forests).<sup>72</sup> Islands are also key links in connecting terrestrial habitats along coastal and offshore areas. These connections are intricately dependent on the species which have evolved to live on islands. This is typified by the relationship between Wet Tropics rainforests, Great Barrier Reef islands, and the pied imperial pigeon (Figure 9.2). Other island bird species migrate or move between many countries across the South Pacific region and beyond.

In addition to their natural values, many islands have significant heritage values for Aboriginal and Torres Strait Islander peoples. They include a diversity of sites of archaeological and cultural significance such as fish traps, middens, rock quarries, story sites and rock art. Well-known examples are on Lizard, Hinchinbrook, Stanley, Cliff and Clack islands and in the Whitsundays where there are spectacular galleries of rock paintings. Many islands also have significant historic heritage values (historic lightstations, associated shipping and navigational history), while a number are a focus for research and defence training activities.



**Figure 9.2 Relationships between the rainforest, islands and the migratory pied imperial pigeons**

Reproduced from *Informing the outlook for the Great Barrier Reef coastal ecosystems*<sup>44</sup>

Islands in the World Heritage Area are important for tourism and recreation — 27 islands have resorts and/or residential communities, and many more have visitor facilities including campgrounds, walking trails and day use areas. An estimated 40 per cent of the 1.8 million tourists to the Great Barrier Reef in 2011 included an island destination in their visit.<sup>47</sup> Magnetic Island, near Townsville, includes a number of urban, village-type settlements wholly located within the World Heritage Area.

### 9.5.2 Responsibilities for islands

Management of the islands in the World Heritage Area is complex, and a range of tenures and management arrangements apply. Table 9.4 presents the tenures of a sample of 21 islands in the World Heritage Area. The State of Queensland has jurisdiction over approximately 980 islands. About 400 of these are protected areas

(national parks), and tenure on the remainder includes leasehold, freehold, unallocated state land, Commonwealth or Deed-of-Grant in Trust land, or a combination of tenures.

Approximately 70 Commonwealth islands are part of the Great Barrier Reef Marine Park and are within the Commonwealth Islands Zone. Twenty-one of those 70 islands include navigational lights or lightstations, and in 1988 responsibility for management of those islands was transferred from the Australian Maritime Safety Authority to the Great Barrier Reef Marine Park Authority (the Authority). The Australian Maritime Safety Authority remains responsible for the operation of the navigational lights, and leases back this portion of land. The Department of Defence is responsible for all the remaining Commonwealth islands in the World Heritage Area, except Little Fitzroy Island which is owned by the Department of Finance and Deregulation.

**Table 9.4 Tenure of islands in the World Heritage Area**

The Authority manages Kent Island, Pilon Island, Unnamed Island (Pilon Island Group), Russell Island, North Reef Island, Low Isles, Pine Islet, Lady Elliot Island, Albany Rock, Coppersmith Island, Hannah Island, Bailey Islet, Clerke Island, Coquet Island, Eshelby Island, Hannibal Island, High Peak Islet, Rocky Island and South Brook Island, together with parts of Dent and Penrith islands.

1050 islands within the Great Barrier Reef World Heritage Area			Examples of islands																				
			Brampton Island	Curtis Island	Dent Island	Flinders Island	Forbes Island	Great Keppel Island	Green Island	Hamilton Island	Heron Island	Hinchinbrook Island	Keswick Island	Lady Elliot Island	Lizard Island	Magnetic Island	North Keppel Island	Orpheus Island	Quoin Island	Raine Island	Townsend Island	Triangular Island	Whitsunday Island
70 Commonwealth islands	Department of Defence (48 islands)																				●	●	
	Department of Finance (1 island)																						
	Great Barrier Reef Marine Park Authority (21 islands)			●									●										
	National park (400 islands)	●	●					●		●	●			●	●	●	●	●	●				●
	Leasehold	●	●	●			●	●	●	●	●			●	●	●	●						
	Unallocated state land		●					●															
	Esplanade							●															
	Freehold	●	●											●	●								
	Aboriginal land					●																	

### 9.5.3 Key issues

In general, the condition of islands is assessed as good and stable, with the exception that condition is deteriorating on some inner islands that are the subject of development activities in the southern World Heritage Area.<sup>73</sup> Many of the natural resource management issues on islands result from the legacy of past activities. Consequently, effective management needs to address past impacts as well as current and emerging threats to the islands and their surrounding marine ecosystems.

Key past and present impacts on the islands throughout the World Heritage Area include: historical guano mining activities; invasive plants and animal pests (for example goats, rats); altered and unmanaged fire regimes; impacts from visitation; increasing impacts of industrial and residential infrastructure; the degradation of cultural heritage (Indigenous and historic); and the consequences of climate change, leading to declines in cultural, ecological and aesthetic values. The diverse range of issues is illustrated by two examples, Raine Island and Lady Elliot Island.

Raine Island is a Queensland Government island and afforded the highest protection as a national park (scientific). It is one of the most remote islands in the World Heritage Area, located in its far north. Guano mining from 1890 to 1892 caused major changes to the island's geomorphology and hydrology, some of which are only now becoming apparent.<sup>74</sup> While Raine Island remains probably the most important seabird breeding site on the

Great Barrier Reef, the island's seabird populations have been affected by a combination of historic activities and climate change. Eighty-four bird species have been recorded at Raine Island — five of these are considered uncommon or rare in Queensland. Sixteen species are known to breed on the island. A comparison of bird populations between 1979 to 1993 and 1994 to 2003 suggests a population decline in 13 of the 16 species over the 24-year period. The combined averages for all 16 species indicate the total rookery population has declined by about 16,000 birds or about 70 per cent.<sup>75</sup>

The world's largest population of nesting green turtles (*Chelonia mydas*) is found on Raine Island.<sup>76</sup> Nightly nesting activity can involve more than 10,000 turtles. Today, a range of potential impacts are affecting turtle reproductive success including, but not limited to, tidal inundation of nests, changing sand distribution and consistency, and soil-borne pathogens. These can lead to a decline in nesting and hatching success.<sup>77,78,79</sup> Active management intervention is now taking place — a situation likely to become more commonplace as impacts related to climate change become more apparent, the condition of values decline and the capacity for natural recovery diminishes.

The southernmost island in the World Heritage Area, Lady Elliot Island is a Commonwealth island and an important site for seabird nesting. For example, *Pisonia* trees on the island are important as breeding habitat for black noddies, while ground-nesting crested terns, red-tailed tropicbirds and bridled terns are common.

From 1863 to 1873, Lady Elliot Island was cleared of vegetation and mined for guano. In 1969, a revegetation program began and a mixed cover of *Casuarina* and *Pisonia* trees was established, providing habitat for many bird species. The island now provides significant opportunities for visitors to view a large diversity of coastal birds. It is accessed mainly by airplane, with accommodation provided by an eco-certified tourism operation.

The heritage-listed lightstation at Lady Elliott Island contains a lighthouse and a cluster of associated service buildings. Many of the buildings are constructed of asbestos which may pose an environmental and occupational health and safety risk if buildings fall into disrepair. Lady Elliot Island lightstation is on the Commonwealth Heritage List and a management plan has been developed for the heritage conservation of the site.



*Lady Elliot Island, a Commonwealth island in the southern Great Barrier Reef*

#### 9.5.4 Current management and its effectiveness

The degree of active management for islands in the World Heritage Area varies widely. They are covered by a variety of tenures (leasehold, freehold, unallocated state land, Commonwealth, and Aboriginal land) and are managed by a range of agencies. Many that are not part of the protected area estate have important values such as seabird rookeries.

##### **Queensland islands (excluding protected area islands)**

Queensland islands that are not part of the protected area estate are multiple-use areas that support a range of activities and industries, such as tourism and residential. They are subject to an evolving range of Queensland Government regulatory tools, including legislation and ongoing management programs. The Queensland Government is committed to ensuring that planning and development of non-protected area islands is appropriately managed to ensure the World Heritage Area retains the values for which it was declared, and that it continues to be one of the best managed areas in the world.

The Queensland Government's coastal management, planning and development framework applies to its islands. This includes legislation, plans, policies and programs that combine to provide a comprehensive regulatory and management system. Their purpose is to provide Queensland Government oversight and regulation of development on these islands, particularly in regard to activities associated with major developments that are expected to occur on islands, for example tourism developments.

Leasehold islands are subject to specific terms of use which dictate the kind of activities that may occur. Activities or uses that have the potential for negative effects are mitigated through lease conditions. Freehold island owners are subject to their environmental duty of care under Queensland and Australian government legislation, such as protecting native species (for example, the Nature Conservation Act (Qld)) or not undertaking actions that may have an impact on matters of national environmental significance.

### Queensland national park islands and Commonwealth islands

Key challenges in managing national parks and Commonwealth islands include resource limitations, the remote location of many islands (where visits by field staff may total only a few hours per year) and the ecological and jurisdictional complexities. These challenges were recognised and resulted in the establishment of joint management arrangements and a formal partnership between the Australian and Queensland governments through the former 1979 Emerald Agreement and the current *Great Barrier Reef Intergovernmental Agreement 2009*. For more than 30 years, this partnership has delivered on-ground management activities through a field management team comprising Queensland and Australian government agencies. Participants in field management include the Queensland Parks and Wildlife Service, Border Protection Command, Customs National Marine Unit, Commonwealth Director of Public Prosecutions, Australian Federal Police, Australian Maritime Safety Authority, Queensland Boating and Fisheries Patrol, and Queensland Police Service.

A large range of planning tools is available to manage islands, including complementary state and Commonwealth zoning (with specific zoning for all Commonwealth islands and Special Management Areas), joint field management business plans, Great Barrier Reef Marine Park plans of management, and Queensland Government national park management plans and management statements.

However, there are many instances when state and Commonwealth legislation is not complementary for waters adjoining islands (such as for seabird closures). In addition, there are no state plans in place to mirror the Authority's plans of management for intertidal waters adjoining islands. While about 400 islands are protected as national parks, there are some islands with high nature conservation value not under protected area tenure (for example, Douglas Islets).

Monitoring of coastal birds, turtles, pests and weeds is undertaken to assess the condition and trend of key species and habitats. Controlling or eradicating pest plants and animals is risk-based and effective at locations where resources are focused (for example, successful eradication of feral pigs from Wild Duck Island, removal of rats from Boydong Island, and controlling outbreaks of scale insects responsible for the destruction of *Pisonia* forests on Tryon Island). In an island context, this is significant because it shows eradication of introduced species is an achievable outcome — with concerted effort, pest programs can be finite and deliver enduring conservation benefits. However, to remain effective, these programs must be coupled with a strong focus on quarantine and surveillance, as preventing adverse environmental impacts is more cost-effective than managing or reversing them.

The Authority and the Department of Defence have responsibility for the protection of natural, historic and cultural heritage values on the 70 Commonwealth-owned islands. Regimes are in place to provide an overarching level of management for these islands (for example, zoning, regulations, surveillance and enforcement). However, a lack of information in a readily accessible format makes it difficult to monitor and assess the condition of those values and in some cases has contributed to delays in the preparation of heritage management plans.

Some cultural heritage sites on islands are known to be degrading, for example Pine Islet lightstation which is closed to public access due to asbestos issues. Long-term maintenance of heritage sites on islands is expensive; particularly for European heritage (for example the Queensland heritage listed Raine Island tower and the Commonwealth heritage listed Lady Elliot Island lightstation). The agency responsible for Commonwealth heritage sites listed under the *Environment Protection and Biodiversity Conservation Act 1999* must make plans to protect and manage heritage values and must not contravene those plans. Plans have been or are under development for Commonwealth heritage listed sites within the World Heritage Area.

Significant progress in island protection and conservation has been made in recent years in partnerships with Traditional Owners, especially through the collaborative development of Traditional Use of Marine Resources Agreements and Indigenous Land Use Agreements. Specific Indigenous compliance partnerships have also been progressed on Cape York Peninsula. Australian Government Reef Rescue funding has been a key driver of this progress.

The independent review of the Authority's assessment concluded:

- **Context** is mostly effective. *Values underpinning matters of national environmental significance for islands are well documented and understood by managers, with management plans or management statements that articulate the values and threats for the islands developed. The geological basis of islands: mangrove islands, continental islands, and reef islands or coral cays influence the management issues associated with each island. Mapping of regional ecosystems and identification of endangered ecosystems has occurred for most islands. The islands vary significantly in their visitation and development profiles.*
- **Planning** is mostly effective. *A range of planning processes is in place to manage the islands, and vary depending on the jurisdictional responsibilities. Plans include the Zoning Plan (including restricted access areas and public appreciation areas), local government planning, Queensland Parks and Wildlife Service management plans, Cairns, Whitsunday and Hinchinbrook Plans of Management and Traditional Use of*



Marine Resources Agreements. Overall, the planning arrangements on islands to protect the values are in place, and are implemented appropriately.

- **Inputs** are partially effective. Significant long-term financial commitments from both the Australian and Queensland governments are required if priority actions to achieve agreed objectives are to be implemented. This is particularly important for high value locations that are remote and difficult to access.
- **Process** is mostly effective. Good governance through the intergovernmental agreement and joint management program, as well as strong stakeholder engagement is in place. Monitoring programs such as the coastal bird monitoring strategy, turtle monitoring, pest and weed programs, and specific vegetation programs such as the impact of the invasive ants in the *Pisonia* forests on Tryon Island are undertaken to assist management decisions and gauge the impact of management actions.
- **Outputs** are mostly effective. Activities relating to island management have generally progressed well, and in accordance with the respective work programs, such as the Field Management Program. Examples include the successful eradication of feral pigs from Duck Island, and removal of rats from Boydong Island. Limited resourcing is seeing a reduction in time spend on remote and isolated islands, and it is expected that this will also impact on the capacity to deliver the work programs.
- **Outcomes** are mostly effective. Management activities are reducing the short-term immediate risks and threats. However, they are not comprehensively addressing issues such as changes in beach profiles due to climate change and severe weather. **Biodiversity outcomes** are mostly effective. Specific actions to address the impacts of changes to beach profiles that affect high value biodiversity outcomes such as turtle breeding on Raine Island have been implemented. However, this has not been comprehensively considered across all islands.

### 9.5.5 Outcomes and potential improvements

The overall condition for islands is projected to remain good in the coming decades, but pressures will increase from climate change and on inner islands in the southern World Heritage Area that are subjected to ongoing pressures from development and use.

The following potential improvements relate to protected area islands in the World Heritage Area (for example Queensland national parks or other conservation tenures, and islands that are part of the Great Barrier Reef Marine Park), representing up about 30 per cent of the total of 1050 islands. Many of the identified management improvements would also benefit the remaining 700 islands not part of the protected area estate and managed under other tenures.

#### Enhancing protection and restoration

Island protected areas will continue to provide visitor opportunities that support the tourism industry and a broad and expanding range of recreational use. As the coastal population increases, so too will the demand for visitor opportunities and the associated facilities required to support use and minimise its effect on island and reef ecosystems. Improving proactive regional and site-specific planning (including for site infrastructure) would provide certainty for recreational and tourism opportunities and sustain visitor enjoyment, environmental quality and the economic benefits of healthy, well-presented islands.

Indigenous and historic heritage values associated with islands are an increasing focus for protection and restoration activities. Improving managing agency capacity to undertake cultural heritage management activities would assist in the delivery of legislated obligations.

A focussed program for Traditional Owner partnerships specifically related to island protected area management would complement progress already made through partnerships with Traditional Owners.

Encouraging an engaged and supportive community is vital to the long-term protection of the Great Barrier Reef World Heritage Area, including its islands. The Queensland Government has initiated a program to increase the support that community volunteers can provide to management of the State's protected areas.

#### Minimising impacts

On-ground management actions are essential in ensuring effective management of protected area islands, whether this is maintaining visitor infrastructure, managing pest plants and animals and fire, or active adaptive management works to remove or reduce threats to vulnerable species such as marine turtles and seabirds.

Improved field management resources, personnel and fit-for-purpose vessels to deploy to all protected area islands at the frequency and duration required for effective on-ground management would improve the protection of natural, social and economic values of island protected areas.

With increasing visitation to islands, the risk of exotic plant and animal pest incursions is likely to increase. The present management framework is effective at controlling or eradicating pest plants and animals at locations where resources are focussed on resolving specific issues. However, there is a need to increase the focus on quarantine and preventative measures and surveillance activities. Prevention or early detection and treatment for island pests is better and less expensive than restoration after an incursion. While there remain many historic pest legacy issues requiring management action, the factors that drive island pest introductions are escalating.

#### Improving adaptive management

Maintaining the natural, economic and social values of Great Barrier Reef islands relies on a good knowledge of values, identifying changes and trends and managing threats to these values. The key known threats to the

natural, economic and social values of World Heritage Area's islands are: the ecosystem consequences of a changing climate; invasive plant and animal pests; altered and unmanaged fire regimes; impacts to island natural and recreational values if human use is not well-managed; and the degradation of cultural heritage (Indigenous and non-Indigenous).

While managers generally understand most of the potential threats to protected area island ecosystem integrity, the capacity to consistently monitor and report on island condition and trend and to plan response treatments strategically across the World Heritage Area is constrained. Improving the capability to monitor and report on the condition and trend of islands over the long term would support a complete adaptive management approach for islands and deliver more resilient island ecosystems. It would also complement the existing marine ecosystem condition and trend monitoring arrangements, providing a more holistic ecosystem assessment of the World Heritage Area.

A fundamental principle for managing island protected areas is maintaining healthy ecosystems. This approach is consistent with that recommended in *Australia's Biodiversity and Climate Change Vulnerability Assessment* (2009) which states under the strategy of Building Resilience: '....the single most important adaptation strategy is the maintenance of well-functioning ecosystems.' This focus on resilience-based management of islands should continue as an important component. Climate change amplifies the disturbance regime in natural systems, with no exception regarding impacts on island ecosystems. Predicting the impacts on island over the next 25 years is difficult; and the capacity to adapt management arrangements to respond to emerging issues or outbreaks is important.

Continued complementarity of management approaches between the Commonwealth and Queensland protected areas (such as uniformity in any management arrangements and regulations across the Commonwealth and state marine parks and adjoining protected area islands) provides certainty for visitors as well as continuing the long-standing joint field management arrangements in the World Heritage Area.

## 9.6 Princess Charlotte Bay

### 9.6.1 Significance

Princess Charlotte Bay is located at the base of Cape York Peninsula (Figure 9.3). The Lama Lama people are the Traditional Owners for the land and sea country extending from the Massey River in the north, around Princess Charlotte Bay, to the Normanby River in the south. The Authority acknowledges the continuing sea country management and custodianship of the Great Barrier Reef by Traditional Owners.





Figure 9.3 Princess Charlotte Bay

The area is recognised for its high biodiversity values. It contributes to all four of the natural criteria for which the World Heritage Area was listed in 1981: exceptional natural beauty, significant geomorphic features, significant ongoing ecological and biological processes, and significant natural habitats for the conservation of biological diversity. The relatively unspoilt nature of the remote area contributes to its high aesthetic value.<sup>80</sup> The area is of high conservation value for largemouth (previously freshwater) and green sawfish,<sup>81</sup> estuarine crocodiles,<sup>82</sup> dugongs,<sup>36</sup> green turtles<sup>77</sup> and Australian snubfin and Indo-Pacific humpback dolphins<sup>83</sup>. In addition, the last known verified specimen of the critically endangered speartooth shark was recorded in 1982 from the Bizant River which flows into the bay.<sup>81</sup> All these species are matters of national environmental significance.

Princess Charlotte Bay is one of the largest tidal wetland systems in Cape York Peninsula. Extensive and interconnected mangrove forests<sup>84,85</sup> and seagrass meadows<sup>84,86,87</sup> provide important fish habitat. Recreational, commercial and Indigenous fishers target barramundi, king and blue threadfin salmon, grunter, mangrove jack and mud crabs. The bay is also the northern extent of most commercial gill-net and crab fishing activities in the World Heritage Area.

The surrounding national parks, Cape York Peninsula Aboriginal Land, state reserves, nationally important wetlands and other lands are characterised by diverse, expansive and relatively undisturbed natural landscapes, including coastal areas, river systems and floodplains, lowlands and wetlands,<sup>88</sup> — notwithstanding some declines.<sup>89</sup> It is the connectivity between terrestrial, aquatic and marine ecosystems which differentiates the Cape York catchment from other catchments adjacent to the World Heritage Area. Connectivity is required to complete the life cycle of many species, typified by the largemouth (freshwater) sawfish (Figure 9.4), which is now rarely found further south.



The agricultural land uses adjacent to Princess Charlotte Bay are principally grazing of natural areas and forestry.<sup>44</sup> They are considered to be well managed and, if these land use practices continue at the same intensity and spatial extent, they are unlikely to adversely affect biodiversity values in the area.

The area has very high Indigenous values and cultural significance.<sup>90</sup> The Flinders Island group contains Aboriginal rock art sites, shell middens and stone arrangements. The rock art on Stanley, Flinders and Clack islands is of international significance.



### 9.6.2 Key issues

For the values which underpin matters of national environmental significance, Princess Charlotte Bay's remote location has provided protection from many of the impacts acting elsewhere in the World Heritage Area. However, the area is identified as the Field Management Program's highest compliance risk because of the very high potential for interaction between fishing (legal and illegal) and species of conservation concern.

The area's remoteness also poses logistical challenges which restrict research and monitoring activities. Long-term data is only available for dugongs,<sup>19</sup> and to a lesser extent seagrasses<sup>86</sup> and some coral reefs.<sup>91</sup> This makes assessing and responding to changes in the distribution, abundance and status of species and habitats very difficult.

Emerging issues of significant concern are:

- potential development in the catchment and associated facilities
- the intensification of agriculture in northern Australia, which is likely in the next 25 years<sup>92</sup>
- visitation to culturally-sensitive sites.

### 9.6.3 Current management and its effectiveness

The majority of the adjacent area is gazetted National Park or Cape York Peninsula Aboriginal Land (Lama Lama, Rinyirru [Lakefield] and Cape Melville National Parks) and jointly managed under the Running Creek, Lilyvale, Rindoparr, Rinyirru and Kalpower Indigenous Land Use Agreements. Traditional Owners are on-country and are best placed to deploy management strategies to provide strong outcomes. The following section refers to the evaluation of current government management and its effectiveness. However, it is acknowledged that Traditional Owners have managed Princess Charlotte Bay on-country well before western systems were implemented and that they continue to do so through cultural practices and traditional ecological knowledge.

Princess Charlotte Bay is within the Remote Natural Area of the Great Barrier Reef Marine Park, which provides recognition and management of the area for its natural and undeveloped character. The purpose of the Remote Natural Area is to ensure the area remains in a state largely unaltered by works and facilities, and to provide opportunities for quiet appreciation and enjoyment. The Remote Natural Area does not include a small area to the north of Bathurst Head in the vicinity of the Flinders Island group.

Management of fishing is a priority in Princess Charlotte Bay. The area from north of Port Stewart around to Bathurst Heads is a Fish Habitat Area under the *Fisheries Act 1994* (Qld) and *Fisheries Regulation 2008* (Qld), which aims to protect critical fish habitat against physical disturbance from coastal development.

The zoning arrangements for Princess Charlotte Bay include a Special Management Area to reduce the threat to dugongs and other species of conservation concern from drowning in commercial set mesh nets (with flow-on benefits to other marine animals) (Figure 9.3). This unique arrangement requires commercial fishers to obtain written permission to use or enter the area for netting (other than bait netting). Since 2009, the Queensland Government has also closed to commercial netting three waterways which flow into Princess Charlotte Bay (North Kennedy, Bizant and Normanby rivers).

The environmental benefits of these fishery closures depend on compliance. The Field Management Program undertakes land-based patrols, complemented by large vessel patrols from Cairns. Helicopter-based patrols since 2011 are an effective surveillance option, but are expensive and limited. Patrolling is weather dependent and resources for these activities are declining. It is suspected there is significant illegal fishing in the area.

Illegal activities can be reported through the Eyes and Ears incident reporting program, in which Indigenous rangers participate. This is but one example of the increasing ways Traditional Owners are working with government to collaboratively manage natural resources within their country. Another example is the Lama Lama Traditional Owners working with the Authority to develop a Traditional Use of Marine Resources Agreement to enhance sustainable management systems, strategies and practices on country (see Chapter 3).

The independent review generally endorsed the assessment by the Authority and concluded:

- **Context** is effective. *Values underpinning matters of national environmental significance are well documented and understood by managers, due in part to its high ecological, traditional and socio-economic values. Condition and trend of a number of species are monitored. The marine habitats and species have been reasonably well researched and documented.*
- **Planning** is effective. *The demonstration case study provides a number of examples of planning systems and zoning that assist with addressing the protection of the values of Princess Charlotte Bay, including Marine National Park, Habitat Protection, Special Management Areas and Conservation Park zones. Complementary systems of protected area management across the marine and terrestrial interface assist with the protection of the values.*
- **Inputs** are partially effective. *Resources for dedicated training of Indigenous rangers have been significant, however resources to enforce the planning schemes is only partially effective, due to a variety of factors including the isolation and remoteness of the area. In particular, compliance officers face practical and logistical difficulties.*



- **Process** is mostly effective. *Sound governance is evident through the intergovernmental agreement, and complementary terrestrial and marine park zoning. Effective partnerships are in place with Traditional Owners.*
- **Outputs** are mostly effective. *Outputs such as the reduction in net fishers to four would suggest a better result than partially effective. Similarly the largely stable dugong population in the area indicates outputs have been mostly effective. Results from management programs such as the Zoning Plan, the Special Management Areas, the declared fish habitat areas, [and] the Reef Rescue program have been mostly effective in managing Princess Charlotte Bay.*
- **Outcomes** overall are mostly effective. *Management activities have been effective to date in protecting the values of Princess Charlotte Bay, as evidenced [by] ongoing high, intact biodiversity values. The greatest threat to Princess Charlotte Bay is potentially from the proposed Wongai Coal mine project which is yet to be approved. Biodiversity outcomes are effective. The intactness and integrity of Princess Charlotte Bay is evidence of effective biodiversity outcomes, although the remoteness and relative inaccessibility plays a significant role in this outcome.*

#### 9.6.4 Outcomes and potential improvements

The projected increase in economic development of the adjacent catchment area is likely to increase pressure on the biodiversity and heritage values of Princess Charlotte Bay, which are predicted to decline in condition in the coming decades. Contributing factors include the current outbreak of crown-of-thorns starfish in the area; extreme weather events that are likely to affect the condition of coral reefs and other habitats; and likely clearing and modifying of additional land in the catchment for agriculture. The improvements in catchment management of land currently used for agriculture are expected to improve inshore marine water quality through reducing sediment loads in flood events and reducing base loads of contaminants during lower flow periods. However, improvements in marine water quality and associated coral reef condition are likely to be slow, with lags in ecosystem responses and potentially long recovery periods.

Outcomes for biodiversity, heritage values and socio-economic benefits would benefit from the following management improvements.

##### Enhancing protection and restoration

Incorporating cumulative impact considerations into management decision making within the area, especially in relation to integrity and connectivity in line with the Remote Natural Area, would create a more holistic and consistent approach.

Ongoing support and resourcing for the implementation of the Lama Lama Traditional Use of Marine Resources Agreement (and other sea country partnership initiatives in the area) and appropriate training and resourcing for Indigenous rangers to conduct assessments of natural resources within their country would provide benefits to the biodiversity of the area as well as to Indigenous heritage sites and community benefits to Traditional Owners.

##### Minimising impacts

Direct impacts on the values of Princess Charlotte Bay could be addressed by reducing risks to species of conservation concern from both legal and illegal netting activity; ongoing compliance patrols in the Special Management Area, including additional resourcing for night-time surveillance technologies to address illegal fishing activities; and implementing safeguards to manage the impacts of increasing recreational fishing effort on species and habitats including from the harvest of fish, vessel strikes, anchor damage and disturbance to wildlife.

##### Improving adaptive management

The protection of the area's values relies on improved understanding, mapping and systematic evaluation of its biodiversity, and Indigenous and historic heritage values. In addition, an improved understanding of the patterns of commercial and recreational use would allow for proactive planning to meet the needs of those uses.

Integrated monitoring is crucial to inform management about conditions and trends of the area's values and to continuously assess the effectiveness of management actions.

Regular reporting on the status of values and management effectiveness will allow management to be adapted over time and to changing conditions.

## 9.7 Cairns Planning Area

### 9.7.1 Significance

Cairns is the principal city of far northern Queensland. The city is located on the east coast of Cape York Peninsula on the coastal strip between the Coral Sea and the Great Dividing Range. The area offshore Cairns contributes to all four of the natural criteria for which the World Heritage Area was listed in 1981: exceptional natural beauty, significant geomorphic features, significant ongoing ecological and biological processes, and significant natural habitats for the conservation of biological diversity. It is well known for its spectacular reefs, abundant wildlife (including whales, dolphins, turtles, dugong and seabirds) and the neighbouring Wet Tropics World Heritage Area, all of which contribute to its high aesthetic value.<sup>80</sup>

This unique environment 'where the rainforest meets the Reef' is an important part of the lifestyle and identity of many of the residents, including the Traditional Owners who have ongoing connections with land and sea country.

Recreational activities such as boating, fishing, scuba diving, snorkelling and yachting are widespread. The area also supports commercial fishing, a growing cruise shipping industry and commercial ports servicing far northern communities.<sup>93</sup>

Tourism is the major economic driver for the Cairns region. The outstanding universal value of the coastal and marine environment, along with the proximity of the reefs and islands to the coast, draws about 50 per cent of the Great Barrier Reef Marine Park's almost two million tourist visits to the area each year. A wide array of nature-based visitor experiences is provided by tourism operators, ranging from cruise ships and live-aboard vessels to day trips on high speed catamarans, kayaking tours and game fishing.

Recognising the intensive tourism activity around Cairns, the Authority prepared the Cairns Area Plan of Management<sup>94</sup> in 1998. The plan's objective is 'to protect and conserve this environmentally significant area, while allowing reasonable opportunities for access and use'.

The area covered by the plan is referred to as the Cairns Planning Area. It extends from Lizard Island in the north to the Frankland Islands in the south (Figure 9.5). The planning area includes many places, biological communities and species of high biodiversity, heritage and socio-economic value. The plan identifies the values that require protection, the main impacts affecting them and the management strategies required to address these issues.

This demonstration case study highlights the complexity of managing cumulative impacts from tourism in an intensively used area in a way that protects biodiversity and provides certainty for a variety of tourism activities.

### 9.7.2 Key issues

Tourism plays a key role in presenting the outstanding universal value of the Great Barrier Reef to visitors. Tourism operators that help protect and present the World Heritage Area to a consistently high standard are recognised by the Authority through the High Standard Tourism program. These operators are independently certified under Ecotourism Australia's ECO Certification Program to the Ecotourism or Advanced Ecotourism levels.

The plan of management recognises the sustainability of this Reef-dependent industry relies on maintaining the environmental and socio-economic values of the area's reefs, cays and waters, as well as the biodiversity these habitats support. A healthy ecosystem also underpins commercial fishing and recreational activities. A focus of the plan is to minimise cumulative impacts predominantly associated with intensive tourism. In particular, the plan seeks to address impacts on heritage, wildlife and the environment associated with intensive tourism use and the potential conflicts between tourism operations and other users.

The suite of management arrangements employed within the planning area is complex, and understanding and applying these management tools is challenging for users. As the plan only applies to a small portion of the Great Barrier Reef Region, management tools and intent do not, on occasion, align well with those in adjoining areas.

### 9.7.3 Current management and its effectiveness

The Cairns Area Plan of Management operates in conjunction with a broad range of management tools, but focuses on issues specific to the area, its species and communities at a regional scale. While the plan addresses the area's intensive tourism activity, it complements the suite of management tools used to manage tourism throughout the Marine Park.

Implementation of the plan began in 1998. Since then it has been reviewed every two to three years, although the last review was in 2008. The plan addresses potential impacts and conflicts in the planning area associated with intensive use through:

- limiting group sizes at locations
- establishing special management arrangements at 11 sensitive locations
- limiting the number of private moorings, pontoons and other structures
- limiting noisy activities
- establishing reef anchorages, no-anchoring areas and public moorings
- capping the number of tourism operations (vessels and aircraft).

Joint management arrangements with the Queensland Government (such as permitting, site infrastructure management and compliance) in place since 1979 and current under the *Great Barrier Reef Intergovernmental Agreement 2009*<sup>95</sup> significantly increased jurisdictional consistency across land and sea. The Queensland Government and the Authority have jointly approved some site management plans, which are non-statutory policy documents; however the Cairns Area Plan of Management has not been approved by the Queensland Government. Reciprocal recognition of planning tools by the Authority and the Queensland Government would improve consistency of management.




Figure 9.5 Cairns Planning Area

The plan and the permitting system employed to manage tourism operations partially address conflicts between tourism and other activities. However, conflicts between recreational and commercial fishers, and between Traditional Owners and other users, are now emerging in the planning area.<sup>93</sup>

The plan has been mostly effective in managing cumulative use by the tourism sector.<sup>6</sup> However, the cumulative impacts from all uses (including recreational use, fishing and shipping) in the planning area are neither the intended focus nor the intended outcome of the plan. The protection of the planning area's values from these impacts is reliant upon other management tools.

Poor water quality (as a result of land management practices in the catchment) correlates with an increased frequency of crown-of-thorns starfish outbreaks.<sup>66</sup> In the past, these outbreaks have severely affected coral cover, a key value of the Cairns Planning Area upon which the tourism industry is heavily reliant. While significant improvements have been made in halting and reversing the decline in water quality, an accelerated program is necessary for the future.

The independent review of the Authority's assessment concluded:

- **Context** was effective — *Values underpinning matters of national environmental significance are well documented and understood by managers, as are the regional and local risks and threats which are the subject of ongoing engagement with regional stakeholders.*
- **Planning** is mostly effective. *A robust planning system is in place but its effectiveness is constrained by the complexity of governance issues with all levels of government needing to be critically involved as well as an array of narrowly-focused industry and interest groups. Agreeing on clear and measurable objectives which give rise to priority actions implemented consistently across jurisdictions in these circumstances remains a challenge. Note that the plan of management addresses overall level of use (within context of an already heavily-used region) but there is little evidence in the plan that it considers or manages cumulative impacts — little or no discussion of this in the plan, no identification of which potential cumulative impacts are of concern and no consideration of associated monitoring to manage this issue — but the demonstration case text suggests that the [Plan] does address cumulative impacts.*
- **Inputs** are mostly effective. *Achieving cross-jurisdictional consistency is always going to be resource intensive. Collaborative governance requires significant long-term financial commitments from all parties if priority actions to achieve agreed objectives are to be implemented.*
- **Process** is partially effective. *The individual components of governance are sound with the individual responsibilities of the different levels of government clear, but the complexity which derives from the number of levels of government involved makes consistent delivery of on-ground results difficult.*
- The delivery of **outputs** assessed by the Authority as effective was not endorsed by the reviewers. *'Mostly effective' would seem to be a more accurate assessment given that while individual programs or activities may have progressed in accordance with timeframes and plans, there is only limited evidence presented that the results have achieved their stated objectives. The fact that [the Authority] no longer provides tourism industry training in a key tourism region, that the field management program is significantly reduced, that planning and policy tools for tourism have not been updated all suggest that a rating of effective is too high.*
- **Outcomes** overall are mostly effective. *Management activities are certainly reducing the major risks and threats but fall short of categorically ensuring either the economic or environmental sustainability of the use of the Reef. Finally, biodiversity outcomes are only partially effective. While risks and threats are reduced, management is only part of the way to being on track to ensure that the values of the Great Barrier Reef are protected.*

Much of the Great Barrier Reef catchment is forecast to experience annual population growth rates of 1.6 per cent or higher in the near future,<sup>93</sup> and this is expected to result in increased recreational use in the planning area. In recent years, regional catchment planning has attempted to identify and conserve regional values, while allowing for growth.

Given the likely impacts from climate change, coastal development and water quality, it seems unlikely that the current plan alone will be effective in protecting the area's values in the future and in providing certainty for Reef-dependant uses such as tourism, commercial fishing, traditional use and some recreational activities.

#### 9.7.4 Outcomes and potential improvements

The Cairns Area Plan of Management has been mostly effective in managing the cumulative impacts of the most intensive tourism area in the Marine Park. However, the condition of biodiversity, heritage values and community benefits are projected to decline in the planning area in the absence of improved management arrangements to address regional impacts (particularly from water quality and coastal development, which are outside the scope of the plan); protection of at-risk marine and terrestrial habitats; and maintenance or restoration of landscape connectivity to ensure functional terrestrial and marine ecosystems.

To improve management of Reef-related activities and resilience in the planning area, potential measures could include the following actions.

##### Enhancing protection and restoration

The resilience of the ecosystem would be improved and values better protected by:



- Improving protection for at-risk marine and terrestrial habitats and maintaining or restoring landscape connectivity to ensure functional terrestrial and marine ecosystems
- Strengthening proactive planning to enhance environmental protection, ecosystem resilience and tourism and recreation values, and to respond to projected changes in patterns of use as a result of population growth in the catchment.
- Improving understanding and presentation of world heritage values, including outstanding universal value, by the tourism industry and visitors to the planning area.

### **Minimising impacts**

Impacts on values would be reduced by:

- Continuing to reduce crown-of-thorns starfish outbreaks, through targeted control activities at source reefs and high value tourism sites and improving water quality.
- Increasing awareness and application of best practices by recreational users, tourism operators, commercial fishers and Traditional Owners.
- Expanding engagement and stewardship programs including a mechanism for auditing and reviewing identified minimum operating standards and requirements.
- Strengthening field management, compliance and enforcement.
- Reducing impacts on biodiversity through the installation of site infrastructure (for example, no-anchoring areas, public moorings, reef protection markers, signs).

### **Improving adaptive management**

The capacity of the Authority to adapt its management to changing circumstances in the planning area would be improved by:

- Regularly reviewing and aligning management tools across the Australian and Queensland governments to ensure currency, consistency, effectiveness and simplicity.

## **9.8 Mackay Whitsunday — water quality improvement**


### **9.8.1 Significance**

The decline in the quality of water entering the Great Barrier Reef from adjacent catchments is one of the most significant impacts on the health of the Reef.<sup>5</sup> This demonstration case study explores how the Queensland and Australian governments are working to halt and reverse the decline in the quality of water entering the Reef. Specifically it highlights some of the work being done to improve the quality of water flowing from the Mackay Whitsunday region (Figure 9.6).

It documents actions delivered through the joint Australian and Queensland government's Reef Water Quality Protection Plan (Reef Plan),<sup>40</sup> including the Commonwealth's Reef Rescue initiative, Queensland's development of best management practices, and management of land use to minimise run-off. It also includes the work of the regional natural resource management bodies in developing a regional water quality improvement plan with water quality targets linked to the Authority's water quality guidelines and Queensland's Environmental Protection Act and policies.


### **9.8.2 Key issues**

Many of the pressures on water quality reflect the land use pattern in the catchment. The dominant land uses are grazing (56 per cent), sugarcane production (19 per cent) and national parks and reserves (17 per cent) (Figure 9.7).<sup>96</sup> In the last decade, understanding of the impact of water quality on the Reef has changed. It has moved from a debate on whether an impact is actually occurring to a consensus that declining water quality has degraded many inshore reef areas. There is evidence that parts of the Great Barrier Reef are now experiencing nutrient enrichment,<sup>53,97,98</sup> high phytoplankton biomass,<sup>53,99</sup> potential changes in the phytoplankton food web structure, increased competition at the coral reef level, and changes in food webs as demonstrated by the proliferation of crown-of-thorns starfish in areas regularly exposed to nutrient loads stemming from human activities.<sup>58,66</sup>



**Figure 9.6 Coastal ecosystems in the Mackay Whitsunday natural resource management region, 2009**

The majority of vegetation in the catchment is classed as 'non-remnant', that is it has been modified to the extent that it is no longer recognised as a functioning habitat. Much has been changed from forest to grassland for grazing purposes (based on *Informing the outlook for Great Barrier Reef coastal ecosystems*<sup>44</sup>).



**Figure 9.7 Conceptual diagram of land use in the Mackay Whitsunday region**

From Reef Plan First Report Card<sup>96</sup>

### 9.8.3 Current management and its effectiveness

There is a range of policies, plans and programs which identify the environmental values that rely on good water quality. Each of these tools — brought together under Reef Plan — performs a different function as shown in Table 9.5.

**Table 9.5 Mechanisms for identifying and protecting water quality values**

Water quality improvement plan	Environment protection (water) policy	Great Barrier Reef water quality guidelines	Queensland water quality guidelines
<ul style="list-style-type: none"> <li>identifies water quality values important to the community</li> <li>informs regional natural resource management</li> </ul>	<ul style="list-style-type: none"> <li>identifies water quality objectives to enhance or protect environmental values</li> <li>links to land use planning and development assessment</li> </ul>	<ul style="list-style-type: none"> <li>sets guidelines for water quality in the Reef to maintain marine ecosystem health</li> </ul>	<ul style="list-style-type: none"> <li>sets guidelines for water quality that maintain freshwater ecosystems health</li> </ul>

The Mackay Whitsunday water quality improvement plan provides the mechanism for identifying high risk areas for the range of pollutants, the targets to achieve the environmental values and water quality objectives identified by the community, and the links to the actions needed to achieve them. The plan guides investment into critical areas for improving water quality linked to aquatic and marine assets.<sup>100</sup> It also defines and uses best management practice frameworks to guide a continuous improvement process. These frameworks, based on the ABCD land condition framework, were developed with each of the industry partners and implemented by community stakeholder working groups.

Reef Plan was established by the Australian and Queensland governments in 2003 to halt and reverse the decline in water quality from diffuse agricultural sources by 2013. It was updated in 2009 after the release of a scientific consensus statement on water quality in the Great Barrier Reef which stated that current management interventions were not effectively solving the problem and that accelerated action was needed. It was updated again in 2013 with a renewed focus on best practice; more coordinated capacity building services and continued collaborative effort by both governments in close partnership with industry and landholders.

The long-term goal of Reef Plan 2013 is to ensure that, by 2020, the quality of water entering the Reef from broadscale land use has no detrimental impact on the health and resilience of the Great Barrier Reef. It includes targets to reduce anthropogenic end-of-catchment dissolved inorganic nitrogen by at least 50 per cent, anthropogenic end-of-catchment loads of sediment and particulate nutrients by at least 20 per cent and end-of-catchment pesticide loads by at least 60 per cent by 2018. It includes a number of direct actions to avoid and mitigate the impacts of catchment run-off from agricultural sources, which is the greatest contributor of pollutant loads entering the Great Barrier Reef lagoon. This includes water quality grants under the Australian Government's Reef Rescue program, the Queensland Government's Reef Protection Regulations, education and extension services, and management of public land (Figure 9.8).






Figure 9.8 Key components of the Reef Water Quality Protection Plan<sup>40</sup>

The reviewers assessed the effectiveness of the Authority's management of water quality across the whole Reef; this is provided in detail in the preceding chapter (Chapter 8). The assessment, which is also relevant to the Mackay Whitsunday region, includes the following key points: *The most significant contribution to water quality decline in the Great Barrier Reef is from activities outside the Marine Park associated with agricultural practices. This limits the Authority's capacity to take direct action with respect to water quality decline.*

- **Context** is effective. *The values that underpin the matters relevant to water quality are well understood by the Authority. While many of the direct and indirect impacts of poor water quality are well known, knowledge is not as comprehensive concerning the consequential and cumulative impacts of water quality.*
- **Planning** is mostly effective. *Since the Outlook Report 2009, a review of Reef Plan has provided better focus and direction for managers including the Authority, including targets for water quality and land management improvement. Water quality guidelines [for the Great Barrier Reef Marine Park]<sup>101</sup> and the development of a Coastal Ecosystems Assessment Framework<sup>102</sup> by the Authority set limits for water entering the Marine Park and provide a framework for assessment of ecosystem services within the basins located in the catchment with a focus on improving the health and resilience of the Reef.*
- **Inputs** are mostly effective. *The Authority has allocated significant resources into understanding the water quality issues from a biophysical aspect, though information is still limited with respect to the socio-economic impact of loss of ecosystem services from poor water quality.*
- The delivery of **outputs** is mostly effective. *The Authority monitors the long-term health of key marine ecosystems and the condition of water quality in the inshore lagoon.*
- **Outcomes** overall and **biodiversity outcomes** are partially effective. *The Authority's management of water quality is through legislation and permits for point source discharges into the Marine Park, as well as undertaking the marine monitoring functions of Reef Plan<sup>39</sup>. The Authority engages with key stakeholders and has expanded the Reef Guardian program (which commenced with schools and local councils), to include farmers, graziers and fishers. However, as the management of water quality leaving the Great Barrier Reef catchment is significantly within the Queensland Government's jurisdiction, the Authority can only work with the community and stakeholders to encourage best practice land management.*



## 9.8.4 Outcomes and potential improvements

The development of the Mackay Whitsunday Water Quality Improvement Plan, and its environmental values and water quality objectives, is critical to restoring and then maintaining the ecological values of the area's waterways and the Reef, and to identifying actions to achieve this. Measures to improve water quality and environmental values are linked to the water quality guidelines developed for the Great Barrier Reef Marine Park<sup>103</sup> and Queensland waters.<sup>104</sup> This establishing of thresholds linked to management tools has been shown to be a very effective measure for addressing the cumulative pressures and impacts of development on water quality. They establish a baseline for healthy systems against which development can be assessed and actions can be taken to halt further decline. They can also be used as a guide in restoring a system to a healthy state.

Recent research on coral decline on the Great Barrier Reef<sup>64</sup> reinforces the need to continue improving water quality to help restore outbreaks of crown-of-thorns starfish to natural cycles. A range of laws and policies are in place to regulate the various sources of pollutants. The package of water quality management tools, coordinated through Reef Plan, demonstrates how government, working with industry and the community, can address the very complex issue of poor water quality. The second Reef Plan report card, which represents the first year (2009–10) of significant full investment in on-ground actions, shows reasonable progress towards Reef Plan's targets.<sup>105</sup> As such, there is an expectation that the target of halting and reversing the decline in the quality of water entering the Reef is achievable; however, this is just the first step towards achieving the much more ambitious target that the quality of water entering the Reef from broad scale land use has no detrimental impact on the health and resilience of the Great Barrier Reef. This would effectively meet the requirement of the Authority's water quality guidelines for maintaining a healthy and resilient Great Barrier Reef.

Initial progress towards achieving healthy waters in the area covered by the Mackay Whitsunday Water Quality Improvement Plan is encouraging. The most recent Reef Plan 2011 report card, released in July 2013, showed that in the Mackay Whitsunday region, total nitrogen, sediment and pesticide loads have been reduced by 13 per cent, 6 per cent and 31 per cent respectively. This is a significant achievement since 2008–09, with the area demonstrating the biggest percentage reduction of all regions for nutrients and pesticides, the two most important pollutants in this region. A diverse range of stakeholders has demonstrated the expertise and commitment to achieving critical water quality improvements. However, ultimate success will require a range of ongoing actions.

### Enhancing protection and restoration

Taking action to improve water quality represents one of the best opportunities to improve the condition of the Great Barrier Reef World Heritage Area, while providing improved resilience to other impacts such as those related to a changing climate. There is a need for increasing emphasis on a whole-of-ecosystem approach to restoring catchment ecosystem health and connectivity, including wetland ecological functions. The Reef 2050 Long-term Sustainability Plan will integrate and coordinate management actions needed to restore, maintain and enhance the Region's values.

### Minimising impacts

A long-term commitment by the Australian and Queensland governments to resourcing water quality improvement initiatives is a key requirement in minimising water quality impacts on the values of the Region. Recognising that emphasis remains on reducing the impacts of broadscale land use, an increasing focus on programs aimed at improving all sources of pollutants (such as urban and industrial discharges) affecting water quality in the catchment and the Region would further reduce impacts.

### Improving adaptive management

A commitment to ongoing monitoring and review of the overarching Reef Plan and the more regionally-based Mackay Whitsunday Water Quality Improvement Plan would improve the ability to measure effectiveness of management and adapt the relevant programs.

## 9.9 Keppel Bay

### 9.9.1 Significance

Keppel Bay is a broad bay on the Capricorn Coast in central Queensland, at the mouth of the Fitzroy River in the southern Great Barrier Reef World Heritage Area (Figure 9.9). The Keppel islands are located in the shallow basin in the north of Keppel Bay, 15 to 20 kilometres off the coastal town of Yeppoon. The Capricorn Coast is the traditional land of the Dharumbal people, and the Woppaburra people are the Traditional Owners of the Keppel Bay islands.

The area contributes to all four of the natural criteria for which the World Heritage Area was listed in 1981: exceptional natural beauty, significant geomorphic features, significant ongoing ecological and biological processes, and significant natural habitats for the conservation of biological diversity. Unusually high coral cover and fish diversity, a vast array of habitats, strong cultural value for Traditional Owners, significant historic heritage, a growing coastal population and extensive access by the community to the local environment make Keppel Bay and islands ecologically, culturally and socially significant, and of high aesthetic value.<sup>80</sup>

The reefs in the area are primarily fringing inshore coral reefs, surrounding the continental islands and rocky outcrops. The islands are under the jurisdiction of the Queensland Government. Fifteen of them (excluding Great

Keppel, Pumpkin and Wedge islands) are surrounded by the Marine National Park zone, while one (Peak island) is within the Preservation zone (Figure 9.9).

Within the area Peak Island, Outer Rock and Egg Rock were identified as 'special and unique sites' during the Representative Areas Program conducted to inform the 2003 rezoning of the Marine Park.<sup>106</sup>

The area is adjacent to the internationally recognised Shoalwater and Corio Bay Area wetland — a matter of national environmental significance.

The bay and the islands are home to 34 listed threatened species and 54 listed migratory species. Its fauna and flora includes:

- three species of inshore dolphin (Australian snubfin, Indo-Pacific humpback and Indo-Pacific bottlenose). The population of the endemic Australian snubfin dolphin is one of the largest known to occur in the World Heritage Area and is at the southernmost extent of its range<sup>107</sup>
- humpback whales, sharks and stingrays
- four species of turtle (flatback, green, loggerhead and hawksbill). Peak Island is one of four major nesting sites for the flatback turtle along the Great Barrier Reef<sup>108</sup>
- one of only a handful of continental island groups on the Reef where sea snakes occur in significant numbers.<sup>109</sup>

Keppel Bay and islands was chosen as a demonstration case study to examine the cumulative impacts from climate change and extreme weather, rapid coastal development, and high recreational and commercial use.

### 9.9.2 Key issues

Many of the habitats in the adjacent Fitzroy River catchment have been modified. Overall, vegetation loss from clearing within the Fitzroy natural resources management region totals more than 89,000 square kilometres.<sup>44</sup>

The Capricorn Coast region, which includes Keppel Bay and islands, is developing rapidly. Since 2000, there has been an increase in tourism (although activity is still below the peaks of the late 1980s and 1990s), and an increase in recreational boating.<sup>110</sup> The Keppel islands are famous for their white sandy beaches and fringing coral reefs and are extensively used for recreational purposes. The largest island is Great Keppel Island, which covers an area of 14.5 square kilometres. It has had a small tourist resort for many years. The Australian and Queensland governments have approved a new marina and expanded tourism development on the island. It is proposed to include a hotel, villas and apartment accommodation, an 18-hole golf course, an upgrade and expansion of the current airstrip, a 250-berth marina at Putney Beach, the restoration of an historic homestead to a research and exhibition centre, and a range of services and utilities to support the resort including a submarine power cable, waste collection and a water treatment plant.


Assessments are currently underway on applications seeking approval to develop new port infrastructure and loading facilities at Port Alma at the mouth of the Fitzroy estuary, including transshipping facilities offshore near Peak Island. These proposals sit in the context of the very high ecological and heritage values of Keppel Bay; the cumulative impacts in the bay of major floods over successive years which delivered significant loads of sediments, nutrients and pesticides from existing agricultural areas; population growth and expanding urban activity in the catchment; an increase in coastal development; and proximity to a major and rapidly expanding port (Port of Gladstone).

The reef communities of Keppel Bay are exposed to a range of environmental pressures. They have been affected by cyclones, flooding and bleaching events over the past 20 years. Prior to the 2011 flood, the Keppel Bay reefs retained an extraordinarily high coral cover — greater than 50 per cent average hard coral cover in 2010<sup>111</sup> — especially given their southern inshore location.

Sea surface temperature anomalies caused severe coral bleaching in 1998, 2002 and 2006.<sup>61,111</sup> The remarkable recovery from the 1998 and 2002 bleaching events and from historic floods demonstrates a high resilience.<sup>112</sup> However, expectations of recovery are less optimistic after almost 40 per cent of corals were lost from the 2006 bleaching event and subsequent, almost annual, flooding between 2007 and 2011.<sup>70</sup>

The discharge from the Fitzroy River exposes affected marine habitats to elevated sediments, nutrients and pesticides as well as large volumes of freshwater, which reduces the salinity of the affected marine waters. Freshwater inundation from a flood in 2011 affected a large proportion of inshore reefs to a depth of at least two metres. The proportion of macroalgae has also increased significantly on several of the inner reefs, while large reef areas have lost the majority of their hard coral cover.

In 2012, the condition and trend of reefs in the Keppel Bay varied greatly. To date, there are no signs of recovery of affected coral communities, and the regional assessment of coral reef condition in this area is poor and declining.<sup>70</sup>



**Figure 9.9 Keppel Bay and its islands**

Perforated and Flat islands are not shown

### 9.9.3 Current management and its effectiveness

Management in the World Heritage is underpinned by the best available science, much of which is conducted in the Keppel area. These studies have provided evidence that the Zoning Plan is an effective ecosystem-based management tool — when there is high compliance, there is an increase in the abundance of coral trout and stripy snapper in Marine Park zones which then act as a source of fish for nearby General Use zones.<sup>113</sup>

In addition to the Zoning Plan, specific threat abatement actions (such as no-anchoring areas, declared fish habitat, a Traditional Use of Marine Resources Agreement and two Public Appreciation Special Management Areas) are in place to maintain biodiversity, heritage values and community benefits in the area.

The Authority is also partnering with the Australian and Queensland governments in delivering Reef Plan to improve the water quality of the Fitzroy catchment through actions such as incentives, education and extension (see Mackay Whitsunday case study, Section 9.8).

The independent review of the Authority's assessment concluded:

- **Context** is mostly effective. *The Authority has reasonable knowledge of the values and key impacts affecting the Keppel Bay region, such as tourism and recreation, fishing, water quality and coastal development. Knowledge of the consequential and cumulative impacts of increased shipping and proposed port development is still being developed.*
- **Planning** is mostly effective. *The demonstration case study provides a number of examples of planning that support the protection of the Keppel Bay area, including zoning plans, specific threat abatement plans with no-anchorage areas, declared fish habitat, Reef Plan; a Traditional Use of Marine Resources Agreement is in place, and two Public Appreciation [Special Management] Areas are in place. Stakeholders are strongly engaged, and the joint Field Management Program is in place. Development of a plan of management could address the increased use of the area. Work to manage wastewater from non-agricultural sources (such as mine discharges) is also required to complement the progress made with the Reef Plan.*
- **Inputs** are partially effective. *Resources for key management actions (for example mitigating impacts of coastal land management), field management (surveillance and compliance, emergency response) and planning for increased recreational use and access are either currently inadequate or not secure in the long term.*

- **Process** is mostly effective. *Effective partnerships with stakeholders and the community are in place, and the joint management arrangements with the Queensland Parks and Wildlife Service ensure consistency in permitting and compliance. The ways in which the plans and strategies are being implemented is appropriate, and there is a sound biophysical and socio-economic knowledge base for decision making.*
- **Outputs** are mostly effective. *Outputs which influence the management effectiveness of the Keppel Bay region include strong stakeholder partnerships, the Recreational Management Strategy and the Great Barrier Reef Biodiversity Conservation Strategy. The strategy emphasises the importance of the Keppel Bay region for the protection of the Australian snubfin dolphin population.*
- **Outcomes** are mostly effective overall. *Issues of multiple use and development within the area are reasonably well managed, and effective partnerships are in place to address impacts of activities on matters of national environmental significance. Biodiversity outcomes are partially effective. Severe weather and flooding have limited the recovery of corals, although the populations of the vulnerable flatback turtles and Australian snubfin dolphin are stable.*

#### 9.9.4 Outcomes and potential improvements

The condition of matters of national environmental significance in Keppel Bay is projected to decline in coming decades due to the cumulative impacts of climate change, water quality and increasing use.

##### Enhancing protection and restoration

Building resilience by reducing cumulative impacts will improve the capacity of the ecosystem to withstand the projected effects of a changing climate. While further degradation is unlikely to be avoided altogether, the extent of damage over coming decades is likely to be influenced by the effectiveness of local management to reduce stress to reefs from degraded water quality, anchor damage, overfishing or other pressures. This requires the maintenance of strong linkages between people of the Keppel region and the surrounding marine environment and strong local support for initiatives that aim to build the resilience of reefs in the area.

Further initiatives to manage pollutants from non-agricultural sources (such as urban and industrial discharges, dumping and resuspension of dredge material) are required to complement the progress made by Reef Plan and to improve conditions for ecosystem recovery.

The area's diversity of habitats, ease of accessibility and range of activities, combined with the projected significant increases in future use, point to the need for a plan of management. Plans of management for other areas in the Region have provided greater certainty around where specific uses may occur, the types of activities allowed, the conditions under which activities may proceed and circumstances where impacts are likely to be unacceptable.

##### Minimising impacts

The installation and maintenance of site infrastructure will be fundamental to protecting biodiversity values and community benefits derived from this unique natural area.

##### Improving adaptive management

The protection of the area's biodiversity values relies on an improved understanding and spatial assessment of the distribution and abundance of species and habitats and supporting ecosystem processes. Better mechanisms for aligning and sharing information are also essential to ensuring management actions are considered using the best available information.

There is a need to identify more fully the area's Indigenous and historic heritage values. There is also a need for better long-term socio-economic information and an improved understanding of the patterns of commercial and recreational use in the Keppel Bay area, especially to inform planning and site infrastructure improvements.

Ongoing monitoring is crucial to effective management, providing information on conditions and trends, supporting assessments of management effectiveness and informing adaptive management.

## 9.10 East Coast Trawl Fishery

### 9.10.1 Significance

The East Coast Trawl Fishery is an example of a fishery making good progress in addressing environmental impacts to Great Barrier Reef values.

The Queensland trawl fishery has a gross value of production of approximately \$100 million, supplying about 8000 tonnes of seafood including prawns, scallops, Moreton Bay bugs and squid. It also directly employs more than 1500 people and provides social and economic benefits to coastal communities along the Queensland east coast.

Trawl fisheries globally and in Australia have been a focus of environmental concern — largely due to the physical impacts when fishing gear is towed on or near the seabed, and because the small mesh used in prawn trawl gear results in bycatch of animals which are discarded, often dead or with a low chance of survival. In the Region, other concerns include the incidental catch of species of conservation concern (for example sea snakes).<sup>5,114</sup>



In the Region, the trawl fleet operates in coastal, lagoon floor and upper continental slope habitats, which support a great diversity of marine life. Trawling interacts with hundreds of non-target species to varying degrees. Some productive prawn grounds are trawled intensely each year; other areas are trawled much less frequently. If not appropriately managed, this fishing activity has the potential to harm inshore and offshore habitats, species and ecological processes. However, improved practices and current trawl fishing effort levels — which are less than half those of the maximum in 1997 — mean overall ecological risks from trawling in the Region have been reduced.

### 9.10.2 Key issues

The commercial trawl fishery developed in the 1950s and has undergone a succession of changes (Figure 9.10).

Over the past 20 years, annual trawl fishing effort has decreased, the fleet has become smaller, and habitat protection through zoning has increased. Trawling is allowed in 34 per cent of the Marine Park. Only about seven per cent of the Marine Park is trawled more than once per year. In addition, the trawl industry has adopted improved fishing practices such as bycatch reduction devices and turtle excluder devices which allow large animals, including marine turtles and some sharks, to escape. These changes have significantly reduced ecological risks.

At the same time, research has provided a better understanding of seabed habitats in non-reef areas, trawling impacts and the recovery for seabed habitats and species; the status of some resource species; trawl bycatch; and protected species interactions with the fishery.<sup>114,115,116,117,118,119,120</sup> However, there remain some challenges for the trawl fishery such as excess fishing capacity of approximately 40 per cent, a few high ecological risks for some species and habitats, and low profitability.<sup>115,121</sup> These remaining challenges put pressure on fishing businesses and regional communities.

From an economic perspective, the trawl fishery currently has too many operators for all individual businesses to be viable. This is reflected in both the profitability of businesses operating in the fishery and in the number of businesses that have chosen to simply not work at all in recent years. While the economic circumstances are unlikely to change into the future, there is a risk that if all operators chose to fish then pressure on individual stocks may become unsustainable.

Better control of trawl fishing effort would likely improve the economics of the fishery and the ecological impacts, since effort levels are a key driver of future ecological risk and impacts from trawling.<sup>115,120</sup>

The current management rules cap overall possible effort and restrict fishing efficiency, such as maximum horsepower for engines and net length restrictions, as well as setting fishing season opening and closing dates. However, improvements are possible. Options for change have been in development since late 2009 and will be considered by the public, fishers and all levels of government prior to approval by the Queensland Government which is expected in the near future.

Several aspects of the trawl fishery may be vulnerable to climate change.<sup>122</sup> Rising sea temperatures and altered ocean currents may send some species southwards, while challenges are posed by the prospect of changing rainfall patterns (and thus changing nutrient inputs), as well as floods, ocean acidification and more intense cyclones.

The trawl fishery may be affected by coastal development and port expansions. Some trawl fishing grounds (which are all in General Use zones, as are ports) have become inaccessible to trawling because they are now used for anchoring many large ships<sup>3</sup>. Fishers report the seabed topography and productivity may be changed by anchoring and that changes in bathymetry may compromise the safe and optimal operation of trawl equipment<sup>3</sup>. Also, port expansions (including reclamation and dredging) may lead to direct loss of grounds. Reduced water quality may be an additional factor, such as through resuspension of sediments and contaminants, potentially affecting productivity for filter-feeding species like scallops. The seafood industry is dependent on access to resources, and the productivity of seafood resources and their market value are influenced by the health and reputation of the Great Barrier Reef.


### Ecological risks from trawling in the Great Barrier Reef Region

The trawl industry has reduced overall risks to the Great Barrier Reef over the past two decades.

Under current practices and trawl fishing effort levels (which are less than half those of the maximum in 1997), overall ecological risks from trawling in the Great Barrier Reef Region are relatively low, with most species, habitat types and the broader ecosystem at low or intermediate–low risk from trawling.

- This low risk profile largely resulted from a combination of: 1) those species, habitats and broader ecosystems exposed to prawn trawling in the Great Barrier Reef having characteristics that give them a degree of resilience to impacts, including from fishing; 2) the relatively low effort exerted by the trawl fishery in recent years leading to relatively low pressure at the scale of the Great Barrier Reef ecosystem under current circumstances; and 3) the extensive knowledge base for the fishery and Great Barrier Reef.
- The low risk profile does not infer that trawling *per se* represents a low risk to marine habitats. The assessment took into account current fishing effort levels and the full suite of management arrangements in place, including protected area management and very fine spatial scale data on the distribution of fishing effort.
- There is no certainty that this level of impact will not increase in the future because much higher levels of fishing effort are allowable under current fishery management arrangements. Even though there is a cap on trawl effort levels within the World Heritage Area, this mechanism would be ineffective at preventing a very substantial increase in fishing pressure, as the amount of effort allowed under the cap is double the amount actually used in 2011.
- Investment in environmental research has provided a robust basis for assessing ecological risks. Access to fishery observer and research information, including previously unpublished data on discards and incidental catches in the trawl fishery, improved confidence in the ecological risk assessment.
- There are some remaining ecological risks posed by the otter trawl fishery. In particular, high risks were identified for 11 species of skates and rays, two species of sea snakes, a poorly-known upper continental slope habitat (90 to 300 metres depth) in the southern Great Barrier Reef and the plant and animal communities there. The by-product species, Balmain bugs, were at intermediate to high risk.
- The fishery still has a substantial interaction with sea snakes. Research and extension work has identified improvements to bycatch reduction devices that can be highly effective at excluding sea snakes, but these are yet to be fully adopted across relevant areas of the fishery.
- No sharks or shark products are allowed to be retained from the Queensland trawl fishery for conservation reasons, and turtle excluder devices allow large animals including some sharks to escape. However, smaller sharks and those with a flattened body form (such as many rays) are still being caught in trawl nets, landed on deck and then discarded. Larger animals are also occasionally caught (for example, green sawfish through entanglement).
- Survival of sharks and rays after trawling is often poor, with fishery observer data indicating immediate survival is only 35 per cent on average, while only 17 per cent are released alive in the deepwater trawl sector.
- The high risk rating for the deepwater habitat was precautionary due to the lack of knowledge about this area. About half of this area receives consistently high levels of trawl fishing effort. Additional ecological and biological information is required to more confidently assess the risks posed by the fishery in this area.
- The trawl fishery is diverse and while some of the remaining ecological risks were found to be fishery-wide, others are specific to fishery sectors. For example, all sectors interact with one or more of the sharks and rays that were assessed as being at high risk, and the red-spot king prawn sector accounted for most of the incidental trawl fishing mortality for sea snakes.

Data from *Ecological risk assessment of the East Coast Otter Trawl Fishery in the Great Barrier Reef Marine Park: summary report*.



**Figure 9.10 Chronology of the Queensland East Coast Trawl Fishery**

Data from *Ecological risk assessment of the East Coast Otter Trawl Fishery in the Great Barrier Reef Marine Park*<sup>1</sup>

### 9.10.3 Current management and its effectiveness

The trawl fishery is directly managed by Fisheries Queensland within the Queensland Department of Agriculture, Fisheries and Forestry. In addition to fisheries legislation, trawling activities in the Great Barrier Reef Region must comply with the Zoning Plan and other environmental laws. The operation of the fishery has been assessed against national environment law since 2004.<sup>123</sup> An independent assessment of management effectiveness for commercial fishing in the Region was completed in 2009,<sup>124</sup> and aspects of this were examined again in 2013 to inform the strategic assessment.

A major review of trawl fishery management arrangements has been underway since 2010; completing this review is a priority of the current Queensland Government. This review process has two advisory groups that provide advice to Fisheries Queensland. The technical advisory group consists of representatives from the commercial fishing industry, recreational fishing, the Authority, conservation organisations, trawl fisheries management, and institutions with science and economic expertise. The scientific advisory group consists of members with expertise in trawl fishing, seafood processing and marketing, economics, science and management. Between 2010 and 2012, these groups examined in detail the pressures facing the industry and considered what the different stakeholders want for the future of the fishery, including legislative requirements.<sup>125,126</sup>

All advisory group members reached a consensus position<sup>127</sup> on the need for change and the type of management change required to develop a profitable, ecologically and socially sustainable trawl fishery for Queensland into the future. The agreed solution was:

- a significant reduction in trawl fishing effort (structural adjustment) to improve profitability and manage ecological risks
- new management rules that change or remove rules that restrict efficiency and that control fishing effort to profitable levels and at optimal times.

It was also agreed that total allocated effort in the fishery must be reduced below current levels of real effort before restrictions on efficiency can be lifted, and that environmental safeguards must be maintained.

Use of a satellite-based vessel monitoring system for the last decade has enabled close monitoring of the trawl fleet's activities, resulting in improved compliance and major benefits for fishery assessments. This detailed spatial information was used to confirm that Great Barrier Reef Marine Park zoning, complemented by Queensland zoning, effectively protected representative seabed habitats and associated biodiversity at levels consistent with the conservation targets underlying the 2003 rezoning of the Marine Park.<sup>128,129</sup> Protection through zoning is also an important measure which acts to limit spatial expansion of the trawl fishery and potential risk to the ecosystem.<sup>115</sup> Risks and impacts have been reduced over the last decade or more, but there is no certainty that the level of impact will not increase again in the future because current fishery effort controls are ineffective.<sup>115</sup>

A number of fishers have voluntarily adopted fishing practices that have less of an environmental impact and which go beyond those required by the Regulations. For example, new prawn trawling otter boards, designed and adopted by some fishers, reduce physical impacts on the seabed and also reduce towing resistance and fuel usage. Additionally, many fishers have been actively working with the Queensland Department of Agriculture, Fisheries and Forestry to trial and improve bycatch reduction devices. This work is helping to inform the optimum selection of these devices for different sectors of the fishery.<sup>114,130</sup>

The Authority is actively helping the industry face future challenges from climate change and extreme weather through measures such as adaptation planning workshops.<sup>131</sup> Some of the responses proposed by stakeholders through these workshops included modernising the trawl fleet and its operations, creating a more flexible fisheries management policy framework, diversifying income streams, developing more forward-looking business strategies, and introducing real-time fisheries data collection.

Effective management of trawling and public confidence in the sustainability of the fishery requires good information, including fishery observer programs, improved reporting via logbooks and ongoing risk monitoring (including analysis of trawl effort patterns).

The effectiveness of the Authority's management of commercial fishing was addressed by the independent reviewers, and is provided in detail in Chapter 8. In brief, the management effectiveness of commercial fishing (including trawl) of the Authority was rated as partially effective for all assessment criteria except context, where it was mostly effective.

*The Authority has a regulatory role in the management of commercial fishing through its Zoning Plan, Regulations and permits for a limited number of Queensland-managed commercial and developmental fisheries. While the Authority has a direct role in ensuring that fishing is ecologically sustainable in the Marine Park it has an advisory role on the management of commercial fisheries and works with other agencies to improve fisheries management. Commercial fisheries arrangements in the Reef are accredited against national sustainability guidelines. The Authority is involved in this role in an informal advisory capacity only. As identified in the Outlook Report 2009, the Authority has a good understanding of commercial catch information. Cumulative impacts associated with commercial fishing are reasonably well known. The recent ecological risk assessment of the East Coast Otter Trawl Fishery in the Great Barrier Reef found the overall ecological risks from trawling are relatively low, but there are some high ecological risks for deep water skates, several rays, and sea snakes.*<sup>115</sup>



*A reduction in trawl fishing effort of more than 40 per cent between 2005 and 2009, driven by prevailing economic conditions rather than management intervention, has significantly reduced the ecological risk from trawling. However, higher effort levels are still allowable under existing management arrangements. Further reductions in trawl bycatch and other efforts to reduce risks for species of conservation concern are important for the sustainability of the fishery.*

#### **9.10.4 Outcomes and potential improvements**

There has been a significant reduction in the risks and impacts to the values of the Great Barrier Reef since the introduction of a management plan for the trawl fishery in 1999 and subsequent management actions (Figure 9.10).

Challenges facing the fishery are increasingly better understood, and an ecosystem-based approach to management is being put into practice. Some successful trawl fishery initiatives, such as the vessel monitoring system and improved gear design to reduce bycatch, have resulted in substantial environmental benefits.

Notwithstanding the significant progress made by the fishery, current management arrangements for trawling are only partially effective at delivering environmental, economic and social outcomes.

##### **Enhancing protection and restoration**

Reaching consensus on desirable changes was an important step in the review of trawl fishery management arrangements. As a result, an opportunity has been identified for future arrangements that would provide net environmental benefits, make a major contribution to protecting the Great Barrier Reef and maintain biodiversity. This will also set the industry on a better path for the future and support a number of coastal and regional communities.

##### **Minimising impacts**

Further actions are required to mitigate the remaining ecological risks — a high priority is reducing incidental catch and death rates of sea snakes and smaller individuals of sharks and rays.<sup>115</sup> Potential risk mitigation strategies have been considered through the recent review processes and may include managing fishing effort, strengthening compliance, supporting research to address knowledge gaps, optimising the use of current bycatch reduction devices, supporting other initiatives to reduce interactions with protected species, and measures that limit fishing pressure on other by-product species.

##### **Improving adaptive management**

The trawl industry is starting to plan for future challenges such as climate change, and some fishers are preparing their businesses. Such future planning may also help with other challenges facing the fishery and help support a healthy marine environment into the future.

### **9.11 Summary of conclusions and guidance for future management**

Along the length of the Region, values, impacts and risks vary. The case studies identify a suite of issues common across the Region (for example, impacts from extreme weather, catchment run-off, degradation of coastal ecosystems and direct use). They also show that the way these issues interact with other activities and values varies from place to place. Regional and local approaches, based on local knowledge and actions, will therefore be central to protecting and managing the values relevant to matters of national environmental significance.

The demonstration case studies also highlight that connectivity within and between terrestrial and marine ecosystems are critical to the health and resilience of the World Heritage Area.

The finer detail assessment of the effectiveness of current management arrangements to protect and manage the relevant matters of national environmental significance demonstrates the need for a suite of management actions to address impacts affecting values across the Region. For each of the case studies, enhancing protection and restoration, minimising direct impacts and improving adaptive management are common themes for potential improvements. These actions range from initiatives that apply across the whole Region, to more regional and local approaches.

Collectively, the demonstration case studies highlight the following pathways for future management.

Being proactive:

- Tackling climate change is fundamental to protecting key habitats such as coral reefs and species such as dugongs which rely on seagrass meadows.
- Clear but adaptable management approaches are required to deal with extreme weather such as flooding and cyclones; and coral bleaching.
- Greater investment and planning is required, particularly in areas of high growth. Aligning, rationalising and updating management tools such as plans of management, policies and permits will make management more outcome-focused, collaborative, responsive and adaptive.
- Increased field management capacity is required to address impacts and improve compliance (both education and enforcement). This recognises that the remoteness of much of the Region protects some values, while simultaneously making management intervention difficult.

Partnerships are the key:

- Enhancing effectiveness of partnerships between policy makers, regulators and researchers, to ensure maximum value of research investments.
- Strengthening efforts to improve water quality entering the Reef from its catchment remains a priority for restoring fundamental processes, such as sedimentation and nutrient cycling, which in turn affect the health and recovery of many habitats and species.
- Strong partnerships across all levels of government are vital to addressing the many cross-jurisdictional issues associated with the protection and management of marine ecosystems.
- Cooperative management approaches between governments, Traditional Owners, stakeholders and the community are an effective way to address impacts and declines in condition at a local scale.
- Cooperative management arrangements with Traditional Owners and programs to support their involvement in protecting and managing Indigenous heritage values will enhance overall protection and management of the Region's values.

Foundational activities underpin management:

- Long-term integrated monitoring and reporting across all the Region and for a range of habitats and species is required to track the condition of the Region's values and inform adaptive management.
- Reef-wide standards for data collection and sharing are required to improve data use across a range of scales and to make the information more accessible for managers and the public.
- Improved understanding about Indigenous and historic heritage values and community benefits derived from the Region is needed to improve their consideration in decision making.

---

## References

1. Pears, R.J., Morison, A.K., Jebreen, E.J., Dunning, M.C., Pitcher, C.R., Courtney, A.J., Houlden, B. and Jacobsen, I.P. 2012a, *Ecological risk assessment of the East Coast Otter Trawl Fishery in the Great Barrier Reef Marine Park: summary report*, Great Barrier Reef Marine Park Authority, Townsville.
2. Sinclair Knight Merz Pty Ltd and Asia-Pacific Applied Science Associates 2013, *Improved dredge material management for the Great Barrier Reef Region: Synthesis report*, Great Barrier Reef Marine Park Authority, Townsville.
3. GHD Pty Ltd 2013, *Ship anchorage management in the Great Barrier Reef World Heritage Area*, Great Barrier Reef Marine Park Authority, Townsville.
4. GHD 2013, *Environmental best practice port development: an analysis of international approaches*, Department of Sustainability, Environment, Water, Population, and Communities, Canberra, Australia.
5. Great Barrier Reef Marine Park Authority 2009a, *Great Barrier Reef Outlook Report 2009*, GBRMPA, Townsville.
6. Hockings, M., Gilligan, B. and Leverington, A. 2013, *Assessment of management effectiveness for the strategic assessment of the Great Barrier Reef Region*, Great Barrier Reef Marine Park Authority, Townsville.
7. Great Barrier Reef Marine Park Authority 1981, *Nomination of the Great Barrier Reef by the Commonwealth of Australia for inclusion in the World Heritage List*, GBRMPA, Townsville.
8. Marsh, H. 2008, *Dugong dugon: IUCN Red List of Threatened Species Version 2013.1*, IUCN The International Union for Conservation of Nature, Cambridge, UK, <<http://www.iucnredlist.org/details/6909/0>>.
9. Marsh, H., Eros, C., Corkeron, P. and Breen, B. 1999, A conservation strategy for dugongs: implications of Australian research, *Marine and Freshwater Research* 50(8): 979-990.
10. Marsh, H., Penrose, H., Eros, C. and Hugues, J. 2002, *Dugong: status report and action plans for countries and territories*, United Nations Environment Program, IUCN, Nairobi.
11. Marsh, H., Channells, P.W., Heinsohn, G.E. and Morrissey, J. 1982, Analysis of stomach contents of dugongs from Queensland, *Australian Wildlife Research* 9(1): 55-67.
12. Marsh, H., Heinsohn, G.E. and Marsh, L.M. 1984, Breeding cycle, life history and population dynamics of the dugong *Dugong dugon* (Sirenia: Dugongidae), *Australian Journal of Zoology* 32: 767-788.
13. Marsh, H., Lawler, I.R., Kwan, D., Delean, S., Pollock, K.H. and Alldredge, M. 2004, Aerial surveys and the potential biological removal technique indicate that the Torres Strait dugong fishery is unsustainable, *Animal Conservation* 7(435): 443.
14. Marsh, H. 1995, The life history, pattern of breeding and population dynamics of the dugong, in *Population and Biology of the Florida Manatee*, eds T. O'Shea, B.B. Ackerman and H. Percival, US Fish and Wildlife Service, Washington, DC, pp. 75-83.
15. Daley, B., Griggs, P. and Marsh, H. 2008, Exploiting marine wildlife in Queensland: the commercial dugong and marine turtle fisheries 1847-1969, *Australian Economic History Review* 48(3): 227-265.
16. Marsh, H., De'ath, G., Gribble, N.A. and Lane, B. 2001, *Shark control records hindcast serious decline in dugong numbers off the urban coast of Queensland*, Great Barrier Reef Marine Park Authority, Townsville.
17. Marsh, H., De'ath, G., Gribble, N.A. and Lane, B. 2005, Historical marine population estimates: triggers or targets for conservation? The dugong case study, *Ecological Applications* 15(2): 481-492.
18. Kidston, W., Dwyer, K., Buhk, A., Waldock, J. and Anderson, G. 1992, *Review of the operation and maintenance of shark meshing equipment in Queensland waters: Report of the Committee of Inquiry*, Department of Primary Industries, Brisbane.
19. Marsh, H., Hodgson, A., Lawler, I.R., Grech, A. and Delean, S. 2007, *Condition, status and trends and projected futures of the dugong in the northern Great Barrier Reef and Torres Strait: Including identification and evaluation of the key threats and evaluation of available management options to improve its status*, Report to the Marine and Tropical Sciences Research Facility, Reef and Rainforest Research Centre, Cairns.
20. Marsh, H. and Lawler, I.R. 2006, *Dugong distribution and abundance on the urban coast of Queensland: A basis for management: Final report to Marine and Tropical Research Sciences Facility interim projects 2005-6*, James Cook University, Townsville.
21. Marsh, H. and Lawler, I.R. 2001, *Dugong distribution and abundance in the southern Great Barrier Reef Marine Park*

and Hervey Bay: Results of an aerial survey in October-December 1999, Great Barrier Reef Marine Park Authority, Townsville.

22. Grech, A. and Marsh, H. 2010, *Conditions, trends and predicted futures of dugong populations in the Great Barrier Reef World Heritage Area: Including an evaluation of the potential cost-effectiveness of indicators of the status of these populations*, Reef and Rainforest Research Centre, Cairns.

23. McKenzie, L.J., Collier, C. and Waycott, M. 2012, *Reef Rescue Marine Monitoring Program: Inshore seagrass, annual report for the sampling period 1st July 2010 – 31st May 2011*, Fisheries Queensland, Cairns.

24. Martin, K., Schaffelke, B., Thompson, A., McKenzie, L.J., Muller, J., Bentley, C., Paxman, C., Collier, C., Waycott, M. and Brando, V. 2014, *Reef Rescue Marine Monitoring Program Synthesis Report 2010/11*, Great Barrier Reef Marine Park Authority, Townsville.

25. Biddle, T.M., Boyle, M. and Limpus, C.J. 2011, *Marine wildlife stranding and mortality database annual report 2009 and 2010: Dugong*, Department of Environment and Resource Management, Brisbane.

26. Sobotzick, S., Hagihara, R., Grech, A. and Marsh, H. 2012, *Aerial survey of the urban coast of Queensland to evaluate the response of the dugong population to the widespread effects of the extreme weather events of the summer of 2010-11, Final report to the Australian Marine Mammal Centre and the National Environmental Research Program*, James Cook University, Townsville.

27. Marsh, H. and Saalfeld, W.K. 1990, The distribution and abundance of dugongs in the Great Barrier Reef Marine Park south of Cape Bedford, *Australian Wildlife Research* 17: 511-524.

28. Marsh, H. and Saalfeld, W.K. 1989, Distribution and abundance of dugongs in the northern Great Barrier Reef Marine Park, *Australian Wildlife Research* 16: 429-440.

29. Marsh, H. and Lawler, I.R. 2002, *Dugong distribution and abundance in the northern Great Barrier Reef Marine Park - November 2000*, Great Barrier Reef Marine Park Authority, Townsville.

30. Marsh, H., Corkeron, P., Lawler, I.R., Lanyon, J. and Preen, A.R. 1996, *The status of the dugong in the Southern Great Barrier Reef Marine Park*, Research Publication No. 41, Great Barrier Reef Marine Park Authority, Townsville.

31. Marsh, H. 2000, Evaluating management initiatives aimed at reducing the mortality of dugongs in gill and mesh nets in the Great Barrier Reef World Heritage Area, *Marine Mammal Science* 16(3): 684-694.

32. Department of Primary Industries and Fisheries 2006, *A report on the Queensland Shark Safety Program*, DPIF, Brisbane.

33. O'Neill, P. 2009, *State of the Environment Report for Shoalwater Bay Training Area 2008*, Department of Defence, Canberra.

34. Dobbs, K., Fernandes, L., Slegers, S., Jago, B., Thompson, L., Hall, J., Day, J., Cameron, D., Tanzer, J., Macdonald, F., Marsh, H. and Coles, R. 2008, Incorporating dugong habitats into the marine protected area design for the Great Barrier Reef Marine Park, Queensland, Australia, *Ocean and Coastal Management* 51(4): 368-375.

35. Grech, A. and Marsh, H. 2008, Rapid assessment of risks to a mobile marine mammal in an ecosystem-scale marine protected area, *Conservation Biology* 22(3): 711-720.

36. Grech, A. and Marsh, H. 2007, Prioritising areas for dugong conservation in a marine protected area using a spatially explicit population model, *Applied GIS* 3(2): 1-14.

37. Chin, A. 2005, Seagrass, in *State of the Great Barrier Reef on-line*, ed. A. Chin, Great Barrier Reef Marine Park Authority, Townsville.

38. Schaffelke, B., Waterhouse, J. and Christie, C. 2002, *A review of water quality issues influencing the habitat quality in Dugong protection areas*, Great Barrier Reef Marine Park Authority, Townsville.

39. Department of the Premier and Cabinet 2009, *Reef Water Quality Protection Plan 2009 for the Great Barrier Reef World Heritage Area and Adjacent Catchments*, Reef Water Quality Protection Plan Secretariat, DPC, Brisbane.

40. Department of the Premier and Cabinet 2013, *Reef Water Quality Protection Plan 2013*, Reef Water Quality Protection Plan Secretariat, Brisbane.

41. Anthony, K., Dambacher, J.M., Walsh, T. and Beeden, R. 2013, *A framework for understanding cumulative impacts, supporting environmental decisions and informing resilience-based management of the Great Barrier Reef World Heritage Area*, Great Barrier Reef Marine Park Authority, Canberra.

42. Wade, P.R. 1998, Calculating limits to the allowable human-caused mortality of cetaceans and pinnipeds, *Marine Mammal Science* 14(1): 1-37.

43. Meager, J.J. and Limpus, C.J. 2012, *Marine wildlife stranding and mortality database annual report 2011*, I Dugong, *Conservation and Technical Data Report* 2011(1): 1-30.

44. Great Barrier Reef Marine Park Authority 2012a, *Informing the outlook for Great Barrier Reef coastal ecosystems*, GBRMPA, Townsville.

45. Department of Agriculture, Fisheries and Forestry and Department of Sustainability, Environment, Water, Population and Communities 2012, *Caring for our Country - Reef Rescue*, DAFF and DSEWPaC, Canberra, <<http://www.nrm.gov.au/about/key-investments/reef-rescue.html>>.

46. Spalding, M.D., Ravilious, C. and Green, E.P. 2001, *World atlas of coral reefs*, University of California, Berkeley.

47. Deloitte Access Economics 2013, *Economic contribution of the Great Barrier Reef*, Great Barrier Reef Marine Park Authority, Townsville.

48. Burke, L., Reyter, K., Spalding, M. and Perry, A. 2011, *Reefs at risk revisited*, World Resources Institute, Washington DC.

49. Wilkinson, C. (ed) 2008, *Status of coral reefs of the world: 2008*, Global Coral Reef Monitoring Network and Reef and Rainforest Research Centre, Townsville.

50. Roff, G., Clark, T.R., Raymond, C.E., Zhao, J., Feng, Y., McCook, L.J., Done, T.J. and Pandolfi, J.M. 2013, Palaeoecological evidence of a historical collapse of corals at Pelorus Island, inshore Great Barrier Reef, following European settlement, *Proceedings of the Royal Society B: Biological Sciences* 280(1750): 2012-2100.

51. Hughes, T.P., Bellwood, D.R., Baird, A.H., Brodie, J., Bruno, J.F. and Pandolfi, J.M. 2011, Shifting base-lines, declining coral cover, and the erosion of reef resilience: Comment on Sweatman et al (2011), *Coral Reefs* 30(3): 653-660.

52. Fabricius, K.E. 2011, Factors determining the resilience of coral reefs to eutrophication: A review and conceptual model, in *Coral Reefs: An Ecosystem in Transition*, eds Z. Dubinsky and N. Stambler, Springer, Dordrecht, pp. 493-508.

53. Fabricius, K.E., De'ath, G., McCook, L.J., Turak, E. and Williams, D.M. 2005, Changes in algal, coral and fish assemblages along water quality gradients on the inshore Great Barrier Reef, *Marine Pollution Bulletin* 51(1-4): 384-398.

54. De'ath, G., Fabricius, K.E., Sweatman, H. and Puotinen, M. 2012, The 27-year decline of coral cover on the Great Barrier Reef and its causes, *Proceedings of the National Academy of Sciences* 109(44): 17995-17999.

55. Bridge, T.C.L., Fabricius, K.E., Bongaerts, P., Wallace, C.C., Muir, P., Done, T.J. and Webster, J.M. 2012, Diversity of scleractinian and octocorallia in the mesophotic zone of the Great Barrier Reef, Australia, *Coral Reefs* 31(1): 179-189.

56. Osborne, K., Dolman, A.M., Burgess, S.C. and Johns, K.A. 2011, Disturbance and the dynamics of coral cover on the Great Barrier Reef (1995-2009), *PLoS ONE* 6(3): e17516.

57. Birkeland, C. and Lucas, J.S. 1990, *Acanthaster planci: major management problem of coral reefs*, CRC Press, Boca Raton.

58. Brodie, J., Fabricius, K.E., De'ath, G. and Okaji, K. 2005, Are increased nutrient inputs responsible for more outbreaks of crown-of-thorns starfish? An appraisal of the evidence, *Marine Pollution Bulletin* 51(1-4): 266-278.

59. Brodie, J.E., Kroon, F.J., Schaffelke, B., Wolanski, E.C., Lewis, S.E., Devlin, M.J., Bohnet, I.C., Bainbridge, Z.T., Waterhouse, J. and Davis, A.M. 2012, Terrestrial pollutant runoff to the Great Barrier Reef: An update of issues, priorities and management responses, *Marine Pollution Bulletin* 65: 81-100.



60. Thompson, A., Costello, P., Davidson, J., Logan, M., Schaffelke, B., Uthicke, S. and Takahashi, M. 2011, *Reef Rescue Marine Monitoring Program, Report of AIMS Activities – Inshore coral reef monitoring 2011, Report for Great Barrier Reef Marine Park Authority*, Australian Institute of Marine Science, Townsville.
61. Berkelmans, R., De'ath, G., Kininmonth, S. and Skirving, W.J. 2004, A comparison of the 1998 and 2002 coral bleaching events on the Great Barrier Reef: Spatial correlation, patterns and predictions, *Coral Reefs* 23(1): 74-83.
62. Done, T., Whetton, P., Jones, R.J., Berkelmans, R., Lough, J.M., Skirving, W. and Wooldridge, S. 2003, *Global climate change and coral bleaching on the Great Barrier Reef*, Department of Natural Resources and Mines, Brisbane.
63. Williams, K.J. and Crimp, S. 2012, *Queensland's biodiversity under climate change: An overview of climate change in Queensland, Climate Adaptation Flagship Working Paper No 12A*, CSIRO Climate Adaptation Flagship, Canberra.
64. Brodie, J.E. 1992, Enhancement of larval and juvenile survival and recruitment in *Acanthaster planci* from the effects of terrestrial runoff: a review, *Australian Journal of Marine and Freshwater Research* 43: 539-554.
65. Brodie, J.E. and Mitchell, A.W. 2005, Nutrients in Australian tropical rivers: Changes with agricultural development and implications for receiving environments, *Marine and Freshwater Research* 56(3): 279-302.
66. Fabricius, K.E., Okaji, K. and De'ath, G. 2010, Three lines of evidence to link outbreaks of the crown-of-thorns seastar *Acanthaster planci* to the release of larval food limitation, *Coral Reefs* 29: 593-605.
67. Wooldridge, S.A. and Done, T.J. 2009, Improved water quality can ameliorate effects of climate change on corals, *Ecological Applications* 19: 1492-1499.
68. Bruno, J.F., Selig, E.R., Casey, K.S., Page, C.A., Willis, B., Harvell, C.D., Sweatman, H. and Melendy, A.M. 2007, Thermal stress and coral cover as drivers of coral disease outbreaks, *PLoS Biology* 5(6): e124.
69. Harvell, C.D., Jordan-Dahlgren, E., Merkel, S., Rosenberg, E., Raymundo, L., Smith, G., Weil, G. and Willis, B. 2007, Coral disease, environmental drivers and the balance between coral and microbial associates, *Oceanography* 20(1): 36-59.
70. Thompson, A., Costello, P., Davidson, J., Schaffelke, B., Uthicke, S. and Liddy, M. 2013, *Reef Rescue Marine Monitoring Program, Report of AIMS Activities – Inshore coral reef monitoring 2011, Report for Great Barrier Reef Marine Park Authority*, Australian Institute of Marine Science, Townsville.
71. McCook, L.J., Folke, C., Hughes, T., Nyström, M., Obura, D. and Salm, R. 2007, Ecological resilience, climate change and the Great Barrier Reef, in *Climate Change and the Great Barrier Reef: A vulnerability assessment*, eds J.E. Johnson and P.A. Marshall, Great Barrier Reef Marine Park Authority and Australian Greenhouse Office, Townsville, pp. 75-96.
72. Turner, M. and Batianoff, G.N. 2007, Vulnerability of island flora and fauna of the Great Barrier Reef to climate change, in *Climate change and the Great Barrier Reef: a vulnerability assessment*, eds J.E. Johnson and P.A. Marshall, Great Barrier Reef Marine Park Authority and Australian Greenhouse Office, Townsville, pp. 621-666.
73. Great Barrier Reef Marine Park Authority 2013, *Great Barrier Reef Region strategic assessment: Strategic Assessment Report. Draft for public comment*, GBRMPA, Townsville.
74. Daley, B. 2005, *Changes in the Great Barrier Reef since European settlement: Implications for contemporary management*, PhD thesis, James Cook University, Townsville.
75. Batianoff, G.N. and Cornelius, N.J. 2005, Birds of Raine Island: population trends, breeding behaviour and nesting habitats, *Proceedings of the Royal Society of Queensland* 112: 1-29.
76. Stokes, T., Dobbs, K., Mantel, P. and Pierce, S. 2004, *Fauna and flora of the Great Barrier Reef World Heritage Area*, Great Barrier Reef Marine Park Authority, Townsville.
77. Limpus, C.J. 2008, *A biological review of Australian marine turtle species, 2 Green turtle, Chelonia mydas (Linnaeus)*, Environmental Protection Agency, Brisbane.
78. Limpus, C.J., Limpus, D.J., Munchow, M. and Barnes, P. 2005, *Queensland turtle conservation project: Raine Island turtle study, 2004-2005*, Environmental Protection Agency, Brisbane.
79. Limpus, C.J., Miller, J.D., Parmenter, C.J. and Limpus, D.J. 2003, The green turtle, *Chelonia mydas*, population of Raine Island and the northern Great Barrier Reef: 1843-2001, *Memoirs of the Queensland Museum* 49(1): 349-440.
80. Context Pty Ltd 2013, *Defining the aesthetic values of the Great Barrier Reef*, Department of Sustainability, Environment, Water, Population and Communities, Canberra.
81. Stevens, J.D., Pillans, R.D. and Salini, J.P. 2005, *Conservation assessment of Glyptis sp A (speartooth shark), Glyptis sp. C (northern river shark), Pristis microdon (freshwater sawfish) and Pristis zijsron (green sawfish): final report to the Department of the Environment and Heritage*, CSIRO Marine Research, Hobart.
82. Read, M.A., Miller, J.D., Bell, I.P. and Felton, A. 2004, The distribution and abundance of the estuarine crocodile, *Crocodylus porosus*, in Queensland, *Wildlife Research* 31(5): 527-534.
83. Parra, G.J. 2005, *Behavioural ecology of Irrawaddy, Orcaella brevirostris (Owen in Gray, 1866), and Indo-Pacific humpback dolphins, Sousa chinensis (Osbeck, 1765), in northeast Queensland, Australia: A comparative study*, PhD thesis, James Cook University, Cairns.
84. Danaher, K.F. 1995, *Marine Vegetation of Cape York Peninsula*, Cape York Peninsula Land Use Strategy, Office of the Coordinator General of Queensland, Brisbane, Department of the Environment, Sport and Territories, Canberra, and Queensland Department of Primary Industries, Brisbane, Brisbane.
85. Duke, N.C. and Larkum, A.W.D. 2008, Mangroves and seagrasses, in *The Great Barrier Reef: biology, environment and management*, eds P. Hutchings, M. Kingsford and O. Hoegh-Guldberg, CSIRO Publishing, Collingwood, pp. 156-170.
86. McKenzie, L.J., Yoshida, R., Grech, A. and Coles, R. 2010, *Queensland seagrasses: Status 2010 - Torres Strait and East Coast*, Department of Employment, Economic Development and Innovation, Brisbane.
87. Lee Long, W.J., Mellors, J.E. and Coles, R.G. 1993, Seagrasses between Cape York and Hervey Bay, Queensland, Australia, *Australian Journal of Marine and Freshwater Research* 44(1): 19-31.
88. Department of Environment and Heritage Protection 2012a, 'Aquatic Conservation Assessment Cape York Catchments V1.1, Riverine and non-riverine Summary Report'.
89. Department of Environment and Heritage Protection 2012b, 'Aquatic Conservation Assessment Cape York Catchments V1.1., Riverine and non-riverine Flora, Fauna and Ecology Expert Panel Report'.
90. Beaton, J. 1985, Evidence for a coastal occupation time-lag at Princess Charlotte Bay (North Queensland) and implications for coastal colonization and population growth theories for Aboriginal Australia, *Archaeology in Oceania* 20: 1-20.
91. Sweatman, H., Cheal, A.J., Coleman, G.J., Emslie, M.J., Johns, K., Jonker, M., Miller, I.R. and Osborne, K. 2008, *Long-term monitoring of the Great Barrier Reef: Status Report 8*, Australian Institute of Marine Science, Townsville.
92. Northern Australia Land and Water Taskforce 2009, *Sustainable development of northern Australia: A report to Government from the Northern Australia Land and Water Taskforce*, Department of Infrastructure, Transport, Regional Development and Local Government, Canberra.
93. Government Statistician 2013, *Queensland Regional Profile: GBR LGAs Region*, Queensland Treasury and Trade, Brisbane.
94. Great Barrier Reef Marine Park Authority 2008, *Cairns Area Plan of Management: reprinted as in force December 2008 and includes Amendment 2008 (No.1)*, GBRMPA, Townsville.
95. Commonwealth of Australia and State of Queensland 2009, *Great Barrier Reef Intergovernmental Agreement*, Commonwealth of Australia and State of Queensland, Canberra and Brisbane.



96. Reef Water Quality Protection Plan Secretariat 2011, *Great Barrier Reef First Report Card: 2009 baseline, Reef Water Quality Protection Plan*, Reef Water Quality Protection Plan Secretariat, Brisbane.
97. De'ath, G. and Fabricius, K.E. 2010, Water quality as a regional driver of coral biodiversity and macroalgae on the Great Barrier Reef, *Ecological Applications* 20(3): 840-850.
98. Wooldridge, S., Brodie, J. and Furnas, M. 2006, Exposure of inner-shelf reefs to nutrient enriched runoff entering the Great Barrier Reef lagoon: Post European changes and the design of water quality targets, *Marine Pollution Bulletin* 52(11): 1467-1479.
99. Cooper, T.F., Uthicke, S., Humphrey, C. and Fabricius, K.E. 2007, Gradients in water column nutrients, sediment parameters, irradiance and coral reef development in the Whitsunday Region, central Great Barrier Reef, *Estuarine, Coastal and Shelf Science* 74(3): 458-470.
100. Drewry, J., Higham, W. and Mitchell, C. 2008, *Water quality improvement plan, Final report for the Mackay Whitsunday region, Mackay Whitsunday Natural Resource Management Group*, Mackay Whitsunday Natural Resource Management Group, Mackay.
101. Great Barrier Reef Marine Park Authority 2009b, *Water quality guidelines for the Great Barrier Reef Marine Park*, GBRMPA, Townsville.
102. Great Barrier Reef Marine Park Authority 2012b, *Great Barrier Reef Coastal Ecosystems Assessment Framework*, GBRMPA, Townsville.
103. Great Barrier Reef Marine Park Authority 2010, *Water quality guidelines for the Great Barrier Reef Marine Park*, GBRMPA, Townsville.
104. Department of Environment and Heritage Protection 2009, *Queensland water quality guidelines 2009*, DEHP, Brisbane.
105. Reef Water Quality Protection Plan Secretariat 2013, *Great Barrier Reef Report Card 2011: Reef Water Quality Protection Plan*, Reef Water Quality Protection Plan Secretariat, Brisbane.
106. Dobbs, K. 2011, *Identifying special or unique sites in the Great Barrier Reef World Heritage Area for inclusion in the Great Barrier Marine Park Zoning Plan 2003*, Great Barrier Reef Marine Park Authority, Townsville.
107. Cagnazzi, D., Parra, G.J., Westley, S. and Harrison, P.L. 2013, At the heart of the industrial boom: Australian snubfin dolphins in the Capricorn Coast, Queensland, need urgent conservation action, *PLoS ONE* 8(2): e56729.
108. Limpus, C.J. 2007, *A biological review of Australian marine turtle species, 5 Flatback turtle, Natator depressus (Garman)*, Environmental Protection Agency, Brisbane.
109. Lukoschek, V., Heatwole, H., Grech, A., Burns, G. and Marsh, H. 2007, Distribution of two species of sea snakes, *Aipysurus laevis* and *Emydocephalus annulatus*, in the southern Great Barrier Reef: metapopulation dynamics, marine protected areas and conservation, *Coral Reefs* 26(2): 291-307.
110. Queensland Government(Unpublished data), 'Vessel Registration Data by LGA for December 2000, 2006 and 2011. Vessel Registration Data by Postcode for 2007, 2009, 2011. Data retrieved from Department of Transport and Main Roads (DTMR) Transport Registration and Integrated Licensing System (TRAILS)'.
111. Oliver, J.K., Berkelmans, R. and Eakin, C.M. 2009, Coral bleaching in space and time, in *Coral bleaching: Patterns, processes, causes and consequences*, eds M.J.H. van Oppen and J.M. Lough, Springer, Berlin, pp. 21-40.
112. Jones, A. and Berkelmans, R. 2010, Marine 'refugia' in the Keppel region of the Great Barrier Reef, *Headmark* 136: 19-27.
113. Harrison, H.B., Williamson, D.H., Evans, R.D., Almany, G.R., Thorrold, S.R., Russ, G.R., Feldheim, K.A., van Herwerden, L., Planes, S., Srinivasan, M., Berumen, M.L. and Jones, G.P. 2012, Larval export from marine reserves and the recruitment benefit for fish and fisheries, *Current Biology* 22(11): 1023-1028.
114. Courtney, A.J., Schemel, B.I., Wallace, R.M., Campbell, M.J. and Mayer, D.J. 2010, *Reducing the impact of Queensland's trawl fisheries on protected sea snakes*, Fisheries Research and Development Corporation and Department of Employment, Economic Development and Innovation, Brisbane.
115. Pears, R.J., Morison, A.K., Jebreen, E.J., Dunning, M.C., Pitcher, C.R., Courtney, A.J., Houlden, B. and Jacobsen, I.P. 2012b, *Ecological risk assessment of the East Coast Otter Trawl Fishery in the Great Barrier Reef Marine Park: Technical report*, Great Barrier Reef Marine Park Authority, Townsville.
116. Courtney, A.J., Haddy, J.A., Campbell, v., Roy, D.P., Tonks, M.L., Gaddes, S.W., Chilcott, K.E., O'Neill, M.F., Brown, I.W., McLennan, M., Jebreen, E.J., van der Geest, C., Rose, C., Kistle, S., Turnbull, C.T., Kyne, P.M., Bennett, M.B. and Taylor, J. 2007, *Bycatch weight, composition and preliminary estimates of the impact of bycatch reduction devices in Queensland's trawl fishery*, Department of Primary Industries and Fisheries, Brisbane.
117. Queensland Department of Employment, Economic Development and Innovation 2011, *Annual status report 2011: East Coast Otter Trawl Fishery*, DEEDI, Brisbane.
118. Pitcher, C.R., Austin, M., Burrridge, C.Y., Bustamante, R.H., Cheers, S.J., Ellis, N., Jones, P.N., Koutsoukos, A.G., Moeseneder, C.H., Smith, G.P., Venables, W. and Wassenberg, T.J. 2008, *Recovery of seabed habitat from the impact of trawling in the far northern section of the Great Barrier Reef Marine Park: CSIRO final report to GBRMPA*, CSIRO Marine and Atmospheric Research, Cleveland, Queensland.
119. Poiner, I.R., Glaister, J., Pitcher, C.R., Burrridge, C.Y., Wassenberg, T.J., Gribble, N.A., Hill, B.J., Blaber, S.J.M., Milton, D., Brewer, D. and Ellis, N. 1998, *Environmental effects of prawn trawling in the far northern section of the Great Barrier Reef: 1991-96*, CSIRO Division of Marine Research, Cleveland.
120. Pitcher, C.R., Doherty, P., Arnold, P., Hooper, J., Gribble, N.A., Bartlett, C., Browne, M., Campbell, N., Cannard, T., Cappel, M., Carini, G., Chalmeres, S., Cheers, S., Chetwynd, D., Colegax, A., Coles, R., Cook, S., Davie, P., De'ath, G., Devereux, D., Done, B., Donovan, T., Ehrke, B., Ellis, N., Ericson, G., Fellegara, I., Forcey, K., Furey, M., Gledhill, D., Good, N., Gordon, S., Haywood, M., Hendricks, M., Jacobsen, I., Johnson, J., Jones, M., Kininmonth, S., Kistle, S., Last, P., Leite, A., Amrks, S., McLeod, I., Oczkowicz, S., Robinson, M., Rose, C., Seabright, D., Sheils, J., Sherlock, M., Skelton, P., Smith, D., Smith, G., Speare, P., Stowar, M., Strickland, C., Van der Geest, C., Venables, W., Walsh, C., Wassenberg, T.J., Welna, G. and Yearsley, G. 2007a, *Seabed biodiversity on the continental shelf of the Great Barrier Reef World Heritage Area. AIMS/CSIRO/QM/QDPI CRC Reef Research Task final report*, CSIRO Marine and Atmospheric Research, Hobart.
121. Dichmont, C.M., Pascoe, S.D., Jebreen, E.J., Pears, R.J., Brooks, K.J. and Perez, P. 2012, *Providing social science objectives and indicators to compare management options in the Queensland trawl planning process - Final Report (FRDC 2009/100)*, CSIRO Marine and Atmospheric Research, Hobart.
122. Morison, A.K. and Pears, R.J. 2012, *Assessment of the ecological vulnerability of the East Coast Otter Trawl Fishery to climate change: a brief synthesis of information and results of an expert workshop*, Great Barrier Reef Marine Park Authority, Townsville.
123. Department of Sustainability, Environment, Water, Population and Communities 2010, *Assessment of the East Coast Otter Trawl Fishery*, DSEWPac, Canberra.
124. Hockings, M. and Gilligan, B. 2009, *Assessment of management effectiveness for the 2009 Great Barrier Reef Outlook Report*, Great Barrier Reef Marine Park Authority, Townsville.
125. Pascoe, S., Dichmont, C.M., Brooks, K., Pears, R.J. and Jebreen, E. 2012, Management objectives of Queensland fisheries: putting the horse before the cart, *Marine Policy* 37: 115.
126. Dichmont, C.M., Pascoe, S., Jebreen, E., Pears, R.J., Brooks, K. and Perez, P. 2013, Choosing a fishery's governance structure using data poor methods, *Marine Policy* 37: 123.
127. Queensland Department of Employment, Economic Development and Innovation 2012, *East Coast Otter Trawl Fishery 2011: Progress against SEWPaC conditions and recommendations*, DEEDI, Brisbane.

128. Pitcher, C.R., Venables, B., Browne, M., Doherty, P. and De'ath, G. 2007b, *Indicators of protection levels for seabed habitats, species and assemblages on the continental shelf of the Great Barrier Reef World Heritage Area*, Reef and Rainforest Research Centre, Cairns.

129. Fernandes, L., Day, J., Lewis, A., Slegers, S., Kerrigan, B., Breen, D., Cameron, D., Jago, B., Hall, J., Lowe, D., Innes, J., Tanzer, J., Chadwick, V., Thompson, L., Gorman, K., Simmons, M., Barnett, B., Sampson, K., De'ath, G., Mapstone, B., Marsh, H., Possingham, H., Ball, I., Ward, T., Dobbs, K., Aumend, J., Slater, D. and Stapleton, L. 2005, Establishing representative no take areas in the Great Barrier Reef: Large scale implementation of theory on marine protected areas, *Conservation Biology* 19(6): 1733-1744.

130. Roy, D. and Jebreen, E.J. 2011, *Extension of Fisheries Research and Development Corporation funded research results on improved bycatch reduction devices to the Queensland East Coast Otter Trawl Fishery*, Fisheries Research and Development Corporation and Department of Employment, Economic Development and Innovation, Brisbane.

131. Great Barrier Reef Marine Park Authority 2012c, *Climate change adaptation: Outcomes from the Great Barrier Reef Climate Change Action Plan 2007-2012*, GBRMPA, Townsville.



An underwater photograph of a mangrove forest. The water is a deep teal color. In the foreground, several dark, vertical mangrove roots are visible. A small, bright yellow fish is swimming near the surface in the upper right quadrant. The background shows more roots and some green foliage at the water's edge.

## Chapter 10

---

## Resilience and risk



10.12 Summary of outcomes.....	10–21
References .....	10–25

## Table of contents

<b>10.1</b>	<b>Understanding ecosystem resilience ...</b>	<b>10–3</b>
10.1.1	Factors that affect ecosystem resilience.....	10–3
<b>10.2</b>	<b>Resilience of key values of biodiversity.....</b>	<b>10–4</b>
10.2.1	Coral reefs and corals.....	10–4
10.2.2	Seagrass meadows .....	10–4
10.2.3	Lagoonal habitats .....	10–4
10.2.4	Deep water and continental slope habitats .....	10–5
10.2.5	Inshore habitats .....	10–5
10.2.6	Bony fishes .....	10–5
10.2.7	Dugongs .....	10–5
10.2.8	Marine turtles .....	10–5
10.2.9	Sharks and rays.....	10–6
10.2.10	Seabirds.....	10–6
<b>10.3</b>	<b>Overall ecosystem resilience .....</b>	<b>10–6</b>
<b>10.4</b>	<b>Active restoration of ecosystem resilience .....</b>	<b>10–7</b>
<b>10.5</b>	<b>Social and cultural resilience .....</b>	<b>10–7</b>
<b>10.6</b>	<b>Assessing risks on the Region's values.....</b>	<b>10–8</b>
<b>10.7</b>	<b>Risks to biodiversity.....</b>	<b>10–9</b>
<b>10.8</b>	<b>Risks to geomorphological features... ..</b>	<b>10–14</b>
<b>10.9</b>	<b>Risks to Indigenous heritage values .. ..</b>	<b>10–15</b>
<b>10.10</b>	<b>Risks to historic heritage values.....</b>	<b>10–19</b>
<b>10.11</b>	<b>Summary of risks to matters of national environmental significance .....</b>	<b>10–20</b>

Cover page image: Rhizophora mangrove roots, Lizard Island



## Extract from Great Barrier Reef Region Strategic Assessment terms of reference

### 5. Projected condition of matters of national environmental significance

#### 5.1 Describe the projected condition of the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area, based on an evaluation of...:

.....

- d) an understanding of ecosystem resilience
- e) an assessment of overall risks to the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area.

## 10.1 Understanding ecosystem resilience

Ecosystem resilience refers to the capacity of an ecosystem to withstand or recover from disturbances and impacts, and maintain key functions without collapsing to a different state. Resilience is not about the single, static ecological state, but rather the capacity of an ever-changing, dynamic system to return to a healthy state.<sup>1,2,3</sup>

Tropical marine ecosystems such as the Great Barrier Reef are subject to a wide range of natural and human-related impacts, such as cyclones, crown-of-thorns starfish outbreaks and pollution, which may damage components of the system. Resilient ecosystems are able to resist or recover from those impacts, if given sufficient time. Understanding the capacity of the ecosystem to absorb or recover from these threats is crucial to the long-term protection of the Reef's values.<sup>4</sup>

The resilience of an ecosystem is determined by a range of variables. Loss of ecosystem resilience is rarely attributed to a single cause, but is typically the consequence of impacts from different activities and drivers, and their accumulation through time or space. Different impacts may combine or even exacerbate each other (synergise), so that the cumulative impacts may be far greater than any individual impact.<sup>5,6</sup> This has two important consequences: first, the need to manage all impacts to reduce cumulative impacts, and that reductions in one impact may reduce the effects of other impacts, potentially providing time to address less accessible impacts such as those caused by climate change.<sup>7,8</sup>

There is no comprehensive information on the resilience of the Great Barrier Reef Region (the Region), largely due to the vast extent and complexity of the ecosystem, and because resilience is a complex property that is difficult to measure. The following description of the current understanding of resilience is based on available information. Although many elements of the ecosystem probably remain resilient, the emergent picture is that the resilience of the ecosystem as a whole is being significantly and incrementally eroded, and that this erosion is increasing.

### 10.1.1 Factors that affect ecosystem resilience

The extensive body of work<sup>1,9,10,11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26</sup> on ecosystem resilience highlights a number of factors important in understanding the concept and its implications:

- Resilience includes the capacity of the system to either resist (absorb) an impact, or to recover from that impact.
- Although the resilience of the ecosystem is distinct from its health or state, they are related — a degraded system may have less capacity to recover if populations are too depleted to supply enough new recruits (for example, larvae or seeds).
- Systems with high levels of diversity, key functional species and a level of functional redundancy are likely to have greater resilience.<sup>12,27</sup> High levels of diversity provide more diverse capacities for adapting to or recovering from new threats. The key functional species play an important role in maintaining the ecological processes underlying resilience (for example herbivorous fishes on coral reefs).
- Networks of elements, for example habitats, may have greater resilience because the connectivity between elements enhances recovery after disturbances.
- A critical aspect of resilience are thresholds or tipping points, whereby an ecosystem that is subject to ongoing impacts may undergo sudden large changes in response to relatively small increases in impacts. For example, a reef subject to excess nutrients and sediments may retain high coral cover, but lose recruitment of new corals. If a storm reduces the coral cover, the lack of recruitment may mean the reef is not able to recover, and so it suddenly changes state. Such tipping points are intrinsically very difficult to predict, making them very difficult to avoid.
- The resilience of a system is strongly linked to the spatial and temporal scales being considered — an individual reef may have lost its resilience, but as part of a connected network, the overall system may

retain resilience and be able to maintain its functions. If sufficient reefs become degraded, the loss of resilience may transfer up to the larger scale, degrading the resilience of the overall system. Thus, a regional scale collapse may become detectable only gradually, as an emergent property of smaller scale changes. Changes like this are very susceptible to the 'shifting baselines' phenomenon (that is, changes occur gradually and piecemeal, and hence are not recognised and responded to).

- Resilience is a property of complex systems, and is itself a complex property, making it very difficult to measure or monitor directly (in contrast to monitoring the state of the system).
- Together, the difficulty of measuring resilience, the nature of tipping points and scale dependence means loss of resilience and environmental degradation are easy to deny and hard to respond to. Thus, managing for resilience requires a sufficiently precautionary approach. Resilience is especially important for risks that are difficult to predict and/or where there is insufficient knowledge of management responses.<sup>28</sup>
- The resilience of ecosystems is very strongly linked to the resilience of social and economic systems that depend on them, and vice versa. For example, fishing and tourism on the Great Barrier Reef depend closely on healthy and resilient ecosystems for their businesses, just as resilient industries have the capacity to operate in ways that support the health of the ecosystem. Also, the resilience of the Region's Indigenous and historic heritage values and community benefits are tightly linked to the resilience of its biodiversity.
- Building the resilience of the ecosystem, such as by reducing local and regional impacts, is a key strategy in addressing the effects of climate change on the ecosystem, but it will not be sufficient. Although it may buy time for the Reef, there is no doubt that direct actions to reduce global carbon emissions are essential for the long-term health of the Great Barrier Reef ecosystems.

## 10.2 Resilience of key values of biodiversity

### 10.2.1 Coral reefs and corals

Coral reefs, the iconic habitat of the Great Barrier Reef, are showing major declines in resilience. Naturally, coral reefs, and corals specifically, have an impressive ability to recover from disturbances such as cyclones, crown-of-thorns starfish outbreaks or coral bleaching. There is recent evidence for quite spectacular recovery of coral reefs at local scales,<sup>20</sup> and reefs in the northern part of the Reef have shown substantial recovery.<sup>29</sup>

However, emerging evidence is increasingly clear that, over the long-term and over a broad scale, coral reefs in the southern two-thirds of the Great Barrier Reef are being gradually but seriously damaged by disturbances and are not recovering sufficiently between disturbances, resulting in the overall long-term (decadal scale) decline in coral cover.<sup>15,29</sup> This loss of resilience is most profound on inshore reefs, where recent evidence suggests reefs are failing to recover from disturbances over the past few decades.<sup>30</sup> This is despite evidence that those reefs had maintained their resilience over many hundreds of years. The loss of resilience is strongly correlated in time with degraded water quality after European settlement.<sup>30</sup>

The causes of this loss of resilience are likely to include direct and indirect effects of a range of impacts, such as catchment run-off, crown-of-thorns starfish outbreaks, and impacts related to climate change, such as increased sea temperatures and ocean acidification, and are almost certainly exacerbated by the broadscale loss of abundance, leading to reduced reproduction and connectivity (see the demonstration case study on corals, (see Section 9.4). Significantly, it is likely that this loss of resilience reflects degradation of both resistance and recovery.

### 10.2.2 Seagrass meadows

Available evidence suggests a serious loss of resilience for many inshore seagrass meadow habitats, especially for the area south of the Daintree River. Not only has there been major loss of area and abundance of seagrasses in recent years; but low or negligible reproductive capacity (loss of seed banks within some meadows) and high tissue nutrient levels (indicating reduced growth due to eutrophic conditions) also suggest loss of resilience.<sup>31</sup> This is further exacerbated by a loss of species diversity and a dominance of rapid growing species over habitat-forming species of seagrass. The severity of the losses in abundance is likely to have led to a loss of connectivity between seagrass meadows and consequently a probable loss of genetic diversity. There is very little information about the resilience of deepwater seagrass meadows, although there is concern about the effects of recent extreme weather events.<sup>31</sup>

### 10.2.3 Lagoonal habitats

There is limited information on the resilience of many lagoonal habitats (such as shoals, *Halimeda* banks, sponge gardens, and muddy and sandy seafloor areas), although it is reasonable to assume that resilience varies considerably among habitat types. Some evidence can be gathered from Great Barrier Reef studies of the habitats' resistance to and recovery from trawling as a proxy for physical disturbance generally.<sup>32,33,34,35,36</sup> Many habitat elements, especially soft sediments, show fairly good resistance to, and recovery from damage. However, structural elements of some other habitat types, such as sponge gardens, are relatively fragile (low resistance) and very slow to recover, taking decades.<sup>35,36</sup> Even relatively deep areas may be strongly affected in extreme weather events: for example, during cyclone Yasi in 2011, the wreck of the SS *Yongala*, at more than 20 metres

deep, was all but stripped bare of a century of growth of sponges and sea fans. Importantly, these more fragile habitat patches may be disproportionately important in ecological terms, as critical links for cross-shelf connectivity of fish or other animal populations, making them in turn critical to the overall resilience of the Region.<sup>37</sup> Notwithstanding these lines of evidence for resilience to physical disturbance, there is very little knowledge of resilience to other impacts, such as excess nutrients related to run-off, herbicides, sedimentation and turbidity, or climate impacts such as increased temperature, acidification or changes to ocean currents.

#### **10.2.4 Deep water and continental slope habitats**

The deep areas of the Region are much less understood than the lagoonal areas, and there is little knowledge of their resilience. Evidence from studies in other parts of the world suggests that deepwater benthos is very slow growing and hence slow to recover from physical damage (decades). Many taxa are also relatively fragile and easily depleted by activities such as trawling.<sup>38,39,40,41</sup> Although the exposure to direct impacts (such as trawling) may be relatively limited for much of the continental slope, some deeper areas are intensively trawled (such as in the south-east of the Region). In the longer term, such deep water and continental slope habitats may be vulnerable to climate-related changes (such as increased temperature), changes to ocean currents (such as upwelling patterns) and ocean chemistry including acidification. Recent evidence suggests deepwater habitats may provide a refuge from disturbances acting on shallow water systems, thus potentially aiding the resilience and recovery of these systems.<sup>42</sup>

#### **10.2.5 Inshore habitats**

The inshore habitats of the Region, especially adjacent to the developed coast south of about Port Douglas, are subject to the most intensive and diverse impacts (see Chapter 5 and 6). This has resulted in more intense degradation in those areas, as well as undermining the resilience of many of the habitats.<sup>43</sup> However, the extent of habitat loss varies considerably between locations and habitat types. Inshore habitats are most affected by terrestrial run-off of nutrients, sediments and pesticides; are most directly affected by coastal development; and are subject to the most intense direct uses such as recreational and commercial fishing. Coastal, estuarine and inshore habitats provide a wide range of ecological and ecosystem services, such as providing nursery areas and supporting key stages in the life cycles of various commercially-harvested fish and crustacean species.<sup>43</sup> Many of those services depend critically on connectivity between coastal waterbodies and the sea, yet many of those connections have been seriously damaged or lost due to coastal development and intensive use of land and marine ecosystems. Further, many of the animals that use these inshore habitats are especially vulnerable to loss of these habitats and to the impacts that cause that loss (for example, dugongs feeding on inshore seagrass meadows).

#### **10.2.6 Bony fishes**

Evidence suggests some targeted fish species such as coral trout can recover relatively quickly after protection from fishing impacts,<sup>44,45</sup> largely due to their fast growth and high rates of reproduction. Other target fish species (for example flowery cod) have characteristics such as late maturity, aggregation spawning and long life spans (over 40 years) which make them more susceptible to depletion by fishing and slower to recover.<sup>46,47</sup> Additionally, recent evidence suggests resilience may be significantly eroded if the species is subject to intense and large-scale disturbance, such as the recent series of extreme weather events.<sup>48</sup> This erosion is likely to be exacerbated by cumulative impacts, especially climate change impacts,<sup>49,50,51,52</sup> and emphasises the need to manage fisheries to protect their resilience.

#### **10.2.7 Dugongs**

Some species of conservation concern have life cycles which make them less resilient to impacts. For example, dugongs are relatively slow growing, have low reproductive rates, have restrictive dietary requirements (almost exclusively seagrasses) and are highly vulnerable to a wide range of direct impacts (refer to the demonstration case study on dugong, (see Section 9.3). Historic depletion has strongly exacerbated this vulnerability, as has the species' dependence on inshore habitats where these impacts are most intense. In the inshore areas of the southern two-thirds of the Region, this low resilience is now further exacerbated by serious declines in seagrass habitats due to poor water quality and extreme weather events.<sup>53</sup> It is unlikely that dugong populations will be able to recover over reasonable time scales without very strong management intervention to improve water quality and seagrass habitat, while also reducing all other sources of mortality.<sup>54</sup>

#### **10.2.8 Marine turtles**

Marine turtles have life histories which result in relatively high vulnerability to many impacts.<sup>55</sup> In particular, reproduction is dependent on beach nesting, a vulnerable part of their life cycle, especially in areas of intensive coastal development.<sup>56</sup> Several species are also vulnerable as bycatch in fisheries, although recent management is proving effective at reducing this mortality.<sup>57</sup> Although some species have been seriously compromised by these and other impacts, green turtles are considered sufficiently abundant (at least in the southern Region) to be able to recover relatively well from population declines when major threats to their populations are removed or addressed.<sup>54,58</sup> This is important given the very high mortality rates from loss of seagrass meadows during extreme weather in recent years (see Section 7.1.3).

## 10.2.9 Sharks and rays

Most species of sharks and rays have life-histories which make them less resilient than many bony fishes.<sup>59</sup> Some species, especially those that inhabit inshore areas such as sawfishes and coral habitats such as grey reef and white tip reef sharks, have declined severely due to fishing and habitat loss; their low reproductive rates mean populations are not resilient in the face of these impacts.<sup>60</sup> Some highly-mobile species use inshore habitats at critical parts of their life cycle, making them vulnerable. However, other species of sharks are thought to be at lower risk because their growth rates are able to balance the effects of the impacts on their populations.<sup>59,62</sup>

## 10.2.10 Seabirds

Life-history traits of offshore and pelagic-foraging seabirds make them susceptible to a number of pressures. These traits include being long-lived; slow growth rates of young with high parental care; low reproductive output; high habitat and trophic specificity (most significantly, these species rely on a particular foraging behaviour that determines specificity in their diet and limits their ability to increase the rate of provision when food resources become scarce).<sup>63</sup> Seabirds have a major influence on island ecosystems, including the transfer of nutrients from pelagic and offshore areas to islands and reefs. Seabirds are also key upper trophic level predators in the marine ecosystem and their demographics and reproduction are strongly influenced by changing marine conditions.<sup>64</sup> Furthermore, seabirds are expected to be directly or indirectly impacted by a majority of the impacts identified in Table 10.3.<sup>65,66</sup>

## 10.3 Overall ecosystem resilience

While the Great Barrier Reef Region may be one of the most resilient tropical marine ecosystems in the world<sup>16,67</sup>, its resilience is being seriously, and increasingly rapidly, eroded. Due largely to the size and complexity of the ecosystem and because resilience is difficult to measure, there is no comprehensive information on the resilience of the Region. However, the available information is of concern: there is increasing evidence of loss of resistance and recovery capacity, although the extent of that loss varies considerably between ecosystem components (for example, dugong versus fish populations) and between localities (for example, inshore southern region compared to offshore northern region).

The loss of ecosystem resilience cannot be attributed to any single cause, but is almost certainly the consequence of cumulative impacts from all the different activities and drivers, and their accumulation through time. The loss of resilience is of serious concern, given the critical role of resilience in allowing the Region's ecosystem to resist the increasing effects of climate change-related impacts.

The integrity of the ecosystem as a whole is a key aspect of both its resilience and its outstanding universal value, yet that integrity is progressively undermined as increasing examples of degraded resilience accrue at smaller scales: that is, as increasing numbers of system sub-components lose their resilience, the likelihood increases of the overall system undergoing a critical change of state.<sup>23,24,68</sup>

It is of particular concern that several of the most significant impacts on ecosystem resilience, and those least effectively managed (see Chapter 8), operate at broad scales. For example, degraded water quality affects all the habitats and species in much of the southern region, so that the impacts interact across those different elements of biodiversity. Further, as the state of the system changes, attributes that confer resilience to a healthy ecosystem may come to undermine the resilience of the compromised system. For example, while the nature of the coral reefs on the Great Barrier Reef as a network of reefs distributed across the Region allows connectivity for corals and fishes, it also provides connectivity for pest species such as crown-of-thorns starfish.

System resilience also depends on the frequency of disturbances (and hence, the length of the intervening recovery periods) as this affects the extent of recovery between disturbances. If disturbances are too frequent, their impacts will accumulate through time. Two of the major causes of coral loss on the Great Barrier Reef, cyclones and crown-of-thorns starfish outbreaks, are being exacerbated by human activities. Cyclone intensity in the Region is expected to increase as a result of climate change,<sup>69,70,71,72,73</sup> and there is strong evidence that increased nutrients in catchment run-off is contributing to the increased frequency of crown-of-thorns starfish outbreaks.<sup>74</sup>

The emerging loss of ecosystem resilience is particularly critical in the context of the projected major increase in severe effects of climate change impacts. As these effects worsen, it is very likely that interactions between climate and other impacts will have increasingly serious consequences. Further, current evidence suggests climate change trajectories remain on course for increasingly serious impacts. Managing for resilience is most important in situations where there is uncertainty about risks and appropriate management responses<sup>28</sup> — the combined consequences of climate change and local and regional impacts on the Great Barrier Reef present such a situation. Maintaining the resilience of the Great Barrier Reef ecosystem will require major increases in effort to reduce local impacts and global climate change.



## 10.4 Active restoration of ecosystem resilience

Overall, major improvements are needed to restore and protect the resilience of the Great Barrier Reef. Previously, the approach to managing resilience has been to focus on supporting the natural resilience of the ecosystem and allowing it to restore its own processes and functions. However, available evidence strongly suggests this strategy has underestimated the extent of the cumulative impacts on the system, and that additional new and stronger approaches will be required.

This is especially true in the face of climate changes already occurring, and the further changes that will result without more effective mitigation of emissions.

Such approaches include more proactive intervention to restore ecosystem functions after disturbances, but there are several important cautions to be noted:

- It is critical that the causes of degradation, including climate change, are addressed in combination with interventions to restore functions. Without addressing the causes, restoration is unlikely to be effective in the long term, whereas combined actions may provide synergistic benefits.
- The costs of restoration are likely to be far greater than the costs of prevention.<sup>75</sup> The clear consequence is that major investment in resilience now is very likely to dramatically reduce future costs.
- At present there is limited capacity and knowledge for such restoration interventions, and little knowledge of the potential risks. Investment in significant research and development would be required to develop the expertise for such interventions in a timely and environmentally responsible manner.

Future attention could focus on major causes of ecosystem decline and on restoring the positive aspects of connectivity within the ecosystem, especially at the land–sea (coastal) interface, and on reducing the frequency and/or impact of major disturbances, wherever feasible. For example, one of the most direct paths to rebuilding coral populations and their dependent communities would be to suppress further crown-of-thorns starfish outbreaks by improving water quality and by developing more effective direct control measures.<sup>29,76</sup> Restoring the positive aspects may include coral transplantation at high value tourism sites.

Finally, active restoration of resilience will need to explicitly address the cumulative impacts on the Reef, and to do so in the context of predicted climate change. This requires:

- a broad portfolio approach, to comprehensively and effectively address all impacts. Such an approach will provide the most effective combined benefits for resilience, whereas only addressing one or two key impacts is not likely to be successful
- a major increase in management efforts across that broad portfolio
- consideration by management of how to best manage resilience while also preparing to manage a system that may well experience climate-driven transformational change in some or all of its elements.

## 10.5 Social and cultural resilience

Social resilience enables societies and individuals to adapt to changing circumstances by learning, storing knowledge and being creative in problem solving.<sup>77</sup> Indigenous systems of heritage conservation and management, including the continuation of cultural practices and retaining and creating traditional ecological knowledge, maintain cultural resilience. Understanding feedbacks between people and nature is critically important in protected area management, enabling appropriate decisions to be made about managing the Region's natural system, while at the same time maintaining options for current and future generations of Reef-dependent industries and communities.<sup>76</sup>

The condition and future of the Great Barrier Reef ecosystem is important for many people including Traditional Owners whose dynamic heritage integrates nature, society and culture. Reef-dependent tourism and commercial fishing are dominated by people who work in their industries because they feel strongly connected to the Reef.<sup>78</sup> Commercial fishing on the Great Barrier Reef is a lifestyle choice for most fishers, who have been in the industry for a long time and cannot imagine doing anything else.<sup>79</sup>

The resilience of the Great Barrier Reef ecosystem is closely linked to the resilience of the communities and industries of the Region, especially for the fishing and tourism industries. There has been a significant increase in available information in this area, and considerable effort to help Reef-dependent industries improve their resilience in the face of climate change.<sup>80</sup> Importantly, many of the impacts on the Reef are influenced strongly by social and economic drivers, such as population changes. Increasing our understanding of these mechanisms is important to our capacity to manage for resilience.

By understanding social resilience and Reef dependency, managers can better promote the Region as an integral part of community life. For example, community benefits derived from the Region, such as a sense of wellbeing associated with living in the Reef catchment, may inspire active stewardship which in turn helps reduce impacts such as habitat clearing and water quality decline.<sup>76</sup>

## 10.6 Assessing risks on the Region's values

To assess the risks to the Great Barrier Reef ecosystem posed by the 40 impacts set out in Chapter 6, the Australian Standard for Risk Assessment (AS/NZS ISO 31000:2009) was followed.

The likelihood and consequence of each predicted impact are ranked on the five-point scale set out in Table 10.1. An overall risk level for each impact over the next 25 years is determined, based on a combination of its likelihood and consequence (Table 10.2).

The assessment is based on the information previously presented in this report, including the current trends in drivers and activities (Chapter 5), the effects of the identified impacts (Chapter 6), the current state of the Region's values (Chapter 7), the effectiveness of current management (Chapters 8) and current resilience of the ecosystem (see above). For this reason, references have not been included in the following tables as they are cited in the chapters referred to above.

The assessments are presented in relation to the broad categories of values relevant to matters of national environmental significance outlined in Chapter 4, namely biodiversity (Section 10.7), geomorphological features (Section 10.8), Indigenous heritage values (Section 10.9) and historic heritage values (Section 10.10). The likely risks to future community benefits are not presented. Although community benefits are closely linked to biodiversity values, the risks to future community benefits are difficult to predict, particularly because societal attitudes, values and expectations (see 'shifting baselines' in Section 7.1) and the ways communities adapt in response to changes in the environment are constantly evolving. Only the impacts relevant to each broad category are included in the assessment.

The implications for each matter of national environmental significance is summarised in Section 10.11.

**Table 10.1 Gratings for likelihood and consequence**

A standard set of criteria allows the comparison of different types of threats within the one risk assessment, based on the likelihood and consequence of each threat. Adapted from the *Great Barrier Reef Outlook Report 2009*.

Likelihood	Expected frequency of a given threat	
Almost certain	Expected to occur more or less continuously throughout a year	
Likely	Not expected to be continuous but expected to occur one or more times in a year	
Possible	Not expected to occur annually but expected to occur within a 10-year period	
Unlikely	Not expected to occur in a 10-year period but expected to occur in a 100-year period	
Rare	Not expected to occur within the next 100 years	

Consequence	Extent of the impact based on current management	
	Broad scale	Local scale
Catastrophic	Impact is clearly affecting, or would clearly affect, the nature of the value over a wide area. Recovery periods greater than 20 years likely.	
Major	Impact is, or would be, significant at a wider level. Recovery periods of 10 to 20 years likely.	Impact is, or would be, extremely serious and possibly irreversible to the condition of a value. Condition of the affected value possibly irretrievably compromised.
Moderate	Impact is, or would be, present at a wider level. Recovery periods of 5 to 10 years likely.	Impact is, or would be, extremely serious to the condition of a value and possibly irreversible over a small area. Recovery periods of 10 to 20 years likely.
Minor	Impact is, or would be, not discernible at a wider level. Impact would not impair the overall condition of the value, including sensitive populations or communities, over a wider level.	Impact is, or would be, significant to the condition of value at a local level. Recovery periods of 5 to 10 years likely.
Insignificant	No impact or if impact is, or would be, present then only to the extent that it has no discernible effect on the overall condition of the value.	No impact or if impact is, or would be, present then only to the extent that it has no discernible effect on the overall condition of the value.

**Table 10.2 Combining likelihood and consequence for overall risk**





































Likelihood	Consequence				
	Insignificant	Minor	Moderate	Major	Catastrophic
Almost Certain	Low	Medium	High	Very high	Very high
Likely	Low	Medium	High	High	Very high
Possible	Low	Low	Medium	High	Very high
Unlikely	Low	Low	Low	Medium	High
Rare	Low	Low	Low	Medium	High

## 10.7 Risks to biodiversity








































The future risks of each of the 40 impacts identified in Chapter 6 to biodiversity in the Region are presented in Table 10.3.









































**Table 10.3 Projected risks to biodiversity in the Region over the next 25 years**































Summary of potential impacts	Risk			
	Low	Medium	High	Very high
<b>Climate change</b>				
<b>Altered ocean currents:</b> A major change in oceanic currents of the Great Barrier Reef over the next 25 years is unlikely. However, an increase in the speed and southern extent of the East Australian Current has already been observed. Major changes to ocean currents would have widespread and potentially irreversible implications for biodiversity. <b>Likelihood:</b> Unlikely ..... <b>Consequence:</b> Major				
<b>Cyclone activity:</b> Cyclones, a natural process in tropical regions, are predicted to become more severe, but less frequent, under current climate change scenarios. Severe cyclones have significant broadscale effects, with recovery times of at least 10 to 20 years. <b>Likelihood:</b> Possible..... <b>Consequence:</b> Major				
<b>Increased sea temperature:</b> The average annual sea surface temperature is almost certain to continue to rise over the next 25 years. Regardless of the variation in climate scenarios, it is predicted that by 2035 the average sea surface temperature will be warmer than any previously recorded. Higher temperatures will affect the nature of the entire ecosystem over a broad scale. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Catastrophic				
<b>Ocean acidification:</b> If trends in global levels in carbon dioxide concentrations continue, in 25 years concentrations could reach more than 450 parts per million. This is anticipated to have widespread effects on coral reef systems. Over the next 25 years, projections suggest the waters of the Great Barrier Reef are almost certainly going to become more acidic. Regardless of the rate of change, recent evidence indicates that even relatively small changes in ocean acidity reduce the capacity of corals and other calcifying organisms to build skeletons and shells, which in turn reduce their capacity to create habitat for reef biodiversity in general. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Catastrophic				
<b>Rising sea level:</b> Projected increases in sea level are almost certain over the next 25 years. This will have a noticeable effect on coastal and shallow water habitats and species at a broad scale. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Major				
<b>Catchment run-off</b>				
<b>Increased freshwater inflow:</b> Future climate projections for northern Queensland suggest an increase in rainfall variability, that is, wetter wet seasons and drier dry seasons. While average annual rainfall may not vary greatly, there may be one or more flood events a year with larger flood plumes. This would have significant consequences for inshore areas exposed to flood plumes and inshore productivity cycles. <b>Likelihood:</b> Likely ..... <b>Consequence:</b> Major				

Summary of potential impacts	Risk			
	Low	Medium	High	Very high
<p><b>Nutrients from catchment run-off:</b> Ongoing improvements in catchment management are likely to reduce nutrient loads in catchment run-off in the future. However, there is likely to be a significant lag time between changes in agricultural practice and measurable water quality improvements in the Region. It is projected that nutrients will continue to enter and remain in the Region over the next 25 years with potentially catastrophic consequences on biodiversity. Risks associated with changes to the Vegetation Management Act include the potential intensification of coastal agricultural development, with subsequent increases in pollutant loads: however, the full impact of these changes is unknown as the Queensland regulatory reform process is continuing.</p> <p><b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Major</p>				
<p><b>Outbreak of crown-of-thorns starfish:</b> Reductions in nutrient loads in catchment run-off may reduce the number of juvenile crown-of-thorns starfish that reach adulthood. Regardless, the presence of an active outbreak on the Reef at any given time is considered likely over the next 25 years, resulting in continued coral mortality. The cumulative effects of a range of impacts are severely compromising the ability of coral reefs to recover from outbreak events.</p> <p><b>Likelihood:</b> Likely..... <b>Consequence:</b> Catastrophic</p>				
<p><b>Pesticides from catchment run-off:</b> Ongoing use of pesticides in the catchment means the Region will almost certainly experience pesticides from catchment run-off over the next 25 years. Continued progress towards targets for reducing pesticides in catchment run-off may reduce the consequences in the Region itself, but pesticides are expected to have extremely serious consequences to some estuarine, seagrass and freshwater ecosystems, resulting in flow-on impacts to biodiversity in the Region.</p> <p><b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Moderate</p>				
<p><b>Sediments from catchment run-off:</b> Although improved practices and restoration of riparian vegetation in many catchment areas has reduced sediment load, sediment will continue to be transported to and remain in the Region. Similar to nutrients from catchment run-off, improvements in agricultural practices may take some time to become evident in water quality within the Region due to the lag time of sediments passing through the system and into sinks within the marine system. Projected increased rainfall variability may also contribute to sediment loads through the erosion of top soils during flood events. Consequences of sedimentation for marine life will depend on the concentration and duration of exposure, however there are likely to be major effects on biodiversity. Risks associated with changes to the Vegetation Management Act include the potential intensification of coastal agricultural development, with subsequent increases in pollutant loads: however, the full impact of these changes is unknown as the Queensland regulatory reform process is continuing.</p> <p><b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Major</p>				
<p><b>Urban discharge:</b> Projected increases in urban development will make discharges such as sewage and stormwater almost certain over the next 25 years. As regulations require sewage to be tertiary treated, sewage discharge is likely to have only minor effects.</p> <p><b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Minor</p>				
<p><b>Industrial discharge:</b> Discharges of wastewater from industrial development and mining that could have irreversible effects over a small area of the Region are possible, but are not expected to occur annually.</p> <p><b>Likelihood:</b> Possible..... <b>Consequence:</b> Moderate</p>				
<b>Degradation of coastal ecosystems</b>				
<p><b>Acid sulphate soils:</b> The projected continuation of coastal development makes the future risk of exposure of potential acid sulphate soils possible. Once disturbed, if not treated, acidic water and heavy metals would continue to be released during rain events over decades if not longer, causing effects that may be irreversible in a small area.</p> <p><b>Likelihood:</b> Possible..... <b>Consequence:</b> Moderate</p>				
<p><b>Artificial barriers to flow:</b> Artificial barriers in the catchment will continue to affect estuarine systems and connectivity.</p> <p><b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Moderate</p>				
<p><b>Atmospheric pollution:</b> Projected increases in urban and industrial development are likely to increase the local contribution of atmospheric pollution, including the potential for more frequent impacts from coal dust at loading ports. Within the next 25 years, atmospheric pollution may start to affect some values; however effects are expected to be only minor.</p> <p><b>Likelihood:</b> Possible..... <b>Consequence:</b> Minor</p>				



Summary of potential impacts	Risk			
	Low	Medium	High	Very high
<b>Coastal reclamation:</b> Projected increases in industrial and urban development make the risk of further coastal reclamation possible. While effects would be localised, the ecosystem in that area would be irretrievably compromised. <b>Likelihood:</b> Possible..... <b>Consequence:</b> Moderate				
<b>Light impacts (artificial):</b> Growth in shipping and urban and industrial development is likely to increase the amount of artificial light. The main known risk of artificial light is its effect on turtle hatchlings' orientation. Other minor effects include the orientation of pelagic species around vessel lights. <b>Likelihood:</b> Likely..... <b>Consequence:</b> Minor				
<b>Modifying supporting terrestrial habitats:</b> An expected increase in and intensification of grazing and projected growth in urban and industrial development makes the likelihood of modifying supporting terrestrial habitats almost certain. However, understanding has increased about the need to ensure protection of coastal ecosystems and their ecosystem services. The consequence to the Region's values is likely to be major over a broad scale. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Major				
<b>Direct use</b>				
<b>Dredging:</b> Continued development of new ports and port expansions would require capital and ongoing maintenance dredging. While maintenance dredging is expected to occur at least one or more times in a year, capital dredging is not expected to occur annually. Whereas the consequence of dredging on a broad scale is considered minor, the consequence for biodiversity within the footprint of the dredging site would be serious and possibly irreversible and pose a high risk at a very local scale. <b>Likelihood:</b> Likely..... <b>Consequence:</b> Minor				
<b>Dumping and resuspension of dredge material:</b> The dumping of dredge material is not continuous; however the frequency of dumping and resuspension of dredge material (from both capital and maintenance dredging) is likely to increase with continued development of new ports and port expansions. The resuspension of significant volumes of sediment could affect the condition of values over a broad scale adding further pressure to already declining inshore ecosystems. <b>Likelihood:</b> Likely..... <b>Consequence:</b> Moderate				
<b>Exotic species and diseases:</b> Despite technological improvements for better detection, projected increases in shipping makes the transport of exotic species and diseases possible. The consequence would depend on the species or disease but is likely to be serious in a small area such as a marina or port. <b>Likelihood:</b> Possible..... <b>Consequence:</b> Moderate				
<b>Extraction — death of discarded species (species of conservation concern):</b> Bycatch reduction devices have significantly reduced the incidental catch of turtles in the trawl fishery. Death of discarded and incidentally caught species of conservation concern across all fisheries and the Queensland Shark Control Program is almost certain, with major consequences for their populations. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Major				
<b>Extraction — death of discarded species (species not of conservation concern):</b> The discard of non-retained catch from fishing activities and the Queensland Shark Control Program are predicted to occur continuously throughout the next 25 years with broadscale consequences for populations of species commonly caught. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Moderate				
<b>Extraction — fishing spawning aggregations:</b> While a number of fish spawning aggregations are currently protected, some fishing effort targets unprotected aggregations. Targeting spawning aggregations can have implications for future population sizes of the species. <b>Likelihood:</b> Likely..... <b>Consequence:</b> Moderate				
<b>Extraction — herbivores (dugong and turtle):</b> Traditional hunting of marine turtles and dugongs is currently managed in a number of areas under Traditional Use of Marine Resources Agreements, and there is the aim of implementing more agreements in the next 25 years. Although not continuous through the year, traditional hunting is likely to occur several times a year with potential effects at a small scale. <b>Likelihood:</b> Likely..... <b>Consequence:</b> Minor				

Summary of potential impacts	Risk			
	Low	Medium	High	Very high
<b>Extraction — herbivores (take of herbivorous fish):</b> Current take of herbivorous fish is low and is unlikely to become very common in the next 25 years. The consequence of removing herbivorous fish from reef ecosystems would be major if extraction was sufficient to cause an ecological phase shift from a coral-dominated system, especially given the predicted effects of climate change. <b>Likelihood:</b> Unlikely..... <b>Consequence:</b> Major				
<b>Extraction — lower order predators:</b> Trends in fishing effort are predicted to remain stable with effects not discernible at a broad scale. There could be greater consequences for slow-breeding species. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Minor				
<b>Extraction — lower trophic orders:</b> Commercial, recreational and traditional fisheries that extract lower trophic orders are projected to continue throughout the next 25 years, with the potential for the trawl fishery effort to increase under current management arrangements. The resilience and biology of these species generally allows them to be sustainably extracted if appropriately managed. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Minor				
<b>Extraction — top order predators:</b> Some shark species extracted through the East Coast Inshore Fin Fish Fishery have life history traits that lend themselves to sustainable exploitation. Other top predators are slow breeding and extraction is likely to have at least moderate consequences. The network of no-take zones has already benefited populations of some predators. However, for larger, more mobile predators, benefits are limited. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Moderate				
<b>Illegal fishing and poaching:</b> Declining global fish stocks are likely to increase the demand on Australian fisheries. This, in turn, will increase the incentive for illegal foreign and domestic fishing activity. The consequence is likely to be major at a broad scale. Trends in illegal fishing and poaching are poorly known, but increasing illegal activity could have major consequences, particularly for sensitive areas and species. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Major				
<b>Marine debris:</b> Ocean currents transport debris around the world's oceans making the Reef vulnerable to debris from both local and more distant sources. Given the rapid increase in plastic production globally, the longevity of this material and the disposable nature of plastic items, plastic marine debris is likely to persist over the next 25 years and to be present at a broad scale within the Region. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Moderate				
<b>Noise pollution:</b> Projected increases in shipping and the continuation of increases in port development and recreational boat ownership mean underwater man-made noise is likely to be continuous over the next 25 years. Little is known about the effects of noise on the Region's species but evidence from elsewhere indicates that effects can be combined with serious consequences close to some sources. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Moderate				
<b>Outbreak of disease:</b> The causes of disease are difficult to ascertain but are likely to be varied. For example, outbreaks of coral disease have been linked to increased sea temperature, making further outbreaks possible. Consequences will vary depending on the disease and duration of outbreak but could have moderate effects at a broad scale. <b>Likelihood:</b> Possible..... <b>Consequence:</b> Moderate				
<b>Outbreak or bloom of other species:</b> Changes in ecological processes as a result of other impacts may cause population explosions of some species. Considering outbreaks and blooms to date, the risk would be significant to a sensitive population or community at a local scale. However, there is a high level of uncertainty. <b>Likelihood:</b> Likely..... <b>Consequence:</b> Minor				
<b>Physical damage — fishing:</b> Current levels of trawling activity pose low risk to shallow (<90m) habitats at a Reef-wide scale, given existing protection through zoning, but local effects may be higher in intensely trawled areas. Consequences could increase if trawl fishing effort increases under more favourable economic conditions. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Minor				

Summary of potential impacts	Risk			
	Low	Medium	High	Very high
<p><b>Physical damage — other:</b> If recreational vessel ownership continues to increase without a corresponding increase in supporting infrastructure and education, it is likely anchor damage will increase. Over the next 25 years, there is likely to be damage from anchors, small vessel groundings, diving and snorkelling throughout the year. The impact could be significant at the areas of impact.</p> <p><b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Minor</p>				
<p><b>Physical damage — ship grounding:</b> Despite projected increases in shipping and reports of skipper fatigue, it is considered that current management of shipping, including the vessel tracking service, significantly reduces the risk of groundings. They are, therefore, not predicted to occur every year but possibly once in 10 years. Groundings can have severe impacts on biodiversity at the site with long recovery periods, and long-term and broadscale effects due to dispersal of anti-fouling paint.</p> <p><b>Likelihood:</b> Possible..... <b>Consequence:</b> Moderate</p>				
<p><b>Spill — large chemical:</b> Although a large chemical spill is unlikely, the effects on biodiversity could be extremely serious and possibly irreversible at a local scale. Consequences would vary depending on the type and amount of spill and are considered major given current management and response plans.</p> <p><b>Likelihood:</b> Unlikely ..... <b>Consequence:</b> Major</p>				
<p><b>Spill — large oil:</b> While shipping is projected to increase, recent improvements in management make the potential for a large oil spill unlikely. The physical smothering of plants and animals, combined with oil toxicity and its chemical reactions with water, mean a large spill is likely to have serious and persistent effects for several years.</p> <p><b>Likelihood:</b> Unlikely ..... <b>Consequence:</b> Major</p>				
<p><b>Spill — small chemical and oil:</b> Small chemical and oil spills are likely to occur frequently in the Region. Projected increases in the number of ships and other vessels are likely to increase the risk in the next 25 years. Effects on sensitive marine life could be significant in the area of the spill, with consequences depending on size and type of spill.</p> <p><b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Minor</p>				
<p><b>Vessel strike on wildlife:</b> Continuing growth in shipping and recreational boating increases the potential for vessel strikes on wildlife. Surface-breathing animals are most at risk but the impact would not be discernible at the ecosystem level.</p> <p><b>Likelihood:</b> Likely ..... <b>Consequence:</b> Minor</p>				
<p><b>Waste discharge from a vessel:</b> Increases in vessel traffic will mean there is likely to be more vessel-based waste discharge over the next 25 years. Effects on biodiversity are anticipated to be minor under current management arrangements.</p> <p><b>Likelihood:</b> Almost certain ..... <b>Consequence:</b> Minor</p>				
<p><b>Wildlife disturbance:</b> Projected increases in population and a continuation of current increases in recreational vessel ownership, particularly in southern and central areas, is likely to lead to an increase in disturbance of wildlife from the presence of boats, snorkelling and diving activities and access to islands. The increase may cause some localised effects, for example on nesting seabirds.</p> <p><b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Minor</p>				

























## 10.8 Risks to geomorphological features

The future risks of relevant impacts to geomorphological features in the Region (see Chapter 6) are presented in Table 10.4.

**Table 10.4 Projected risks to geomorphological features in the Region over the next 25 years**

Summary of potential impacts	Risk			
	Low	Medium	High	Very high
<b>Climate change</b>				
<b>Altered ocean currents:</b> Altered ocean currents could influence the sediment transport pathways with flow-on effects to geomorphological features. Increased velocity or change in direction of currents could alter erosion processes on islands, shorelines and river deltas. Changes to tidal jets that deliver nutrients from deeper waters will control the location of <i>Halimeda</i> banks. <b>Likelihood:</b> Unlikely..... <b>Consequence:</b> Minor	■	■	■	■
<b>Cyclone activity:</b> Cyclone-induced damage is of concern to coral reefs as a geomorphological feature. Increased severity of cyclones reduces the resilience of corals and affects their capacity to recover from cumulative impacts. Cyclones are also likely to affect islands, shorelines and river deltas. <b>Likelihood:</b> Possible..... <b>Consequence:</b> Moderate	■	■	■	■
<b>Increased sea temperature:</b> Increased sea temperatures are predicted to be of minor consequence to the affected geomorphological features, such as coral reefs, seagrass and <i>Halimeda</i> banks. Reef building may slow or halt and erosive processes may begin. Loss of seagrass may lead to increased sediment movement. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Minor	■	■	■	■
<b>Ocean acidification:</b> Ocean acidification is likely to affect the geomorphological features reliant upon calcifying species, such as coral reefs and <i>Halimeda</i> banks. Erosional processes may dominate, and features that are protected by reefs may be indirectly impacted such as shorelines. Sediments may be produced at lower rates and with less durability, affecting Reef islands. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Moderate	■	■	■	■
<b>Rising sea level:</b> Rising sea level could potentially affect islands, shorelines and coral reefs that are unable to grow fast enough to maintain their shallow depth. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Insignificant	■	■	■	■
<b>Catchment run-off</b>				
<b>Increased freshwater inflow:</b> Flood events have the potential to rearrange river deltas and affect the health of reef-building corals and seagrass meadows. <b>Likelihood:</b> Likely..... <b>Consequence:</b> Minor	■	■	■	■
<b>Nutrients from catchment run-off:</b> Increased nutrients from catchment run-off will have minor effects on some geomorphological features, such as coral reefs and <i>Halimeda</i> banks. Increased nutrients can promote macroalgae growth on coral reef features which can, in turn, affect, slow or halt the reef-building process. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Minor	■	■	■	■
<b>Outbreak of crown-of-thorns starfish:</b> Coral mortality caused by a crown-of-thorns starfish outbreak will have an effect on coral reef features, halting growth and promoting erosional processes. <b>Likelihood:</b> Possible..... <b>Consequence:</b> Minor	■	■	■	■
<b>Sediments from catchment run-off:</b> Increased sediment will have an effect on geomorphological features reliant upon photosynthetic species such as coral reefs and seagrass meadows. It could also infill palaeochannels, burying the feature. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Moderate	■	■	■	■
















































Summary of potential impacts	Risk			
	Low	Medium	High	Very high
<b>Degradation of coastal ecosystems</b>				
<b>Coastal reclamation:</b> Reclaimed areas experience changes in sediment composition and can alter ocean currents in the surrounding area, which affect sediment transport. There is also potential that geomorphological features in reclaimed areas are directly affected. <b>Likelihood:</b> Possible..... <b>Consequence:</b> Moderate				
<b>Direct use</b>				
<b>Dredging:</b> Dredging can result in direct effects on geomorphological features at the site of dredging. It can also have flow-on effects to features and geomorphological processes by changing sediment transport pathways through dredged channels. <b>Likelihood:</b> Likely..... <b>Consequence:</b> Minor				
<b>Dumping and resuspension of dredge material:</b> Dumping and resuspension of dredge material can affect geomorphological features such as coral reefs by restricting light and smothering them. Sediment may also settle in palaeochannels. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Minor				
<b>Physical damage — fishing:</b> Little is known of the current impacts of deep-sea trawling on deeper water geological features, but the consequences could be moderate. <b>Likelihood:</b> Possible..... <b>Consequence:</b> Moderate				
<b>Physical damage — other:</b> Anchors can cause physical damage to shallow water geomorphological features at the site of contact, with potentially long-term consequences on their shape and function. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Minor				
<b>Physical damage – ship grounding:</b> The grounding of a large vessel has the potential to damage geomorphological features at the site with long-term effects on the function and shape of the feature. Given the scale of the potential impact relative to the size of most geomorphological features, the overall consequence would be minor. <b>Likelihood:</b> Possible..... <b>Consequence:</b> Minor				

















































## 10.9 Risks to Indigenous heritage values

The future risks of each of the 40 impacts identified in Chapter 6 to Indigenous heritage values in the Region are presented in Table 10.5.

**Table 10.5 Projected risks to Indigenous heritage values in the Region over the next 25 years**

Summary of potential impacts	Risk			
	Low	Medium	High	Very high
<b>Climate change</b>				
<b>Altered ocean currents:</b> Any effects on biodiversity will have direct effects on cultural values. Many species are totems for Traditional Owners and have cultural significance for customary practice, lore, story and songlines. <b>Likelihood:</b> Unlikely ..... <b>Consequence:</b> Major				
<b>Cyclone activity:</b> Severe cyclones can affect culturally important sites (including sacred sites) and places of historic Indigenous significance. <b>Likelihood:</b> Possible ..... <b>Consequence:</b> Major				

Summary of potential impacts	Risk			
	Low	Medium	High	Very high
<b>Increased sea temperature:</b> The effect of increased sea temperature on the entire ecosystem over a broad scale will also have a major effect on Indigenous cultural values. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Major				
<b>Ocean acidification:</b> The effects on coral reefs and related ecosystems from ocean acidification will change the environment and will consequently affect Indigenous heritage values. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Major				
<b>Rising sea level:</b> Rising sea level could affect coastal and shallow water historical cultural sites, as well as cause changes to custom. Loss of access to fish traps, burial sites (which may be in coastal sand dunes), or rock art located in beach caves will have adverse consequences to cultural practices. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Moderate				
<b>Catchment run-off</b>				
<b>Increased freshwater inflow:</b> Increased flood events are likely to affect seagrass meadows, driving animals such as turtle and dugong to leave areas in search of food. These movements would have significant localised effects on cultural values. This is in addition to the changes in the environment that occur with major flooding events. <b>Likelihood:</b> Likely..... <b>Consequence:</b> Major				
<b>Nutrients from catchment run-off:</b> Any impacts that affect the Region's water quality, and consequently marine plants and animals, will have an impact on Indigenous heritage values. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Moderate				
<b>Outbreak of crown-of-thorns starfish:</b> Little is known of the potential effects of outbreaks of crown-of-thorns starfish to Indigenous cultural values, but they are likely to be similar to those on biodiversity. <b>Likelihood:</b> Possible..... <b>Consequence:</b> Moderate				
<b>Pesticides from catchment run-off:</b> Any impacts that affect the Region's water quality, and consequently marine plants and animals, will have an impact on Indigenous heritage values. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Minor				
<b>Sediments from catchment run-off:</b> Any impacts that affect the Region's water quality, and consequently marine plants and animals, will have an impact on Indigenous heritage values. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Moderate				
<b>Urban and industrial discharge:</b> Urban and industrial discharge is already having an adverse effect on cultural practices. High concentrations of heavy metals due to polluted water are found in the liver of culturally significant species such as dugong and turtle. Any increase in discharge will continue to place cultural values and practices at further risk. <b>Likelihood:</b> Likely..... <b>Consequence:</b> Moderate				
<b>Degradation of coastal ecosystems</b>				
<b>Acid sulphate soils:</b> Localised effects of run-off from exposed acid sulphate soils can have significant effects on Traditional Owner cultural values. Impacts on fish and other marine resources from acid water and heavy metals would impact on the subsistence lifestyles that many Traditional Owners still practice today. <b>Likelihood:</b> Possible..... <b>Consequence:</b> Moderate				
<b>Artificial barriers to flow:</b> Change caused by construction of artificial barriers continues to alter and disrupt cultural connections for Traditional Owners. Physical changes to the environment not only disrupt species composition and abundance, but can affect cultural sites (e.g. burial sites, birthing sites). <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Moderate				
<b>Atmospheric pollution:</b> Atmospheric pollution may have minor effects on cultural values overall, but could have more significant effects in localised areas. <b>Likelihood:</b> Possible..... <b>Consequence:</b> Minor				

Summary of potential impacts	Risk			
	Low	Medium	High	Very high
<b>Coastal reclamation:</b> Without adequate consultation with Traditional Owners, reclamation on culturally significant sites would be possible and the values would be irretrievably compromised. <b>Likelihood:</b> Possible..... <b>Consequence:</b> Major				
<b>Light impacts (artificial):</b> Artificial lighting can reduce survival rates of turtle hatchlings which could have flow-on impacts on cultural values such as traditional hunting. <b>Likelihood:</b> Likely..... <b>Consequence:</b> Minor				
<b>Modifying supporting terrestrial habitats:</b> Even relatively small changes to land and seascapes have very significant consequences for cultural values. Cultural observances, customs, storylines and songlines can be lost by changes to terrestrial habitats. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Major				
<b>Direct use</b>				
<b>Dredging:</b> Dredging near coastal islands and in port areas can disturb culturally significant sites. The disturbance of burial sites or sacred sites has catastrophic effects on Indigenous cultural values. <b>Likelihood:</b> Likely..... <b>Consequence:</b> Major				
<b>Dumping and resuspension of dredge material:</b> Dumping or resuspension of dredge material in areas with undiscovered heritage sites and features is possible but would have only minor consequences. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Moderate				
<b>Exotic species and diseases:</b> The consequence would depend on the species or disease but could have serious effects to Indigenous heritage values in the local area. <b>Likelihood:</b> Possible..... <b>Consequence:</b> Moderate				
<b>Extraction—death of discarded species:</b> Many discarded species have a cultural significance to Traditional Owners as either a food source, totem or for customary practice. Any decrease in species populations has a significant effect for Traditional Owners. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Major				
<b>Extraction—fishing spawning aggregations:</b> Changes to population sizes of species targeted by fishing spawning aggregations would have subsequent effects on Indigenous heritage values connected to those species. <b>Likelihood:</b> Likely..... <b>Consequence:</b> Moderate				
<b>Extraction—herbivores:</b> Traditional hunting for turtle and dugong is likely to continue and has a positive impact on Indigenous heritage values. <b>Likelihood:</b> Likely..... <b>Consequence:</b> (Positive)				
<b>Extraction—lower order predators:</b> Localised effects from the extraction of lower order predators by commercial and recreational fishing can cause changes to customary practice if Traditional Owners have to fish or collect in non-traditional areas. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Moderate				
<b>Extraction—lower trophic orders:</b> Localised effects from the extraction of lower trophic orders by commercial and recreational fishing extraction can cause changes to customary practice if Traditional Owners have to fish or collect in non-traditional areas. <b>Likelihood:</b> Likely..... <b>Consequence:</b> Moderate				
<b>Extraction—top order predators:</b> Top order predators are totems for many Traditional Owners. The exploitation of these animals and localised impacts on populations will affect the cultural values of Traditional Owners with sea country estates. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Moderate				
<b>Illegal fishing and poaching:</b> Changes in species numbers and biodiversity directly affect Traditional Owners' ability to practice customary lore, use their cultural tools and technology, and follow cultural observances. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Major				

Summary of potential impacts	Risk			
	Low	Medium	High	Very high
<b>Marine debris:</b> As well as entangling and killing culturally important marine animals and birds, marine debris washes up in culturally important and sacred sites. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Moderate				
<b>Noise pollution:</b> Little is known of the potential effects of increased noise on Indigenous cultural values, but they are likely to be similar to those on biodiversity. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Minor				
<b>Outbreak of disease:</b> Outbreaks of disease such as coral disease and fibropapilloma in turtles can affect cultural practices, customs and lore. Outbreaks that may seem moderate at a broad scale could have significant impacts at a smaller, more local level. <b>Likelihood:</b> Possible..... <b>Consequence:</b> Moderate				
<b>Outbreak or bloom of other species:</b> Little is known of the potential effects of outbreaks or blooms of other species to Indigenous cultural values, but they are likely to be similar to those on biodiversity. <b>Likelihood:</b> Likely..... <b>Consequence:</b> Minor				
<b>Physical damage — fishing:</b> Any damage caused to culturally significant sites would seriously affect Indigenous heritage values. Customs and story and songlines for those damaged areas would be lost or have to change. <b>Likelihood:</b> Possible..... <b>Consequence:</b> Moderate				
<b>Physical damage — other:</b> Culturally significant sites (including sacred sites, burial sites and sites that have storylines associated with them) can be affected by damage to the seafloor and its habitats. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Minor				
<b>Physical damage — ship grounding:</b> Many reefs have strong cultural value to Traditional Owners. Song and storylines are connected to them and in some cases they are sacred sites. The destruction and damage caused by ship grounding would be major and long term. <b>Likelihood:</b> Possible..... <b>Consequence:</b> Moderate				
<b>Spill — large chemical:</b> A large chemical spill that affects biodiversity would have a flow-on effect to the cultural values of Traditional Owners. Localised extinctions could have extremely serious and possibly irreversible effects on cultural practice, observances, story and songlines and places of cultural significance at a local scale. <b>Likelihood:</b> Unlikely..... <b>Consequence:</b> Major				
<b>Spill — large oil:</b> Impacts on animals and land and seascapes from a large oil spill would have a similarly negative effect on Indigenous heritage values. <b>Likelihood:</b> Unlikely..... <b>Consequence:</b> Major				
<b>Spill — small chemical and oil:</b> This localised effect could have significant impacts on Indigenous heritage values for those connected to that sea country. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Minor				
<b>Vessel strike on wildlife:</b> As the species most affected are culturally significant, the impact on cultural values is significant at a local scale. <b>Likelihood:</b> Likely..... <b>Consequence:</b> Minor				
<b>Waste discharge from a vessel:</b> The minor effects of waste discharge from a vessel on biodiversity will have flow-on effects on Indigenous heritage values. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Minor				
<b>Wildlife disturbance:</b> Changes to animal behaviour caused by the presence of boats or people can change the nature of customary practice, affect sites of cultural importance and change storylines. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Moderate				















































## 10.10 Risks to historic heritage values

The future risks of relevant impacts to historic heritage values in the Region (see Chapter 6) are presented in Table 10.6.

**Table 10.6 Projected risks to historic heritage values in the Region over the next 25 years**

The risk assessment applies to buildings and structures of historic heritage value.

Summary of potential impacts and their risk parameters	Risk			
	Low	Medium	High	Very high
<b>Climate change</b>				
<b>Altered ocean currents:</b> A major change in ocean currents is likely to have only minor effects on heritage values such as shipwrecks. <b>Likelihood:</b> Possible..... <b>Consequence:</b> Minor				
<b>Cyclone activity:</b> Severe cyclones can have major impacts on heritage sites and features such as shipwrecks, lighthouses and World War II sites. The severity of cyclones is likely to increase in the future. <b>Likelihood:</b> Possible..... <b>Consequence:</b> Major				
<b>Increased sea temperature:</b> Increased sea temperatures could accelerate the natural degradation of historic heritage sites. <b>Likelihood:</b> Possible..... <b>Consequence:</b> Minor				
<b>Ocean acidification:</b> Ocean acidification could have an effect on shipwrecks, but it is likely to be insignificant. <b>Likelihood:</b> Almost certain..... <b>Consequence:</b> Insignificant				
<b>Rising sea level:</b> Rising sea level is likely to have some minor effects on coastal and shallow water historic heritage sites. <b>Likelihood:</b> Likely..... <b>Consequence:</b> Minor				
<b>Degradation of coastal ecosystems</b>				
<b>Atmospheric pollution:</b> Atmospheric pollution may have minor effects on heritage such as lighthouses and World War II sites. <b>Likelihood:</b> Possible..... <b>Consequence:</b> Minor				
<b>Coastal reclamation:</b> Coastal reclamation on or close to a heritage site is not likely to happen in a 100-year period. If it were to happen, for example on an undeclared heritage site, it would have major consequences. <b>Likelihood:</b> Rare..... <b>Consequence:</b> Major				
<b>Modifying supporting terrestrial habitats:</b> Modifying terrestrial habitats could affect undiscovered heritage sites. <b>Likelihood:</b> Possible..... <b>Consequence:</b> Moderate				
<b>Direct use</b>				
<b>Dredging:</b> Dredging in areas with undiscovered heritage sites and features is unlikely but would have major consequences. <b>Likelihood:</b> Unlikely..... <b>Consequence:</b> Major				
<b>Dumping and resuspension of dredge material:</b> Dumping of dredge material in areas with undiscovered heritage sites and features is possible but would have only minor consequences. <b>Likelihood:</b> Possible..... <b>Consequence:</b> Minor				
<b>Marine debris:</b> Marine debris such as discarded fishing nets can potentially become entangled on submerged historic sites, potentially degrading their heritage value. <b>Likelihood:</b> Rare..... <b>Consequence:</b> Minor				

Summary of potential impacts and their risk parameters	Risk			
	Low	Medium	High	Very high
<b>Physical damage — fishing:</b> It is possible for trawlers to accidentally damage undiscovered heritage sites and features, which could have serious effects for the site or feature. <b>Likelihood:</b> Possible..... <b>Consequence:</b> Moderate				
<b>Physical damage — other:</b> Small vessel groundings and anchor damage to heritage sites or features is possible over the next 25 years, with moderate consequences. <b>Likelihood:</b> Possible..... <b>Consequence:</b> Moderate				
<b>Physical damage — ship grounding:</b> A ship grounding in an area with a heritage site is not likely to happen in a 100-year period but would have major consequences. <b>Likelihood:</b> Rare ..... <b>Consequence:</b> Major				
<b>Spill — large chemical:</b> A large chemical spill would present a risk to heritage sites and features if the spill was close to the site. <b>Likelihood:</b> Unlikely..... <b>Consequence:</b> Moderate				
<b>Spill — large oil:</b> It is not likely in the next 25 years that an oil spill will have an effect on heritage sites or features. <b>Likelihood:</b> Rare ..... <b>Consequence:</b> Insignificant				
<b>Spill — small chemical and oil:</b> The risk of small chemical and oil spills affecting heritage sites is considered insignificant. <b>Likelihood:</b> Rare ..... <b>Consequence:</b> Insignificant				
<b>Waste discharge from a vessel:</b> Waste from vessels could possibly affect historic heritage sites over the next 25 years; however the effects would not be discernible. <b>Likelihood:</b> Possible..... <b>Consequence:</b> Insignificant				

## 10.11 Summary of risks to matters of national environmental significance

A summary of the likely future risks affecting each matter of national environmental significance is provided in Table 10.7.

**Table 10.7 Summary of projected risks to matters of national environmental significance in the Region**

Matter of national environmental significance	Implications of highest risk impacts
World heritage properties	The implications for the Great Barrier Reef World Heritage Area are the same as for the Marine Park. In addition, risks that occur in the World Heritage Area but outside the Marine Park (such as Queensland islands and internal waters and port areas) will have implications on the property's resilience, outstanding universal value and integrity. The Region's ecological and biological processes; natural beauty and phenomenon; and habitats for conservation of biodiversity are dependent on the condition of its biodiversity values. Therefore projected high and very high risks for biodiversity values will present an equivalent for the attributes underpinning the property's outstanding universal value. The sources of these high and very high risks continue to be climate change, catchment run-off, degradation of coastal ecosystems and direct use. While overall risk to attributes underpinning the major stages of Earth's evolutionary history are relatively low, projected increases in the intensity of cyclones, increased sediment, dredging and dumping of dredge material, physical impacts from fishing and ocean acidification are likely to pose the greatest risks. In addition, the risks identified would likely affect the property's integrity as its overall condition declines.
Great Barrier Reef Marine Park	About three-quarters of the major to catastrophic risks assessed as almost certain to affect the Marine Park's values over the next 25 years originate outside the Region, including impacts on a global scale (for example those related to climate change) and those originating in the catchment (for example from catchment run-off and the degradation of coastal ecosystems). While management of risks that originate within the Marine Park may help respond to impacts and restore resilience, the future health of the values relevant to the Marine Park depends to a large

Matter of national environmental significance	Implications of highest risk impacts
	degree on avoiding and mitigating risks arising beyond its boundaries. Some of the most significant risks arising from direct use— illegal fishing and poaching, and death of discarded species — have implications for the population health of target species and recovery of species of conservation concern. Other very high risks — such as dumping and resuspension of dredge material and outbreak of crown-of-thorns starfish — affect keystone species such as corals, which support a variety of life in the Region.
National heritage places	The implications for Great Barrier Reef as a national heritage place are the same as for the Great Barrier Reef World Heritage Area.
Commonwealth marine area	The implications for the Commonwealth marine area are the same as for the Marine Park.
Listed migratory and listed threatened species	Many listed migratory and listed threatened species rely on coastal and island habitats for feeding and breeding (e.g. marine turtles, dugong, some sharks, inshore dolphins, sawfish, some seabirds and shorebirds). This makes them more vulnerable to high risks such as modifying supporting terrestrial habitats, dredging-related impacts and illegal fishing and poaching. The migratory nature of many of these species means they may spend some of their life outside the Region, beyond the protection of the Marine Park.
Wetlands of international importance	The size and integrity of the Shoalwater and Corio Bays Area and its location within a defence practice area means it is less likely to be affected by many of the high risks (except climate change). Over the next 25 years, the area may provide an important refuge to many species (including listed and migratory species).

## 10.12 Summary of outcomes

- The Great Barrier Reef Region remains one of the most resilient tropical marine ecosystems in the world, however there is increasing evidence that its resilience is being seriously eroded.
- The loss of ecosystem resilience cannot be attributed to any single cause, but is almost certainly the consequence of impacts from the different activities and direct drivers, and their accumulation through time.
- Coral reefs, seagrass meadows and other inshore habitats have lost some of their resilience due to declines in condition, continuing impacts and a loss of connectivity. Dugongs have lower resilience due to historic depletion, low rates of reproduction and loss of seagrass habitats.
- The emerging loss of ecosystem resilience is particularly critical given the projected major increase in the severity of impacts related to climate change.
- The Region faces a range of increasing risks into the future. The principal sources of the impacts assessed as high and very high risk remain climate change, catchment run-off, degradation of coastal ecosystems and direct use.
- For all the Region's values, impacts associated with climate change such as ocean acidification, increased sea temperatures and rising sea level are likely to become more severe into the future. Therefore, these impacts pose an increasing risk to the Region's values.
- Legacy impacts, such as the extraction of herbivores (for example, the dugong harvest fishery which stopped operating in 1969), and many of the impacts of fishing have a reduced future risk due to improved management.
- Although there have been few large oil spills and no large chemical spills recorded in the Region and these events remain unlikely in the future, the consequence of a large spill would be major.
- Geomorphological features are most likely to be at risk from ocean acidification, sediments from catchment run-off, dumping and resuspension of dredge material, and physical impacts of fishing.
- The close connections between biodiversity values and Indigenous heritage values mean the projected risk for many impacts is equivalent for both sets of values. Examples include illegal fishing and poaching, nutrients from catchment run-off and rising sea level which are rated as high risk for both biodiversity and Indigenous heritage values. However, in some cases the assessments of risk differ because of differences in perspective.
- None of the impacts assessed were rated as presenting a very high risk to historic heritage values. Only two present a high risk — cyclone activity and ocean acidification.

A summary of the likely future risks affecting the Region's values is provided in Table 10.8 to Table 10.11 for biodiversity, geomorphological features, Indigenous heritage values and historic heritage values respectively.

Table 10.8 Summary of projected risks to biodiversity values over the next 25 years

Likelihood	Consequence				
	Insignificant	Minor	Moderate	Major	Catastrophic
Almost certain		<ul style="list-style-type: none"> <li>Extraction — lower order predators</li> <li>Extraction — lower trophic orders</li> <li>Physical damage — fishing</li> <li>Physical damage — other</li> <li>Spill — small chemical and oil</li> <li>Urban discharge</li> <li>Waste discharge from a vessel</li> <li>Wildlife disturbance</li> </ul>	<ul style="list-style-type: none"> <li>Artificial barriers to flow</li> <li>Extraction — death of discarded species (most species)</li> <li>Extraction — top order predators</li> <li>Marine debris</li> <li>Noise pollution</li> <li>Pesticides from catchment run-off</li> </ul>	<ul style="list-style-type: none"> <li>Extraction — death of discarded species (species of conservation concern)</li> <li>Illegal fishing and poaching</li> <li>Modifying supporting terrestrial habitats</li> <li>Nutrients from catchment run-off</li> <li>Sediments from catchment run-off</li> <li>Rising sea level</li> </ul>	<ul style="list-style-type: none"> <li>Increased sea temperature</li> <li>Ocean acidification</li> </ul>
Likely		<ul style="list-style-type: none"> <li>Extraction — herbivores (dugong and turtle)</li> <li>Light impacts (artificial)</li> <li>Outbreak or bloom of other species</li> <li>Vessel strike on wildlife</li> <li>Dredging</li> </ul>	<ul style="list-style-type: none"> <li>Extraction — fishing spawning aggregations</li> <li>Dumping and resuspension of dredge material</li> </ul>	<ul style="list-style-type: none"> <li>Increased freshwater inflow</li> </ul>	<ul style="list-style-type: none"> <li>Outbreak of crown-of-thorns starfish</li> </ul>
Possible		<ul style="list-style-type: none"> <li>Atmospheric pollution</li> </ul>	<ul style="list-style-type: none"> <li>Acid sulphate soils</li> <li>Coastal reclamation</li> <li>Exotic species and diseases</li> <li>Industrial discharge</li> <li>Outbreak of disease</li> <li>Physical damage — ship grounding</li> </ul>	<ul style="list-style-type: none"> <li>Cyclone activity</li> </ul>	
Unlikely				<ul style="list-style-type: none"> <li>Altered ocean currents</li> <li>Extraction — herbivores (fish)</li> <li>Spill — large chemical</li> <li>Spill — large oil</li> </ul>	
Rare					
Projected risk		Low risk	Medium risk	High risk	Very high risk



**Table 10.9 Summary of projected risks to geomorphological features over the next 25 years**

Likelihood	Consequence				
	Insignificant	Minor	Moderate	Major	Catastrophic
<b>Almost certain</b>	<ul style="list-style-type: none"> <li>Rising sea level</li> </ul>	<ul style="list-style-type: none"> <li>Dumping and resuspension of dredge material</li> <li>Increased sea temperature</li> <li>Nutrients from catchment run-off</li> <li>Physical damage — other</li> </ul>	<ul style="list-style-type: none"> <li>Ocean acidification</li> <li>Sediments from catchment run-off</li> </ul>		
<b>Likely</b>		<ul style="list-style-type: none"> <li>Dredging</li> <li>Increased freshwater inflow</li> </ul>			
<b>Possible</b>		<ul style="list-style-type: none"> <li>Outbreak of crown-of-thorns starfish</li> <li>Physical damage — ship grounding</li> </ul>	<ul style="list-style-type: none"> <li>Coastal reclamation</li> <li>Cyclone activity</li> <li>Physical damage — fishing</li> </ul>		
<b>Unlikely</b>		<ul style="list-style-type: none"> <li>Altered ocean currents</li> </ul>			
<b>Rare</b>					
<b>Projected risk</b>		Low risk	Medium risk	High risk	Very high risk

Table 10.10 Summary of projected risks to Indigenous heritage values over the next 25 years

Likelihood	Consequence				
	Insignificant	Minor	Moderate	Major	Catastrophic
Almost certain		<ul style="list-style-type: none"> <li>Noise pollution</li> <li>Pesticides from catchment run-off</li> <li>Physical damage — other</li> <li>Spill — small chemical and oil</li> <li>Waste discharge from a vessel</li> </ul>	<ul style="list-style-type: none"> <li>Artificial barriers to flow</li> <li>Dumping and resuspension of dredge material</li> <li>Extraction — lower order predators</li> <li>Extraction — top order predators</li> <li>Marine debris</li> <li>Nutrients from catchment run-off</li> <li>Rising sea level</li> <li>Sediments from catchment run-off</li> <li>Wildlife disturbance</li> </ul>	<ul style="list-style-type: none"> <li>Extraction — death of discarded species</li> <li>Illegal fishing and poaching</li> <li>Increased sea temperature</li> <li>Modifying supporting terrestrial habitats</li> <li>Ocean acidification</li> </ul>	
Likely		<ul style="list-style-type: none"> <li>Light impacts (artificial)</li> <li>Outbreak or bloom of other species</li> <li>Vessel strike on wildlife</li> </ul>	<ul style="list-style-type: none"> <li>Extraction — lower trophic orders</li> <li>Extraction — fishing spawning aggregations</li> <li>Urban and industrial discharge</li> </ul>	<ul style="list-style-type: none"> <li>Dredging</li> <li>Increased freshwater inflow</li> </ul>	
Possible		<ul style="list-style-type: none"> <li>Atmospheric pollution</li> </ul>	<ul style="list-style-type: none"> <li>Acid sulphate soils</li> <li>Exotic species and diseases</li> <li>Outbreak of crown-of-thorns starfish</li> <li>Outbreak of disease</li> <li>Physical damage — fishing</li> <li>Physical damage — ship grounding</li> </ul>	<ul style="list-style-type: none"> <li>Coastal reclamation</li> <li>Cyclone activity</li> </ul>	
Unlikely				<ul style="list-style-type: none"> <li>Altered ocean currents</li> <li>Spill — large chemical</li> <li>Spill — large oil</li> </ul>	
Rare					
Projected risk		Low risk	Medium risk	High risk	Very high risk

Table 10.11 Summary of projected risks to historic heritage values over the next 25 years

Likelihood	Consequence				
	Insignificant	Minor	Moderate	Major	Catastrophic
Almost certain	• Ocean acidification				
Likely		• Rising sea level			
Possible	• Waste discharge from a vessel	• Altered ocean currents • Atmospheric pollution • Dumping and resuspension of dredge material • Increased sea temperature	• Modifying supporting terrestrial habitats • Physical damage — other • Physical damage — fishing	• Cyclone activity	
Unlikely			• Spill — large chemical	• Dredging	
Rare	• Spill — small chemical and oil • Spill — large oil	• Marine debris		• Coastal reclamation • Physical damage — ship grounding	

Projected risk	Low risk	Medium risk	High risk	Very high risk
----------------	----------	-------------	-----------	----------------

## References

- McCook, L.J., Folke, C., Hughes, T., Nyström, M., Obura, D. and Salm, R. 2007, Ecological resilience, climate change and the Great Barrier Reef, in *Climate Change and the Great Barrier Reef: A Vulnerability Assessment*, eds J.E. Johnson and P.A. Marshall, Great Barrier Reef Marine Park Authority and Australian Greenhouse Office, Townsville, pp. 75-96.
- Walker, B., Carpenter, S., Anderies, J., Abel, N., Cumming, G., Janssen, M., Lebel, L., Norberg, J., Peterson, G.D. and Pritchard, R. 2002, Resilience Management in Social-ecological Systems: a working hypothesis for a participatory approach, *Conservation Ecology* 6(1): 14.
- Holling, C.C. 1973, Resilience and stability of ecological systems, *Annual Review of Ecological Systems* 4: 1-23.
- Hughes, T.P., Baird, A.H., Bellwood, D.R., Card, M., Connolly, S.R., Folke, C., Gosberg, R., Hoegh-Guldberg, O., Jackson, J.B.C., Kleypas, J.A., Lough, J.M., Marshall, P.A., Nyström, M., Palumbi, S.R., Pandolfi, J.M., Rosen, B. and Roughgarden, J. 2003, Climate change, human impacts, and the resilience of coral reefs, *Science* 301: 929-933.
- Anthony, K.R.N., Maynard, J.A., Diaz-Pulido, G., Mumby, P.J., Marshall, P.A., Cao, L. and Hoegh-Guldberg, O. 2011, Ocean acidification and warming will lower coral reef resilience, *Global Change Biology* 17: 1798-1808.
- McCook, L.J., Wolanski, E. and Spagnol, S. 2001, Modelling and visualizing interactions between natural disturbances and eutrophication as causes of coral reef degradation, in *Oceanographic processes of coral reefs: physical and biological links in the Great Barrier Reef*, ed. E. Wolanski, CRC Press, Boca Raton, Florida, USA, pp. 113-125.
- Hoegh-Guldberg, O., Mumby, P.J., Hooten, A.J., Steneck, R.S., Greenfield, P., Gomez, E., Harvell, C.D., Sale, P.F., Edwards, A.J., Caldeira, K., Knowlton, N., Eakin, C.M., Iglesias-Prieto, R., Muthiga, N., Bradbury, R.H., Dubi, A. and Hatzios, M.E. 2007, Coral reefs under rapid climate change and ocean acidification, *Science* 318: 1737-1742.
- Wooldridge, S.A. and Done, T.J. 2009, Improved water quality can ameliorate effects of climate change on corals, *Ecological Applications* 19: 1492-1499.
- ResilienceAlliance 2013, *Resilience*, Resilience Alliance, <<http://www.resalliance.org/576.php>>.
- Nyström, M., Folke, C. and Moberg, F. 2000, Coral reef disturbance and resilience in a human dominated environment, *Trends in Ecology and Evolution* 15(10): 413-417.
- McCook, L.J. 1999, Macroalgae, nutrients and phase shifts on coral reefs: scientific issues and management consequences for the Great Barrier Reef, *Coral Reefs* 18: 357-367.
- Hughes, T.P., Rodrigues, M.J., Bellwood, D.R., Ceccarelli, D., Hoegh-Guldberg, O., McCook, L.J., Moltschanivskyj, N.A., Pratchett, M.S., Steneck, R.S. and Willis, B. 2007a, Phase shifts, herbivory, and the resilience of coral reefs to climate change, *Current Biology* 17(4): 360-365.
- McCook, L.J., Almany, G.R., Berumen, M.L., Day, J.C., Green, A.L., Jones, G.P., Leis, J.M., Planes, P., Russ, G.R., Sale, P.F. and Thorrold, S.R. 2009, Management under uncertainty: guidelines for incorporating connectivity into the protection of coral reefs, *Coral Reefs* 28: 353-366.
- Hughes, T.P. 1994, Catastrophes, phase shifts, and large-scale degradation of a Caribbean coral reef, *Science* 265(5178): 1547-1551.
- Bellwood, D.R., Hughes, T.P., Folke, C. and Nyström, M. 2004, Confronting the coral reef crisis, *Nature* 429: 827-833.
- Pandolfi, J.M., Bradbury, R.H., Sala, E., Hughes, T.P., Bjorndal, K.A., Cooke, R.G., McArdle, D., McClenachan, L., Newman, M.J.H., Paredes, G., Warner, R.R. and Jackson, J.B.C. 2003, Global trajectories of the long-term decline of coral reef ecosystems, *Science* 301(5635): 955-958.
- Szmant, A.M. 1997, Nutrient effects on coral reefs: A hypothesis on the importance of topographic and trophic complexity to reef nutrient dynamics, in *Proceedings of the 8th international coral reef symposium*, eds. H. A. Lessios and I. G.

Macintyre, Smithsonian Tropical Research Institute, Panama City, Panama, pp.1527-1532.

18. Done, T.J. 1999, Coral community adaptability to environmental change at the scales of regions, reefs and reef zones, *American Zoologist* 39: 66-79.

19. Fabricius, K.E., De'ath, G., McCook, L.J., Turak, E. and Williams, D.M. 2005, Changes in algal, coral and fish assemblages along water quality gradients on the inshore Great Barrier Reef, *Marine Pollution Bulletin* 51(1-4): 384-398.

20. Diaz-Pulido, G., McCook, L.J., Dove, S., Berkelmans, R., Roff, G., Kline, D.I., Weeks, S., Evans, R.D., Williamson, D.H. and Hoegh-Guldberg, O. 2009, Doom and boom on a resilient reef: Climate change, algal overgrowth and coral recovery, *PLoS ONE* 4(4): e5239.

21. Sweatman, H., Cheal, A.J., Coleman, G.J., Emslie, M.J., Johns, K., Jonker, M., Miller, I.R. and Osborne, K. 2008, *Long-term monitoring of the Great Barrier Reef: Status Report 8*, Australian Institute of Marine Science, Townsville.

22. McClanahan, T.R., Donner, S.D., Maynard, J.A., MacNeil, M.A., Graham, N.A.J., Maina, J., Baker, A.C., Alemu, J.B., Beger, M., Campbell, S.J., Darling, E.S., Eakin, C.M., Heron, S.F., Jupiter, S.D., Lundquist, C.J., McLeod, E., Mumby, P.J., Paddock, M.J., Selig, E.R. and van Woesik, R. 2012, Prioritizing key resilience indicators to support coral reef management in a changing climate, *PLoS ONE* 7(8): e42884.

23. Mumby, P.J. and Steneck, R.S. 2008, Coral reef management and conservation in light of rapidly evolving ecological paradigms, *Trends in Ecology and Evolution* 23(1): 555-563.

24. Nyström, M., Graham, N.A.J., Lokrantz, J. and Norstrom, A.V. 2008, Capturing the cornerstones of coral reef resilience: Linking theory to practice, *Coral Reefs* 27: 795-809.

25. Nyström, M., Norström, A.V., Blenckner, T., de la Torre-Castro, M., Eklof, J.S., Folke, C., Österblom, H., Steneck, R.S., Thyresson, M. and Troell, M. 2012, Confronting feedbacks of degraded marine ecosystems, *Ecosystems* 15(5): 695-710.

26. Hughes, T.P., Graham, N.A.J., Jackson, J.B.C., Mumby, P.J. and Steneck, R.S. 2010, Rising to the challenge of sustaining coral reef resilience, *Trends in Ecology and Evolution* 25(11): 633-642.

27. Hoey, A.S. and Bellwood, D.R. 2009, Limited functional redundancy in a high diversity system: single species dominates key ecological process on coral reefs, *Ecosystems* 12(8): 1316-1328.

28. Howell, L. 2013, *Global Risk 2013: An initiative of the Risk Response Network*, World Economic Forum, Geneva, Switzerland.

29. De'ath, G., Fabricius, K.E., Sweatman, H. and Puotinen, M. 2012, The 27-year decline of coral cover on the Great Barrier Reef and its causes, *Proceedings of the National Academy of Sciences* 109(44): 17995-17999.

30. Roff, G., Clark, T.R., Raymond, C.E., Zhao, J., Feng, Y., McCook, L.J., Done, T.J. and Pandolfi, J.M. 2013, Palaeoecological evidence of a historical collapse of corals at Pelorus Island, inshore Great Barrier Reef, following European settlement, *Proceedings of the Royal Society B-Biological Sciences* 280(1750): 2012-2100.

31. McKenzie, L.J., Collier, C. and Waycott, M. 2012, *Reef Rescue Marine Monitoring Program: Inshore seagrass, annual report for the sampling period 1st July 2010 – 31st May 2011*, Fisheries Queensland, Cairns.

32. Poiner, I.R., Glaister, J., Pitcher, C.R., Burrridge, C.Y., Wassenberg, T.J., Gribble, N.A., Hill, B.J., Blaber, S.J.M., Milton, D., Brewer, D. and Ellis, N. 1998, *Environmental effects of prawn trawling in the far northern section of the Great Barrier Reef: 1991-96*, CSIRO Division of Marine Research, Cleveland, Qld.

33. Pitcher, C.R., Burrridge, C.Y., Wassenberg, T.J., Hill, B.J. and Poiner, I.R. 2009, A large scale BACI experiment to test the effects of prawn trawling on seabed biota in a closed area of the Great Barrier Reef Marine Park, Australia, *Fisheries Research* 99(3): 168-183.

34. Burrridge, C.Y., Pitcher, C.R., Wassenberg, T.J., Poiner, I.R. and Hill, B.J. 2003, Measurement of the rate of depletion of benthic fauna by prawn (shrimp) otter trawls: An

experiment in the Great Barrier Reef, Australia, *Fisheries Research* 60(2-3): 237-253.

35. Pitcher, C.R., Wassenberg, T.J., Cappo, M.C., Smith, G.P., Austin, M., Gordon, S.R., Bustamante, R.H., Moeseneder, C.H., Speare, P.J., Kennedy, J.A., Doherty, P.J. and Hooper, J.N.A. 2004, *Dynamics of large sessile seabed fauna, important for structural fisheries habitat and biodiversity of marine ecosystems and the use of these habitats by key finfish species, Final report to the FRDC on project 1997/205*, CSIRO Marine Research, Cleveland, Qld.

36. Pitcher, C.R., Austin, M., Burrridge, C.Y., Bustamante, R.H., Cheers, S.J., Ellis, N., Jones, P.N., Koutsoukos, A.G., Moeseneder, C.H., Smith, G.P., Venables, W. and Wassenberg, T.J. 2008, *Recovery of seabed habitat from the impact of trawling in the far northern section of the Great Barrier Reef Marine Park: CSIRO final report to GBRMPA*, CSIRO Marine and Atmospheric Research, Cleveland, Queensland.

37. Cappo, M. and Kelley, R. 2001, Connectivity in the Great Barrier Reef World Heritage Area: An overview of pathways and processes, in *Oceanographic processes of coral reefs: Physical and biological links in the Great Barrier Reef*, ed. E. Wolanski, CRC Press, Boca Raton, Florida, pp. 161-187.

38. Sainsbury, K.J., Campbell, R.A., Lindholm, R. and Whitelaw, A.W. 1997, Experimental management of an Australian multi-species fishery: examining the possibility of trawl-induced habitat modification, in *Global trends in fisheries management*, eds E.K. Pititch, D.D. Huppert and M.P. Sissenwine, American Fisheries Society, Bethesda, USA, pp. 107-112.

39. Bax, N.J. and Williams, A. 2001, Seabed habitat on the south-eastern Australian continental shelf: context, vulnerability and monitoring, *Marine and Freshwater Research* 52: 491-512.

40. Williams, A., Schlacher, T.A., Rowden, A.A., Althaus, F., Clark, M.R., Bowden, D.A., Stewart, R., Bax, N.J., Consalvey, M. and Kloser, R.J. 2010, Seamount megabenthic assemblages fail to recover from trawling impacts, *Marine Ecology* 31(Suppl 1): 183-199.

41. Asch, R.G. and Collie, J.S. 2008, Changes in a benthic megafaunal community due to disturbance from bottom fishing and the establishment of a fishery closure, *Fishery Bulletin* 106(4): 438-456.

42. Bridge, T. and Guinotte, J. 2013, *Mesophotic coral reef ecosystems in the Great Barrier Reef World Heritage Area: Their potential disturbance and possible role as refugia from disturbance. Research Publication No.109*, Great Barrier Reef Marine Park Authority, Townsville.

43. Great Barrier Reef Marine Park Authority 2012a, *Informing the outlook for Great Barrier Reef coastal ecosystems*, GBRMPA, Townsville.

44. Russ, G.R., Cheal, A.J., Dolman, A.M., Emslie, M.J., Evans, R.D., Miller, I.R., Sweatman, H. and Williamson, D.H. 2008, Rapid increase in fish numbers follows creation of world's largest marine reserve network, *Current Biology* 18(12): R514-R515.

45. McCook, L.J., Ayling, T., Cappo, M., Choat, J.H., Evans, R.D., DeFreitas, D.M., Heupel, M., Hughes, T.P., Jones, G.P., Mapstone, B., Marsh, H., Mills, M., Molloy, F.J., Pitcher, C.R., Pressey, R.L., Russ, G.R., Sutton, S., Sweatman, H., Tobin, R., Wachenfeld, D. and Williamson, D. 2010, Adaptive management of the Great Barrier Reef: A globally significant demonstration of the benefits of networks of marine reserves, *Proceedings of the National Academy of Sciences* 107(43): 18278-18285.

46. Pears, R.J., Choat, J., H., Mapstone, B., D. and Begg, G., A. 2006, Demography of a large grouper *Epinephelus fuscoguttatus*, from the Great Barrier Reef: implications for harvest limits and size regulations, *Marine Ecology Progress Series* 307: 259-271.

47. Pears, R., Choat, J., H., Mapstone, B., D. and Begg, G., A. 2007, Reproductive biology of a large, aggregation-spawning serranid, *Epinephelus fuscoguttatus*: management implications, *Journal of Fish Biology* 71: 795-817.

48. Tobin, A.J., Schlaff, R., Tobin, R., Penny, A., Ayling, T., Ayling, A., Krause, B., Welch, D., Sutton, S., Sawynok, B., Marshall, N.A., Marshall, P.A. and Maynard, J.A. 2010a, *Adapting to change: minimising uncertainty about the effects of rapidly-*



changing environmental conditions on the Queensland Coral Reef Fin Fish Fishery, Final Report to the Fisheries Research and Development Corporation, Project 2008/103, James Cook University, Townsville.

49. Munday, P.L., Dixon, D.L., McCormick, M.I., Meekan, M., Ferrari, M.C.O. and Chivers, D.P. 2010, Replenishment of fish populations is threatened by ocean acidification, *Proceedings of the National Academy of Sciences* 107(29): 12930.

50. Munday, P.L., Donelson, J.M., Dixon, D.L. and Endo, G.G.K. 2009a, Effects of ocean acidification on the early life history of a tropical marine fish, *Proceedings of the Royal Society B: Biological Sciences* 276(1671): 3275-3283.

51. Munday, P.L., Cheal, A.J., Graham, N.A.J., Meekan, M.G., Pratchett, M.S., Sheaves, M., Sweatman, H. and Wilson, S.K. 2009, Tropical coastal fish, in *Marine climate change impacts and adaptation report card for Australia 2009*, eds E.S. Poloczanska, A. Hobday and A.J. Richardson, www.oceanclimatechange.org.au edn, Southport, pp. 186-204.

52. Munday, P.L., Jones, G.P., Pratchett, M.S. and Williams, A. 2008, Climate change and the future of coral reef fishes, *Fish and Fisheries* 9(3): 261-285.

53. Great Barrier Reef Marine Park Authority 2012b, *A vulnerability assessment for the Great Barrier Reef: Seagrass*, GBRMPA, Townsville.

54. Chaloupka, M.Y., Bjørndal, K.A., Balazs, G.H., Bolten, A.B., Ehrhart, L.M., Limpus, C.J., Suganuma, H., Trøeng, S. and Yamaguchi, M. 2008, Encouraging outlook for recovery of a once severely exploited marine megaherbivore, *Global Ecology and Biogeography* 17(2): 297-304.

55. Lutz, P.L., Musick, J.A. and Wyneken, J. (eds) 2003, *The biology of sea turtles*, CRC Press, Boca Raton, Florida, USA.

56. Fuentes, M.M.P.B., Limpus, C.J. and Hamann, M. 2011, Vulnerability of sea turtle nesting grounds to climate change, *Global Change Biology* 17(1): 140-153.

57. Brewer, D., Heales, D., Milton, D., Dell, Q., Fry, G., Venables, B. and Jones, P. 2006, The impact of turtle excluder devices and bycatch reduction devices on diverse tropical marine communities in Australia's northern prawn trawl fishery, *Fisheries Research* 81(2-3): 176-188.

58. Limpus, C.J. 2008, *A biological review of Australian marine turtle species, 2 Green turtle, Chelonia mydas (Linnaeus)*, Environmental Protection Agency, Brisbane.

59. Cortés, E. 2000, Life history patterns and correlations in sharks, *Reviews in Fisheries Science* 8: 299-344.

60. Robbins, W.D., Hisano, M., Connolly, S.R. and Choat, J.H. 2006, Ongoing collapse of coral reef shark populations, *Current Biology* 16: 2314-2319.

61. Chin, A., Tobin, A., Simpfendorfer, C. and Heupel, M. 2012, Reef sharks and inshore habitats: patterns of occurrence and implications for vulnerability, *Marine Ecology Progress Series* 460: 115-125.

62. Tobin, A.J., Simpfendorfer, C.A., Mapleston, A., Currey, L., Harry, A.V., Welch, D.J., Ballagh, A.C., Chin, A., Szczecinski, N., Schlaff, A., White, J. and Moore, B. 2010b, *A quantitative ecological risk assessment of sharks and finfish of the Great Barrier Reef World Heritage Area inshore waters: a tool for fisheries and marine park managers. Identifying species at risk and potential mitigation strategies*, Marine and Tropical Sciences Research Facility, Cairns.

63. Congdon, B.C. 2008, Seabirds, in *The Great Barrier Reef: biology, environment and management*, eds P.A. Hutchings, M.J. Kingsford and O. Hoegh-Guldberg, CSIRO, Collingwood, pp. 359-368.

64. Congdon, B.C., Erwin, C.A., Peck, D.R., Baker, B.G., Double, M.C. and O'Neill, P. 2007, Vulnerability of seabirds on the Great Barrier Reef to climate change, in *Climate change and the Great Barrier Reef: a vulnerability assessment*, eds J.E. Johnson and P.A. Marshall, Great Barrier Reef Marine Park Authority and Australian Greenhouse Office, Townsville, pp. 427-463.

65. Chambers, L.E., Dann, P., Deveney, C., Dunlop, N. and Woehler, E. 2012, Seabirds, in *Marine Climate Change Impacts and Adaptation Report Card for Australia 2012*, eds E.S. E.S. Poloczanska, A.J. Hobday and A.J. Richardson, .

66. Chambers, L.E., Congdon, B.C., Dunlop, N., Dann, P. and Devney, C.A. 2009, Seabirds and climate change, in *Marine climate change: impacts and adaptation responses: 2009 report card*, eds E.S. Poloczanska, A.J. Hobday and A.J. Richardson, National Climate Change Adaptation Research Facility, Southport, pp. 18 p.

67. Hughes, T.P., Gunderson, L.H., Folke, C., Baird, A.H., Bellwood, D.R., Berkes, F., Crona, B., Helfgott, A., Leslie, H., Norberg, J., Nyström, M., Olsson, P., Osterblom, H., Scheffer, M., Schuttenberg, H., Steneck, R.S., Tengö, M., Troell, M., Walker, B., Wilson, J. and Worm, B. 2007b, Adaptive management of the Great Barrier Reef and the Grand Canyon World Heritage Areas, *Ambio* 36(7): 586-592.

68. Mumby, P.J., Steneck, S. and Hastings, A. 2013, Evidence for and against the existence of alternate attractors on coral reefs, *Oikos* 122(4): 481-491.

69. Abbs, D. 2012, *The impact of climate change on the climatology of tropical cyclones in the Australia region*, Working Paper No. 11, CSIRO Climate Adaptation Flagship.

70. Kuleshov, Y., Fawcett, R., Qi, L., Trewin, B., Jones, D., McBride, J. and Ramsay, H. 2010, Trends in tropical cyclones in the South Indian Ocean and the South Pacific Ocean, *Journal of Geophysical Research* 115: doi:10.1029/2009JD012372.

71. Williams, K.J. and Crimp, S. 2012, *Queensland's biodiversity under climate change: An overview of climate change in Queensland, Climate Adaptation Flagship Working Paper No 12A*, CSIRO Climate Adaption Flagship.

72. Knutson, T.R., McBride, J.L., Chan, J., Emanuel, K., Holland, G., Landsea, C., Held, I., Kossin, J.P., Srivastava, A.K. and Sugi, M. 2010, Tropical cyclones and climate change, *Nature Geoscience* 3: 157-163.

73. Steffen, W. and Hughes, L. 2013, *The critical decade 2013: Climate change science, risks and responses*, Climate Commission Secretariat, Canberra.

74. Fabricius, K.E., Okaji, K. and De'ath, G. 2010, Three lines of evidence to link outbreaks of the crown-of-thorns seastar *Acanthaster planci* to the release of larval food limitation, *Coral Reefs* 29: 593-605.

75. Moberg, F. and Ronnback, P. 2003, Ecosystem services of the tropical seascape: interactions, substitutions and restoration, *Ocean and Coastal Management* 46(1-223310): 27-46.

76. Anthony, K., Marshall, P., Abdullah, A., Beeden, R., Bergh, C., Black, R., Eakin, M., Game, E., Gooch, M., Graham, N., Green, A., Heron, S., van Hooidonk, R.J., Knowland, C., Mangubhai, S., Marshall, N., Maynard, J., McGinnity, P., McLeod, E., Mumby, P.J., Nyström, M., Obura, D., Oliver, J., Possingham, H., Pressey, B., Rowlands, G., Tamelander, J., Wachenfeld, D. and Wear, S. Unpublished report, 'Operationalising resilience for marine conservation'.

77. Gunderson, L. and Holling, B. (eds) 2002, *Panarchy: Understanding transformations in human and natural systems*, Island Press, London.

78. Marshall, N.A. and Tobin, R. 2012, *"More than meets the eye": The social and economic impacts of recent natural disasters on marine resource dependent industries of the Great Barrier Reef region*, Great Barrier Reef Marine Park Authority, Townsville.

79. Tobin, R., Sutton, S., Penny, A., Williams, L., Maroske, J. and Nilsson, J. 2010c, *Baseline socio-economic data for Queensland east-coast inshore and rocky reef fishery stakeholders. Part A: commercial fishers*, FRDC Project No. 2007/048, Technical Report No. 5, Fishing and Fisheries Research Centre, Townsville.

80. Great Barrier Reef Marine Park Authority 2012c, *Climate change adaptation: Outcomes from the Great Barrier Reef Climate Change Action Plan 2007-2012*, GBRMPA, Townsville.







## Chapter 11

---

**Projected condition**



## Table of contents

11.1	<b>Current condition and trend of values .....</b>	<b>11-3</b>
11.2	<b>Effectiveness of current management .....</b>	<b>11-3</b>
11.2.1	Broadscale assessment .....	11-4
11.2.2	Outcomes of demonstration case studies .....	11-4
11.3	<b>Ecosystem resilience .....</b>	<b>11-4</b>
11.4	<b>Overall risks to values .....</b>	<b>11-5</b>
11.5	<b>Projected condition .....</b>	<b>11-5</b>
11.5.1	The future of the Great Barrier Reef .....	11-5
11.5.2	Projected condition of the Region's values .....	11-6
11.6	<b>Future scenarios .....</b>	<b>11-15</b>
11.6.1	Coral reef scenarios.....	11-16
11.6.2	Seagrass meadow scenarios .....	11-18
11.6.3	Dugong scenarios.....	11-18
	<b>References .....</b>	<b>11-20</b>



**Extract from Great Barrier Reef Region Strategic Assessment terms of reference**

**5. Projected condition of matters of national environmental significance**

**5.1 Describe the projected condition of the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area, based on an evaluation of their:**

- a) *current status and trends*
- b) *actual and potential impacts*
- c) *the effectiveness of the Program to protect the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area, and manage impacts*
- d) *an understanding of ecosystem resilience*
- e) *an assessment of overall risks to the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area.*

The projected condition of matters of national environmental significance in the Region is affected by a wide range of variables and is difficult to predict. Information presented in the previous chapters provides a basis for analysing likely condition into the future:

- The assessment of condition and trend of values (Chapter 7) provides an understanding of the current state of each of the values underpinning matters of national environmental significance and their recent trend. The systematic assessment undertaken, which takes into account the past and present impacts (Chapter 6), allows the values of most concern to be identified.
- The independent assessment of the effectiveness of current management (Chapter 8) provides an understanding of the strengths and weaknesses of current management arrangements. It forms an important contribution to developing projected condition as the future predictions assume that these arrangements remain unchanged.
- An understanding of ecosystem resilience (Chapter 10) informs consideration of how strongly the values of most concern are likely to bounce back from current and future impacts.
- The assessment of risk (Chapter 10) analyses the future likelihood and consequence of the identified impacts on values relevant to matters of national environmental significance. The risk of each impact is assessed separately, allowing the impacts that are likely to be of highest risk to be identified. This analysis is a vital component in developing projected condition and focusing future management activities.

In this chapter, a summary of the findings of the previous chapters is provided, followed by an assessment of the likely future condition of the Great Barrier Reef Region and values and attributes relevant to matters of national environmental significance.

## **11.1 Current condition and trend of values**

The current condition and trend of the values and attributes relevant to matters of national environmental significance are assessed in Chapter 7. Information on condition and trend is limited to a relatively small number of attributes. For many, the assessment is based on limited evidence and anecdotal information or is inferred from the condition of supporting habitats and processes. Most values and attributes are assessed as being in good or very good condition at the scale of the Region. However, in the southern two-thirds of the Region, a number of habitats (such as coral reefs, seagrass meadows and open waters) and environmental processes which are critical in supporting a range of other values are assessed as being in poor or very poor condition. There are four examples of species showing good recovery after past serious declines: humpback whales, estuarine crocodiles, loggerhead turtles and green turtles (southern stock).

The values of most concern generally occur in the southern two-thirds of the Great Barrier Reef Region, and coincide with areas where impacts on the Region's environment are compounding.

## **11.2 Effectiveness of current management**

An independent assessment considered the effectiveness of the Authority's existing measures to protect and manage the Region, examining both 15 major management topics addressed by the Authority (Chapter 8) and a series of demonstration case studies (Chapter 9).

### 11.2.1 BROADSCALE ASSESSMENT

The broadscale assessment recognised the difficulties in achieving positive outcomes on the ground, given the complex nature, large extent and long timescale of the threats and the increasing gap between the growth in management resources and the growth in issues to be addressed.

Management effectiveness was assessed to be strongest on issues limited in scale or intensity and presenting only minor or moderate complexity, such as defence and research activities. Tourism, which operates across much of the Region and is moderately complex, was also found to be effectively managed.

Management effectiveness challenges were most evident for those broadscale issues which are complex socially, biophysically and jurisdictionally. These included Indigenous heritage, climate change and extreme weather, water quality protection, coastal development, port activities, shipping, and commercial and recreational fishing.

The independent assessment reported many of the issues and key values are well understood and that stakeholder engagement was consistently strong across the management topics. Having adequate resources, implementing adequate processes and achieving desired outcomes were the elements where the Authority's current arrangements were generally assessed as least effective.

Addressing cumulative and consequential impacts and applying Indigenous knowledge to management were the weakest indicators of effectiveness across the entire assessment.

### 11.2.2 Outcomes of demonstration case studies

In addition to the broadscale assessment that considered the entire range of the Authority's management activities, a more detailed understanding of effectiveness was achieved through a series of demonstration case studies. Outcomes particularly relevant to developing the projected condition for the Region include:

- the value of addressing the many cross-jurisdictional issues through strong partnerships across all levels of government, especially having a common goal
- the importance of using all the management tools available to address complex use and environmental issues
- the effectiveness of cooperative management approaches between governments, Traditional Owners, stakeholders and the community in addressing impacts and declines in condition at a local scale
- the importance of cooperative management arrangements with Traditional Owners
- the significance of remoteness — protecting some of the Region's values, while at the same time making management intervention difficult.

## 11.3 Ecosystem resilience

The capacity of an ecosystem to withstand or recover from disturbances and impacts, and maintain key functions, is an important component in determining its future. Resilience is not about the single, static condition of an ecosystem, but its capacity to absorb or recover from impacts.

A key aspect of resilience is that different impacts may combine or even exacerbate each other so that the cumulative impacts may be far greater than any individual impact. This has two important consequences: first, the need to manage all impacts to reduce cumulative impacts; and second, reductions in one impact may reduce the effects of other impacts, potentially 'buying time' to address less accessible impacts, such as those related to climate change.

There is limited information on ecosystem resilience for the Great Barrier Reef Region, largely due to the vast extent and complexity of the ecosystem, and because resilience is a complex property which is difficult to measure. The resilience of the Great Barrier Reef ecosystem is discussed in Chapter 10, based on the best available information.

Overall, while the Region may be one of the most resilient tropical marine ecosystems in the world, there is increasing concern that its resilience is being seriously, and increasingly rapidly, eroded. The extent of loss varies considerably between components and localities. It is almost certainly the consequence of many different impacts accumulating through time and in different locations.

It is of particular concern that several of the most significant pressures on resilience, and those least effectively managed, are broad in scale. For example, degraded water quality significantly affects most of the habitats and species in inshore areas in the southern two-thirds of the Region and, to a lesser extent, in the northern third. Further, as the state of the system changes, attributes that confer resilience to a healthy ecosystem may begin to undermine resilience in a compromised system. For example, the network of coral reefs distributed across the Region confers vital connectivity for corals and fishes, but it also provides similar connectivity for destructive outbreaks of other species such as crown-of-thorns starfish.

Ecosystem resilience is also affected by the frequency of disturbances. The length of time available for recovery before the next disturbance occurs is a factor in the amount of recovery between disturbances. If disturbances are too frequent, their effects will accumulate through time. Two of the major causes of coral loss on the Great Barrier Reef — cyclones and crown-of-thorns starfish outbreaks — are thought to be increasing in intensity and frequency, respectively.

The emerging loss of resilience is particularly critical in the context of the projected major increase in severity of impacts related to climate change. As these impacts increase, it is very likely that interactions between other impacts and those related to climate change will have increasingly serious consequences for the resilience of the Region's ecosystem.

## 11.4 Overall risks to values

A detailed assessment of the values relevant to the matters of national environmental significance is presented in Chapter 10. This includes consideration of the likelihood and consequence of each predicted impact, in relation to biodiversity, geomorphological features, and Indigenous and historic heritage values.

The impacts likely to present a high or very high risk to the matters of national environmental significance have their origin in a number of factors. These may originate well beyond the boundaries of the Region:

- climate change
  - cyclone activity
  - increased sea temperature
  - ocean acidification
  - rising sea level
- catchment run-off
  - increased freshwater inflow (also linked to climate change)
  - nutrients in catchment run-off
  - outbreaks of crown-of-thorns starfish, linked to higher concentrations of nutrients
  - sediments in catchment run-off
  - urban and industrial discharge
- degradation of coastal ecosystems
  - artificial barriers to flow
  - modifying supporting terrestrial habitats
  - coastal reclamation
- direct use of the Region
  - extraction — death of discarded species
  - wildlife disturbance
  - dredging
  - dumping and resuspension of dredge material
  - extraction — lower trophic orders, lower order predators and top order predators
  - extraction — fishing spawning aggregations
  - illegal fishing and poaching
  - marine debris
  - noise pollution.

## 11.5 Projected condition

### 11.5.1 The future of the Great Barrier Reef

The health of the Great Barrier Reef is declining, particularly in inshore areas south of Cooktown, and management is not keeping pace with the cumulative impacts that are acting on the system. The causes of decline are known, and the potential for restoration is strong provided more than a 'business-as-usual' approach is adopted. Improving the Reef's resilience now is also the best approach to managing the effects of climate change in the future.

The past decade of extreme weather events has taken its toll on the Reef. The frequency of cyclones and floods has reduced the capacity of the Reef ecosystem to recover from these and other disturbances such as outbreaks of crown-of-thorns starfish. Legacy issues, such as broadscale catchment clearing and commercial harvesting of iconic species, are still affecting the Reef. Some of these issues go back decades, even to the 1800s, and their impacts are likely to continue long into the future.

The impacts outlined in this report do not operate in isolation, but overlap and interact with each other. Their accumulation through time, and over an ever-increasing area, is diminishing the ecosystem's ability to bounce back. There is increasing evidence that both the ecosystem's resistance and its capacity to recover is being lost, although the extent of that loss varies considerably between ecosystem components (for example, dugong compared to some fishes) and between localities (for example, the inshore southern two-thirds of the Region compared to places offshore and further north).

The declining condition of the Great Barrier Reef and its loss of resilience cannot be attributed to any single cause — it is almost certainly the result of cumulative impacts.

It is of particular concern that the most serious risks to the Region's values, and those least effectively managed, operate at broad scales. They originate well beyond the Region and affect an area much larger than the Region.

Climate change remains the most serious long-term risk facing the Reef and is likely to have far reaching consequences for the Region's environment. Future climate change predictions indicate sea level rises and temperature increases will continue, and the ocean will become gradually more acidic. Extreme weather events are predicted to increase in severity. These changes will have dramatic effects on the health and resilience of the Reef. The impacts of climate change will be amplified by the Reef's declining resilience and the accumulation of other impacts. In turn, the effects of climate change will exacerbate the effects of other impacts, potentially accelerating the decline in the condition of the Region's values.

The urgent need to limit global warming to two degrees Celsius above pre-industrial levels has been recognised by almost 200 nations.<sup>1</sup> At present, global emissions are not on track to achieve such a target,<sup>2</sup> and even a two degree Celsius rise would be a very dangerous level of warming for coral reef ecosystems, including the Great Barrier Reef, and the people who derive benefits from them.<sup>3</sup> To ensure the Reef remains a coral-dominated system, the latest science indicates global average temperature rise would have to be limited to 1.2 degrees Celsius.<sup>3,4</sup>

Water quality in the Region has declined markedly, especially in inshore areas adjacent to the developed coast. Agricultural practices in the Great Barrier Reef catchment are improving, leading to reduced nutrient and sediment loads entering the Reef through catchment run-off. However, there is likely to be a significant lag time before there are measurable water quality improvements in the Region, with sediments and nutrients projected to continue affecting biodiversity for the next 25 years or so. The continuation of Reef Rescue funding until 2018, to support ongoing reductions in the amount of nutrients and sediments entering the Reef from the catchment, will assist continued improvement in the Region's water quality. Potential changes to the present broadscale land clearing legislation could affect the gains made in the catchment and add to the loads entering the Region. With continued pesticide use in the catchment, it is almost certain they will also be a component of catchment run-off over the next quarter of a century. Of major concern, is the effect that pesticides will have on freshwater and estuarine systems in the catchment that support the biodiversity of the Reef.

There is increasing evidence that current trends for crown-of-thorns starfish outbreaks are likely to continue over the next 25 years, contributing further to coral mortality. The Reef's ability to recover from outbreak events has been severely compromised by cumulative impacts. Current understanding of the relationship between crown-of-thorns starfish and water quality indicates future improvements in the Region's water quality will likely reduce the frequency of outbreaks in the future.

Modifying terrestrial habitats that support the Great Barrier Reef is likely to continue, based on the expected increase and intensification of agriculture and projected increases in urban and industrial development. It is predicted the flow-on effects, especially in areas close to the coast, will continue to present a very high risk to the Region's values; for example, through changes to water quality and connectivity.

The impacts of incidental catch continue to have a major impact on species of conservation concern, although the trawl fishery has significantly reduced its incidental catch of marine turtles and other non-retained bycatch species by using improved equipment. Death of discarded and incidentally caught species of conservation concern across all fisheries and the Queensland Shark Control Program is almost certain, with major consequences for their populations. As most species discarded are significant for Traditional Owners, either as food, a totem or for customary practice, this impact is likely to have a major effect on their cultural values.

The operation and expansion of ports requires initial capital and ongoing maintenance dredging. Approval of all port development proposals currently being assessed would significantly increase the volume of dredge material disposal in the Great Barrier Reef World Heritage Area. Of key concern are the effects of redistribution and resuspension of sediments which affects habitats such as coral reefs and seagrass meadows and a range of species.

The cumulative effects of impacts highlight the need to manage all of them to reduce stresses on the system. However, many of the activities and drivers having the most effect are outside of the Authority's control — climate change being an obvious example — which limits its capacity to respond directly to the threats.

Managing the multiple pressures that are affecting the Reef requires a multi-pronged and multi-disciplinary approach. It is clear that a 'business-as-usual' approach to managing the impacts affecting the Great Barrier Reef will not be enough. North of Cooktown there is a need to focus on safeguarding the Reef's health. In contrast, restoring the condition of values of concern and halting and reversing impacts on those values will be the key themes for restoring Reef health in the southern two-thirds of the Region.

A concerted international effort to reduce global climate change, combined with action at the national and local levels to build the Reef's resilience by reducing direct and indirect impacts, is our best insurance for protecting one of nature's most outstanding masterpieces.

### **11.5.2 Projected condition of the Region's values**

The projected condition of the Region's values is assessed in Table 11.1 to Table 11.3, based on the assessments of condition, impacts, resilience, risk and management effectiveness presented in previous chapters. This analysis is used to inform the recommendations for improvement to management presented in























































Chapter 12. It also forms the basis for a new outcome-based approach to management by the Authority — one which is focused on working with all parties to collectively achieve the outcomes needed to ensure the future of the Great Barrier Reef.

























The projected condition of community benefits will, in part, be based on the condition of the Region’s biodiversity, geomorphological features, and Indigenous and historic heritage values; however, community benefit values are also dependent on social, cultural and economic drivers that are difficult to predict. Given these dependencies and variability, projected condition has not been assessed for community benefits.

Table 11.4 presents a summary of the projected condition for each of the matters of national environmental significance relevant to the Region.

Table 11.1 Projected condition of biodiversity values				
The projected condition grade for each key value and attribute relevant to matters of national environmental significance (MNES) is a grade of best fit across all elements of the value. If a number of the elements are likely to have a 'poor' projected condition then the group is assigned this grade, even if some are likely to have a better projected condition. Note that the projected condition of some values is based on limited data and knowledge and, as such, should be treated with caution.				
Understanding the table				
<b>Very good:</b> The values are likely to remain healthy and resilient for the foreseeable future with strong recovery in threatened species and at damaged locations. Additional management intervention is not required to maintain the values.	<b>Good:</b> With only minor additional management intervention, the values are likely to remain generally healthy and resilient for the foreseeable future, with only some values showing signs of significant deterioration.	<b>Poor:</b> Without significant additional management intervention, some of the values will deteriorate in the next 25 years and only a few values are likely to be healthy and resilient in the longer term.	<b>Very poor:</b> Without urgent and effective additional management intervention, the values are likely to deteriorate rapidly with the loss of most values in the longer term.	
			<div><div>Projected Condition</div><div><div>Very good</div><div>Good</div><div>Poor</div><div>Very poor</div></div></div>	
<b>Overview:</b> The health of the Great Barrier Reef ecosystem is declining, particularly in inshore areas south of Cooktown. Without urgent and effective additional management intervention, the Region's biodiversity values are likely to continue to deteriorate. A decade of extreme weather events has severely affected many habitats and species, and has reduced the capacity of the Reef to recover from these and other ongoing impacts. Legacy issues continue to affect the Reef.			<div><div></div><div></div><div></div><div></div></div>	
			<div><div>Projected Condition</div><div><div>Very good</div><div>Good</div><div>Poor</div><div>Very poor</div></div></div>	
<b>Key values and attributes</b>				
<b>Coral reefs and corals:</b> Coral reefs in the northern third of the Region are currently in good condition. Those in the southern two-thirds of the Region are in poor or very poor condition. Ongoing impacts from cyclones, bleaching events associated with increased sea temperatures, poor water quality and increased frequency of crown-of-thorns starfish outbreaks mean they are not able to recover between disturbances. Their condition is likely to deteriorate further given the increasing risks from climate change, coastal ecosystem degradation and timeframes for achieving water quality improvements, combined with their lack of resilience.			<div><div></div><div></div><div></div><div></div></div>	
<b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; national heritage place; Commonwealth marine area; listed migratory and threatened species; wetlands of international importance.				

Key values and attributes	Projected Condition			
	Very good	Good	Poor	Very poor
<p><b>Seagrass meadows and seagrasses:</b> Seagrass meadows in the northern third of the Region are currently believed to be in very good condition. Those in the southern two-thirds of the Region are known to be in poor or very poor condition. Increased nutrients, sediments and freshwater, combined with recent cyclone damage, have seriously affected their condition and reduced their resilience. Timeframes to realise improvements in water quality and increased effects from climate change impacts are likely to cause further declines in condition.</p> <p><b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; Commonwealth marine area; national heritage place; listed migratory and threatened species; wetlands of international importance.</p>				
<p><b>Islands:</b> While islands are generally in good condition, the introduction of pest species has affected many and those close to population centres are under pressure. All islands are likely to be significantly affected by rising sea levels and cyclone activity. Cays are particularly vulnerable to altered ocean currents, with ramifications on species using cays to roost and nest. Modified fire regimes have also altered island ecosystems.</p> <p><b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; national heritage place; Commonwealth marine area; listed migratory and listed threatened species; wetlands of international importance.</p>				
<p><b>Beaches and coastlines:</b> Many beaches and coastlines adjacent to the developed coast have been modified, affecting coastal habitats and processes. Economic and population growth is likely to drive further changes in adjacent beaches. Predicted increases in sea level and cyclones will affect the coastline, and recovery from these disturbances is likely to be poor where natural coastal processes have been disrupted.</p> <p><b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; national heritage place; Commonwealth marine area; listed migratory and threatened species; wetlands of international importance.</p>				
<p><b>Mangrove forests and mangroves:</b> Mangrove forests and mangroves are currently stable and their abundance is being maintained; however, as the population increases, there is likely to be further pressure on undeveloped coastal mangrove habitats.</p> <p><b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; national heritage place; Commonwealth marine area; listed migratory and threatened species; wetlands of international importance.</p>				
<p><b>Lagoon floor:</b> Although it is likely that the lagoon floor is in good condition and stable, there has been no long-term monitoring and there is likely to be some effects of trawling and extreme weather.</p> <p><b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; national heritage place; Commonwealth marine area; listed migratory and listed threatened species.</p>				
<p><b>Shoals:</b> Shoals remain in good condition throughout the Region due to their relative isolation from human impacts and are unlikely to be exposed to increased risks in the future. Shoals have been affected previously by ship groundings; however, the current management of shipping is considered effective at minimising this risk of groundings. The effect of climate change on shoals is unknown. Shoals are likely to remain in good condition.</p> <p><b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; national heritage place; Commonwealth marine area; listed migratory and threatened species.</p>				
<p><b>Halimeda banks:</b> This habitat is poorly studied, but is likely to be in very good condition given its isolation from land-based impacts and its level of protection from trawling. However, it is at risk from climate change as increased ocean acidity is likely to reduce the rate of calcification.</p> <p><b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; national heritage place; Commonwealth marine area; listed migratory and threatened species.</p>				
<p><b>Continental slope:</b> Much of the habitat remains undisturbed and is, therefore, likely to be in very good condition. A deepwater trawl fishery operating in the south-east of the Region is likely to have some physical impacts on the continental slope, but little is known of the interaction.</p> <p><b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; national heritage place; Commonwealth marine area.</p>				
<p><b>Open waters:</b> The inshore open water habitat in the southern two-thirds of the Region is in poor condition, particularly as a result of catchment run-off. Time lags in sediments and nutrients reaching the Region, and sinks of such pollutants within the Region mean it may be decades before the positive effects of improved land management in the catchment are realised. Resuspension of pollutants, for example during cyclones and after dumping of dredge material, degrades the quality of the open water habitat.</p> <p><b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; national heritage place; Commonwealth marine area; listed migratory and threatened species; wetlands of international importance.</p>				

Key values and attributes	Projected Condition			
	Very good	Good	Poor	Very poor
<p><b>Terrestrial habitats that support the Great Barrier Reef:</b> Losses and modification of terrestrial habitats in the catchment have affected the Region's environment and coastal integrity. The profound nature of many of these changes mean that natural recovery of ecosystem function is unlikely, and significant intervention would be required to restore key components. Rising sea level and cyclones will affect tidal habitats, and recovery from these disturbances is likely to be poor where terrestrial habitats have been degraded.</p> <p><b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; national heritage place; Commonwealth marine area; listed migratory and threatened species; wetlands of international importance.</p>				
<p><b>Macroalgae:</b> Based on the limited information about macroalgae in the Region, it is considered to be in very good condition. As coral reefs become increasingly degraded by cumulative impacts, fleshy macroalgae is likely to become more dominant. There may be enhanced rates of photosynthesis and growth, and increased carbon storage in some fleshy macroalgae species. Projected increases in ocean acidity are likely to affect calcification rates in calcareous algae, limiting reef growth and strength. A shift to an algal-dominated system would have major implications to future coral reef recovery.</p> <p><b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; national heritage place; Commonwealth marine area; listed migratory and threatened species.</p>				
<p><b>Benthic microalgae:</b> There is limited information about the current and projected future condition of benthic microalgae; however, they are assumed to have been undisturbed and in good condition and are likely to remain stable.</p> <p><b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; national heritage place; Commonwealth marine area.</p>				
<p><b>Other invertebrates:</b> Rising sea temperature and ocean acidification will have significant effects on the condition and functioning of other invertebrates (i.e. those other than coral). The majority of invertebrates are likely to be in good condition with strong resilience, including fisheries species. The changing inshore environment in the southern two-thirds of the Region will continue to affect some invertebrates, including through dumping of dredge material and physical damage to the seafloor. There is evidence that reduced water quality is a factor in the increased frequency of crown-of-thorns starfish outbreaks, with serious effects on coral reefs. Consequential impacts from increased shipping (for example, cargo spills and groundings) have the potential to affect invertebrates on a local scale. There is also a potential for trawl effort to increase under current management arrangements.<sup>5</sup></p> <p><b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; national heritage place; Commonwealth marine area; listed migratory and threatened species; wetlands of international importance.</p>				
<p><b>Plankton and microbes:</b> Little is known about the current and projected future status of plankton and microbes in the Region, although there are concerns about the effects of water quality, marine debris (particularly microplastics) and climate change. Improvements to land practices are likely to improve water quality, although there may be a significant lag period before improvements are observed. Altered ocean currents have the potential to change the abundance and distribution of plankton, which would have flow-on effects to higher trophic levels. Increased sea temperatures and ocean acidification are also likely to affect the abundance, distribution and calcification of some plankton.</p> <p><b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; national heritage place; Commonwealth marine area; listed migratory and threatened species.</p>				
<p><b>Bony fish:</b> While most fish populations are currently considered to be in very good or good condition, some targeted species are under significantly more pressure in southern areas, with some especially vulnerable. The combination of fishing pressure, coastal habitat degradation, climate change and extreme weather may undermine this for some species. Also, considering the predicted poor or very poor condition of key habitats for bony fish, it is likely that the future condition of this group is poor.</p> <p><b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; national heritage place; Commonwealth marine area; listed migratory and threatened species; wetlands of international importance.</p>				
<p><b>Sharks and rays:</b> Overall, sharks and rays are considered to be in poor condition. Some species are vulnerable to impacts with slow recovery rates, or their migratory patterns expose them to impacts outside the Region. Coastal and inshore specialists are the most affected by activities within and adjacent to the Region (for example, dwarf sawfish, freshwater sawfish and green sawfish — all of which are listed threatened species). Further habitat modification, illegal fishing, death of discarded catch and any unsustainable extraction are risks into the future.</p> <p><b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; national heritage place; Commonwealth marine area; listed migratory species and threatened species.</p>				

Key values and attributes	Projected Condition			
	Very good	Good	Poor	Very poor
<p><b>Sea snakes:</b> Some sea snake species are taken as incidental bycatch in the trawl fishery, but the effects on the populations are poorly understood. Continued incidental take of some species in this fishery, and predicted declines in habitats such as coral reefs, are likely to affect sea snake populations in the future.</p> <p><b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; national heritage place; Commonwealth marine area.</p>				
<p><b>Estuarine crocodile:</b> Estuarine crocodiles occur in most coastal waters in the Region. They are also regularly reported at mid-shelf and some offshore islands. The species is steadily recovering from previous population declines, with no recorded expansions in its range. Its projected condition is likely to be very good.</p> <p><b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; national heritage place; Commonwealth marine area; listed migratory species.</p>				
<p><b>Marine turtles:</b> Each species of marine turtle has a different status. They are vulnerable to a range of impacts, from marine debris and incidental take to declines in seagrass habitats. Some are affected by impacts occurring outside the Region (for example, coastal development adjacent to flatback turtle nesting beaches at Peak Island). Green turtle populations have been stressed by recent declines in seagrass meadows, but the population is expected to absorb the impact. Past population losses, declines in supporting habitats and loss of nesting habitat reduce the resilience of marine turtle populations. Continuing pressures on the populations, including beyond the Region, make it likely that the condition of some species will continue to deteriorate.</p> <p><b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; national heritage place; Commonwealth marine area; listed migratory and listed threatened species (all species).</p>				
<p><b>Seabirds:</b> Some serious declines have been recorded in seabird nesting populations. Although all seabirds are vulnerable to impacts of climate change, it is the offshore and pelagic-foraging seabirds that are most at risk. Listed migratory seabirds are particularly vulnerable. Marine debris will continue to be a risk to many species. Changes to coastal habitats affect some coastal nesting species. Given seabird nesting and feeding habits, they are likely to have a poor capacity to recover after impacts.</p> <p><b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; national heritage place; Commonwealth marine area; listed migratory and listed threatened species.</p>				
<p><b>Shorebirds:</b> Shorebird populations are thought to be in poor condition throughout the Region, and are seriously affected by changes in supporting coastal habitats, including changes in tidal flow. Sea level rise will increasingly affect habitats used by shorebirds. Many are listed migratory species and are subject to impacts well beyond the Region. Shorebird resilience is likely to be poor and any recovery is likely to be slow.</p> <p><b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; national heritage place; listed migratory species.</p>				
<p><b>Whales:</b> Listed species such as the blue, fin and sei whales are rarely sighted within the Region, and very little is known about their movements or use patterns there. Humpback whales are the most commonly sighted listed marine mammal in the Region. The east Australian population is recovering from severe depletion by commercial whaling in the 20th century. The greatest risks for listed migratory whale species over the next 25 years are likely to come from outside the Region. Nevertheless, risks within the Region such as direct disturbance from defence activities, collision with large vessels, entanglement and marine debris, may affect the species.</p> <p><b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; national heritage place; Commonwealth marine area; listed threatened and listed migratory species.</p>				
<p><b>Dolphins:</b> The Indo-Pacific humpback and Australian snubfin dolphin are listed migratory species and are likely to be in decline. The resilience of these species is assessed as poor due to continuing pressures on their inshore habitats combined with low reproductive rates. Continued development within and adjacent to their habitat is likely to increase future risks to these species. Little is known about other dolphin species. Their lesser reliance on inshore habitats is likely to mean they are less affected by future impacts to those habitats.</p> <p><b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; national heritage place; Commonwealth marine area; listed migratory species.</p>				
<p><b>Dugongs:</b> Dugong populations in the southern two-thirds of the Region have declined substantially in recent decades, most recently as a result of extreme weather events. The species' ability to recover is very poor due to the massively reduced population, loss of seagrass habitats and low reproduction rates. Recent management has reduced man-made impacts on the southern population, but it is unlikely to recover unless all such mortality is eliminated.</p> <p><b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; national heritage place; Commonwealth marine area; listed migratory species.</p>				



















**Table 11.2 Projected condition of geomorphological features**

The projected condition grade for each key value and attribute relevant to matters of national environmental significance (MNES) is a grade of best fit across all elements of the value. If a number of the elements are likely to have a 'poor' projected condition then the group is assigned this grade, even if some are likely to have a better projected condition.

### Understanding the table

<b>Very good:</b> The values are likely to remain healthy and resilient for the foreseeable future with strong recovery at damaged locations. Additional management intervention is not required to maintain the values.	<b>Good:</b> With only minor additional management intervention, the values are likely to remain generally healthy and resilient for the foreseeable future, with only some values showing signs of significant deterioration.	<b>Poor:</b> Without significant additional management intervention, some of the values will deteriorate in the next 25 years and only a few values are likely to be healthy and resilient in the longer term.	<b>Very poor:</b> Without urgent and effective additional management intervention, the values are likely to deteriorate rapidly with the loss of most values in the longer term.

	Projected Condition			
	Very good	Good	Poor	Very poor
<b>Overview:</b> The long time periods associated with changes to geomorphological features means most changes in the condition of the related biological systems may take many decades to be reflected in the Region's geomorphological features. However, some features, such as reef islands, can change and disappear in relatively short periods. Overall, it is likely the condition of some features, such as coral reefs, islands, shorelines and river deltas will deteriorate over the next 25 years. Deeper, more distant features, such as submarine canyons and channels, are likely to be less affected by cumulative human impacts, but may still be impacted by climate change-related impacts such as ocean acidification.				

	Projected Condition			
	Very good	Good	Poor	Very poor
<b>Key values and attributes</b>				
<b>Coral reefs:</b> Increasing risks from climate change and coastal habitat degradation and continuing impacts from catchment run-off, combined with their lack of resilience, are likely to result in further declines in coral cover. As a result, coral reef features are likely to change in their appearance and become more susceptible to erosion. Their overall structure is unlikely to change significantly. <b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; national heritage place; Commonwealth marine area.				
<b>Islands and shorelines:</b> Rising sea level and intense cyclones will continue to affect islands and shorelines, and recovery from these disturbances is likely to be poor where natural coastal processes have been disrupted. High islands are likely to suffer the least from impacts. Unvegetated sand cays are likely to experience the greatest changes, but it will be difficult to determine the degree of change as they are naturally dynamic features. <b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; national heritage place; Commonwealth marine area; listed migratory species and threatened species; wetlands of international importance.				
<b>Channels and canyons:</b> Although little is known about submarine canyons, karstic channels and blue holes, their depth and distance from shore mean they are likely to be in very good condition and stable into the future. Palaeochannels are found closer to the coast and are, therefore, more exposed to any impacts from human activity. The future effects of climate change on channels and canyons are unknown. <b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; national heritage place; Commonwealth marine area.				

















Key values and attributes	Projected Condition			
	Very good	Good	Poor	Very poor
<b>River deltas:</b> It is likely that some river deltas have been affected by coastal development, changed sediment loads from the catchment, and artificial barriers to flow. These effects are likely to continue into the future; however, river deltas are dynamic features and likely to remain in fairly good condition. <b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; national heritage place; Commonwealth marine area.				
<b>Halimeda banks:</b> <i>Halimeda</i> banks are poorly studied, but it is likely the features are in good condition, given the isolation from land-based impacts and level of protection from trawling. Increases in ocean acidification are likely to have an effect on the calcification rates of <i>Halimeda</i> , with long-term implications for the future of the geomorphological feature. <b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; national heritage place; Commonwealth marine area.				
<b>Seagrass meadows:</b> Seagrass meadows in the northern third of the Region are currently believed to be in very good condition. Those in the southern two-thirds of the Region are in poor or very poor condition. Increased nutrients, sediments and freshwater, combined with recent cyclone damage, have seriously affected their condition and reduced their resilience. Continuing poor water quality and increased effects from climate change impacts are likely to cause further declines in condition, which will have an impact on seagrass meadows as a geomorphological feature and their associated role in trapping and stabilising sediment. <b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; national heritage place; Commonwealth marine area; listed migratory species and threatened species; wetlands of international importance.				





**Table 11.3 Projected condition of Indigenous and historic heritage values**









The projected condition grade for each key value and attribute relevant to matters of national environmental significance (MNES) is a grade of best fit across all elements of the value. If a number of the elements are likely to have a 'poor' projected condition then the group is assigned this grade, even if some are likely to have a better projected condition.

Understanding the table			
<b>Very good:</b> The values are likely to remain healthy and resilient for the foreseeable future with strong recovery at damaged locations. Additional management intervention is not required to maintain the values.	<b>Good:</b> With only minor additional management intervention, the values are likely to remain generally healthy and resilient for the foreseeable future, with only some values showing signs of significant deterioration.	<b>Poor:</b> Without significant additional management intervention, some of the values will deteriorate in the next 25 years and only a few values are likely to be healthy and resilient in the longer term.	<b>Very poor:</b> Without urgent and effective additional management intervention, the values are likely to deteriorate rapidly with the loss of most values in the longer term.

Indigenous heritage values				
	Projected Condition			
	Very good	Good	Poor	Very poor
<b>Overview:</b> Indigenous heritage values are assessed as currently being in poor condition overall. It is likely these values will continue to deteriorate, based on inextricable connections between Indigenous culture and the Reef environment and the increasing pressures of coastal development.				

Key values and attributes	Projected Condition			
	Very good	Good	Poor	Very poor
<b>Cultural practices, observances, customs and lore:</b> Traditional Owners with connections to the Region maintain their cultural practices and customs in relation to the Region, and this is likely to continue into the future. <b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; national heritage place; Commonwealth marine area; listed migratory and threatened species; wetlands of international importance.				
<b>Indigenous sacred sites, sites of particular significance, places important for cultural tradition:</b> Sacred sites and other sites of cultural significance are under pressure in many coastal areas in and adjacent to the Region. These values are likely to continue to be affected, primarily through coastal ecosystem degradation, direct use of the Region and climate change. Others are intact and likely to continue to be well managed by Traditional Owners into the future. <b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; national heritage place; Commonwealth marine area; wetlands of international importance.				
<b>Indigenous stories, songlines, totems and languages:</b> Direct use of the Region and other activities are likely to continue to affect Indigenous stories, songlines, totems and languages (for example, shipping and modifying supporting terrestrial habitats). <b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; national heritage place; Commonwealth marine area; listed migratory and threatened species; wetlands of international importance.				
<b>Indigenous structures, technology, tools and archaeology:</b> It is expected that Traditional Owners will continue to fulfil their traditional responsibilities to use and manage specific Indigenous structures, tools and sites. While Traditional Owners may have adopted new tools and technologies, the cultural practice of activities such as hunting and gathering of resources will continue to be a major component of Indigenous heritage. <b>Relevant MNES:</b> World Heritage Area; Great Barrier Reef Marine Park; national heritage place; Commonwealth marine area; wetlands of international importance.				

Historic heritage values				
	Projected Condition			
	Very good	Good	Poor	Very poor
<b>Overview:</b> Historic heritage values are assessed as currently being in good condition overall. Some historic values, such as shipwrecks, are well studied and are likely to remain in good condition. However, limited knowledge about other historic heritage values means their projected condition is difficult to predict.				

Key values and attributes	Projected Condition			
	Very good	Good	Poor	Very poor
<b>Places of historic significance — historic shipwrecks:</b> There are more than 1300 known historic shipwrecks (i.e. those greater than 75 years) in the Region. Of these, only six are protected. Known wrecks have been systematically recorded as part of the Australian National Shipwrecks Database. However, the majority are poorly recorded or their locations remain unknown. <b>Relevant MNES:</b> Great Barrier Reef Marine Park; Commonwealth marine area.				
<b>Places of historic significance — World War II features and sites:</b> The features and sites of significance in relation to World War II include shipwrecks, aircraft wrecks, unexploded ordnances and features on islands. Generally, they have been identified but little is known of their current and, hence, projected condition. <b>Relevant MNES:</b> Great Barrier Reef Marine Park; Commonwealth marine area.				

Key values and attributes	Projected Condition			
	Very good	Good	Poor	Very poor
<b>Places of historic significance — lightstations:</b> The four Commonwealth heritage-listed lightstations are being maintained and restored under heritage management plans. There is strong ongoing management for these sites. Increased cyclones and rising sea level are likely to present an increasing risk to these structures. <b>Relevant MNES:</b> Great Barrier Reef Marine Park; Commonwealth marine area.				
<b>Places of historic significance — other:</b> Other places of historic significance in the Region include Endeavour Reef, sites of turtle and dugong factories, and Mrs Watson's cottage on Lizard Island. Most are poorly recorded and their current and, hence, projected condition is not well understood. <b>Relevant MNES:</b> Great Barrier Reef Marine Park; Commonwealth marine area.				
<b>Places of scientific significance:</b> Areas of scientific significance are generally well recorded and maintained. However, as with the overall condition of the Great Barrier Reef, the condition of some sites continues to decline. <b>Relevant MNES:</b> Great Barrier Reef Marine Park; Commonwealth marine area.				
<b>Places of social significance — iconic sites:</b> The condition of some iconic sites has deteriorated. As many are associated with coral reefs, future risks to that habitat are likely to also affect their condition. The defined area of such sites makes it more feasible to undertake management intervention to maintain their condition. <b>Relevant MNES:</b> Great Barrier Reef Marine Park; Commonwealth marine area.				

























**Table 11.4 Projected condition of matters of national environmental significance**

The projected condition grade for each matter of national environmental significance (MNES) is a grade of best fit across all elements of each matter. If a number of the elements are likely to have a 'poor' projected condition, then the group is assigned this grade, even if some are likely to have a better projected condition.

Understanding the table			
Very good	Good	Poor	Very poor
<b>Very good:</b> The values are likely to remain healthy and resilient for the foreseeable future with strong recovery at damaged locations. Additional management intervention is not required to maintain the values.	<b>Good:</b> With only minor additional management intervention, the values are likely to remain generally healthy and resilient for the foreseeable future, with only some values showing signs of significant deterioration.	<b>Poor:</b> Without significant additional management intervention, some of the values will deteriorate in the next 25 years and only a few values are likely to be healthy and resilient in the longer term.	<b>Very poor:</b> Without urgent and effective additional management intervention, the values are likely to deteriorate rapidly with the loss of most values in the longer term.

Key values and attributes	Projected Condition			
	Very good	Good	Poor	Very poor
<b>World heritage properties, including outstanding universal value:</b> Many elements that make up the outstanding universal value of the Great Barrier Reef World Heritage Area are in good condition, but declining. While the condition of some attributes such as humpback whales has improved, others have experienced serious declines since 1981. The natural beauty of most of the Region remains intact; however, its underwater aesthetic value has declined in southern inshore areas. Recognising the increasing threats facing the Great Barrier Reef environment, the future condition of the world heritage property is projected to be poor.				



Key values and attributes	Projected Condition			
	Very good	Good	Poor	Very poor
<b>The Great Barrier Reef Marine Park:</b> The Marine Park is exposed to impacts that occur both within and adjacent to its boundaries (for example, climate change, catchment run-off, coastal ecosystem degradation). While direct use of the Marine Park, in most cases, is considered ecologically sustainable, the cumulative effect of all impacts from within the Marine Park and beyond has resulted in this matter being in poor condition in its southern two-thirds. Its condition is likely to deteriorate further in the future. The close connections between the natural environment and Indigenous heritage values mean that these are also likely to deteriorate into the future. As a multiple-use protected area, there are diverse and numerous social and economic (community) benefits derived from the Marine Park environment. Their future condition will depend, in large part, on the condition of the area's natural environment.				
<b>National heritage places:</b> National heritage listing of the Great Barrier Reef is based on its recognition as a world heritage property. The values that underpin this matter of national environmental significance are the same as those identified for the world heritage property, and the overall projected condition is the same.				
<b>Commonwealth marine areas:</b> The Commonwealth marine area extends beyond the Region, into the Torres Strait and Coral Sea, and to the south of the Region. The values, risks and projected condition relevant to this matter of national environmental significance in the Region are the same as those for the Great Barrier Reef Marine Park.				
<b>Listed migratory species:</b> The very nature of migratory species is that they move, often large distances. This means that most of the Region's listed migratory species spend significant amounts of time outside the Region and, hence, are exposed to impacts that may not occur within the Region. Within the Region, migratory species are likely to be at risk from climate change, degradation of coastal ecosystems, catchment run-off and commercial fishing. Catastrophic nesting failure has already been recorded for some seabirds. Australia-wide declines in shorebirds of between 70 and 80 per cent have been recorded in the past 24 years and the poor resilience of most species, combined with continuing impacts, means the projected condition of these species in the Region is poor.				
<b>Listed threatened species:</b> All of the listed threatened species found within the Region move outside the Region, often over large distances, including internationally, and are, therefore, exposed to impacts beyond the Region. The greatest risks for the listed threatened species over the next 25 years are likely to come from: commercial fishing, climate change, coastal development, habitat loss and catchment run-off. Their current poor status, combined with the generally low resilience, means that the projected condition for the Region's threatened species is poor.				
<b>Wetlands of international importance:</b> The Shoalwater and Corio Bays wetland area is in very good condition, which can be attributed mainly to its restricted access as a defence training area. It is likely that its current condition will mean that the area is better able to recover from future broadscale impacts that may affect the area, such as those from climate change and catchment run-off.				

## 11.6 Future scenarios

As outlined in previous chapters, drivers, activities, past and current impacts and future risks do not operate independently, but are intertwined in a complex web of cumulative effects. The qualitative models used in Chapter 6 to assess cumulative impacts on coral reefs, seagrass and dugong can also be employed to account for and predict how future drivers and activities are likely to have an effect throughout the entire network of ecosystem interactions. Such models can guide the development of targets for management actions required to halt and reverse deteriorating conditions of values. Recognising the models are preliminary and subject to further expert review, they are a first step in an examination of the likelihood of changes in condition as a result of future changes in some of the most serious impacts. This type of analysis can enhance understanding of likely projected condition and inform recommendations to improve future management.

As part of the Sustainable Regional Development Program project *Great Barrier Reef resilience decision framework*<sup>6</sup>, the method of Hosack *et al.* (2008)<sup>7</sup> has been used to consider cumulative impacts from multiple sources, as well as to compare alternative models of ecosystem function.

The resultant Bayesian Belief Networks, shown later in this section, follow a standard format. The top rows contain the ecosystem variables (for example, *Acropora* coral species, crown-of-thorns starfish and seagrass meadows) in either a state of increase, unchanged or decrease. The bottom rows contain the inputs — the drivers, activities or impacts that may influence the system (for example, ocean warming or nutrients). These also have the three states: increase, unchanged or decrease. The likely response of an ecosystem variable is affected by the state of each input — that is, whether or not there is a significant change to that input in the system. For instance, in the first scenario (Figure 11.1 A), the model variables are set so that there is no significant change in

the concentrations of nutrients, turbidity and sedimentation, or the frequency of intense storms (that is, it is likely the variables will remain unchanged). However, for ocean warming the likelihood has been altered, with an equal likelihood given to both a significant increase and no change.

### 11.6.1 Coral reef scenarios

Coral reefs are part of a complex set of interactions, affected by a range of activities and drivers. A simplified qualitative representation of the interactions and linkages is presented in Chapter 6, Figure 6.27. To reflect this complexity and the different hypotheses of ecosystem function in relation to coral reefs, four alternate models of coral reefs were examined as part of the resilience decision framework project, and predictions of likely responses generated. A more detailed explanation of the models, including a comparison of outcomes from the four alternate models, is presented in the technical report supporting the coral demonstration case study available at [www.gbrmpa.gov.au](http://www.gbrmpa.gov.au).

Figure 11.1 presents the outcomes of one of the models — assuming a positive link between nutrients and crown-of-thorns starfish populations, and between sediments and herbivorous fish.


The outcomes of the modelling can be summarised as:

- **Scenario A:** A 50 per cent chance of an increase in ocean warming is likely to lead to an approximate 50 per cent chance of a decrease in coral cover and coral dependent species as a result of bleaching impacts. This symmetry of input and response is a result of the relatively simple cause-and-effect pathway between ocean warming (bleaching) and coral cover.
- **Scenario B:** A 50 per cent chance of an increase in the frequency of intense storms is likely to lead to an approximate 50 per cent reduction in coral cover. The responses of corals, fish and invertebrates are predicted to be similar to the previous scenario, with crown-of-thorns starfish, macroalgae and herbivorous fishes having an equal likelihood of increasing or not changing, and turf algae and coral recruitment having an equal likelihood of decreasing or not changing. The pathways for this cause-and-effect relationship are more numerous and diffuse than scenario A, but nonetheless result in a similar response for corals and coral dependent fishes and invertebrates.
- **Scenario C:** A simultaneous increase in ocean warming and storm events intensifies the predicted decline in coral cover and fish and invertebrates. All other predictions for variables remain unchanged from scenarios A and B.
- **Scenario D:** Even with simultaneous increases in the likelihood of ocean warming and intensity of storm events, improvements to water quality can make a significant difference to the likely future condition of coral reef variables. Under this scenario, there is a considerable increase in the probability of increases in corals, coral recruitment and coral dependent fish and invertebrates, and a decrease in macroalgae. The modelled improvements to water quality are based on the *Reef Water Quality Protection Plan*<sup>8</sup> targets for strong reductions in nutrients and moderate reductions in sediments. These are expressed as an 80 per cent likelihood of a significant decrease in nutrients and 50 per cent likelihood of a significant decrease in turbidity and sediments. Additional modelling shows that these results are robust, regardless of the coral reef model used.


Testing these four scenarios for coral reefs adds weight to the projections that climate change variables, such as ocean warming and the increased intensity of storm events, are likely to substantially affect the future condition of reefs in the Region. Importantly, it also demonstrates the likely effectiveness of improving water quality in the Region as a means of improving the resilience of coral reefs to climate change and other impacts.

Although coral reefs on the Great Barrier Reef remain among the most well managed in the world, it is important to understand that simply removing pressures will not automatically lead to recovery, especially if the Reef degrades to a point beyond its natural resilience level (a phase shift).<sup>9,10</sup> A large proportion of coral reefs around the world are degraded to such a degree that recovery is highly unlikely, including many of the reefs in the Caribbean.<sup>11,12</sup>


### A: Increased likelihood of ocean warming




### B: Increased likelihood of storms



### C: Increased likelihood of ocean warming and storms



### D: Increased likelihood of ocean warming and storms plus improvements in water quality



**Figure 11.1 Future modelled changes in the condition of coral reefs**

Four major factors affecting coral reefs are varied (blue circles) to illustrate likelihood of change in outcomes for the key components of a coral reef ecosystem (red and green arrows).

### 11.6.2 Seagrass meadow scenarios

As with coral reefs, seagrass meadows are influenced by a complex set of interactions, involving a multitude of species and affected by a range of activities and drivers. They are especially important as a habitat for listed species such as dugongs and green turtles. A simplified qualitative representation of the interactions and linkages for seagrass meadows is presented in Chapter 6, Figure 6.28. Again, to reflect this complexity and the different hypotheses of ecosystem function in relation to seagrass meadows, four alternate models were tested in the resilience decision framework project.<sup>6</sup>

Figure 11.2 presents the outcomes of the modelling for the seagrass meadow model that includes a negative effect from epiphytes (a plant which grows upon another but does not derive food or water from it) to seagrass (from shading effects on seagrass growth), and dugong populations regulated by additional factors not specified in the model.

The outcomes of the modelling can be summarised as:

- **Scenario A:** A simultaneous increase in structural damage, erosion and ocean warming results in an ambiguous response for many variables. There is an increased likelihood that seagrass species with higher growth rates and shorter life spans (known as *r* strategy species) will be more abundant; and that there will be a likely decrease in marine turtles as a result of impacts on their nesting beaches and changes to incubation conditions, rather than food availability.
- **Scenario B:** The cumulative effects of an increase in structural damage, erosion and ocean warming, operating simultaneously with improvements to water quality, results in predictions for many variables similar to those in scenario A. There is more variability in the outcomes, with dugongs having a slightly greater likelihood for either increasing or decreasing, mirroring the likelihoods for change in seagrass abundance. The assumptions made about water quality improvement were the same as those in scenario D for coral reefs (Section 11.6.1).
- **Scenario C** relates to dugong populations and is discussed below.

Recognising the relative simplicity of the model and the level of understanding of the interactions taking place between all the components of the seagrass meadow habitat, testing of the three scenarios illustrates the complex interactions within this habitat and the variability in the likely responses to changes in the factors affecting it. The results suggest climate change impacts such as ocean warming and increased structural damage and erosion may increase the likelihood of a shift in seagrass species away from slow-growing, long-lived species (known as *K* strategy species) to shorter-lived, faster growing pioneer species (*r* strategy species).

### 11.6.3 Dugong scenarios

The same qualitative model was employed to investigate future implications for dugong populations as for seagrass meadows, recognising the close interconnections between the species and its habitat.


In summary, the outcomes of the modelling depicted in Figure 11.2, with respect to dugongs, are:

- **Scenario A:** A simultaneous increase in structural damage, erosion and ocean warming results in variable outcomes for dugongs with only a 60 per cent chance of increasing or not changing.
- **Scenario B:** The cumulative effects of an increase in structural damage, erosion and ocean warming, operating simultaneously with improvements to water quality, results in even more variability in the outcomes for dugongs.
- **Scenario C:** An increased likelihood of dugongs surviving other causes of mortality combined with changes in the other variables, similar to scenario B, results in a reversal in the predicted fate of dugongs, which now have a much greater chance of increasing.


While the outcomes of testing these three scenarios for dugong populations are ambiguous, the outcomes of scenario C suggest improving the condition of seagrass meadows on which dugong depend, and continuing to reduce the mortality of dugongs from man-made sources, such as incidental take and vessel strike, is important.




### A: Increased likelihood of ocean warming and storms



### B: Increased likelihood of ocean warming and storms plus improvements in water quality



### C: Increased likelihood of ocean warming and storms plus improvements in water quality and increased dugong survival




**Figure 11.2 Future modelled changes in the condition of seagrass meadows and dugongs**

Five major factors affecting seagrass meadows are varied (blue circles) to illustrate likelihood of change in outcomes for the key components of the ecosystem (red and green arrows). The effect of an increased likelihood of dugongs surviving other causes of mortality (abbreviated as Input\_to\_Dugong) is an additional variable tested in scenario C.

## References

1. United Nations Climate Change Commission Secretariat 2011, *Durban conference delivers breakthrough in international community's response to climate change*, UNFCCC, Bonn, Germany.
2. International Energy Agency 2012, *World Energy Outlook 2012*, Organisation for Economic Co-operation and Development, Paris, France.
3. Frieler, K., Meinhausen, M., Golly, A., Mengel, M., Lebek, K., Donner, S.D. and Hoegh-Guldberg, O. 2012, Limiting global warming to 2C is unlikely to save most coral reefs, *Nature Climate Change* 3(2): 165-170.
4. Hoegh-Guldberg, O., Mumby, P.J., Hooten, A.J., Steneck, R.S., Greenfield, P., Gomez, E., Harvell, C.D., Sale, P.F., Edwards, A.J., Caldeira, K., Knowlton, N., Eakin, C.M., Iglesias-Prieto, R., Muthiga, N., Bradbury, R.H., Dubi, A. and Hatzioiols, M.E. 2007, Coral reefs under rapid climate change and ocean acidification, *Science* 318: 1737-1742.
5. Pears, R.J., Morison, A.K., Jebreen, E.J., Dunning, M.C., Pitcher, C.R., Courtney, A.J., Houlden, B. and Jacobsen, I.P. 2012, *Ecological risk assessment of the East Coast Otter Trawl Fishery in the Great Barrier Reef Marine Park: Technical report*, Great Barrier Reef Marine Park Authority, Townsville.
6. Anthony, K., Dambacher, J.M., Walsh, T. and Beeden, R. 2013, *A framework for understanding cumulative impacts, supporting environmental decisions and informing resilience-based management of the Great Barrier Reef World Heritage Area*, Great Barrier Reef Marine Park Authority, Canberra.
7. Hosack, G.R., Hayes, K.R. and Dambacher, J.M. 2008, Assessing model structure uncertainty through an analysis of system feedback and Bayesian networks, *Ecological Applications* 18: 1070-1082.
8. Department of the Premier and Cabinet 2009, *Reef Water Quality Protection Plan 2009 for the Great Barrier Reef World Heritage Area and Adjacent Catchments*, Reef Water Quality Protection Plan Secretariat, DPC, Brisbane.
9. Norström, A.V., Nyström, M., Lokrantz, J. and Folke, C. 2009, Alternative states on coral reefs: Beyond coral-macroalgal phase shifts, *Marine Ecology Progress Series* 376: 295-306.
10. Hughes, T.P., Rodrigues, M.J., Bellwood, D.R., Ceccarelli, D., Hoegh-Guldberg, O., McCook, L.J., Moltschaniwskyj, N.A., Pratchett, M.S., Steneck, R.S. and Willis, B. 2007, Phase shifts, herbivory, and the resilience of coral reefs to climate change, *Current Biology* 17(4): 360-365.
11. Wilkinson, C. (ed) 2008, *Status of coral reefs of the world: 2008*, Global Coral Reef Monitoring Network and Reef and Rainforest Research Centre, Townsville.
12. Bellwood, D.R., Hughes, T.P., Folke, C. and Nyström, M. 2004, Confronting the coral reef crisis, *Nature* 429: 827-833.






## Chapter 12

### **Recommended changes to management**





## Table of contents

12.1	How recommended improvements were identified .....	12-3
12.2	The need to strengthen the Authority's management .....	12-3
12.3	Recommended improvements .....	12-4
12.3.1	Recommended improvements to the Authority's management arrangements.....	12-4
12.3.2	Recommended improvements to related local, state and national government programs.....	12-9
12.4	Recommendations of the coastal zone assessment.....	12-12



## Extract from Great Barrier Reef Region Strategic Assessment terms of reference

### 6.1 Recommendations for changes to the Program

- 6.1.1 *Recommend changes to the current Program to improve its effectiveness to deliver its objectives, including outcomes that protect the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area. Recommendations for improvements should specifically address the matters listed in Section 4.1.1.*
- 6.1.2 *Consistent with Section 6.1.1 above, recommend improvements to related local, state and national government programs.*

Note: In this report 'current Program' is referred to as 'current management arrangements'.

This Chapter provides the recommended improvements to the management arrangements of the Great Barrier Reef Marine Park Authority (the Authority), in partnership with other government agencies, Traditional Owners and stakeholders, to protect and manage matters of national environmental significance relevant to the Great Barrier Reef Region (the Region).

The recommendations are designed to strengthen protection of matters of national environmental significance and guide improvements to the management of impacts, including actions required to avoid, mitigate, offset and adaptively manage.

The recommendations guide development of the accompanying Program Report for the Region which sets out the Authority's future management. Their subsequent implementation would be subject to normal government legislative and policy development processes.

## 12.1 How recommended improvements were identified

The recommended improvements are based on the findings of the strategic assessment. They include specific consideration of the outcomes from:

- the assessment of impacts (past and present, plus future risk) to matters of national environmental significance (see Chapters 6 and 10)
- the assessment of the condition and trend of values and attributes relevant to matters of national environmental significance (current and projected) (see Chapters 7 and 11)
- the independent review of management effectiveness (see Chapter 8)
- outcomes from demonstration cases (see Chapter 9)
- the identification of key information gaps and processes to address them (see Sections 6.10 and 7.7)
- the independent peer review of, and public submissions on, the draft Strategic Assessment Report and draft Program Report (see Supplementary Report).

Their development has also been informed by the results of recent Sustainable Regional Development Program projects (see Chapter 2) and relevant recent recommendations of the World Heritage Committee (see Chapter 1).

## 12.2 The need to strengthen the Authority's management

Through the *Great Barrier Reef Marine Park Act 1975*, the Authority has a clear and long-standing responsibility to protect and manage the Great Barrier Reef. Over the past three decades the Authority has established a strong and comprehensive set of management arrangements to protect the Reef, adapting them in response to emerging issues and improved understanding.

The independent assessment of management effectiveness presented in Chapter 8 found the Authority's management of activities within the Region for which it has direct responsibility is effective and its actions have delivered benefits for the Reef's resilience. The assessment highlighted that more is required to halt and reverse observed declines in Reef health, especially in addressing impacts arising outside the Region.

Effective future protection of matters of national environmental significance in the Region relies on the integration and strengthening of management measures, with a focus on protecting the Great Barrier Reef environment as a whole.

Managing the multiple impacts affecting the Reef requires a comprehensive and coordinated approach across all levels of government. A concerted international effort to reduce global climate change combined with action at national, state and local levels to build the Reef's resilience by reducing impacts is considered the best insurance for protecting the outstanding universal value of the Great Barrier Reef. By working together with an ecosystem

approach, the Region's values, including its outstanding universal value, can be protected, conserved, presented and transmitted to future generations.

The strategic assessment highlights the success of cooperative actions to 'halt and reverse the decline' of water quality entering the Region, coordinated through the Reef Water Quality Protection Plan (Reef Plan). Extension and adaption of this model across a range of high risk impacts and key values is likely to be an effective approach to improving environmental outcomes for the Region.

There is also a need to actively restore habitats and ecosystem processes which support the Reef's biodiversity, recognising regional and local differences in the condition of values and in the type and severity of the impacts affecting them. In the northern third of the Region, the values relevant to matters of national environmental significance are generally in good and very good condition, and the impacts affecting them are less severe. For that area, the aim should be to maintain and enhance the Reef's health. In contrast, in the southern two-thirds, where values are mostly in poorer condition and impacts are more severe, restoring the condition of values and reducing impacts should be the primary aim. Critically, there is a need to improve management of cumulative impacts and develop mechanisms which will deliver net conservation benefits across the Region.

Recognising the varied sources of impacts on the Region's values and the range of parties contributing to their reduction, it is considered that the definition of a comprehensive set of desired outcomes for the Region's values would provide a strong basis for identifying and prioritising actions to improve environmental condition. It would also guide decision making on development proposals likely to affect the Reef. The outcomes could be supported by measurable targets to gauge success. A clear target-driven management framework was one of the key recommendations of the World Heritage Committee in 2012 (see Chapter 1).

There is a need to improve understanding and protection of heritage values within the Region, together with the community benefits derived from the Region's environment. Consideration of these values needs to be integrated into planning and assessment decision-making.

## **12.3 Recommended improvements**

In line with the terms of reference, recommended improvements to the Authority's management arrangements are presented, as well as recommended improvements to related local, state and national government programs.

### **12.3.1 Recommended improvements to the Authority's management arrangements**

Based on the findings of the strategic assessment presented in the preceding chapters, the recommended improvements to the Authority's management arrangements are focused on:

- improving identification and consideration of the values relevant to matters of national environmental significance, including Indigenous and historic heritage values and the community benefits derived from the Region's environment
- improving the assessment of impacts on the Region's values, particularly understanding and managing cumulative impacts
- better avoiding impacts on the Region's values, including through improving compliance with the Authority's management arrangements and strengthening best practice and stewardship activities
- improving actions to mitigate impacts on the Region's values
- establishing arrangements for offsetting those impacts which cannot be avoided or mitigated so that there is a net benefit to the Region's values
- providing a management framework that sets out outcomes and targets for the Region's values and progressively incorporates ecosystems' thresholds as understanding improves
- supporting protection and restoration activities that enhance the Region's values and build resilience, including an emphasis on cooperative regional actions linked to adjacent catchments
- coordinating and integrating relevant monitoring activities to better inform adaptive management
- improving governance arrangements for key development activities
- adapting to climate change.

The recommended improvements to the Authority's management arrangements are outlined in Table 12.1. The Authority will work in partnership with Traditional Owners and stakeholders in implementing recommendations.

**Table 12.1 Recommended improvements to the Authority's management arrangements**

The identified strengths and weaknesses and the resulting recommended improvements are based on the findings of the strategic assessment. Recommendations take into consideration comments received by independent peer reviewers and public submissions on the draft Strategic Assessment Report and draft Program Report.

Strengths	Weaknesses	Recommended improvements
<b>Identifying matters of national environmental significance</b>		
The Authority's focus on the Marine Park's matters of national environmental significance means that values relevant to other matters of national environmental significance, in particular biodiversity values, are implicitly considered in decision making	Explicit consideration of all matters of national environmental significance in decision making, and the need to strengthen consideration of heritage values and community benefits	<b>REC1:</b> Explicitly incorporate consideration of all matters of national environmental significance, including attributes of the property's outstanding universal value, into the Authority's programs, plans and policies
Bioregional mapping provides a strong foundation for management. Development activities and commercial use are well mapped	The spatial distributions of most species and habitats are poorly collated, synthesised and mapped. Limited understanding of connectivity and ecological processes operating at local, regional and Reef-wide scales	<b>REC2:</b> Improve spatial mapping capabilities to support planning and assessment decision-making, including the range of values mapped and public availability
Monitoring and research programs have provided a good understanding of the status and trend of many biodiversity values	Further work is required to identify and understand the aesthetic and superlative natural phenomena which make up the world heritage property	<b>REC3:</b> Work closely with, Queensland and local government agencies to help identify values of the Great Barrier Reef World Heritage Area that are not easily represented and measured, such as aesthetic values
Ecological knowledge held by Traditional Owners  Strong collaborative relationships with Traditional Owners, including through the development of Traditional Use of Marine Resources Agreements	Existing information is restricted to a limited number of locations within the Region	<b>REC4:</b> Collaborate with Traditional Owners to undertake an assessment of the Indigenous heritage values of the Region
Ecological knowledge held by Traditional Owners  Strong collaborative relationships with Traditional Owners, including through the development of Traditional Use of Marine Resources Agreements	There is a need for policies and systems to guide the collection, handling and sharing of culturally sensitive information	<b>REC5:</b> Develop and implement knowledge management systems for Indigenous and historic heritage information, including a protocol for managing culturally sensitive information and improved information sharing arrangements
Marine Park has been managed as a multiple-use area since its inception	There is little systematic monitoring of community benefits derived from the Region	<b>REC6:</b> Improve understanding of the role the Great Barrier Reef plays in the life of the community
<b>Assessing impacts on matters of national environmental significance</b>		
There is a rigorous and well established process in place, supported by legislation, for assessing projects that may impact matters of national environmental significance	There are no established standards or guidelines to assist proponents and decision makers consider and address cumulative impacts	<b>REC7:</b> Work closely with Australian and Queensland government agencies to improve understanding and management of cumulative impacts from activities within and adjacent to the Region, and provide clearer guidance on how proponents and decision makers should address cumulative impacts in assessments
Intergovernmental Agreement provides a strong foundation for complementary joint permit assessment and approval processes	There is currently significant duplication in assessment processes across jurisdictions. The complexity of processes presents significant challenges to public engagement in decision making	<b>REC8:</b> Streamline assessment processes across jurisdictions and seek to have a more coordinated approach to community consultation

Strengths	Weaknesses	Recommended improvements
A focus on the Marine Park as a matter of national environmental significance means that values relevant to other matters of national environmental significance are implicitly considered in decision making	There is a need for clearer alignment of the Authority's permission system with the requirements of the EPBC Act	<b>REC8A:</b> Implement measures to enhance alignment between the Authority's permission system and EPBC Act assessment and approval processes, to ensure that activities will not result in unacceptable impacts to matters of national environmental significance
<b>Avoiding impacts on matters of national environmental significance</b>		
Complementary zoning arrangements provide a strong basis for coordinated management and protection of matters of national environmental significance  Tourism impacts are well understood and managed	Tourism management is administratively complex. Plans of management require updating, and complementary arrangements across jurisdictions (particularly intertidal areas) need improved alignment	<b>REC9:</b> Improve alignment between the Authority's and the Queensland Government's protected area and tourism management arrangements, and look for opportunities to streamline
Existing plans of management for high use areas	Absence of site planning in high growth areas to manage potentially conflicting use and increasing pressures on values	<b>REC10:</b> Develop and implement plans of management in areas of the Great Barrier Reef Marine Park that have high growth for recreation and other uses
Existing incidence response plans are in place for ship groundings, oil and chemical spills, coral disease and crown-of-thorns starfish outbreaks and major coral bleaching events	There is no dedicated role within the Authority to coordinate a response in the event of a ship grounding, large oil or chemical spills, or major natural disasters	<b>REC10A:</b> Improve and strengthen the Authority's arrangements for coordinating the response to major incidents, such as ship groundings, large oil or chemical spills, or major natural disasters
Technical expertise in coastal ecosystems and water quality  Policy on dredge spoil disposal and hydrodynamic modelling	No regulatory provisions that set out where ports can be developed within the Region	<b>REC11:</b> Support development of a Queensland ports strategy that concentrates port development around long-established major ports in Queensland, and encourage port master planning
The Authority has an existing Dredging and Spoil Disposal Policy for the Great Barrier Reef Marine Park	There is a need for a cross-jurisdictional policy consistent with the findings of the strategic assessment, including consideration of cumulative impacts and delivery of net benefits	<b>REC11A:</b> Facilitate the development of a whole of government policy to provide a strategic and consistent approach to the sustainable management of dredging and dredge spoil disposal in the Great Barrier Reef World Heritage Area
The Zoning Plan provides the basis for managing activities within the Region	There is a lack of strategic planning for marinas along the Great Barrier Reef coast	<b>REC12:</b> Promote a strategic approach to the development and operation of marinas and other access infrastructure along the Great Barrier Reef coast
The Great Barrier Reef's natural, Indigenous and historic heritage values are recognised as an important component of the Region, as reflected in the main object of the Great Barrier Reef Marine Park Act	Lack of recognition and statutory protection of some heritage values  Current heritage strategy is outdated  Need for improved alignment and complementarity in statutory protection arrangements across jurisdictions	<b>REC13:</b> Review and update the Great Barrier Reef Marine Park Heritage Strategy to guide management actions to strengthen recognition and protection of heritage values
Existing Reef Guardian and High Standard Tourism programs provide a strong platform for uptake of stewardship activities	Consistency in defining best management practice standards and implementation of continuous improvement measures  Inconsistent uptake of practices across and within sectors	<b>REC14:</b> Promote, recognise and encourage stewardship and best practice efforts by community, industry and government
Existing network of moorings and reef protection markers protect values in some high use locations	Inadequate resources available to meet increasing demand for Reef protection infrastructure	<b>REC15:</b> Support increased investment in site infrastructure to protect matters of national environmental significance in the Great Barrier Reef Region



Strengths	Weaknesses	Recommended improvements
Effective cross-jurisdictional arrangements are in place through the joint Australian and Queensland government compliance program	Lack of real time vessel tracking. Current penalties are not providing sufficient deterrent for repeat offenders	<b>REC16:</b> Improve compliance through more effective surveillance and compliance activities, access to latest technology, increased coordination across jurisdictions, and strengthened powers to prevent repeat offending
Effective systems and frameworks for managing protected area estate in the World Heritage Area	Lack of sufficient controls for island biosecurity and on-ground management actions to protect and maintain island natural integrity	<b>REC17:</b> Support a collaborative, Reef-wide management strategy for islands, and contribute to its development and implementation
All commercial fisheries have made significant progress towards achieving better sustainability outcomes in the Region over recent years	Some issues remain, and there is a need to continue to work with the Queensland Government and other relevant partners to improve environmental outcomes	<b>REC17A:</b> Work with the Queensland Government to provide technical and policy advice on actions to secure the long-term ecological, social and economic sustainability of Great Barrier Reef Region fisheries
<b>Mitigating impacts on matters of national environmental significance</b>		
Water quality guidelines provide a sound basis for the assessment of impacts on plants and animals in the Marine Park	Water quality guidelines do not address regional specific requirements or account for cumulative impacts  Primary research on the effects of cumulative impacts is not currently available	<b>REC18:</b> Update and strengthen the Great Barrier Reef Marine Park water quality guidelines to address a broader range of habitats and species and account for cumulative impacts
Hydrodynamic guidelines provide a sound basis for the assessment of potential impacts associated with dredging	Current guidelines do not account for longshore drift, resuspension and inter-annual variability in currents  Absence of an agreed model to predict hydrodynamic conditions	<b>REC19:</b> Improve the effectiveness of the Authority's hydrodynamic guidelines as a decision-making tool by requiring consideration of a greater range of environmental factors, and regularly reviewing them to reflect improvements in understanding
Some ecosystem thresholds for coral reef ecosystems have been defined	Limited understanding of critical ecosystem thresholds for a range of habitats and species  Primary research on ecosystem thresholds is not currently available	<b>REC20:</b> Support research on critical ecosystem thresholds, with a focus on inshore biodiversity and associated ecosystems
Existing environmental impact assessment framework for the Great Barrier Reef	Potential impacts of underwater noise on Great Barrier Reef species are not well understood  No standards or guidelines have been developed for the management of underwater noise impacts in the Region	<b>REC21:</b> Improve understanding and the Authority's management of the impacts of noise on species, particularly at-risk and inshore species
Reef Plan is successfully reducing nutrient loads from the catchment to the Reef  Crown-of-thorns starfish control program has been successfully trialled and implemented locally	The time lags expected before improvements in catchment run-off are reflected in Reef water quality	<b>REC22:</b> Reduce crown-of-thorns starfish outbreaks by continuing to improve water quality, and through a long-term control program
<b>Offsetting impacts on matters of national environmental significance</b>		
Sound regulatory basis for requiring offsets	Lack of strategic approach and coordination between jurisdictions  Uncertainty for proponents and the public regarding offsetting requirements	<b>REC23:</b> Develop a policy and supporting mechanisms to facilitate strategic and collaborative implementation of offsets across jurisdictions
Improved knowledge of actions likely to deliver environmental benefits	As above	<b>REC24:</b> Inform implementation of Australian and Queensland government offsets policies and restoration programs by identifying actions that will maximise the delivery of environmental benefits to the Region

Strengths	Weaknesses	Recommended improvements
<b>Enhancing matters of national environmental significance</b>		
Systematic framework to identify values and impacts	Primary research to support the setting of targets	<b>REC25:</b> Establish a management framework with clear outcomes and targets for the protection of values and the management of impacts, including cumulative impacts
Two strategic assessments have been undertaken  There is international attention on the Reef's future	No explicit overall program to coordinate efforts to enhance matters of national environmental significance	<b>REC26:</b> Develop and implement a long-term sustainability plan for the Great Barrier Reef in cooperation with Australian and Queensland governments
Existing partnership arrangements provide a strong foundation for effective engagement with other government regulators	Many of the high risk impacts affecting the Reef are outside of the Authority's jurisdictional control	<b>REC27:</b> Strengthen engagement with all relevant partners to facilitate actions that maintain and enhance the condition of values and reduce impacts, particularly in relation to climate change, catchment run-off, degradation of coastal ecosystems and direct use (Section 12.3.2)
Strong collaborative relationships between the Authority and Traditional Owners, including through the development of Traditional Use of Marine Resources Agreements	Lack of an overarching framework to promote and coordinate the contributions of Traditional Owners to management, and to guide management of Indigenous heritage values in the Region	<b>REC28:</b> Develop a comprehensive management framework and an Indigenous heritage strategy for Traditional Owner use and management of the Great Barrier Reef
Network of Local Marine Advisory Committees and strong working relationships with local government, industry and experts	Lack of regional data on condition and trend of many values	<b>REC29:</b> Adopt regionally-based cooperative approaches to protect inshore biodiversity hotspots — supporting local actions and encouraging cooperation
Significant expertise is available to undertake research	Despite significant progress in the alignment of management and research needs, gaps remain in a number of key areas	<b>REC30:</b> Improve alignment and coordination of strategic research priorities, and strengthen partnerships between the Authority and research institutions to facilitate the delivery of critical research needs
<b>Monitoring and evaluation</b>		
Significant expertise in monitoring available from a range of sources	Lack of overarching strategy and coordination of monitoring across the Great Barrier Reef  Lack of standardised procedures and protocols for data collection, handling and interpretation	<b>REC31:</b> Establish an integrated monitoring, reporting and adaptive management program for the Great Barrier Reef World Heritage Area, including more explicit reporting on the condition and trend of matters of national environmental significance to allow the performance of activities to be monitored against regulatory objectives, outcomes for matters of national environmental significance and relevant guidelines and standards, including ecosystem thresholds
Baseline level of understanding of human and natural impacts to marine megafauna	Limited data available on species of conservation concern	<b>REC32:</b> Maintain and improve monitoring, investigation and data management relating to critical species and habitats
A long-term socio-economic program is currently under development	A lack of systematic monitoring of social and economic values	<b>REC33:</b> Support implementation of a long-term social and economic monitoring program to improve understanding of changing use, investment and values
<b>Governance</b>		

Strengths	Weaknesses	Recommended improvements
Intergovernmental Agreement provides a strong foundation for an integrated and collaborative approach to management	Cross jurisdictional governance arrangements for development activities which span the catchment and the Region (for example, ports and associated activities)	<b>REC34:</b> Contribute to the development of improved governance arrangements for the management and coordination of development activities that affect the Great Barrier Reef
Existing partnerships and cooperative arrangements with all levels of government, as well as Traditional Owners, industry sectors and other stakeholders	There is currently no independent advisory body for the Great Barrier Reef that brings together the diverse cross-section of stakeholders in one group	<b>REC34A:</b> Establish a peak Great Barrier Reef Advisory Group made up of Traditional Owners, scientific, conservation and industry experts, to provide high level advice on the implementation of the Authority's management program
<b>Adapting to climate change</b>		
Wide body of expert information available on potential climate change impacts	Lack of clear explanations of the implications of climate change for the Reef's future	<b>REC35:</b> Communicate the implications of climate change impacts for the Great Barrier Reef, and the critical need to halt increasing concentrations of global greenhouse gases and restore them to levels that will support growth, recruitment and recovery processes of the Great Barrier Reef ecosystem
Data is available on the likely impacts of climate change, including increased magnitude of extreme weather events  Initiatives such as tourism contingency planning	Policies require updating to reflect improved knowledge	<b>REC36:</b> Ensure the impacts of climate change and extreme weather are appropriately considered in the Authority's management decisions
Industry initiatives to reduce emissions; for example, in the tourism industry	Limited uptake of initiatives across and within sectors	<b>REC37:</b> Encourage reduction of greenhouse gas emissions in the Great Barrier Reef Region in partnership with industry and communities
Industry initiatives to scope and adapt to climate change; for example, the tourism industry	Limited understanding among users of the likely effects of climate change on their businesses and activities	<b>REC38:</b> Support initiatives to build the capacity of management agencies and Reef users to adapt and respond to climate change and extreme weather events

### 12.3.2 Recommended improvements to related local, state and national government programs

One of the key findings of the strategic assessment is that most of the serious risks to the Region's values operate at broad scales, originating well beyond the Region or outside the Authority's management responsibilities. The interconnected and overlapping jurisdictional framework for managing these and other issues relevant to the Region (see Chapter 3), means that much of what is needed to protect and restore the Region's values relies on the actions of other government agencies and bodies. As a result, one of the key roles of the Authority is collaborating with and influencing its management partners to improve environmental outcomes in the Region.

The above recommendation for the Authority — to '*strengthen engagement with all relevant partners to facilitate actions that maintain and enhance the condition of values and reduce impacts, particularly in relation to climate change, catchment run-off, degradation of coastal ecosystems and direct use*' — requires the Authority to enhance its partnerships and cooperative arrangements with all levels of government, as well as Traditional Owners, industry sectors and other stakeholders. It requires the Authority to more clearly communicate the outcomes required for a healthy Great Barrier Reef, and to provide strong support and advice to those taking decisions and undertaking actions that may affect the Region's future.

Consistent with the terms of reference, the following is a description of potential avenues for improvements in related local, state and national government programs. It is noted that any such recommended improvements are the view of the Authority and not necessarily those of the other relevant agencies.

#### Climate change

Severe degradation of key Great Barrier Reef values, especially coral reefs, is forecast to occur before the middle of the century unless global concentrations of atmospheric carbon dioxide are reduced towards 350 parts per million. This would prevent temperature, ocean acidity, sea level and extreme weather events exceeding the

resilience capacity of the ecosystem. Ultimately, the outlook for the Great Barrier Reef is likely to be determined by the concentration of greenhouse gases in the Earth's atmosphere.

The enormity of issues relating to climate change means actions are required at all levels across the globe. It is important to continue to work with other Australian Government agencies and relevant partners to highlight the critical need to halt increasing concentrations of greenhouse gases and the implications for the Reef of impacts related to climate change. It is also important to lead by example, and encourage industries and the community to be active partners in reducing greenhouse gas emissions in the Region and adapting and responding to the effects of climate change.

#### **Catchment run-off**

Catchment run-off, particularly from rural diffuse sources, is one of the key sources of impacts on the Region's values, and much is being done under the joint *Reef Water Quality Protection Plan 2013* (Reef Plan) to reduce the loads of pollutants entering the Region. These management interventions are showing positive trends, and actions to reduce water quality impacts and improve ecosystem health need to continue and be accelerated where possible. It is recommended that the Authority continue, to work with the Queensland Government and other relevant partners in relation to catchment run-off, including seeking to enhance outcomes through:

- actively promoting the uptake of improved land management practices to more rapidly achieve Reef Plan targets
- responding to sub-regional differences in water quality impacts in the Region by supporting development and implementation of further regionally-based water quality improvement plans for the catchment
- encouraging strong linkages between water quality improvement initiatives and actions to protect and restore inshore biodiversity in the Region
- supporting consideration of expanding the scope of Reef Plan to incorporate other sources of pollutants (for example, urban and industrial activities) to provide a sound coordination mechanism
- encouraging the strengthened application of water quality guidelines across the Great Barrier Reef World Heritage Area, so that there is a consistent approach in the application of standards and best practice guidelines.

#### **Degradation of coastal ecosystems**

Coastal ecosystems play a vital role in supporting the values of the Region. In the southern two-thirds of the catchment, historic land use changes have resulted in the loss and degradation of many coastal ecosystems. This has seriously changed hydrological and ecological processes — in particular, affecting the Reef's water quality and species that rely on coastal areas. There is evidence that thresholds for ecosystem health have been exceeded in much of the southern two-thirds of the Region, particularly inshore, and there is a need to further reduce impacts from all sources. The northern Great Barrier Reef provides a refuge for many species and a buffer to pressures experienced further south.

It is recommended the Authority continues to enhance collaborative actions with Queensland and local governments and other relevant partners in relation to protecting and restoring supporting coastal ecosystems, including through:

- communicating the findings of the Authority's coastal ecosystem basin assessments and management case studies and encouraging their use in determining priorities for protecting and restoring coastal ecosystems and in taking actions likely to improve Great Barrier Reef health and resilience. This would help ensure available resources are employed most effectively; for example, by informing the Australian Government when developing offsets, and by guiding natural resource management bodies when planning basin restoration actions
- promoting and supporting protection of remaining intact systems within the Great Barrier Reef catchment, as identified through the outcomes of the basin assessments and the management case studies
- communicating the importance of functioning coastal ecosystems to Reef health through a position statement on the maintenance and restoration of critical coastal ecosystems, including identifying actions likely to have significant effects, and promoting a whole of ecosystem approach to management of the Reef and its catchment
- better informing relevant Queensland and local government coastal planning and decision-making frameworks about the findings of the Authority's technical reports and coastal ecosystem basin assessments, with the aim of achieving clearer linkages between catchment development and ecosystem health outcomes.

#### **Direct use — port activities**

Port activities can directly and indirectly affect the Region's values on a local or wider scale. While most port infrastructure along the Great Barrier Reef coast is outside the Region, the environmental footprint associated with access channels and ship anchorages extends into the Region.

In order to improve environmental outcomes for port activities, there is a need to adopt a strategic approach to port development and to enhance collaboration with the Queensland Government, other Australian Government agencies, port corporations and relevant partners, including through:



- improving certainty regarding the location of ports, and reducing further fragmentation of coastal ecosystems through a Queensland ports strategy that concentrates port development to around long-established major ports in Queensland
- supporting a strategic approach to the planning, assessment and management of port development, including long-term integrated planning for the network of trading ports on the Great Barrier Reef coast, with a view to achieving environmental, social and economic sustainability at a Reef-wide scale
- contributing to the development of improved governance arrangements across Great Barrier Reef ports aimed at strengthening coordination between responsible agencies across all jurisdictions, focusing on efficient and sustainable development of ports and associated activities
- improving understanding and management of environmental impacts from dredging and dredge material disposal in the Great Barrier Reef World Heritage Area, recognising the current uncertainty surrounding the duration, intensity and extent of predicted dredge material plumes, and their impacts on the Region's values. In particular by:
  - exploring with proponents and government agencies all alternatives which may avoid and reduce the need for dredging and dredge material disposal, and provide better environmental outcomes
  - ensuring dredging and dredge material disposal decisions take account of the Great Barrier Reef hydrodynamic and water quality guidelines, and do not exceed ecosystem thresholds
- working with port corporations, other agencies and proponents to identify and address critical information needs, and to promote research and development into innovative best practice arrangements to mitigate cumulative impacts of port activities within the Region.

### **Direct use — shipping**

Vessel visitation to ports within the Great Barrier Reef World Heritage Area is predicted to substantially increase in coming decades. The draft North-East Shipping Management Plan sets out a strong basis for collaboratively managing shipping and reducing potential risks in the Region. The Authority supports the actions outlined in the plan and will work with the Australian Maritime Safety Authority, Maritime Safety Queensland and other relevant partners to implement the final plan.

### **Direct use — fishing**

All commercial fisheries have made significant progress towards achieving better sustainability outcomes in the Region over recent years. Some issues remain, and the need to continue to work with the Queensland Government and other relevant partners to improve environmental outcomes is recognised, including:

- improving compliance arrangements to reduce illegal fishing — an impact that continues to undermine the benefits derived from the Great Barrier Reef Marine Park Zoning Plan 2003 and the sustainability of the Region's fisheries. In particular, the Authority recognises the importance of implementing real-time vessel tracking of the commercial fishing fleet and improving the effectiveness of penalties for illegal activity as a deterrent
- promoting further research and development into better gear technologies designed to reduce interactions with protected species and other bycatch, and examining options to legislate use of new bycatch reduction devices and strategies that display significant reductions in bycatch
- supporting actions to reduce effort in Great Barrier Reef fisheries to ensure long-term ecological, social and economic sustainability. There is evidence of excess fishing capacity in some Reef-related fisheries, and current management controls for some fisheries in the Region cannot prevent overfishing at a regional scale or a stock level, or control ecological risks
- improving protection of key herbivorous fish. This is based on the key ecological role they have in preventing a shift from coral reefs to algal-dominated systems after disturbances, as demonstrated in other coral reef systems where reduced populations of these fish have contributed to reef decline. While herbivorous fish are not currently generally targeted by fishers in the Region, it is considered that their protection would provide valuable insurance for reef recovery following future extreme weather events and incidents (for example, coral bleaching, cyclones, crown-of-thorns starfish outbreaks)
- improving collection, access and sharing of fisheries data across all relevant agencies
- fostering further uptake of best practice and stewardship within the Reef's fisheries, including in the recreational fishing sector.

### **Direct use — tourism and recreation**

Continuing the close and effective partnerships that have been developed with relevant government agencies, the tourism industry and the community is recognised as an important ingredient in maintaining effective management, including actions aimed at:

- streamlining tourism management arrangements across all jurisdictions, recognising the opportunities to improve alignment between existing management tools (for example, plans of management, policies and permits), and the value to both managing agencies and the industry of a contemporary and business-focused management approach
- promoting the presentation of the Reef's values in partnership with the tourism industry, including through training and development of educational material. The tourism industry has a central role in meeting Australia's international obligation to present the values of the world heritage property. At the same time,

increased understanding and appreciation of the Reef's outstanding universal value will assist in building support for the Region's protection

- fostering further uptake of best practice and stewardship within the Reef's tourism industry, building on the continuing success of the High Standard Tourism program
- improved planning and site infrastructure in existing and emerging high use areas to avoid potential conflicts of use across a range of uses, and manage predicted increase in recreational use
- promoting a strategic approach to the development and operation of marinas and other access infrastructure along the Great Barrier Reef coast
- improving understanding of tourism and recreational use and the benefits derived, including through supporting implementation of a long-term socio-economic monitoring program.

## 12.4 Recommendations of the coastal zone assessment

As part of the complementary strategic assessment of the adjacent coastal zone, the Queensland Government has made a number of commitments to improve management. Some of these overlap and are consistent with the Authority's recommendations; for example, development of an outcomes-based framework, a long-term sustainability plan, an integrated monitoring program, and improved understanding and management of cumulative impacts. The alignment between the two sets of recommendations is presented in Table 12.2.

As illustrated in the recommended improvements set out in Section 12.3 above, the Authority is committed to continuing its close partnership with Queensland Government agencies in achieving improved outcomes for the Great Barrier Reef.

**Table 12.2 Alignment between the recommended improvements of the Authority and the Queensland Government**

There is strong alignment between the recommended improvements of the Authority, as set out in this Chapter, and those of the Queensland Government in the strategic assessment of the adjacent coastal zone. The different jurisdictions and different terms of reference mean that some recommendations are relevant to only one assessment.

Authority recommended improvements	Queensland Government recommended improvements
<b>Identifying matters of national environmental significance</b>	
<b>REC1:</b> Explicitly incorporate consideration of all matters of national environmental significance, including attributes of the property's outstanding universal value, into the Authority's programs, plans and policies	<p><b>COM1:</b> The Queensland Government will complete regional plans in the Great Barrier Reef coastal zone where there is a gap, and continue to update other regional plans to ensure they respond to the latest information and pressures</p> <p><b>COM10:</b> The Queensland Government will use the Australian Government's 'Protected Matters Search Tool' in conducting planning and making environmental impact statement (EIS) assessment decisions related to EPBC Act protected matters</p>
<b>REC2:</b> Improve spatial mapping capabilities to support planning and assessment decision making, including the range of values mapped and public availability	<p><b>COM6:</b> The Queensland Government is committed to working with the Australian Government, including the Authority, to develop MNES guidelines for proponents to consider when assessing impacts on MNES during the EIS processes under the Program</p> <p><b>COM8:</b> The Queensland Government will develop an offsets register to spatially identify areas used as offsets under Queensland legislation, and priority areas for future offsets</p>
<b>REC3:</b> Work closely with Australian and Queensland government agencies to help identify values of the Great Barrier Reef World Heritage Area that are not easily represented and measured such as aesthetic values	<b>COM6:</b> The Queensland Government is committed to working with the Australian Government, including the Authority, to develop MNES guidelines for proponents to consider when assessing impacts on MNES during the EIS processes under the Program
<b>REC4:</b> Collaborate with Traditional Owners to undertake an assessment of the Indigenous heritage values of the Region	<b>COM14:</b> The Queensland Government will require project proponents to apply the Australian Government's guidelines for consulting with Indigenous people in relation to cultural heritage and the management of traditional use. The Australian Government's guidelines will be developed in cooperation with Queensland's and the state will also explore ways to streamline Indigenous consultation processes between the two governments
<b>REC5:</b> Develop and implement knowledge management systems for Indigenous and historic heritage information, including a protocol for managing culturally sensitive information and improved information sharing arrangements	

Authority recommended improvements	Queensland Government recommended improvements
<b>REC6:</b> Improve understanding of the role that the Great Barrier Reef plays in the life of the community	
<b>Assessing impacts on matters of national environmental significance</b>	
<b>REC7:</b> Work closely with Australian and Queensland government agencies to improve understanding and management of cumulative impacts from activities within and adjacent to the Region, and provide clearer guidance on how proponents and decision makers should address cumulative impacts in assessments	<b>COM15:</b> The Queensland Government will work with the Australian Government, including the Authority, to develop guidelines for proponents to consider when assessing cumulative impacts on matters of national environmental significance in the Great Barrier Reef World Heritage Area
<b>REC8:</b> Streamline assessment processes across jurisdictions, and seek to have a more coordinated approach to community consultation	<b>COM9:</b> The Queensland Government will develop a Direct Benefit Management Plan for the Great Barrier Reef World Heritage Area consistent with the accredited Queensland Offsets Framework  <b>COM23:</b> The Queensland Government will continue to work closely with the Authority to increase the implementation of complementary actions across protected area jurisdictions, including the streamlining of assessment and joint permitting processes, the formulation of joint park user policies, and discouraging repeat offending
<b>REC8A:</b> Implement measures to enhance alignment between the Authority's permission system and EPBC Act assessment and approval processes, to ensure that activities will not result in unacceptable impacts to matters of national environmental significance	
<b>Avoiding impacts on matters of national environmental significance</b>	
<b>REC9:</b> Improve alignment between the Authority's and the Queensland Government's protected area and tourism management arrangements, and look for opportunities to streamline	<b>COM2:</b> The Queensland Government will maintain and work to add to its protected area estate, and continue to provide funding for protected area management in the Great Barrier Reef coastal zone
<b>REC10:</b> Develop and implement plans of management in areas of the Great Barrier Reef Marine Park that have high growth for recreation and other uses	<b>COM11:</b> The Queensland Government will ensure that stringent conditions addressing matters of national environmental significance and outstanding universal value will be incorporated into approval recommendations  <b>COM24:</b> The Queensland Government will continue to fund and support ongoing joint field management activities with the Australian Government, including the Authority
<b>REC10A:</b> Improve and strengthen the Authority's arrangements for coordinating the response to major incidents, such as ship groundings, large oil or chemical spills or major natural disasters	<b>COM32:</b> The Queensland Government is committed to reducing the risk of shipping incidents and potential pollution of the marine environment, including implementing its responsibilities as part of the North-East Shipping Management Group
<b>REC11:</b> Support development of a Queensland ports strategy that concentrates port development around long-established major ports in Queensland, and encourage port master planning	<b>COM4:</b> The Queensland Government will introduce legislation to implement key actions of the Queensland Ports Strategy. The legislation will concentrate development at five Priority Port Development Areas and introduce port master planning which will incorporate environmental considerations and community engagement. The Queensland Ports Strategy also prohibits dredging within and adjoining the Great Barrier Reef World Heritage Area for the development of new, or the expansion of existing port facilities outside Priority Port Development Areas, for the next 10 years
<b>REC11A:</b> Facilitate the development of a whole of government policy to provide a strategic and consistent approach to the sustainable management of dredging and dredge spoil disposal in the Great Barrier Reef World Heritage Area	
<b>REC12:</b> Promote a strategic approach to the development and operation of marinas and other access infrastructure along the Great Barrier Reef coast	<b>COM11:</b> The Queensland Government will ensure that stringent conditions addressing matters of national environmental significance and outstanding universal value will be incorporated into approval recommendations
<b>REC13:</b> Review and update the Great Barrier Reef Marine Park Heritage Strategy to guide management actions to strengthen recognition and protection of heritage values	

Authority recommended improvements	Queensland Government recommended improvements
<b>REC14:</b> Promote, recognise and encourage stewardship and best practice efforts by community, industry and government	<p><b>COM18:</b> The Queensland Government is providing \$12 million over three years in grants under the Everyone's Environment Grants program</p> <p><b>COM19:</b> The Queensland Government will provide \$30 million of natural resource management funding to the Reef for biodiversity, wetlands, water quality, coastal risk, sustainable agriculture, and weeds and pest management projects over the next five years. This will support the sustainable management of natural resources and help protect significant natural assets</p>
<b>REC15:</b> Support increased investment in site infrastructure to protect matters of national environmental significance in the Great Barrier Reef Region	<b>COM2:</b> The Queensland Government will maintain and work to add to its protected area estate, and continue to provide funding for protected area management in the Great Barrier Reef coastal zone
<b>REC16:</b> Improve compliance through more effective surveillance and compliance activities, access to latest technology, increased coordination across jurisdictions, and strengthened powers to prevent repeat offending	<b>COM23:</b> The Queensland Government will continue to work closely with the Authority to increase the implementation of complementary actions across protected area jurisdictions, including the streamlining of assessment and joint permitting processes, the formulation of joint park user policies, and discouraging repeat offending
<b>REC17:</b> Support a collaborative, Reef-wide management strategy for islands and contribute to its development and implementation	<b>COM24:</b> The Queensland Government will continue to fund and support ongoing joint field management activities with the Australian Government, including the Authority
<b>REC17A:</b> Work with the Queensland Government to provide technical and policy advice on actions to secure the long-term ecological, social and economic sustainability of Great Barrier Reef Region fisheries	<b>COM16:</b> The Queensland Government will ensure that fisheries are managed for the purpose of ecological sustainability, supported by the ongoing collection of commercial and recreational data through various monitoring programs
<b>Mitigating impacts on matters of national environmental significance</b>	
<b>REC18:</b> Update and strengthen the Great Barrier Reef water quality guidelines to address a broader range of habitats and species and account for cumulative impacts	<b>COM3:</b> The Queensland Government will undertake on-ground actions which will deliver long-term benefits for threatened species
<b>REC19:</b> Improve the effectiveness of the Authority's hydrodynamic guidelines as a decision-making tool by requiring consideration of a greater range of environmental factors, and regularly reviewing them to reflect improvements in understanding	<b>COM11:</b> The Queensland Government will ensure that stringent conditions addressing matters of national environmental significance and outstanding universal value will be incorporated into approval recommendations
<b>REC20:</b> Support research on critical ecosystem thresholds, with a focus on inshore biodiversity and associated ecosystems	<b>COM12:</b> The Queensland Government will prioritise actions to recover species, taking into account national recovery plans,



Authority recommended improvements	Queensland Government recommended improvements
<p><b>REC21:</b> Improve understanding and the Authority's management of the impacts of noise on species, particularly at-risk and inshore species</p>	<p>threat abatement plans and conservation advice</p> <p><b>COM18:</b> The Queensland Government is providing \$12 million over three years in grants under the Everyone's Environment Grants program</p> <p><b>COM19:</b> The Queensland Government will provide \$30 million of natural resource management funding to the Reef for biodiversity, wetlands, water quality, coastal risk, sustainable agriculture, and weeds and pest management projects over the next five years. This will support the sustainable management of natural resources and help protect significant natural assets</p> <p><b>COM20:</b> The Queensland Government will continue to support programs that improve the outstanding universal value of the Wet Tropics World Heritage Area</p> <p><b>COM21:</b> The Queensland Government will continue to support the Queensland Wetlands Program to deliver a range of new mapping, information and decision-making tools and products to enable local, state and federal government agencies, landowners, regional natural resource management bodies and conservation groups to protect and manage wetlands into the future</p> <p><b>COM22:</b> The Queensland Government is committed to providing 40 new Indigenous Land and Sea Rangers in Queensland over three years, bringing the total number of Indigenous Land and Sea Rangers to 80</p> <p><b>COM24:</b> The Queensland Government will continue to fund and support ongoing joint field management activities with the Australian Government, including the Authority</p> <p><b>COM28:</b> The Queensland Government will work with the Australian Government, including the Authority, to develop a Reef 2050 — Long Term Sustainability Plan for the Great Barrier Reef World Heritage Area by the end of 2014, and ensure its implementation</p> <p><b>COM33:</b> The Queensland Government is committed to funding of \$55 million over the next five years to develop, promote and install best management practice systems to improve Reef water quality</p>
<p><b>REC22:</b> Reduce crown-of-thorns starfish outbreaks by continuing to improve water quality and through a long-term control program</p>	<p><b>COM34:</b> The Queensland Government will continue to fund and support the Reef Water Quality Protection Plan and the associated Paddock to Reef monitoring program to help achieve the long-term goal of no detrimental impact from the quality of water entering the Great Barrier Reef. Consideration will be given to the inclusion of other pollutants, other than broadscale land use, during the Plan's next review in 2018</p>

Authority recommended improvements	Queensland Government recommended improvements
<b>Offsetting impacts on matters of national environmental significance</b>	
<b>REC23:</b> Develop a policy and supporting mechanisms to facilitate strategic and collaborative implementation of offsets across jurisdictions	<b>COM7:</b> The Queensland Government will apply the Australian Government Offsets Policy until the Queensland Offsets Framework is accredited by the Australian Government. Offset guidelines that deliver net benefits will be prepared for application by planning and development decision-makers in consultation with the Australian Government
<b>REC24:</b> Inform implementation of Australian and Queensland government offsets policies and restoration programs by identifying actions that will maximise the delivery of environmental benefits to the Region	<b>COM8:</b> The Queensland Government will develop an offsets register to spatially identify areas used as offsets under Queensland legislation and priority areas for future offsets  <b>COM9:</b> The Queensland Government will develop a Direct Benefit Management Plan for the Great Barrier Reef World Heritage Area consistent with the accredited Queensland Offsets Framework
<b>Enhancing matters of national environmental significance</b>	
<b>REC25:</b> Establish a management framework with clear outcomes and targets for the protection of values and the management of impacts, including cumulative impacts	<b>COM19:</b> The Queensland Government will provide \$30 million of natural resource management funding to the Reef for biodiversity, wetlands, water quality, coastal risk, sustainable agriculture, and weeds and pest management projects over the next five years. This will support the sustainable management of natural resources and help protect significant natural assets
<b>REC26:</b> Develop and implement a long-term sustainability plan for the Great Barrier Reef World Heritage Area in cooperation with Australian and Queensland government agencies to better coordinate programs designed to manage and improve the condition of the Reef	<b>COM21:</b> The Queensland Government will continue to support the Queensland Wetlands Program to deliver a range of new mapping, information and decision-making tools and products to enable local, state and federal government agencies, landowners, regional natural resource management bodies and conservation groups to protect and manage wetlands into the future
<b>REC27:</b> Strengthen engagement with all relevant partners to facilitate actions that maintain and enhance the condition of values and reduce impacts, particularly in relation to climate change, catchment run-off, degradation of coastal ecosystems and direct use	<b>COM28:</b> The Queensland Government will work with the Australian Government, including the Authority, to develop a Reef 2050 — Long Term Sustainability Plan for the Great Barrier Reef World Heritage Area by the end of 2014, and ensure its implementation
<b>REC28:</b> Develop a comprehensive management framework and an Indigenous heritage strategy for Traditional Owner use and management of the Great Barrier Reef	<b>COM29:</b> The Queensland Government will work with the Australian Government, including the Authority, to develop an outcomes-based framework for the Great Barrier Reef World Heritage Area as part of the Reef 2050 — Long Term Sustainability Plan
<b>REC29:</b> Adopt regionally-based cooperative approaches to protect inshore biodiversity hotspots — supporting local actions and encouraging cooperation	<b>COM30:</b> The Queensland Government will work with the Australian Government, including the Authority, to establish an integrated monitoring framework and program for the Great Barrier Reef World Heritage Area as part of the Reef 2050 — Long Term Sustainability Plan
<b>REC30:</b> Improve alignment and coordination of strategic research priorities, and strengthen partnerships between the Authority and research institutions to facilitate the delivery of critical research needs	
<b>Monitoring and evaluation</b>	
<b>REC31:</b> Establish an integrated monitoring, reporting and adaptive management program for the Great Barrier Reef World Heritage Area, including more explicit reporting on the condition and trend of matters of national environmental significance, to enable decisions about use to be made and the performance of activities to be monitored against regulatory objectives, outcomes for matters of national environmental significance and relevant guidelines and standards, including ecosystem thresholds	<b>COM13:</b> The Queensland Government will prioritise actions to recover species, taking into account national recovery plans, threat abatement plans and conservation advice  <b>COM16:</b> The Queensland Government will ensure that fisheries are managed for the purpose of ecological sustainability, supported by the ongoing collection of commercial and recreational data through various monitoring programs
<b>REC32:</b> Maintain and improve monitoring, investigation and data management relating to critical species and habitats	

Authority recommended improvements	Queensland Government recommended improvements
<p><b>REC33:</b> Support implementation of a long-term social and economic monitoring program to improve understanding of changing use, investment and values</p>	<p><b>COM17:</b> The Queensland Government will incorporate reporting on matters of national environmental significance into Queensland State of the Environment reporting</p> <p><b>COM30:</b> The Queensland Government will work with the Australian Government, including the Authority, to establish an integrated monitoring framework and program for the Great Barrier Reef World Heritage Area as part of the Reef 2050 — Long Term Sustainability Plan</p> <p><b>COM31:</b> The Queensland Government will continue to work with industry and other stakeholders in Gladstone Harbour, through the Gladstone Healthy Harbour Partnership, to ensure open and accountable management of Gladstone Harbour, including annual reporting on ecosystem health and future actions underpinned by rigorous monitoring and science</p> <p><b>COM34:</b> The Queensland Government will continue to fund and support the Reef Water Quality Protection Plan and the associated Paddock to Reef monitoring program to help achieve the long-term goal of no detrimental impact from the quality of water entering the Great Barrier Reef. Consideration will be given to the inclusion of other pollutants, other than broadscale land use, during the Plan's next review in 2018</p>
<b>Governance</b>	
<p><b>REC34:</b> Contribute to the development of improved governance arrangements for the management and coordination of development activities that affect the Great Barrier Reef</p>	<p><b>COM26:</b> The Queensland Government will report to the Australian Government regarding proposed developments that may impact upon world heritage properties to ensure Australia's international obligations continue to be met</p> <p><b>COM27:</b> The Queensland Government will report annually to the Great Barrier Reef Ministerial Forum on implementation of the Reef 2050 — Long Term Sustainability Plan</p>
<p><b>REC34A:</b> Establish a peak Great Barrier Reef Advisory Group, made up of Traditional Owners, scientific, conservation and industry experts, to provide high level advice on the implementation of the Authority's management program</p>	
<b>Adapting to climate change</b>	
<p><b>REC35:</b> Communicate the implications of climate change impacts for the Great Barrier Reef, and the critical need to halt increasing concentrations of global greenhouse gases and restore them to levels that will support growth, recruitment and recovery processes of the Great Barrier Reef ecosystem</p>	
<p><b>REC36:</b> Ensure the impacts of climate change and extreme weather are appropriately considered in the Authority's management decisions</p>	
<p><b>REC37:</b> Encourage reduction of greenhouse gas emissions in the Great Barrier Reef Region in partnership with industry and communities</p>	
<p><b>REC38:</b> Support initiatives to build the capacity of management agencies and Reef users to adapt and respond to climate change and extreme weather events</p>	<p><b>COM28:</b> The Queensland Government will work with the Australian Government, including the Authority, to develop a Reef 2050 — Long Term Sustainability Plan for the Great Barrier Reef World Heritage Area by the end of 2014, and ensure its implementation</p>







## Chapter 13

## Adaptive management



13.6.2	Review and public reporting .....	13–34
13.6.3	Adapting to new information .....	13–34
13.6.4	Parties responsible for monitoring and reporting .....	13–35
<b>13.7</b>	<b>Addressing the endorsement criteria .....</b>	<b>13–35</b>
<b>13.8</b>	<b>Program review, modification and continuous improvement .....</b>	<b>13–42</b>
13.8.1	Review and continuous improvement of Program Report Part B .....	13–42
13.8.2	Review and continuous improvement of Program Report Part C .....	13–42
<b>References</b>	<b>.....</b>	<b>13–42</b>

## Table of contents

<b>13.1</b>	<b>Background .....</b>	<b>13–3</b>
<b>13.2</b>	<b>Comprehensive management program (Part B) .....</b>	<b>13–4</b>
<b>13.3</b>	<b>The permission system (Part C) .....</b>	<b>13–6</b>
13.3.1	Benefits of approving the permission system .....	13–6
13.3.2	Activities under the program .....	13–6
13.3.3	permission system objectives and outcomes .....	13–7
13.3.4	permission system process .....	13–8
13.3.5	Summary of program commitments .....	13–20
13.3.6	permission system alignment with the EPBC Act .....	13–21
<b>13.4</b>	<b>Applying the principles of ecologically sustainable development .....</b>	<b>13–25</b>
<b>13.5</b>	<b>Applying adaptive management ....</b>	<b>13–29</b>
13.5.1	Addressing risk .....	13–29
13.5.2	Addressing uncertainty .....	13–29
13.5.3	Implementing adaptive management to protect and manage matters of national environmental significance .....	13–30
13.5.4	Addressing World Heritage Committee recommendations .....	13–31
<b>13.6</b>	<b>Monitoring and reporting .....</b>	<b>13–34</b>
13.6.1	Monitoring .....	13–34

Cover page image: Releasing a turtle rehabilitated at Reef HQ Turtle Hospital



### **Extract from Great Barrier Reef Region strategic assessment terms of reference**

#### **6.2 Principles of ecologically sustainable development**

*Describe how the principles of ecologically sustainable development have been applied in the proposed program. The principles of ecologically sustainable development as described in section 3A of the Environment Protection and Biodiversity Conservation Act 1999 are:*

- a) decision-making processes should effectively integrate both long-term and short-term economic, environmental, social and equitable considerations*
- b) if there are threats of serious or irreversible environmental damage, lack of full scientific certainty should not be used as a reason for postponing measures to prevent environmental degradation*
- c) the principle of intergenerational equity — that the present generation should ensure that the health, diversity and productivity of the environment is maintained or enhanced for the benefit of future generations*
- d) the conservation of biological diversity and ecological integrity should be a fundamental consideration in decision making*
- e) improved valuation, pricing and incentive mechanisms should be promoted.*

#### **6.3 Adaptive management**

- a) identify the key adaptive management measures in the proposed program that address uncertainties and risks inherent in the decision-making process*
- b) describe how the adaptive management measures will be implemented to ensure the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area, are effectively protected and managed over the life of the program.*

#### **6.4 Monitoring and reporting**

- a) describe the monitoring, review and public reporting process that will be used to examine whether the program adequately protects and manages the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area, and the framework for measuring success*
- b) describe the processes for adapting the program in response to new information*
- c) nominate the parties responsible for undertaking the monitoring, review and reporting and for implementing any actions arising.*

#### **6.5 Review, modification or abandonment**

- a) identify and analyse likely circumstances and procedures that may result in the review, modification or abandonment of the program. This is to include a discussion of how any commitments under the program would continue to be met under these situations.*

## **13.1 Background**

The purpose of this chapter is to demonstrate how the Authority's program:

- provides for the protection of the Great Barrier Reef Region, including matters of national environmental significance
- applies the principles of ecologically sustainable development
- utilises an adaptive management approach
- conducts monitoring and reporting.

The final section focuses on program review, modification and continuous improvement.

The constantly changing nature of the Great Barrier Reef requires that the Authority's management program is sufficiently focused to consistently deliver results, yet agile enough to seamlessly shift and respond to issues when necessary. This adaptive management approach is embedded in the operations of the Authority, and forms a basis for our future program.

The Authority's future management arrangements are presented in three parts (Part A, B and C) of the Program Report. The program takes into account comments received from public consultation and independent reviews:

- **Part A** provides an overview of the Authority's program; describes the significance of the Great Barrier Reef; and outlines the strategic assessment process and key findings.
- **Part B** describes the Authority's comprehensive management arrangements and future commitments to protect and manage the environment, biodiversity and heritage values of the Region.
- **Part C** identifies the Authority's environmental assessment and decision-making process (the permission system) and explains how the Marine Park, including matters of national environmental significance, is protected.

## 13.2 Comprehensive management program (Part B)

The *Great Barrier Reef Marine Park Act 1975* (the GBRMP Act) and Regulations govern the protection and management of the Great Barrier Reef Region. The Act establishes an objectives-based regulatory framework, the main object of which is to provide for the long-term protection and conservation of the environment, biodiversity and heritage values of the Great Barrier Reef Region.

The other objects, so far as is consistent with the main object, are to:

- allow ecologically sustainable use of the Great Barrier Reef Region
- encourage engagement in the protection and management of the Great Barrier Reef Region
- assist in meeting Australia's international responsibilities in relation to the environment and protection of world heritage.

These objectives, together with the Authority's commitment to ensure activities carried out do not have an unacceptable impact by avoiding, mitigating and offsetting impacts on matters of national environmental significance protected under the EPBC Act, underpin the Authority's comprehensive management program.

To support attainment of the program's objectives, including the protection of world heritage values, the Authority has committed to the adoption of a comprehensive outcomes-based management approach. This framework explicitly identifies the desired outcomes for the Region's values including each matter of national environmental significance, and attributes and ecosystem processes relevant to matters of national environmental significance (see Chapter 4).

The life of the program is 25 years.

The Authority's comprehensive management program is built around three program areas:

- **Environmental regulation** — the Authority is the primary environmental regulator for the protection of the Great Barrier Reef. It utilises a range of spatially explicit planning instruments and system-based management tools such as Regulations, the Zoning Plan, plans of management, permits, compliance and statutory reporting.
- **Engagement** — the Authority recognises that protection of the Great Barrier Reef requires local, national and international effort. It works with Traditional Owners, the Australian and international community, business, industry and local government, together with Australian, Queensland and international government agencies to influence best practice and find pragmatic solutions to secure the future health of the Reef.
- **Knowledge, integration and innovation** — the Authority strives to access and capture the best available science from a network of science providers, both nationally and internationally, as well as drawing on traditional ecological knowledge and information from the wider community.

Based on the findings of the strategic assessment, the Authority has committed to a number of measures to strengthen its comprehensive management program. These are set out in Table 13.1.



**Table 13.1 Implementation of measures to strengthen management and forward commitments**

Over the next five years, the Authority will focus its management efforts on the implementation of a suite of measures to strengthen foundational management. Measures are designed to strengthen the protection and, where relevant, restoration of the Region's values including matters of national environmental significance.

	Within 12 months	Within two years	Within five years	Ongoing	Annually	Five-yearly
<b>New initiatives</b>						
Outcomes and targets	•					
Offset guidelines and net benefit policy	•					
Cumulative impact assessment guidelines	•					
Reef recovery program				•		
Integrated monitoring and reporting program			•			
<b>Measures to strengthen management</b>						
<b>Environmental regulation</b>						
Regionally-based standards for ecosystem health			•			
Dredging and dredge material disposal policy	•					
Strengthened guidelines (e.g. water quality)		•				
Measures to enhance alignment of permission system with EPBC Act assessment processes	•					
Streamlining, harmonising and enhancing management tools		•				
Improving assessment of matters of national environmental significance				•		
Further develop operational activities that support recovery and build resilience				•		
Improving certainty for planning and management			•			
Strengthening protection of heritage values			•			
Improving compliance				•		
Improving incident response capacity				•		
<b>Engagement</b>						
Influencing drivers and activities affecting the Region				•		
Supporting best practice and stewardship				•		
Improving consultation arrangements		•				
Establishment of a peak Reef advisory committee	•					
<b>Knowledge, innovation and integration</b>						
Improving identification and understanding of matters of national environmental significance				•		
Improving identification of Indigenous heritage values			•			
Developing an historic heritage database		•				
Improving understanding of community benefits			•			
Improving alignment and coordination of research priorities	•					
Increasing emphasis on use of modelling			•			
Better integration of knowledge into management			•			
<b>Forward commitments</b>						
Reef 2050 Long-Term Sustainability Plan	•					
Adaptive management — review cycles					•	•
Governance (Ministerial Forum)					•	

### 13.3 The permission system (Part C)

The Authority's permission system is established under the GBRMP Act. The GBRMP Act and its Regulations govern the protection and management of the Great Barrier Reef Region. The Act provides the Authority's head of power to regulate certain activities in the Region. In managing the Marine Park and carrying out its functions, the Authority must have regard to, and seek to act in a way that is consistent with, the objects of the Act, the principles of ecologically sustainable use and the protection of the world heritage values of the Great Barrier Reef World Heritage Area (subsection 7(3) of the GBRMP Act).

The permission system draws upon and operates within the context of the Authority's spatially explicit and systems-based regulatory tools.

The primary planning instrument under the GBRMP Act is the *Great Barrier Reef Marine Park Zoning Plan 2003* (the Zoning Plan). The Zoning Plan divides the Great Barrier Reef into eight zones, setting out which activities can occur in each zone. Each zone has specific conservation and management objectives for the protection of the environmental values of that zone.

The GBRMP Regulations provide for a system of permits which are required for specified activities likely to impact on the environmental values of each zone. For each permit application, the Authority undertakes an assessment of the nature and scale of activities, and makes a determination based on the acceptability of impacts from those activities on the environment.

As part of the strategic assessment process, the Authority is seeking approval of activities under the program in accordance with the provisions of Part 10 of the *Environment Protection and Biodiversity Conservation Act 1999* (the EPBC Act).

#### 13.3.1 Benefits of approving the permission system

The Marine Park is jurisdictionally complex, with overlapping tenures and responsibilities for management. Following a strategic assessment, the Minister for the Environment may endorse the program and approve actions, or classes of actions to be undertaken in accordance with the endorsed program. As part of the decision about whether to approve actions or classes of actions under the program, the Minister for the Environment must inform other Ministers with administrative responsibilities relating to the actions of the proposed approval, and invite comments including about economic and social matters relevant to those actions.

If approved, activities under the program are deemed to have been assessed and approved under the EPBC Act, and further consideration is not needed under this legislation.

A key benefit of the approval of actions taken in accordance with an endorsed program is the streamlining and reduced duplication of assessment processes while maintaining strong environmental safeguards.

The strategic assessment found that one of the strengths of the Authority's current management arrangements was that a focus on the Marine Park as a matter of national environmental significance means that values relevant to other matters of national environmental significance are implicitly considered in decision making. The implementation of recommendation 1 will further improve this approach:

REC1: Explicitly incorporate consideration of all matters of national environmental significance, including attributes of the property's outstanding universal value, into the Authority's programs, plans and policies.

The strategic assessment also found that although the Intergovernmental Agreement (between the Australian and Queensland governments) provides a strong foundation for complementary joint permit assessment and approval processes, there is currently significant duplication in assessment processes across jurisdictions. The complexity of processes presents significant challenges to public engagement in decision making. An approval of actions under Part 10 of the EPBC Act would support recommendation 8 to:

REC8: Streamline assessment processes across jurisdictions and seek to have a more coordinated approach to community consultation.

#### 13.3.2 Activities under the program

The Great Barrier Reef Marine Park is a multiple-use area. It provides for a range of ecologically sustainable commercial, recreational and research opportunities and for the continuation of traditional activities.

The GBRMP Act includes provisions which **prohibit mining** operations (which includes prospecting or exploration for, as well as recovery of, minerals) in the Great Barrier Reef Region, unless authorisation to carry out the operations is granted by a permission under the Regulations, for the purpose of research or investigations relevant to the conservation of the Marine Park.

Approval of activities under the program will be subject to alignment with any approval bilateral agreement under Part 5 of the EPBC Act that may be in effect for the program area.

An updated assessment bilateral agreement between the Australian and Queensland governments was signed on 13 December 2013 through which the Queensland Government is accredited to undertake environmental assessments of any proposed actions that may impact the Great Barrier Reef. In addition, the two governments

are negotiating an approval bilateral agreement to accredit Queensland Government processes for assessment and approval of proposed actions under the EPBC Act. This agreement aims to reduce duplication of environmental assessment and approval processes between the Commonwealth and Queensland, while maintaining high environmental standards.

The intent of the Authority seeking approval of classes of action is also to maximise opportunities for streamlining and reducing potential regulatory duplication, while maintaining high environmental standards.

Activities covered under the permission system are:

- all activities requiring a permission under the Zoning Plan
  - aquaculture operations
  - harvest and development fisheries
  - research (other than limited impact research)
  - tourism programs and developments
  - educational programs (other than limited impact educational programs)
  - vessel or aircraft charter operations
  - navigating a managed vessel or aircraft
  - operating a facility, including
    - discharging waste from a facility
    - installation, operation and decommissioning of a facility
    - moorings
    - operating a landing area or a facility for aircraft
  - carrying out works, including:
    - dredging
    - dumping of spoil
    - reclamation
    - beach protection works
    - harbour works
  - taking animals and plants that pose a threat to human life or safety, marine ecosystems of the Marine Park or use or amenity of a part of the zone or adjacent area.

The activities under Part 5 of the Zoning Plan are not intended to be covered by this program.

The Authority will continue to implement joint management arrangements for the area of shared jurisdiction with the Queensland Government and in accordance with the Great Barrier Reef Intergovernmental Agreement.

If the Authority receives a permit application which is likely to have a significant impact on a matter of national environmental significance which is not covered by the above classes of activities in the strategic assessment, the Authority will provide applicants with the earliest possible advice of this and direct them to the EPBC Act assessment pathways administered by the Department of the Environment, or the Queensland Government where an accredited process exists or delegated authorities.

### **13.3.3 Permission system objectives and outcomes**

The objectives of the permission system are that all activities permitted under the permission system must be carried out in a way that ensures the impacts to the Marine Park, including matters of national environmental significance, and the broader environment, are of an acceptable level and are carried out in a manner that is consistent with the objectives of the GBRMP Act and the principles of ecologically sustainable use. If a proposed project or activity would have an unacceptable impact on the Marine Park, including a matter of national environmental significance, the Authority would not grant a permit for the project or activity to proceed under the permission system.

The objective of the Permission system is to ensure the values of the Region are transmitted in good and very good condition. In assessing and managing impacts, the approach to date has been to avoid, mitigate and offset significant residual impacts. This would be sufficient if the attributes critical to the Marine Park, including matters of national environmental significance, were in good or very good condition and the ecosystem's ability to recover from disturbance was high.

However, the majority of these attributes in the inshore southern two-thirds of the Region have been assessed to be in either poor or declining condition, and there is increasing evidence that the ecosystem's ability to recover from disturbance is being lost. Furthermore, the strategic assessment found that it is not only significant residual impacts of development actions, but the cumulative effect of all impacts which are affecting the condition of the Marine Park, including matters of national environmental significance and relevant attributes and environmental processes.

Consequently, the Authority's management program includes measures to 'offset' all residual impacts in order to maintain the current condition of attributes and environmental processes supporting matters of national environmental significance. Furthermore, as many of the attributes and environmental processes have been assessed to be in poor or declining condition, additional measures are proposed to not only 'offset' residual impacts but to deliver an overall net improvement in the condition of the Marine Park.

The matters relevant to this strategic assessment and the program's overarching commitment to outcomes of environmental protection are outlined in Table 13.2.

**Table 13.2 The Authority's commitment to safeguard matters of national environmental significance**

Matter of national environmental significance	Outcome
All matters of national environmental significance	Matters of national environmental significance are protected and conserved.
World heritage properties	The outstanding universal value of the Great Barrier Reef World Heritage property is identified, protected, conserved, presented and transmitted to future generations.
Great Barrier Reef Marine Park	The outstanding universal value of the Great Barrier Reef Marine Park, as a world heritage property, is identified, protected, conserved, presented and transmitted to future generations. The environmental, biodiversity and heritage values of the Great Barrier Reef Marine Park are protected and conserved for the long term, consistent with the objects of the Act.
National heritage places	The outstanding value to the nation of national heritage properties is identified, protected, conserved, presented and transmitted to future generations of Australians.
Commonwealth marine areas	The ecosystem functioning and integrity of Commonwealth marine areas are maintained or enhanced in full conformity with relevant marine bioregional plans.
Listed migratory species	The survival and conservation status of listed migratory species and their critical habitat is promoted and enhanced, consistent with Australia's international obligations.
Listed threatened species and ecological communities	The survival and conservation status of listed threatened species and ecological communities is promoted and enhanced, including through the conservation of habitat critical to the survival of a species or community and other measures contained in any recovery plans, threat abatement plans or conservation advices.
Wetlands of international importance	The ecological character of each Ramsar wetland is maintained, and the conservation and wise and sustainable use of the wetland is promoted for the benefit of humanity in a way that is compatible with maintenance of the natural properties of the ecosystem. This is to be achieved through the implementation of ecosystem approaches, within the context of sustainable development.

#### 13.3.4 Permission system process

The Authority assesses permit applications against the requirements of the permission system. The Authority is obliged to take into account all relevant considerations in exercising its decision-making power. Relevant considerations include:


- the Authority's environmental regulation requirements
- the additional commitments outlined in the program to maintain and enhance protection of matters of national environmental significance
- the Authority's commitments to consider and manage impacts on matters of national environmental significance to be implemented via the permission system as set out in Table 10 of the Program Report.

Under the GBRMP Act the protection and conservation of the environment, biodiversity and heritage values of the Great Barrier Reef Region takes primacy over social and economic considerations. Consequently, the decision framework is governed by the requirement to first protect the environment and matters of national environmental significance and, second, to support sustainable use.



The Authority's assessment under the permission system involves experienced assessment teams and specialists evaluating applications with reference to all relevant internal and external sources of information. This includes applicable legislation, scientific reports, international agreements, management plans and relevant policies, guidelines and standards. When permit applications relate to matters of national environmental significance that occur in the Marine Park, the Authority's assessments have regard to and are undertaken consistently with the EPBC Act's statutory and policy documents related to those matters of national environmental significance.

Each permit application is subject to an assessment in line with Authority policy and includes scoping, assessment, decision, and audit and compliance steps as outlined in Figure 13.1.



**Figure 13.1 The four key steps in the permission assessment and decision process**

Examples of how application of the permission system protects matters of national environmental significance are provided in Appendix 7 – Permission system case studies.

An overview of how the permission system ensures the appropriate level of consideration and management of impacts for each matter of national environmental significance is provided in Table 13.3. This table outlines how the steps in the Authority's permission system and the program's commitments align with the EPBC Act assessment process.

**Table 13.3 Steps in the Authority's permission system and EPBC Act assessment and decision-making process**

Permission system — step components	Description	Authority mechanism	EPBC Act, MNES
<b>Step 1: Scoping phase</b>			
<b>Initial inquiry</b>	<p>An applicant may approach the Authority to discuss initial concepts, designs, alternatives and potential mitigation of impacts for any proposed activity or development which may impact on the Great Barrier Reef Marine Park and require a permit.</p> <p>Where permits may be required from other government agencies, the Authority recommends a whole-of-Government meeting be held to identify the scope of the proposed activity. Discussions will include consideration of matters of national environmental significance (MNES). The Authority has committed to developing guidance for applicants which explain requirements relating to MNES. The guidance will assist applicants in meeting the information requirements necessary for MNES.</p>	<p>The Authority's Environmental Impact Management Policy.</p> <p>Under the program, the Authority will:</p> <ul style="list-style-type: none"> <li>• develop guidelines to assist applicants to determine whether an action is likely to have significant impact on the Great Barrier Reef Marine Park MNES. The Authority will seek to have these guidelines approved by the Minister.</li> </ul>	<p>Equivalent to EPBC Act, 'pre-referral meeting' where regulatory arrangements are explained to applicants.</p> <p>MNES and outstanding universal value (OUV) will be specifically included in Authority programs, plans and policies.</p>
<b>Tests of 'significant impact'</b>	<p>The Authority will develop and implement referral guidelines to assist in the determination on whether a proposed action is likely to have a significant impact on the Great Barrier Reef Marine Park and other matters of national environmental significance.</p> <p>The Authority will update its policy documents for permittees and internal assessment officers to ensure relevant EPBC Act guidelines are considered.</p>	<p>GBRMP Act section 7(4), policy documents.</p> <p>Under the program, the Authority will require:</p> <ul style="list-style-type: none"> <li>• consideration of relevant EPBC Act policy documents by applicants, including relevant significant impact guidelines, in the scoping phase of the permission assessment and decision process</li> <li>• consideration of prudent and feasible alternatives, which may have a lower impact on the environment by applicants, and amend its policy documentation to include reference to this requirement.</li> </ul>	<p>The EPBC Act advisory documents include significant impact guidelines for MNES, and OUV of the Great Barrier Reef World Heritage Area.</p>
<b>Permit application</b>	<p>An applicant must complete a Permit Application Form and pay a Permit Application Assessment Fee.</p> <p>In order to be granted a permit to undertake an activity, the applicant is required to establish that the environmental impacts of the proposed activity are acceptable.</p> <p>The application must provide sufficient information to enable a clear understanding of the intended activity.</p> <p>The Authority undertakes the following processes at this stage:</p> <ul style="list-style-type: none"> <li>• The permit application is lodged with the Authority in both hard form and electronically.</li> <li>• The application undergoes an initial check to ensure it meets minimum legal requirements (including EPBC Act</li> </ul>	<p>GBRMP Regulation 88A (standard permission) and Division 2A.3 (special permissions) require applications to be made in a form approved by the Authority.</p> <p>Fees are set out in GBRMP Regulation 128.</p> <p>Under the program, the Authority will amend its Permit Application Form to include a section which specifically addresses MNES.</p>	<p>MNES and OUV will be included within Authority programs, plans and policies.</p>

Permission system — step components	Description	Authority mechanism	EPBC Act, MNES
	<p>requirements relating to MNES).</p> <ul style="list-style-type: none"> <li>• A determination of the Permit Application Assessment Fee is undertaken.</li> <li>• An acknowledgement letter is sent to the applicant; this includes details of assessment procedures and likely timeframes for assessment.</li> <li>• A native title notification is sent if required.</li> </ul>		
<b>Step 2: Assessment phase</b>			
<b>Allocation of assessment level — nature and scale</b>	<p>Permit applications are allocated to an assessment level, depending on the scale of the project, the degree and duration of impacts and other factors such as public interest and sensitivity of the environment.</p> <p>The level of complexity of the permit is determined against the Authority's Environmental Impact Management Policy by the delegate. The level of assessment is also determined at this stage, for example by an Environmental Impact Statement (EIS).</p> <p>The level of permit assessment is determined based on the nature and scale of the activity which is proposed. Considerations may include, but are not limited to:</p> <ul style="list-style-type: none"> <li>• The size of the project.</li> <li>• The duration/timeframe of the project.</li> <li>• The adequacy and completeness of the information in the permit on the relevant impacts of the action.</li> <li>• The duration (temporary or ongoing) and reversibility of impacts.</li> <li>• The current environmental situation at the action location and/or area of impact.</li> <li>• Consideration of outcomes if approved in its current form.</li> <li>• Consideration of potential to carry out activity in a manner that reduces impacts.</li> <li>• The level of knowledge and certainty of involved risks (and how to improve knowledge).</li> <li>• The environmental elements that are likely to be affected by the activity.</li> <li>• The extent to which potential relevant impacts have already been or will be assessed under Queensland Government processes (including through the local or state government planning process).</li> <li>• The degree of public concern associated with the proposal, or with similar proposals having comparable likely impacts on the environment.</li> <li>• What further information is required from the applicant? Are there gaps in information or insufficient information to provide a high level of certainty about the project?</li> <li>• An initial assessment of the costs and benefits, including environmental costs and benefits, to the community and the</li> </ul>	<p>The Authority's Environmental Impact Management Policy.</p> <p>Under the program, the Authority will:</p> <ul style="list-style-type: none"> <li>• update the Great Barrier Reef Marine Park Regulations 1983 to explicitly outline the levels of assessment under the permission system</li> <li>• develop guidance material to outline the levels of assessment and the general requirements for applicants.</li> </ul>	<p>The EPBC Act enables the identification of whether approval of an action is required.</p> <p>Actions likely to have a significant impact on a MNES protected under the EPBC Act would fall into Level 3 or 4 where options include EIS and Public Environmental Review assessments.</p>

Permission system — step components	Description	Authority mechanism	EPBC Act, MNES
	<p>person proposing to take the action of further data collection about, and analysis of, the relevant impacts of the proposal.</p> <ul style="list-style-type: none"> <li>• What type/level of monitoring would be required to ensure compliance?</li> <li>• Would an Environmental Management Plan be required to implement and manage onsite impacts?</li> <li>• Would an Advisory Committee be required to advise on environmental impacts?</li> <li>• Is an Environmental Site Supervisor required to supervise environmental controls?</li> <li>• Would a deed of agreement need to be entered into for the protection of the environment, and would a bond be required to be paid as part of the deed of agreement to cover the risks associated with the activity?</li> </ul> <p>The assessment level is chosen on the basis that it will provide the information needed to satisfactorily answer the assessment questions. Larger scale projects with longer timeframes and a greater nature and scale of impacts will require more detailed assessment.</p> <p>After allocation, all permits are immediately registered on the permit assessment database.</p> <p>Indicatively, the level of environmental impact management tools utilised are as follows:</p> <ul style="list-style-type: none"> <li>• Level 1 assessments may require compliance monitoring.</li> <li>• Level 2 assessments may require a site inspection, standard Deed and a Bond.</li> <li>• Level 3 assessments may require a site inspection (and issues-based site inspections), an Environmental Management Plan (and an Advisory Committee), a standard Deed and a Bond.</li> <li>• Level 4 assessments may require a site inspection (and issues-based site inspections, and reactive site inspections and monitoring consultant(s)), an Environmental Management Plan (and an Advisory Committee and Environmental Site Supervisor), a specific Deed, and a Bond.</li> </ul> <p>All activities with a likely significant impact on MNES will undergo a Level 3 or 4 assessment (or equivalent) which requires an EIS or PER be prepared and assessed for the activity.</p> <p>Deeds are required for permitted commercial activities, and establish terms and conditions related to the permit, and identify indemnities, bonds and insurance requirements.</p> <p>Bonds seek to ensure that costs associated with rectification of environmental damage are borne by the operator.</p>		



Permission system — step components	Description	Authority mechanism	EPBC Act, MNES
<b>Permit assessment</b>	<p>Equivalent to the EPBC significant impact test, a permit assessment determines whether EPBC referral would be triggered by the actions proposed in the Permit Application.</p> <p>The environmental management tools and permit conditions generally become more robust as the assessment level scale increases.</p> <p>Permits may be assessed solely on the permit documentation. This type of assessment is typical for smaller scale assessments. Activities with significant impacts on MNES would not be assessed at this level.</p> <p>During the assessment process, a Public Environment Report may be requested to ensure sufficient information on the impacts of the action on MNES is available to inform decision-making.</p> <p>A Public Environment Report has the same meaning as within Division 5, Part 8 of the EPBC Act.</p>	<p>The Authority's Environmental Impact Management Policy.</p> <p>Each MNES will be identified separately in the assessment process.</p> <p>GBRMP Regulation 88A.</p> <p>GBRMP Regulations 88E and 128(3).</p> <p>The Authority's Environmental Impact Management Policy.</p> <p>Each MNES will be explicitly identified and impacts on them must be explicitly assessed.</p>	<p>The EPBC Act provides for different levels of assessment of controlled actions, commensurate with the nature and scale of the impacts on EPBC protected matters.</p> <p>Smaller scale assessments are the equivalent to assessment of a referral under Part 7 of the EPBC Act where the activity is not determined to have a significant impact on a MNES.</p> <p>Division 5, Part 8 of the EPBC Act.</p>
<b>Environmental Impact Statement</b>	<p>During the assessment process, an EIS may be requested to ensure sufficient information on the impacts of the action on MNES is available to inform decision-making.</p> <p>An EIS has the same meaning as within Division 6, Part 8 of the EPBC Act.</p>	<p>GBRMP Regulations 88E and 128(4).</p> <p>Each MNES will be explicitly identified and impacts on them must be explicitly assessed.</p>	<p>Division 6, Part 8 of the EPBC Act.</p>
<b>Consultation</b>	<p>During the assessment process, the Permit Application may be referred to relevant parties for comment (for example applications with the potential to impact fisheries would be referred to the Commonwealth and State departments responsible for management of fisheries).</p> <p>Applications that may have a significant impact on MNES will be available for public consultation via the Authority's website.</p> <p>All applications with significant impacts will be publically advertised.</p>	<p>The Authority's Environmental Impact Management Policy.</p> <p>Internal procedures and guidelines.</p> <p>Authority's website.</p> <p>Relevant policies will be updated to explicitly state that permit applications for an action where granting a permission may restrict the reasonable use by the public of a part of the Marine Park will be advertised for public comment through either EIS or PER assessment.</p> <p>Relevant policies will be updated to explicitly state that appropriate consultation will be undertaken with persons or organisations, including Australian and Queensland government agencies, whose functions, interests or activities could be impacted by the proposed activity.</p>	<p>CI74 of Division 1, Part 7, of the EPBC Act.</p>
<b>Advertising application</b>	<p>During the assessment process, the permission system provides for public notification and consultation where an action may restrict the reasonable use by the public of a part of the Great Barrier Reef Marine Park.</p>	<p>GBRMP Regulation 88D provides for the advertising of applications.</p> <p>The Authority will publish notification of all permit applications on the internet</p>	<p>Divisions 4, 5 and 6 Part 8 of the EPBC Act.</p>

Permission system — step components	Description	Authority mechanism	EPBC Act, MNES
	All activities likely to have a significant impact on a MNES will be advertised. Consultation will occur on EIS and PER documents for no less than 30 days.	as soon as practicable after receiving a permit application.	
<b>Mandatory and discretionary criteria</b>	<p>The Authority will make decisions consistent with the Objects of the GBRMP Act which includes the main object at section 2A(1) to 'provide for the long-term protection and conservation of the environment, biodiversity and heritage values of the Great Barrier Reef Region'. This necessarily includes MNES.</p> <p>During the assessment process, the permit application is assessed in accordance with mandatory and discretionary criteria outlined within the GBRMP Regulations. These include relevant Australian and Queensland Government environmental management laws, and conservation advices.</p>	<p>GBRMP Regulations 88Q (mandatory considerations) and 88R (discretionary considerations).</p> <p>Some activities (for example the taking of Protected Species, 88S) require additional criteria to be met.</p> <p>Assessments of impacts will have regard to the full range of MNES information sources.</p> <p>Under REC1 of the Great Barrier Reef Strategic Assessment Report, the Authority has committed to explicitly incorporate consideration of all values relevant to matters of national environmental significance, including elements of the property's outstanding universal value, into the Authority's programs, plans and policies</p>	Subdivision b, Division 1, Part 9 of the EPBC Act.
<b>Assessment of risk</b>	<p>The Authority's Environmental Impact Assessment Policy and Environmental Assessment and Management Risk Assessment Guidelines (developed in accordance with the Australian Standards Environmental Risk Management Handbook HB203:2006) provide guidance in the assessment of risk, including analysis of the likelihood and consequence of identified hazards being realised to obtain a Hazard Risk Grade.</p> <p>During the assessment process, hazards are identified and a risk assessment (of consequence and likelihood) is determined.</p> <p>At a minimum, a Risk Profile is completed to document the risk assessment process. The risk assessment process requires sufficient information to be included in order to enable a decision to be made on each MNES, which will be individually identified in the assessment.</p> <p>Additional risk management instruments utilised with permits include Deeds and Bonds.</p>	<p>GBRMP Regulation 88Q(a)</p> <p>Where an action is likely to have a significant impact on a matter protected under the EPBC Act within the Marine Park, the Authority will:</p> <ul style="list-style-type: none"> <li>• require applicants to consider the impacts of actions on MNES and relevant attributes and environmental processes as described in the Strategic Assessment Report and as amended in accordance with the Authority's policy review procedures</li> <li>• require applicants to consider impacts (including direct, indirect and cumulative impacts) arising from the action in the context of all impacts affecting the matter</li> <li>• consider any relevant international conventions and Australian-Queensland Government agreements in assessing and determining permissions under its permission system</li> <li>• consider all relevant plans, policies, documents and guidance</li> </ul>	

Permission system — step components	Description	Authority mechanism	EPBC Act, MNES
		(as specified in Table 10 of the Program Report) in assessing and determining permissions under its permission system.	
<b>Step 3: Decision phase</b>			
<b>Acceptable level</b>	<p>Applicants must demonstrate that the environmental impacts and risks of the activity will be of an acceptable level.</p> <p>The Authority will update its policy documents to provide guidance on the key considerations for determining ‘acceptability’ of direct, indirect and cumulative impacts on MNES.</p> <p>The Authority will comply with the commitments in Table 10 of the Program Report.</p>	<p>GBRMP Act Section 7(4), policy documents.</p> <p>Under the program the Authority will develop and implement:</p> <ul style="list-style-type: none"> <li>• guidelines for the application of Great Barrier Reef offsets to maintain or improve the condition of MNES and relevant attributes and environmental processes, where impacts cannot be avoided or mitigated. The guidelines will seek to deliver an outcome equivalent to, or better than, the outcome that would apply if the EPBC Act Environmental Offsets Policy was applied</li> <li>• a Great Barrier Reef net benefit policy (that will be updated from time to time) to enhance the condition of the Marine Park, including MNES and relevant attributes and environmental processes.</li> </ul>	Includes, but is not limited to, an evaluation of whether an activity has, will have or is likely to have unacceptable impacts on MNES.
<b>Granting and refusing permissions</b>	<p>The Authority may grant or refuse a permission based on the assessment. Permission may be granted with any conditions appropriate to the attainment of the objects of the GBRMP Act and the protection of MNES.</p> <p>The Authority will not grant a permit for an activity inconsistent with regulatory objectives set out under the GBRMP Act, as outlined in Table 10 of the Program Report.</p>	<p>GBRMP Regulations 88X and 88ZE.</p> <p>The Program’s commitments to MNES (see Table 10 of the Program Report)</p> <p>The Authority will strengthen its system for monitoring and managing compliance risks associated with permit conditions. It will review and make any necessary improvements to internal policies, procedures, training protocols and mechanisms to systematically enhance the effectiveness of its compliance and auditing process. This will include specific reference to MNES.</p>	Part 9 of the EPBC Act.

Permission system — step components	Description	Authority mechanism	EPBC Act, MNES
<b>Step 4: Audit and compliance phase</b>			
<b>Permitted activities</b>	An activity that has been granted permission must be carried out in accordance with the GBRMP permit and any conditions made. Conditions will be applied to ensure an unacceptable impact does not occur on MNES.	GBRMP Regulations, Division 2A.5 and 2A.6.	
<b>Environmental Management Plans</b>	For actions which may have a significant impact on the Marine Park, including MNES, an Environmental Management Plan (EMP) will be required to specify the strategies to minimise the potential impacts.  EMPs generally include the identification of environmental impacts, how on-water activities will be managed to reduce these impacts, a monitoring program, emergency response plans and any relevant issue-based plans.  The EMP is developed in close consultation with the Authority's assessing officer and may be modified throughout the assessment phase.	The Authority's Environmental Impact Management Policy will be updated to ensure EMPs address MNES.	
<b>Environmental Site Supervisor</b>	The Environmental Site Supervisor and Monitoring Consultant must be independent of (with no conflicts of interest), and funded by the applicant.  The Authority will select and directly contract the Environmental Site Supervisor and Monitoring Consultant. These positions may be engaged for the whole project or any relevant phases. Supervisors have a direct 'on the ground' role in ensuring activities are carried out in accordance with conditions and EMPs. Supervisors will additionally consider MNES under the endorsed program. Supervisors would be appointed where there is a higher risk of activities to MNES, to ensure mitigation measures are applied.	The Authority's Environmental Impact Management Policy.	MNES and OUV will be specifically included in Authority programs, plans and policies.
<b>Advisory Committee</b>	An Advisory Committee may be established for a project to provide advice to the Authority. The Advisory Committee may include representatives from government, the applicant, community, the Environmental Site Supervisor, and experts in the fields of science and management. Under the program, MNES is an explicit consideration. Advisory Committees in future may include expertise specifically relevant to MNES.	The Authority's Environmental Impact Management Policy.	MNES and OUV will be specifically included in Authority programs, plans and policies.
<b>Performance outcomes and performance standards</b>	The Authority will audit environmental performance of some permitted activities and developments and, if environmental performance or compliance does not meet required standards, then penalties and/or remediation may be required. MNES will be included in audits following the approval of the program.	The Authority's Environmental Impact Management Policy.  The Authority's Outcomes Framework.	Environmental performance outcomes and standards must take account of all relevant information, including (but not limited to) management guidance, standards relevant to MNES under the EPBC Act, and the program's commitments in Table 10 of the Program



Permission system — step components	Description	Authority mechanism	EPBC Act, MNES
			Report. MNES and OUV will be specifically included in Authority programs, plans and policies.
<b>Compliance</b>	The Authority will determine the need for a Bond and/or a Deed to provide ongoing indemnity and, where necessary, to ensure there is an ongoing contractual obligation to protect the environment if the permit lapses. These would be applied where there is a higher risk of activities to MNES.	GBRMP Act, Part VAA The Joint Field Management Program (Great Barrier Reef Intergovernmental Agreement – Schedule C) undertakes compliance activities through both education and enforcement in order to encourage adherence to legal requirements.	If environmental performance or compliance does not meet required standards for protection of MNES and OUV, penalties and/or remediation will be required.
<b>Incident response</b>	Increasing use of the Region and the likelihood of increased extreme weather events will require the Authority and its management partners to have a strong capacity to respond to incidents. MNES will be relevant considerations of any incident response measures.	The Authority has a cooperative compliance management and surveillance program involving Australian and Queensland government agencies. These include the Queensland Department of National Parks, Recreation, Sport and Racing; Border Protection Command; Queensland Boating and Fisheries Patrol; Queensland Police Service; Commonwealth Director of Public Prosecutions; Australian Federal Police; Australian Maritime Safety Authority; and Maritime Safety Queensland.	A partnership approach with the Australian and Queensland governments enables the Authority to promote protection of MNES and OUV to a broader monitoring group, and deliver greater incident management coverage for the Region.

An overview of how the permission system permit application content requirements align with the EPBC Act assessment requirements is described in Table 13.4.

**Table 13.4 Content requirements for Permit Applications**

Required information	Content requirements
<b>Details of the applicant</b>	In order to be granted a permit to undertake an activity, an application must provide details of the applicant, including a history of environmental performance regarding matters of national environmental significance.
<b>Summary of the proposal</b>	<p>GBRMP Regulation 88A (standard) and Regulation 2A.3 (applications for special permissions) require applications to be made in a form approved by the Authority.</p> <p>In order to be granted a permit to undertake an activity, an application must provide enough information so permit assessors can clearly understand what the intended activity is.</p> <p>The application must contain a description of each of the activities that will be carried out, including a description of works to be undertaken and facilities to be used to carry out the activity, the location of the activities, and proposed timetables for carrying out the activities.</p> <p>The application forms will include a section to address impacts on matters of national environmental significance to assist with the assessment.</p>

Required information	Content requirements
<b>Description of prudent and feasible alternatives considered</b>	The Authority recommends applicants and interest groups to discuss initial concepts, designs, alternatives and potential mitigation of impacts for any proposed activity or development which may impact on the Great Barrier Reef Marine Park and require a permit. Alternatives to activities with a significant impact on matters of national environmental significance will be required.
<b>Description of the existing environment that may be affected by the proposal</b>	<p>The application must contain a description of the characteristics of the existing environment that may be affected by the proposal and details of sensitivities of that environment, including attributes and environmental processes relevant to matters protected under the EPBC Act.</p> <p>The description should cover the zone of influence likely to be affected by the proposed activity and must be sufficient to inform consideration of the nature and scale of the impacts by the Authority in its assessment (for example, species/populations that may potentially be affected by upstream and downstream impacts such as water quality or noise effects).</p>
<b>Description of sensitivities</b>	The application must contain a description of sensitivities of the impacted environment, including matters of national environmental significance protected under the EPBC Act and relevant attributes and environmental processes.
<b>Description of environmental impacts and risks</b>	<p>Chapter 6 of the Strategic Assessment Report describes the impacts of activities on values. Consistent with the Authority's Environmental Impact Management Policy, applicants need to consider direct, indirect and cumulative impacts appropriate to the nature and scale of activities for which permission is sought. If a proposal is regarded as having the potential for significant environmental impact under the EPBC Act, then applicants are advised inclusion of an Environmental Impact Statement or Public Environment Report may be required.</p> <p>The Authority has committed to specifically including matters of national environmental significance and outstanding universal value in Authority programs, plans and policies.</p>
<b>Evaluation of impacts (direct, indirect and cumulative) and risks</b>	<p>In order to be approved, an application for an activity under the permission system is required to contain an evaluation of all the impacts, including direct, indirect and cumulative impacts and risks, including relative level of certainty.</p> <p>The application must further demonstrate that impacts and risks are consistent with the desired outcomes for the protection of matters of national environmental significance. Assessment guidelines will include information on condition and trend of all matters of national environmental significance.</p> <p>Chapter 7 of the Strategic Assessment Report assesses the current condition and trend for the Region's values and processes. Part B of the Program Report outlines the comprehensive management program based on outcomes and targets.</p> <p>The desired outcome for each value is informed by the assessment of its current condition and trend as evaluated in the strategic assessment. This approach recognises that the condition and trend of each value and process varies across the Region.</p> <p>The outcome for each value and process is a combination of the strategic assessment findings on condition and trend, and the international and national obligations to protect them.</p> <p>For example, if a value is assessed to be good condition, the outcome is that its condition is maintained and enhanced. However, if a value is assessed to be in poor condition, the outcome is that its condition is restored to good condition. If the trend of the value is assessed to be declining, the outcome is that the decline is halted and reversed.</p> <p>There will be improvement in the collection, management, analysis and interpretation of information derived from the Authority's permission system and compliance monitoring.</p>
<b>Performance outcomes, standards and measurement criteria</b>	<p>The setting of environmental performance outcomes and performance standards must take into account all relevant information which includes, but is not limited to, management guidance and standards relevant to all EPBC Act protected matters.</p> <p>These have to be benchmarked against regulatory objectives, outcomes for matters of national environmental significance and relevant GBRMP Act and EPBC Act guidelines and standards.</p> <p>An Advisory Committee may be established for a project to provide advice to the Authority. The Advisory Committee may include representatives from government, the applicant, community, the Environmental Site Supervisor, and experts in the fields of science and management.</p> <p>The Authority will determine the need for a cash bond or bank guarantee and a Deed of Agreement to provide ongoing indemnity and, where necessary, to ensure there is an ongoing contractual obligation to protect the environment if the permit lapses. Civil</p>

Required information	Content requirements
	<p>penalties are available for offences.</p> <p>The Authority will audit environmental performance of some permitted activities and developments and, if environmental performance or compliance does not meet required standards, then penalties and/or remediation may be required.</p> <p>MNES and OUV will be specifically included in Authority programs, plans and policies.</p>
<b>Description of environmental monitoring and reporting standards</b>	<p>For certain activities, including any that are likely to have a significant impact on MNEES, the Authority requires development of an Environmental Management Plan (EMP). Where an EMP is required for implementation of an activity it is set out as a condition of a Marine Parks Permit.</p> <p>As a minimum, the EMP needs to address:</p> <ul style="list-style-type: none"> <li>• what values of the Marine Park may be impacted by the activities</li> <li>• how the applicant will manage actions to minimise those impacts</li> <li>• what the applicant will do to ensure that all staff and clients know what they should be doing to meet the requirements of the EMP, including who is the responsible person for key actions</li> <li>• how it will be determined if actions are working in accordance with the conditions of the EMP and Permit conditions</li> <li>• how adaptive management principles will be applied to improve the EMP.</li> </ul> <p>At a minimum, the EMP elements should include:</p> <ul style="list-style-type: none"> <li>• <i>Activity or Issue</i> – Description of the activity being managed or considered.</li> <li>• <i>Potential Impact/s</i> - Description of the potential effects of the activity on all environmental issues.</li> <li>• <i>Aim</i> — What the EMP element hopes to achieve for environmental issues.</li> <li>• <i>Management strategies</i> — How all activity will be managed to achieve the aim</li> <li>• <i>Performance indicators</i> — What will be measured to show that the aim is being met</li> <li>• <i>Responsibility</i> — Who will be the person nominated to manage this element</li> <li>• <i>Monitoring and Reporting</i> - How and when the performance indicators will be measured to test whether the aim has been achieved</li> <li>• <i>Corrective Action</i> — The action to be taken and by whom, if a performance requirement is not met</li> <li>• environmental performance standards and measures for the activities to be carried out, benchmarked against regulatory objectives, outcomes for matters of national environmental significance and relevant GBRMP Act and EPBC Act guidelines and standards</li> <li>• environmental monitoring and reporting standards to be employed to measure environmental performance, appropriate to the type of activities.</li> </ul>
<b>Legislative requirements</b>	<p>Permit applications will be considered on a case by case basis, with consideration given to the objectives of the zone in which the proposed conduct will take place, as set out in the Zoning Plan.</p> <p>In making decisions that are consistent with the Objects of the GBRMP Act, the Authority must consider the potential impacts of the conduct proposed to be permitted by the permission on the environment and on the social, cultural and heritage values of the Marine Park or a part of the Marine Park.</p> <p>Content of the EMP must be consistent with the nature and scale of the activity, and have regard to the mandatory and discretionary considerations under GBRMP Regulations 88Q and 88R.</p>
<b>Additional requirements</b>	<p><i>Deed of Agreement</i> — Once a permit is issued, depending on the type and scale of the proposed activity, it is likely to be subject to a Deed of Agreement to cover the risks associated with the proposed activity.</p> <p><i>Bond</i> — If a bond is required, the amount determined will be specified in the Deed of Agreement. The permittee will be required to secure with the Authority a bank guarantee for the bond amount.</p> <p><i>Insurance</i> — The Authority sets out minimum insurance requirements under the Deed of Agreement. The Authority recommends permittees seek independent insurance advice concerning the level and type of cover that is appropriate for its specific operation and its associated activities.</p> <p><i>Site Supervision</i> — Site supervision is an essential component of the management of projects that have the capacity to have significant impacts on the Marine Park and its users. It is also used as a tool to enable the flexible management for certain activities (for example, filming to provide access to or allow activities at a sensitive location in the presence of a nominated site supervisor).</p>

### 13.3.5 Summary of program commitments

As outlined in Table 13.3, the following program commitments are made under the program:

A clear understanding of legislative requirements, including criteria for determining whether an action is likely to have a significant impact, together with consideration of prudent and feasible options at an early stage, will maximise prospects of the proposed development proceeding in an environmentally acceptable way. Accordingly, the Authority will:

- require consideration of relevant EPBC Act policy documents by applicants, including relevant significant impact guidelines, in the scoping phase of the permission assessment and decision process
- require consideration of prudent and feasible alternatives which may have a lower impact on the environment by applicants, and amend its policy documentation to include reference to this requirement
- develop guidelines to assist applicants to determine whether an action is likely to have significant impact on the Great Barrier Reef Marine Park matters of national environmental significance. The Authority will seek to have these guidelines approved by the Minister.

The Authority will amend its Permit Application Form to include a section which specifically addresses matters of national environmental significance.

The Authority will seek to pursue amendments to the Great Barrier Reef Marine Park Regulations 1983 to explicitly outline the levels of assessment under the permission system.

The Authority will update its permissions guidance material to outline the general information requirements for applicants according to the level of assessment.

The Authority will seek to pursue amendments to the GBRMP Regulations (in particular, regulation 128) to ensure continued application of fees for the Authority's assessment of activities that are covered by an approval granted under section 146 of the EPBC Act (Part 10 – Strategic Assessments).

The Authority will publish notification of all permit applications on the internet as soon as practicable after receiving a permit application.

The Authority will seek to pursue amendments to the GBRMP Regulations (in particular, regulation 88D) to require advertising of applications where the granting of a permission may restrict the reasonable use by the public of a part of the Marine Park or is likely to have a significant impact on the Marine Park, including on matters of national environmental significance

Relevant policies will be updated to explicitly state that appropriate consultation will be undertaken with persons or organisations, including Australian and Queensland government agencies, whose functions, interests or activities could be impacted by the proposed activity.

Where an action is likely to have a significant impact on a matter protected under the EPBC Act within the Marine Park, the Authority will:

- require applicants to consider the impacts of actions on matters of national environmental significance and relevant attributes and environmental processes, as described in the Strategic Assessment Report and as amended in accordance with the Authority's policy review procedures
- require applicants to consider impacts (including direct, indirect and cumulative impacts) arising from the action in the context of all impacts affecting the matter
- consider any relevant international conventions and Australian-Queensland Government agreements in assessing and determining permissions under its permission system
- consider all relevant plans, policies, documents and guidance in assessing and determining permissions under its permission system.

The Authority will develop and implement:

- guidelines for the application of Great Barrier Reef offsets to maintain or improve the condition of matters of national environmental significance and relevant attributes and environmental processes, where impacts cannot be avoided or mitigated. The guidelines will seek to deliver an outcome equivalent to, or better than, the outcome that would apply if the EPBC Act Environmental Offsets Policy was applied.
- a Great Barrier Reef net benefit policy (that will be updated from time to time) to enhance the condition of the Marine Park, including matters of national environmental significance and relevant attributes and environmental processes.

Based on these findings, and in addition to previously identified commitments, the Authority commits to progressively strengthening (and, from time to time, amending) its policies, guidance material and support tools for assessing and determining permissions as follows:

- Amend policies and guidance material (that will be updated) used in the determination of permissions so they explicitly require the consideration of impacts on matters of national environmental significance and relevant attributes, including:
  - biodiversity and environmental processes
  - Indigenous and historic heritage values



- community benefits derived from the environment, including those not easily represented or measured (such as aesthetic values).
- Develop and maintain regionally-based standards for ecosystem health (including water quality) that support best management practices, and may be used to inform:
  - the assessment of individual, indirect and cumulative impacts on matters of national environmental significance and relevant attributes
  - actions to mitigate impacts and cumulative impacts on matters of national environmental significance and relevant attributes
  - outcomes-based decision making and considerations for determining 'acceptability' based on the commitments in Table 10 of the Program Report of impacts on matters of national environmental significance.
- Develop guidelines (that will be updated) requiring consideration of regionally-based standards for ecosystem health to improve the assessment of impacts and cumulative impacts to the Region's values, including matters of national environmental significance and outstanding universal value.
- Facilitate development of a dredging and dredge material disposal policy.
- Improve the hydrodynamic modelling guidelines (that will be updated) so they take account of longshore drift, resuspension and inter-annual current variability.
- Develop guidance material (that will be updated) to assist in determining the acceptability of impacts. Criteria shall include consistency of assessment outcomes with regulatory objectives, outcomes for matters of national environmental significance and relevant GBRMP Act and EPBC Act guidelines and standards.
- Develop guidance material (that will be updated) to be used in the determination of permissions to require the consideration of the impacts of underwater noise, particularly on at-risk species and inshore species.
- Develop guidelines (that will be updated) to improve the collection, management, analysis and interpretation of information derived from the Authority's permission system and compliance monitoring. The information will be used to better inform adaptive management strategies.
- Develop and/or strengthen knowledge and information management tools (including spatial mapping tools, hydrological connectivity tools and systems for managing Indigenous and historic heritage information) so their use can enhance understanding of:
  - the effects of impacts of actions on matters of national environmental significance
  - measures to avoid, mitigate and offset
  - actions to deliver net benefits and best practice adaptive management.
- Develop specific advice documents (that will be updated from time to time) for notifications under Part 5 of the Zoning Plan to make reference to consideration of the matters of national environmental significance under the EPBC Act.

The Authority will consider relevant Department of the Environment policy documents, guidelines, plans of management and other online data sources available on the Department's website in its assessment and decision-making process.

The Authority will integrate relevant components of the Department of the Environment's EPBC Act assessment templates and manuals into its assessment documents as they are reviewed and updated.

The Authority will strengthen its system for monitoring and managing compliance risks associated with permit conditions. It will review and make any necessary improvements to internal policies, procedures, training protocols and mechanisms to systematically enhance the effectiveness of its compliance and auditing process. This will include specific reference to matters of national environmental significance.

Update the Environmental Impact Management Policy to enable the effective integration of all the commitments above.

The Authority will work with Queensland Government agencies and the Department of the Environment to establish administrative arrangements including, but not limited to, reporting and communication mechanisms, information and data, environmental assessment and decision-making processes.

The Authority will investigate and implement opportunities through its ongoing regulatory reform processes to:

- enhance the effectiveness of its permission system to achieve objectives and outcomes for the protection and management of the Marine Park, including matters of national environmental significance
- harmonise requirements with relevant Commonwealth and state legislation and reduce regulatory burden.

### 13.3.6 Permission system alignment with the EPBC Act

The strategic assessment considered the *Framework of Standards for Accreditation*, prepared in 2014 by the Department of the Environment. This Framework was developed to assist state and territory governments to streamline their environmental assessment and approval processes with those of the Commonwealth through bilateral agreements. Bilateral agreements are a separate process under the EPBC Act to strategic assessments;

however, the Framework does provide a plain English description of baseline requirements under the EPBC Act, many of which are applicable.

In particular, the Framework provides guidance in relation to protection of matters of national environmental significance under the EPBC Act, which are addressed in this report, and also includes guidance on risk-based assessment, assessment and approvals policy, and transparency of processes and decisions.

Table 13.5 compares the outcomes achieved by the Authority's permission system against the requirements of the EPBC Act, noting that the Authority has committed to explicitly incorporate consideration of all attributes relevant to matters of national environmental significance, including the property's outstanding universal value, into the Authority's programs, plans and policies.

**Table 13.5 Comparison of the EPBC Act requirements and the permission system**

EPBC Act regulatory requirement	The permission system
<p><b>Referral</b></p> <p>A person proposing to take an action that will have or is likely to have a significant impact on a matter of national environmental significance protected under the EPBC Act must refer their proposal to the Department of the Environment, for a decision by the Minister on whether further assessment is required.</p> <p><b>Outcome(s):</b></p> <ul style="list-style-type: none"> <li>Matters that are referred under the EPBC Act will be assessed for a decision on whether further assessment is required. There is a risk that actions with the potential to significantly impact on protected matters may not be referred and may not be assessed for approval.</li> <li>Deliver certainty and efficiency by systematically identifying actions that are likely to have a significant impact on a matter of national environmental significance.</li> <li>Compliance actions may apply to a person who takes (or intends to take) such an action without approval.</li> </ul> <p>Once a proposed action is referred under the EPBC Act the Minister must decide whether the action is:</p> <ul style="list-style-type: none"> <li>not a controlled action (NCA)</li> <li>not controlled provided it is conducted in a particular manner (NCA PM)</li> <li>controlled action (CA — requires assessment)</li> <li>clearly unacceptable (cannot proceed).</li> </ul>	<p><b>All activities</b></p> <p>The Zoning Plan is the primary planning instrument for the conservation and management of the Marine Park. The Zoning Plan has regard to the objects set out in subsection 32(7) of the GBRMP Act, taking into account the World Heritage values of the Marine Park and the principles of ecologically sustainable development.</p> <p>Activities that will have or are likely to have a significant impact on the Marine Park, including MNES, would also require a Marine Park Permit which would be assessed under the permission system.</p> <p>The provisions of the Zoning Plan are enforced by Sections 38A to 38E of the GBRMP Act (which set out offences relating to use or entry of a zone for purposes that are not permitted, or in contravention of permission requirements or conditions) and sections 38M to 38MB of the GBRMP Act (which set out offences in relation to the use of ships). Contravening notification requirements or directions of the Authority is also an offence.</p> <p>Under Regulation 88Q(a) of the GBRMP Regulations, the Authority must consider the potential impacts of the conduct proposed to be permitted by the permission on the environment and on the social, cultural and heritage values of the Marine Park or a part of the Marine Park. This relevantly includes matters of national environmental significance.</p> <p>The Authority has committed to develop specific advice document(s) that stakeholders should consider in the preparation of their Permit Applications, to make reference to consideration of the MNES under the EPBC Act.</p> <p>This advice will include references to relevant International Conventions and Agreements, Commonwealth and State Agreements, and EPBC Act guidance documents to be considered by stakeholders in preparing Permit Applications. The Authority has committed to undertaking an assessment of whether an activity is likely to have a significant impact on matters of national environmental significance. This will be implemented by progressively strengthening (and, from time to time, amending) policies, guidance material and support tools for assessing and determining permissions.</p>
<p><b>Assessment</b></p> <p>Not all referred actions require further assessment (see Approvals below). For those actions that do require further assessment, a level of assessment is set. The Minister for the Environment considers the scale and nature of impacts, the complexity of the issues, and the degree of public concern. The level of assessment can range from assessment on the referral information already submitted to a public inquiry.</p> <p>Once the method of assessment is determined, the EPBC Act and Regulations provide further details about the process, including additional content requirements (if any) and the public comment process.</p> <p>If an action is deemed controlled, further assessment is required, including an opportunity for public comment.</p>	<p><b>Assessment</b></p> <p>The permission system's regulatory objective and outcomes-based approach ensures that the level of assessment is commensurate with the nature and scale of the activity, and its risk-based approach ensures it is appropriate to the potential impacts and risks to the Marine Park, including MNES.</p> <p>The Authority recommends applicants and interest groups discuss initial concepts, designs, alternatives and potential mitigation of impacts for any proposed activity or development which may impact on the Great Barrier Reef Marine Park and require a permit.</p> <p>In order to be granted a permit to undertake an activity, an application must provide enough information so permit assessors can clearly understand what the intended activity is.</p> <p>The Authority has committed to specifically including MNES in</p>

EPBC Act regulatory requirement	The permission system
<p><b><u>Outcome(s):</u></b></p> <ul style="list-style-type: none"> <li>The level of assessment is appropriate to the nature, scale, and potential impacts and risks to matters protected under the EPBC Act.</li> <li>Efficiency and transparency by employing assessment approaches that reflect the risk of the proposed action and provide sufficient information for a decision maker to make an informed decision.</li> </ul>	<p>Authority programs, plans and policies.</p> <p>The Authority assigns an assessment level to each permission application. The assessment level is based on the potential level of impact and the level of information required to adequately address mandatory and discretionary considerations, and is outlined in the Great Barrier Reef Marine Park Authority Environmental Impact Management Policy.</p> <p>The Permit application is assessed in accordance with the GBRMP Regulations which include mandatory and discretionary criteria. These include relevant Australian and Queensland government laws relating to the management of the environment and conservation advices.</p> <p>Some activities (for example take of Protected Species (88S) require additional criteria to be met).</p> <p>Assessments of impacts will have regard to the full range of MNES. Information sources used in assessments will be included in the Authority's assessment report.</p>
<p><b><u>Approvals</u></b></p> <p>The Department of the Environment or relevant state agency prepares a report for the Minister, and the Minister (or delegate) makes a decision on whether or not to approve the action. In making this decision, the Minister for the Environment considers the impacts of the proposed action on matters protected by the EPBC Act, social and economic matters, as well as other matters for consideration as set out under the EPBC Act. If that decision is to approve the action, the Minister for the Environment may place conditions on the approval.</p> <p><b><u>Outcome(s):</u></b></p> <ul style="list-style-type: none"> <li>Actions that could have an unacceptable impact on MNES protected under the EPBC Act are not allowed to proceed.</li> <li>Actions that could have a significant impact on MNES protected under the EPBC Act are assessed and, if approved, are regulated to manage such impacts and ensure development proceeds in accordance with the principles of ecologically sustainable development.</li> <li>Deliver certainty, transparency and legally robust decisions by undertaking environmental assessment that adequately addresses all matters of national environmental significance, and making approval decisions based on recognised principles of environmental policy, as set out in the Intergovernmental Agreement on the Environment 1992.</li> </ul>	<p><b><u>Decision Phase</u></b></p> <p>Applicants must demonstrate that potential impacts and risks to the Marine Park, including MNES and the broader environment, will be consistent with the objects of the GBRMP Act in their permit application and Environmental Management Plan.</p> <p>Content of the EMP must be consistent with the nature and scale of the activity, and have regard to the mandatory and discretionary considerations under GBRMP Regulations 88Q and 88R.</p> <p>The Authority may grant or refuse a permission based on the assessment. A permit may be granted with any conditions appropriate to the attainment of the objects of the GBRMP Act, including the principles of ecologically sustainable use, and the protection of matters of national environmental significance.</p> <p>An Environmental Management Plan (EMP) will be required for activities that may have an unacceptable impact on the Marine Park, including MNES. The EMP outlines the proposed activity, its possible impacts on the Marine Park and specifies strategies to minimise potential impacts.</p> <p>The Authority will audit environmental performance of some permitted activities and developments and, if environmental performance or compliance does not meet required standards, then penalties and/or remediation may be required.</p>
<p><b><u>Transparency</u></b></p> <p>Actions referred to the Department of the Environment under the EPBC Act are published on that Department's website and there is a public invitation to comment. Public comments must be taken into account in both referral and approval decisions.</p> <p>Notifications of key stages of the assessment are posted on the website, including a notice of all decisions (for example referral, approval decisions).</p> <p>All decisions are subject to judicial review.</p> <p>Conditions on approvals are published on the Department of the Environment's website, as are all approved documents. The Minister may condition approvals to require publication of additional documents or reports such as monitoring plans.</p> <p><b><u>Outcome(s):</u></b></p> <ul style="list-style-type: none"> <li>Systems are transparent and offer appropriate opportunities for public engagement, and decisions are legally robust.</li> </ul>	<p><b><u>Transparency</u></b></p> <p>The Great Barrier Reef Marine Park Authority is an independent statutory authority established by the GBRMP Act. As an independent regulator, the Authority's decision-making processes are based entirely in law. This is a robust regulatory system, which provides confidence to the Australian community that the regulator will not make decisions on any basis other than those enshrined in law that has passed both houses of Parliament.</p> <p>Applications that may have a significant impact on MNES will be available for public consultation via the Authority's website. The permit application is assessed by the permit assessor/project manager and referred to relevant parties for comment (for example applications with potential to impact fisheries would be referred to the Department of Agriculture and Fisheries Qld).</p> <p>Regulation 88D of the GBRMP Regulations requires applications to be advertised if the Authority considers that granting of a permission may restrict the reasonable use by the public of a part of the Marine Park.</p> <p>This advertisement must be published in a newspaper circulating in the part of the State of Queensland adjacent to that part of the</p>

EPBC Act regulatory requirement	The permission system
	<p>Marine Park in which the conduct, for which permission is being sought, is to be engaged in.</p> <p>The advertisement is to include an invitation for interested persons to make comment on the application within a period of not less than 30 days. The advertisement will also be published on the Authority's website.</p> <p>The Authority will seek to amend Regulation 88D of the GBRMP Regulations to require advertising of applications where the granting of a permission may restrict the reasonable use by the public of a part of the Marine Park or is likely to have a significant impact on the Marine Park, including on matters of national environmental significance.</p> <p>The Authority's management approach includes ensuring that decision making is transparent and accountable.</p> <p>The Authority's website can be used to search for specific details of permit applications and permit decisions. All current EPBC referrals, where the Authority is providing advice to the Department of the Environment, are available on the Authority's website, with links to relevant documentation currently hosted on the Department website. Following endorsement and approval, this information would be hosted on the Authority's website.</p> <p>Section 183 of the GBRMP Regulations requires for Notice of Certain Decisions on the Authority's website, including decisions on Permissions.</p> <p>GBRMP Regulations 184 allows for persons affected by certain decisions to obtain a statement of reasons from the Authority.</p> <p>Reconsideration of decisions by the Authority is available under GBRMP Regulations 186. Administrative Appeals Tribunal review of decisions after reconsideration is available under GBRMP Regulations 187.</p>
<p><b>Compliance and Enforcement</b></p> <p>The EPBC Act compliance framework includes a broad range of enforcement mechanisms for monitoring and/or managing potential non-compliance of both referred and non-referred actions. Compliance measures include post-approval reporting, monitoring with approval conditions and compliance auditing. Enforcement mechanisms to address non-compliance include revocation of approvals, civil and criminal penalties, remediation orders and enforceable undertakings.</p> <p><b>Outcome(s):</b></p> <ul style="list-style-type: none"> <li>• That all actions approved under the EPBC Act are undertaken in accordance with the relevant legislative and regulatory controls.</li> <li>• The community has confidence that actions with significant impacts are regulated effectively.</li> </ul>	<p><b>Compliance and Enforcement</b></p> <p>Approved activities must be carried out in accordance with a Great Barrier Reef Marine Park Permit and any conditions (GBRMP Regulations Division 2A.5 and 2A.6).</p> <p>The Authority will determine the need for a cash bond or bank guarantee and a Deed of Agreement to provide ongoing indemnity and, where necessary, to ensure there is an ongoing contractual obligation to protect the environment if the permit lapses (GBRMP Regulation 88ZE).</p> <p>Civil penalties are available for offences (GBRMP Act Part VAA).</p> <p>The Authority has a well-established inspection, compliance, monitoring and enforcement regime to ensure compliance with all legislative requirements. It schedules and conducts inspections according to a risk rating of the potential impacts of activities, and implements a graduated enforcement strategy to educate and promote compliance with the legislation (including the EPBC Act and protection of MNES).</p> <p>The Joint Field Management Program (Great Barrier Reef Intergovernmental Agreement — Schedule C) undertakes compliance activities – both through education and enforcement — to encourage adherence to legal requirements such as those contained in permits, plans of management and zoning.</p> <p>A partnership approach with Australian and Queensland government agencies and stakeholders enables the Authority to promote protection of the Marine Park, including MNES and OUV, to a broader monitoring group and deliver greater incident identification and response coverage for the Region.</p>




## 13.4 Applying the principles of ecologically sustainable development

The principles of ecologically sustainable development as described in section 3A of the EPBC Act are:

- Decision-making processes should effectively integrate both long-term and short-term environmental, economic, social and equitable considerations.
- If there are threats of serious or irreversible environmental damage, lack of full scientific certainty should not be used as a reason for postponing measures to prevent environmental degradation.
- The principle of intergenerational equity — that the present generation should ensure the health, diversity and productivity of the environment is maintained or enhanced for the benefit of future generations.
- The conservation of biological diversity and ecological integrity should be a fundamental consideration in decision making.
- Improved valuation, pricing and incentive mechanisms should be promoted.

It is a requirement of the GBRMP Act that, in managing the Marine Park and performing its other functions, the Authority has regard to and seeks to act in a way that is consistent with the principles of ecologically sustainable use. These principles, defined in the GBRMP Act closely match the above principles of ecologically sustainable development (Figure 13.2). The Authority, bound by the requirements of the GBRMP Act, will continue to apply these principles to its protection and management of the environment, biodiversity and heritage values of the Region.



**Figure 13.2 Principles of ecologically sustainable use**

Principles of ecologically sustainable use outlined within the GBRMP Act, and the equivalent principles of ecologically sustainable development within the *EPBC Act*

The ways in which the principles of ecologically sustainable development will continue to be applied in the Authority's comprehensive management program (Part B) and its permission system (Part C) are summarised in Table 13.6.

Table 13.6 Applying the principles of ecologically sustainable development

Principles of ecologically sustainable development	Applying the principles of ESD in the Authority's comprehensive management program (Part B)	Applying the principles of EST in the Authority's permission system (Part C)
a) Decision-making processes should effectively integrate long-term and short-term economic, environmental, social and equitable considerations.	<ul style="list-style-type: none"> <li>Section 7(3) of the Act requires the Authority to perform its functions consistent with the principles of ecologically sustainable use. This includes consideration of these principles in the preparation of zoning plans, plans of management and permit decision-making.</li> <li>The potential impacts on the environment and on social, cultural and heritage values must be considered in all permit decisions.</li> <li>The effects on public appreciation, understanding and enjoyment of the Marine Park, as well as impacts on other activities may be considered in permit decisions.</li> <li>The Authority's Environmental Impact Management Policy requires consideration of impacts on environmental, economic and social values.</li> <li>Decision making is informed by the best available information and, where appropriate, stakeholders are engaged and public comment is sought (for example, during major permit assessments).</li> <li>There is explicit recognition of the importance of the Great Barrier Reef to its Traditional Owners and their cultural heritage. They are increasingly engaged in its management.</li> <li>The community benefits derived from the environment are explicitly recognised in the Program Report and, increasingly, the outcomes of integrated monitoring and modelling will be available for consideration in decision making, providing guidance on socio-economic benefits and impacts and likely future trends.</li> <li>Biodiversity, heritage values and community benefits derived from the environment are the foundations of the Authority's strategic planning.</li> <li>Minimum standards for public consultation are identified under the Act and in the Authority's policies to guide those involved in the Authority's decision-making processes.</li> <li>Cumulative impact assessment guidelines will incorporate this principle.</li> </ul>	<ul style="list-style-type: none"> <li>Section 7(3) of the Act requires the Authority in performing its functions to have regard to, and seek to act in a way that is consistent with, the principles of ecologically sustainable use. This includes consideration of these principles in the preparation of zoning plans, plans of management and permit decision-making.</li> <li>The potential impacts on the environment and on social, cultural and heritage values of the Marine Park or part of the Marine Park must be considered in all permit decisions.</li> <li>The effects on public appreciation, understanding and enjoyment of the Marine Park, as well as impacts on other activities may be considered in permit decisions.</li> <li>The Authority's Environmental Impact Management Policy requires consideration of impacts on environmental, economic and social values.</li> <li>Decision making is informed by the best available information and, where appropriate, stakeholders are engaged and public comment is sought (for example, during major permit assessments).</li> <li>Minimum standards for public consultation are identified under the Act and in the Authority's policies to guide those involved in the Authority's decision-making processes.</li> <li>A cumulative impact assessment policy will incorporate this principle.</li> </ul>
b) If there are threats of serious or irreversible environmental damage, lack of full scientific certainty should not be used as a reason for postponing measures to prevent environmental degradation.	<ul style="list-style-type: none"> <li>Section 7(3) of the Act requires the Authority to perform its functions consistent with the objects of the Act and the principles of ecologically sustainable use. This includes consideration of these principles in the preparation of zoning plans, plans of management and permit decision-making.</li> <li>Consideration of permit applications is based on the best available science.</li> <li>Where there is scientific uncertainty, the Authority seeks to harness</li> </ul>	<ul style="list-style-type: none"> <li>Section 7(3) of the Act requires the Authority in performing its functions to have regard to, and seek to act in a way that is consistent with, the objects of the Act and the principles of ecologically sustainable use. This includes consideration of these principles in the preparation of zoning plans, plans of management and permit decision-making.</li> <li>Consideration of permit applications is based on the best available science.</li> <li>Where there is scientific uncertainty, the Authority seeks to harness</li> </ul>

Principles of ecologically sustainable development	Applying the principles of ESD in the Authority's comprehensive management program (Part B)	Applying the principles of EST in the Authority's permission system (Part C)
	<p>expert opinion, plus knowledge held by Traditional Owners and stakeholders, including through its advisory committees and through seeking public comment.</p> <ul style="list-style-type: none"> <li>• Risks and threats to the Great Barrier Reef are identified and assessed in the <i>Great Barrier Reef Outlook Report 2009</i>, which is updated every five years.</li> <li>• Increasingly, outcomes of integrated monitoring and modelling will be available for consideration in decision making, providing guidance on likely future trends.</li> <li>• Cumulative impact assessment guidelines will incorporate this principle and increase the rigour of environmental assessment processes.</li> </ul>	<p>expert opinion, plus knowledge held by Traditional Owners and stakeholders, including through its advisory committees and through seeking public comment.</p> <ul style="list-style-type: none"> <li>• A cumulative impact assessment policy will incorporate this principle and increase the rigour of environmental assessment processes.</li> </ul>
c) The principle of intergenerational equity — that the present generation should ensure the health, diversity and productivity of the environment is maintained or enhanced for the benefit of future generations.	<ul style="list-style-type: none"> <li>• Section 7(3) of the Act requires the Authority to perform its functions consistent with the principles of ecologically sustainable use. This includes consideration of these principles in the preparation of zoning plans, plans of management and permit decision-making.</li> <li>• The Authority recognises the overall condition of the Great Barrier Reef has declined. Future management is focused on minimising further impacts and enhancing and restoring the values of most concern.</li> <li>• A policy on delivering net benefits, including offsets, will contribute to the maintenance and enhancement of the health, diversity and productivity of the environment for future generations.</li> <li>• Programs and actions are implemented to address remaining risks identified in the Outlook Report and improve the future outlook for the Reef.</li> <li>• Regional and local actions facilitated by the Authority, including Traditional Use of Marine Resources Agreements and the Reef Guardian program, are an important element in enhancing health, diversity and productivity and often involve activities that engage a range of generations.</li> <li>• Increasingly, the outcomes of integrated monitoring and modelling will be available for consideration in decision making, providing guidance on community benefits and likely future trends, and evaluating the delivery of net benefits.</li> <li>• Cumulative impact assessment guidelines will incorporate this principle.</li> <li>• An outcomes-based management approach and a net benefit policy will improve actions to maintain and enhance the condition of the Region's values for future generations.</li> </ul>	<ul style="list-style-type: none"> <li>• Section 7(3) of the Act requires the Authority to perform its functions consistent with the principles of ecologically sustainable use. This includes consideration of these principles in the preparation of zoning plans, plans of management and permit decision-making.</li> <li>• A policy on delivering net benefits, and guidelines for the application of offsets in the Marine Park, will contribute to the maintenance and enhancement of the health, diversity and productivity of the environment for future generations.</li> <li>• Increasingly, the outcomes of integrated monitoring and reporting will be available for consideration in decision making, providing guidance on community benefits and likely future trends, and evaluating the delivery of net benefits.</li> <li>• A cumulative impact assessment policy will incorporate this principle.</li> <li>• An outcomes-based management approach and a net benefit policy will improve actions to maintain and enhance the condition of the Region's values for future generations.</li> </ul>
d) The conservation of biological diversity and ecological integrity	<ul style="list-style-type: none"> <li>• The main object of the Act includes providing for the long-term protection and conservation of the environment, biodiversity and heritage of the</li> </ul>	<ul style="list-style-type: none"> <li>• The main object of the Act includes providing for the long-term protection and conservation of the environment, biodiversity and</li> </ul>

Principles of ecologically sustainable development	Applying the principles of ESD in the Authority's comprehensive management program (Part B)	Applying the principles of EST in the Authority's permission system (Part C)
should be a fundamental consideration in decision making.	<p>Great Barrier Reef Region. This is the basis for the Authority's management arrangements.</p> <ul style="list-style-type: none"> <li>• Zoning arrangements, which include about one-third of the Marine Park in no-take zones, are a vital component in conserving biological diversity and ecological integrity. Connectivity and protecting a representative area of all bioregions were key considerations in designating no-take zones.</li> <li>• A range of policies, including the <i>Great Barrier Reef Biodiversity Conservation Strategy</i>, guide the Authority's management relevant to the conservation of biological diversity. Additionally, the <i>Great Barrier Reef Climate Adaptation Strategy and Action Plan</i> assists the Authority to address the challenges of climate change in relation to biological diversity.</li> <li>• The potential impacts on the environment, including its biodiversity, and the objects of the zone where an activity will occur, are required to be considered in all permit decisions.</li> <li>• Any relevant recovery, conservation threat abatement plans or approved conservation advice under the EPBC Act may be considered in permit decisions.</li> <li>• Cumulative impact assessment guidelines and a net benefit policy will incorporate this principle.</li> </ul>	<p>heritage of the Great Barrier Reef Region. This is the basis for the Authority's management arrangements.</p> <ul style="list-style-type: none"> <li>• Zoning arrangements, which include about one-third of the Marine Park in no-take zones, are a vital component in conserving biological diversity and ecological integrity. Connectivity and protecting a representative area of all bioregions were key considerations in designating no-take zones.</li> <li>• A range of policies, including the <i>Great Barrier Reef Biodiversity Conservation Strategy</i>, guide the Authority's management relevant to the conservation of biological diversity. Additionally, the <i>Great Barrier Reef Climate Adaptation Strategy and Action Plan</i> assists the Authority to address the challenges of climate change in relation to biological diversity.</li> <li>• The potential impacts on the environment, including its biodiversity, and the objects of the zone where an activity will occur, are required to be considered in all permit decisions.</li> <li>• All relevant recovery, conservation threat abatement plans or approved conservation advice under the EPBC Act may be considered in permit decisions.</li> <li>• A cumulative impact assessment policy and a net benefit policy will incorporate this principle.</li> </ul>
e) Improved valuation, pricing and incentive mechanisms should be promoted.	<ul style="list-style-type: none"> <li>• Socio-economic monitoring will be included in long-term core monitoring for the Region.</li> <li>• The Authority's management is informed by economic valuations of Reef-dependent activities.</li> <li>• Tourism operators are recognised and rewarded for being certified as meeting best practice standards, including with longer-term permits.</li> <li>• There is public recognition of participants in the Reef Guardian program and of Traditional Use of Marine Resources Agreements.</li> <li>• Cost recovery principles are applied in the Authority's management arrangements.</li> <li>• A framework will be developed to guide implementation of investments for net conservation benefits designed to increase the resilience of the Great Barrier Reef Region.</li> <li>• Cumulative impact assessment guidelines will incorporate this principle.</li> </ul>	<ul style="list-style-type: none"> <li>• Socio-economic monitoring will be included in long-term core monitoring for the Region.</li> <li>• The Authority's management is informed by economic valuations of Reef-dependent activities.</li> <li>• Tourism operators are recognised and rewarded for being certified as meeting best practice standards, including with longer-term permits.</li> <li>• Cost recovery principles are applied in the Authority's management arrangements.</li> <li>• A framework will be developed to guide implementation of investments for net conservation benefits designed to increase the resilience of the Great Barrier Reef Region. This framework will support the Authority's advice to the Australian and Queensland governments on investments, including for Reef Trust.</li> <li>• A cumulative impact assessment policy will incorporate this principle.</li> </ul>



## 13.5 Applying adaptive management

### 13.5.1 Addressing risk

The Authority adopts a risk-based approach in its management and decision making. The strategic assessment identifies existing and emerging impacts that are predicted to present the highest risks to the values relevant to matters of national environmental significance. The Authority's future management program is clearly focused on the highest risk impacts and the cumulative impact of medium and lower risks.

Section 54(3)(d) of the Act requires 'an assessment of the risks to the ecosystem' within the Region be included in each Outlook Report. This legislative requirement means the assessment of current impacts and future risks will be regularly updated. The risk assessment presented in the first Outlook Report in 2009 was undertaken using the Australian Standard for Risk Assessment, and has subsequently been a principal guide for the Authority in determining management priorities. The Authority also applies the standard in its environmental assessment and management and its permitting processes.

One key component of managing risks is having access to the most up-to-date information. The integrated monitoring and reporting program will substantially improve the knowledge base available to the Authority. Recognising the significance of the Great Barrier Reef to local communities and the nation, and the community benefits derived from the environment, the monitoring and reporting program has an expanded focus to better integrate information on social, cultural and economic values. The Authority also intends to expand the sources of information it uses in making decisions, so that traditional ecological knowledge and information held by the broader community are better considered. A commitment to the development of a heritage database, including a protocol to manage culturally sensitive information, will further improve consideration of cultural, social and economic values relevant to matters of national environmental significance.

Ongoing improvements in linkages with information providers and advances in information technology are likely to continue to greatly improve the ways in which the Authority and its management partners provide, access and integrate information used to address risk.

This strategic assessment has clearly demonstrated that impacts — whether the result of human activities or natural forces — do not operate in isolation or on a single value or in a specific area. It also shows consideration of cumulative impacts is one of the weakest aspects of the Authority's current management arrangements. A commitment in the Authority's management program is to integrate risk management and the consideration of cumulative impacts into regulatory frameworks through development of cumulative impact guidelines.

### 13.5.2 Addressing uncertainty

Inherently, there is a large amount of uncertainty surrounding protection and management of the Great Barrier Reef — an enormous, complex marine protected area which is managed for multiple use.

The Authority will continue to employ a range of tools to provide certainty with regard to where activities may occur and under what conditions they may proceed. These tools are outlined in the Program Report and include zoning plans, plans of management, permits, policies and guidelines. The resulting level of certainty provided for different activities in the Region is presented in Chapter 8 of this report. Where uncertainty exists, the Authority adopts a risk-based approach and applies the principles of ecologically sustainable use to its decision making. The use of these tools, approaches and principles is embedded in the Authority's governing legislation (the Act) and reflected in its policies. The Authority also deals with uncertainty by, where feasible, maintaining flexibility in its management approach, so that changing circumstances can be considered and reflected in management arrangements.

Uncertainty is also addressed by always seeking to improve the knowledge base available for decision making. The Authority will continue to be actively involved in determining research and monitoring priorities for the Region, and maintaining strong connections with the research community (including those undertaking voluntary monitoring) so that emerging information can be understood and applied to management. The Authority will also seek to improve the ways that it is informed by knowledge and understanding held by the Great Barrier Reef's Traditional Owners, stakeholders and the community (see Section 7 of the Program Report). This knowledge will continue to be acquired through ongoing liaison, specific workshops and the Authority's Reef Advisory Committees and Local Marine Advisory Committees.

For key matters that affect the Region, the Authority takes a leading role in improving understanding and defining the priorities for management action. For example, in the case of reducing impacts on the Region's values arising from clearing and modifying coastal habitats, the Authority:

- improved technical understanding of the role that coastal habitats play in providing ecosystem services to the Reef ecosystem, culminating in the report *Informing the outlook for Great Barrier Reef coastal ecosystems*<sup>1</sup>
- is defining priorities for action by undertaking basin assessments to identify the coastal basins that remain intact and those that are slightly to moderately affected by human activity.

These bodies of work will guide the Authority's partnership activities with local and state government agencies, natural resource management bodies, Reef Guardian participants and local communities to actively manage and restore the functioning of coastal ecosystems critical to the health of the Great Barrier Reef.

In its future management, the Authority will adopt the systematic approach used in this report to the identification of the values relevant to matters of national environmental significance, and the drivers, activities and impacts affecting them (see Chapter 4, 5 and 6). It will also adopt a systematic approach to considering the cumulative effect of impacts on values (see Chapter 2, Table 2.1). This will include using models (qualitative or quantitative) to examine the relationships between multiple drivers, activities and impacts; for example, those that can affect key habitats such as coral reefs and seagrass meadows. Such tools can help the Authority deal with uncertainty by allowing consideration of multiple alternate models in situations where the cause-and-effect relationships between values and impacts are uncertain or contentious. They can also help tackle a key source of uncertainty — the cumulative effect of impacts. It is intended that ongoing use of modelling in the Authority's decision-making processes will be formalised through a guideline on cumulative impact assessment (see Section 6 of the Program Report).

### **13.5.3 Implementing adaptive management to protect and manage matters of national environmental significance**

Through the ongoing foundational activities and the future improvements identified in the Program Report and the vital contributions of its partners, the Authority anticipates declines in key components of the Region's ecosystem can be halted and reversed and the desired outcomes achieved for values relevant to matters of national environmental significance. It is recognised that this will require a sustained and ongoing commitment of resources, and that such improvements in environmental condition may take decades to achieve.

Adaptive management has played a central role in the understanding and management of impacts in the Region in the last decade.<sup>2,3,4,5,6,7,8,9</sup> The Authority recognises that to achieve the desired outcomes for a system as dynamic and complex as the Great Barrier Reef Region, an ongoing adaptive approach will be required.

In addition to its management through the permission system outlined above, the Authority will continue to adopt a comprehensive and coordinated approach to its management of impacts on matters of national environmental significance, operating at a number of levels (Figure 13.3) influencing drivers and activities:

- mitigating (reducing) impacts and pressures
- restoring, maintaining and enhancing the state or condition of the environment
- promoting appreciation of the community benefits derived from the environment.

Management responses directed towards drivers and activities are often the most effective as they act on the source of impacts and enable them to be avoided. Management actions that directly address impacts or pressures focus on reducing or mitigating the magnitude of the impact, and are fundamental to allowing use to occur within ecologically sustainable limits. Management actions that work directly to improve the condition of the environment once the impact has occurred are generally least effective and often most costly. Lastly, interventions which target community benefits are focused on promoting understanding and awareness of the benefits derived from the Region (for example, enjoyment and income) or of the World Heritage Area (for example, outstanding universal value).

Adaptive management that responds to changing circumstances requires an up-to-date and comprehensive understanding of the status and trend of values relevant to matters of national environmental significance, the impacts that may be affecting them and the benefits being derived from the environment. One of the Authority's new measures to strengthen management is an integrated monitoring and reporting program to standardise and integrate ecological, social and economic monitoring for the Great Barrier Reef as part of an adaptive management cycle (Section 13.4).

The Authority will continue to use its five-yearly preparation of the statutory *Great Barrier Reef Outlook Report* to examine the current status, future risks and likely outlook for the Region. The outcomes of each report will inform the Authority's future priorities and be the principle guide to adaptive management. Importantly, the 'management effectiveness' section will identify management topics that require improvement, and the 'outlook' section of each report will inform adaptive management by integrating current knowledge and modelling to produce a predicted future condition of the values relevant to matters of national environmental significance.

Annual strategic planning, taking into account emerging information and monitoring results, will inform finer-scale adjustments to management arrangements.

The Authority will also continue to review and update its science information needs every five years as part of the outlook report cycle.

The Authority will systematically and regularly evaluate its performance towards achieving the program's desired outcomes for the protection of matters of national environmental significance. Evaluating performance will allow the Authority to learn from its successes, improve continually as an organisation, adapt its management arrangements and better deliver on its statutory objectives.




Figure 13.3 Pathways of management intervention

### 13.5.4 Addressing World Heritage Committee recommendations

The 35<sup>th</sup> session of the World Heritage Committee requested a monitoring mission be undertaken jointly by the World Heritage Centre (WHC) and the International Union for Conservation of Nature (IUCN). The reactive monitoring mission was undertaken from 6–14 March 2012 with the objective of assessing the state of conservation of the Great Barrier Reef World Heritage Area property and to contribute to the strategic assessment process.

The Authority has taken account of the recommendations of the WHC and IUCN in developing the strategic assessment and in outlining the Authority's future management program. Specific actions addressing each recommendation are provided in Table 13.7. The timeframe of the implementation of a number of these recommendations can be found in Table 13.1.

Table 13.7 Progress on the recommendations of the 2012 World Heritage Centre and International Union Conservation of Nature reactive monitoring mission

2012 World Heritage Centre and IUCN mission recommendations	Great Barrier Reef Region strategic assessment response
<b>R1:</b> Water quality	<ul style="list-style-type: none"> <li>• <b>REC18:</b> Update and strengthen the Great Barrier Reef Marine Park water quality guidelines to address a broader range of habitats and species and account for cumulative impacts</li> <li>• <b>REC19:</b> Improve the effectiveness of the Authority's hydrodynamic guidelines as a decision-making tool by requiring consideration of a greater range of environmental factors, and regularly reviewing them to reflect improvements in understanding</li> <li>• <b>REC22:</b> Reduce crown-of-thorns starfish outbreaks by continuing to improve water quality and through a long-term control program</li> </ul>
<b>R2:</b> No port development outside existing and long-established port areas	<ul style="list-style-type: none"> <li>• <b>REC11:</b> Support development of a Queensland ports strategy that concentrates port development around long-established major ports in Queensland, and encourage port master planning</li> </ul>
<b>R3:</b> Manage development in Gladstone Harbour and on Curtis Island	<ul style="list-style-type: none"> <li>• <b>REC11:</b> Support development of a Queensland ports strategy that concentrates port development around long-established major ports in Queensland, and encourage port master planning</li> </ul>
<b>R4:</b> Ensure that any development, including ports and associated infrastructure, is carried out consistently with the highest international standards of best practice	<ul style="list-style-type: none"> <li>• <b>REC3:</b> Work closely with Australian, Queensland and local government agencies to help identify values of the Great Barrier Reef World Heritage Area that are not easily represented and measured such as aesthetic values</li> <li>• <b>REC7:</b> Work closely with Australian and Queensland government agencies to improve understanding and management of cumulative impacts from activities within and adjacent to the Region, and provide clearer guidance on how applicants and decision makers should address cumulative impacts in assessments</li> </ul>
<b>R5:</b> Strategic assessments and long-term sustainability plan	<ul style="list-style-type: none"> <li>• <b>REC26:</b> Develop and implement a long-term sustainability plan for the Great Barrier Reef in cooperation with Australian and Queensland government agencies to better coordinate programs designed to manage and improve the condition of the Reef</li> </ul>

2012 World Heritage Centre and IUCN mission recommendations	Great Barrier Reef Region strategic assessment response
<b>R6:</b> Include in the future editions of the Outlook Report, commencing with the version to be published in 2014, a specific assessment of the condition, trends, threats and prospects for the outstanding universal value of the Great Barrier Reef World Heritage Area	<ul style="list-style-type: none"> <li>• The Great Barrier Reef Marine Park Regulations have been amended to require future Outlook Reports to include an assessment of heritage values including, but not limited to, the outstanding universal value of the Great Barrier Reef World Heritage Area.</li> <li>• The <i>Outlook Report 2014</i> will contain a specific assessment of the condition, trends, threats and prospects for the outstanding universal value of the Great Barrier Reef World Heritage Area.</li> </ul>
<b>R7:</b> No development to impact individually or cumulatively on the outstanding universal value of the property	<ul style="list-style-type: none"> <li>• <b>REC1:</b> Explicitly incorporate consideration of all matters of national environmental significance, including attributes of the property's outstanding universal value, into the Authority's programs, plans and policies</li> </ul>
<b>R8:</b> Prevent any approval of major projects that may compromise the outcomes of the strategic assessment, until it is completed No development to impact individually or cumulatively on the outstanding universal value of the property	<ul style="list-style-type: none"> <li>• <b>REC1:</b> Explicitly incorporate consideration of all matters of national environmental significance, including attributes of the property's outstanding universal value, into the Authority's programs, plans and policies</li> </ul>
<b>R9:</b> Ensure all components of the outstanding universal value of the Great Barrier Reef are clearly defined and form a central element within the protection and management system for the property	<ul style="list-style-type: none"> <li>• <b>REC1:</b> Explicitly incorporate consideration of all matters of national environmental significance, including attributes of the property's outstanding universal value, into the Authority's programs, plans and policies</li> </ul>
<b>R10:</b> Develop and adopt clearly defined and scientifically justified targets for improving the state of conservation of the outstanding universal value of the Great Barrier Reef World Heritage Area	<ul style="list-style-type: none"> <li>• <b>REC25:</b> Establish a management framework with clear outcomes and targets for the protection of values and the management of impacts, including cumulative impacts</li> </ul>
<b>R11:</b> Commission an independent review of the overall institutional and legal mechanisms that provide coordinated planning, protection and management of the Great Barrier Reef World Heritage Area as a whole	<ul style="list-style-type: none"> <li>• An independent review of management effectiveness was conducted as part of the comprehensive strategic assessment. The program is one of the products of this review.</li> </ul>
<b>R12:</b> Ensure increased resources for the protection and management of the property	<ul style="list-style-type: none"> <li>• <b>REC15:</b> Support increased investment in site infrastructure to protect matters of national environmental significance in the Great Barrier Reef Region</li> </ul>
<b>R13:</b> Take an integrated approach to planning, regulation and management of ports and shipping activity	<ul style="list-style-type: none"> <li>• <b>REC2:</b> Improve spatial mapping capabilities to support planning and assessment decision making, including the range of values mapped and public availability</li> <li>• <b>REC11:</b> Support development of a Queensland ports strategy that concentrates port development around long-established major ports in Queensland, and encourage port master planning</li> <li>• <b>REC11A:</b> Support the development of a whole-of-government policy to provide a strategic and consistent approach to the sustainable management of dredging and dredge spoil disposal in the Great Barrier Reef World Heritage Area</li> <li>• <b>REC12:</b> Promote a strategic approach to the development and operation of marinas and other access infrastructure along the Great Barrier Reef coast</li> </ul>
<b>R14:</b> Share best practices and success stories with other world heritage sites facing similar management challenges	<ul style="list-style-type: none"> <li>• The Authority will continue its international engagement program to share knowledge on marine park management and related issues.</li> </ul>
<b>R15:</b> Inform the committee of developments that may affect outstanding universal value	<ul style="list-style-type: none"> <li>• The Authority will update the administrative agreement with the Department of the Environment to provide relevant information for reporting on OUV</li> </ul>
Other actions to support recommendations	<ul style="list-style-type: none"> <li>• <b>REC4:</b> Collaborate with Traditional Owners to undertake an assessment of the Indigenous heritage values of the Region</li> <li>• <b>REC5:</b> Develop and implement knowledge management systems for Indigenous and historic heritage information, including a protocol for managing culturally sensitive information and improved information sharing arrangements</li> <li>• <b>REC6:</b> Improve understanding of the role the Great Barrier Reef plays in the life of the community</li> </ul>



2012 World Heritage Centre and IUCN mission recommendations	Great Barrier Reef Region strategic assessment response
	<ul style="list-style-type: none"> <li>• <b>REC8:</b> Streamline assessment processes across jurisdictions and seek to have a more coordinated approach to community consultation</li> <li>• <b>REC9:</b> Improve alignment between the Authority's and the Queensland Government's protected area and tourism management arrangements and look for opportunities to streamline</li> <li>• <b>REC10:</b> Develop and implement plans of management in areas of the Great Barrier Reef Marine Park that have high growth for recreation and other uses</li> <li>• <b>REC13:</b> Review and update the <i>Great Barrier Reef Marine Park Heritage Strategy</i> to guide management actions to strengthen recognition and protection of heritage values</li> <li>• <b>REC14:</b> Promote, recognise and encourage stewardship and best practice efforts by community, industry and government</li> <li>• <b>REC17:</b> Support a collaborative, Reef-wide management strategy for islands and contribute to its development and implementation</li> <li>• <b>REC17A:</b> Work with the Queensland Government to provide technical and policy advice on actions to secure the long-term ecological, social and economic sustainability of Great Barrier Reef Region fisheries</li> <li>• <b>REC20:</b> Support research on critical ecosystem thresholds, with a focus on inshore biodiversity and associated ecosystems</li> <li>• <b>REC21:</b> Improve understanding and the Authority's management of the impacts of noise on species, particularly at-risk and inshore species</li> <li>• <b>REC23:</b> Develop a policy and supporting mechanisms to facilitate strategic and collaborative implementation of offsets across jurisdictions</li> <li>• <b>REC24:</b> Inform implementation of Australian and Queensland government offsets policies and restoration programs by identifying actions that will maximise the delivery of environmental benefits to the Region</li> <li>• <b>REC27:</b> Strengthen engagement with all relevant partners to facilitate actions that maintain and enhance the condition of values and reduce impacts, particularly in relation to climate change, catchment run-off, degradation of coastal ecosystems and direct use</li> <li>• <b>REC28:</b> Develop a comprehensive management framework and an Indigenous heritage strategy for Traditional Owner use and management of the Great Barrier Reef</li> <li>• <b>REC29:</b> Adopt regionally-based cooperative approaches to protect inshore biodiversity hotspots — supporting local actions and encouraging cooperation</li> <li>• <b>REC30:</b> Improve alignment and coordination of strategic research priorities and strengthen partnerships between the Authority and research institutions to facilitate the delivery of critical research needs</li> <li>• <b>REC31:</b> Establish an integrated monitoring and reporting program to support adaptive management for the Great Barrier Reef World Heritage Area, including more explicit reporting on the condition and trend of matters of national environmental significance, to enable the performance of activities to be monitored against regulatory objectives, outcomes for matters of national environmental significance and relevant guidelines and standards, including ecosystem thresholds</li> <li>• <b>REC32:</b> Maintain and improve monitoring, investigation and data management relating to critical species and habitats.</li> <li>• <b>REC33:</b> Support implementation of a long-term social and economic monitoring program to improve understanding of changing use, investment and values</li> <li>• <b>REC34:</b> Contribute to the development of improved governance arrangements for the management and coordination of development activities that affect the Great Barrier Reef</li> <li>• <b>REC 34A:</b> Establish a peak Great Barrier Reef Advisory Group made up of Traditional Owners, scientific, conservation and industry experts to provide high level advice on the implementation of the Authority's management program</li> <li>• <b>REC35:</b> Communicate the implications of climate change impacts for the Great Barrier Reef, and the critical need to halt increasing concentrations of global greenhouse gases and restore them to levels that will support growth, recruitment and recovery processes of the Great Barrier Reef ecosystem</li> <li>• <b>REC36:</b> Ensure the impacts of climate change and extreme weather are appropriately considered in the Authority's management decisions</li> <li>• <b>REC37:</b> Encourage reduction of greenhouse gas emissions in the Great Barrier Reef Region in partnership with industry and communities</li> <li>• <b>REC38:</b> Support initiatives to build the capacity of management agencies and Reef users to adapt and respond to climate change and extreme weather events</li> </ul>

## 13.6 Monitoring and reporting

### 13.6.1 Monitoring

Monitoring provides information to track the condition and trend of the Region's values, as well as the status of the impacts and activities affecting those values. It allows early detection of trends and the assessment of future risks, as well as playing a fundamental role in evaluating management effectiveness. It is also critical to informing the development, refinement and application of maps and models that help managers understand trends and patterns in the Region. Monitoring data are used to meet statutory reporting requirements such as the Authority's five-yearly Outlook Report and its annual report, as well as other reporting obligations including communicating the outcomes of *Reef Water Quality Protection Plan* actions.

The integrated monitoring and reporting program, outlined in Section 8 of the Program Report, will underpin the monitoring, evaluation and public reporting process to be employed by the Authority. Once implemented, the program will provide important information for management, support the Authority's outcome-based management approach, and be a vital component of the Authority's adaptive management of the Region.

The integrated monitoring and reporting program identifies three types of monitoring required for management:

- long-term core monitoring programs — to assess the condition and trend of the Region's values and broadscale impacts, such as water quality, over many years
- short to medium-term, issue-specific monitoring — to examine the condition of, extent of impact on and recovery rate of species, habitats or community benefits
- compliance monitoring — to target the impacts of a development action (for example, construction of a marina or a dredging program) in accordance with conditions specified in a permit, licence or approval.

The program will improve the integration and coordination of existing monitoring programs through the development and implementation of standardised protocols for information collection, collation and data sharing. It will improve the scalability of data (from point source or local, to regional and Reef-wide scales) and synthesis of information from different sources. This will provide a more comprehensive and systematic understanding of the condition of values and scale of impacts.

### 13.6.2 Review and public reporting

Research and monitoring as well as community-derived data is captured, synthesised and presented every five years through preparation of the Outlook Report. The Outlook Report is a statutory reporting responsibility for the Authority, providing a regular and reliable assessment of the Great Barrier Reef and its management. The outcomes of the integrated monitoring and reporting program will be synthesised and comprehensively reported in future Outlook Reports. In this way, the Outlook Report will be the primary reporting mechanism of the integrated monitoring and reporting program.

There are good examples of monitoring programs already tightly integrated into adaptive management, such as the marine monitoring program which operates under Reef Plan, and the monitoring of the effectiveness of the Zoning Plan. These programs are reported more regularly than the Outlook Report cycle because they are linked to specific management objectives and feed directly into assessments of management effectiveness. However, for many monitoring programs, linkages to management effectiveness are not as explicit. Through the integrated monitoring and reporting program, links to management will be made clearer and the causal relationships between drivers, pressures, impacts and condition will be better understood. Data from monitoring programs will be able to be synthesised with other data and readily incorporated into adaptive management processes.

### 13.6.3 Adapting to new information

Research is fundamental to interpreting monitoring data and understanding the effects of impacts on values. In addition, research analysis of existing datasets to uncover patterns and relationships and develop unifying concepts such as on ecosystem health, resilience and sustainability, substantially contribute to adaptive management. Synthesising data and other information is also vital to informing ecological and social risk assessments and developing standards and guidelines for the management of impacts.

Information from research and monitoring will be critical to implementing and adapting the new initiatives outlined in the Program Report — for example, the development of thresholds for ecosystem health, and targets for management action. Research will also be needed to conceptualise and, in some cases, model how a system works and how the elements interact and respond to changing pressures.

Traditional ecological knowledge and local community knowledge shared with the Authority by Traditional Owners, stakeholders and members of the community play a central role in informing its adaptive management and decision making. Combined with scientific information, such knowledge informs development of management strategies and policies and, where feasible, will continue to be used to monitor and assess the effectiveness of management arrangements (see Section 7 of the Program Report). Furthermore, the integration of traditional and community knowledge with scientific knowledge can extend the time perspective of scientific knowledge and highlight potential subject areas for future studies.<sup>10</sup>

### 13.6.4 Parties responsible for monitoring and reporting

The Authority will be the lead agency responsible for the implementation of the integrated monitoring and reporting program, in partnership with research institutions, government agencies, Traditional Owners, Reef-based industries and members of the community.

Monitoring activities to inform adaptive management will be conducted by:

- research institutions and government agencies (as identified above) where monitoring is carried out by scientists, technicians and field staff
- Traditional Owners of the Great Barrier Reef who are actively involved in monitoring and protecting the health and biodiversity of their sea country. This monitoring provides a valuable link between traditional ecological knowledge and modern science. Examples include training in scientific and wildlife monitoring as part of a two-way information exchange between Traditional Owners, scientists and management agencies.
- Reef-based industries where monitoring may be a voluntary contribution to management, undertaken to meet a permit condition, or a compulsory reporting obligation
- Members of the community, typically on a voluntary basis and associated with a recognised community monitoring program.

While the Authority will be responsible for ensuring it continues to meet its statutory monitoring and reporting functions, a whole of government approach, across all levels of government, such as that described in the integrated monitoring and reporting program, is required to coordinate and integrate monitoring and reporting functions across the Great Barrier Reef and its catchments.

### 13.7 Addressing the endorsement criteria

When deciding whether to endorse the Program, the Minister must be satisfied that the Strategic Assessment Report adequately addresses the impacts to which the Agreement (Appendix 1) relates and that any recommendations by the Minister to modify the program have been responded to appropriately.

Table 13.8 describes how the Authority's comprehensive management program (Part B) and permission system (Part C) meets the endorsement criteria as described in the Agreement (Appendix 1).

Table 13.8 Addressing endorsement criteria

Endorsement Criteria	Addressing the endorsement criteria in the Authority's comprehensive management program (Part B)	Addressing the endorsement criteria in the Authority's permission system (Part C)
In determining whether or not to endorse the program, the Minister will have regard to the extent to which the program meets the objects of the EPBC Act. In particular, that it:		
a) Protects the environment, especially those aspects of the environment that are matters of national environmental significance.	<ul style="list-style-type: none"> <li>The main object of the GBRMP Act is the long-term protection and conservation of the environment, biodiversity and heritage values of the Great Barrier Reef Region.</li> <li>Management focus on the Marine Park as a MNES means that values relevant to other MNES are implicitly considered in decision making.</li> </ul>	<ul style="list-style-type: none"> <li>The permission system ensures protection of the Great Barrier Reef Region, which necessarily includes matters of national environmental significance.</li> <li>The Authority commits to updating its policies to make explicit reference to consideration of matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area.</li> <li>The commitments in the Program Report, actions approved under permission system, ensure protection of the Great Barrier Reef Region, which necessarily includes protection of matters of national environmental significance.</li> </ul>
b) Promotes ecologically sustainable development through the conservation and ecologically sustainable use of natural resources.	<ul style="list-style-type: none"> <li>Consistent with the main object of the GBRMP Act.</li> <li>Ecologically sustainable use of the Region is allowed for only if it is consistent with the main object of the GBRMP Act.</li> <li>Maintaining sustainable multiple use and community benefits derived from the environment is one of the four goals of the Authority.</li> </ul>	<ul style="list-style-type: none"> <li>The permission system is a key component of the Authority's regulatory approach to achieve the objects of the GBRMP Act.</li> <li>The main object of the GBRMP Act is the long-term protection and conservation of the environment, biodiversity and heritage values of the Great Barrier Reef Region.</li> <li>Ecologically sustainable use of the Region is allowed for only if it is consistent with the main object.</li> </ul>
c) Promotes the conservation of biodiversity.	<ul style="list-style-type: none"> <li>Consistent with the main object of the GBRMP Act.</li> <li>Protecting and restoring the Reef's ecosystem health and biodiversity is one of the four goals of the Authority.</li> </ul>	<ul style="list-style-type: none"> <li>Consistent with the main object of the GBRMP Act.</li> <li>Part 13 of the EPBC Act provides mechanisms, including recovery plans, threat abatement plans, and conservation plans for migratory and marine species to promote the conservation of biodiversity.</li> <li>The commitments in the Program Report for the permission system ensure that assessment of activities that are likely to have a significant impact on matters of national environmental significance will consider all of the above mechanisms.</li> </ul>
d) Provides for the protection and conservation of heritage.	<ul style="list-style-type: none"> <li>Consistent with the main object of the GBRMP Act.</li> <li>Safeguarding the Reef's heritage values is one of the four goals of the Authority.</li> <li>Implementation of: <ul style="list-style-type: none"> <li><b>REC 5</b> to develop and implement knowledge management systems</li> </ul> </li> </ul>	<ul style="list-style-type: none"> <li>The main object of the GBRMP Act is the long-term protection and conservation of the environment, biodiversity and heritage values of the Great Barrier Reef Region.</li> <li>Consistent with 88Q(a) of the GBRMP Regulations, the Authority must consider the potential impacts of the conduct proposed on the environment and on the social, cultural and heritage values of the</li> </ul>



Endorsement Criteria	Addressing the endorsement criteria in the Authority's comprehensive management program (Part B)	Addressing the endorsement criteria in the Authority's permission system (Part C)
	<p>for Indigenous and historic heritage information</p> <ul style="list-style-type: none"> <li>- <b>REC 13</b> to review and update the <i>Great Barrier Reef Marine Park Heritage Strategy</i> to guide management actions to strengthen recognition and protection of heritage values</li> </ul> <p>will support protection and conservation of heritage.</p>	<p>Marine Park or a part of the Marine Park — this includes Indigenous social, cultural and heritage values.</p> <ul style="list-style-type: none"> <li>• The commitments made under the Program Report will require the Authority to amend its policies and guidance material (that will be updated from time to time) which are used in the determination of permissions, so they explicitly require the consideration of matters of national environmental significance and relevant attributes, including Indigenous and historic heritage values; and to develop and/or strengthen knowledge and information management tools, including systems for managing Indigenous and historic heritage.</li> </ul>
e) Promotes a cooperative approach to the protection and management of the environment.	<ul style="list-style-type: none"> <li>• Consistent with object 2A(b) of the GBRMP Act to encourage greater engagement in the protection and management of the Great Barrier Reef Region by interested persons and groups, including Queensland and local governments, communities, Indigenous persons, business and industry.</li> <li>• Engagement is one of the three areas of the Authority's foundational management program.</li> <li>• The Authority's management program focuses on enhancing protection measures within the Region, and influencing drivers and activities which affect the Region but fall outside of the Authority's jurisdictional control.</li> <li>• The Authority recognises that protection of the Great Barrier Reef requires local, national and international effort.</li> <li>• The Authority will continue to work with Traditional Owners, the community, business, industry, local government, plus Australian and Queensland government agencies to influence best practice and find pragmatic solutions to secure the future health of the Reef.</li> <li>• The Authority will continue to undertake engagement through partnerships, providing advice, education and community awareness, stewardship and best practice, and consultation.</li> </ul>	<ul style="list-style-type: none"> <li>• The permission system supports a cooperative approach to management, and includes joint permitting and a Joint Field Management Program with the Queensland Government.</li> <li>• Any permit application which may restrict the reasonable use by the public of a part of the Marine Park will be placed on the Authority's website, and will require public advertising as outlined in Regulation 88D of the GBRMP Regulations.</li> <li>• In accordance with Regulation 88Q(e) of the GBRMP Regulations, any written comments received about the application in response to the public advertisement are a mandatory consideration in the assessment of the permit application.</li> <li>• Consistent with section 2A(2)(b) of the GBRMP Act to encourage engagement in the protection and management of the Great Barrier Reef Region by interested persons and groups, including Queensland and local governments, communities, Indigenous persons, business and industry; where a permit application will likely have significant impacts on matters of national environmental significance, the Authority will publicly advertise for written comments on the application. Advertising would be done in the same manner as required under Regulation 88D of the GBRMP Regulations, and the information sought that would be to inform consideration of applications in accordance with regs 88Q and 88R of the GBRMP Regulations.</li> </ul>
f) Assists in the co-operative implementation of Australia's international environmental responsibilities.	<ul style="list-style-type: none"> <li>• Consistent with the main object of the GBRMP Act.</li> <li>• Consistent with object 2A(c) of the GBRMP Act to assist in meeting Australia's international responsibilities in relation to the environmental management and protection of world heritage (especially Australia's responsibilities under the World Heritage Convention).</li> </ul>	<ul style="list-style-type: none"> <li>• The permission system supports the objects of the GBRMP Act.</li> <li>• The commitments made under the Program Report will require the Authority to update its current administrative arrangements with the Department of the Environment, including for the transfer of information required for reporting to the World Heritage Committee on implementation of Australia's international environmental</li> </ul>

Endorsement Criteria	Addressing the endorsement criteria in the Authority's comprehensive management program (Part B)	Addressing the endorsement criteria in the Authority's permission system (Part C)
	<ul style="list-style-type: none"> <li>The <i>Great Barrier Reef Outlook Report</i> will contain a specific assessment of the condition, trends, threats and prospects for the outstanding universal value of the Great Barrier Reef World Heritage Area.</li> </ul>	responsibilities.
g) Recognises the role of Indigenous people in the conservation and ecologically sustainable use of Australia's biodiversity.	<ul style="list-style-type: none"> <li>Consistent with object 2A(b) of the GBRMP Act to encourage greater engagement in the protection and management of the Great Barrier Reef Region by interested persons and groups, including Queensland and local governments, communities, Indigenous persons, business and industry.</li> <li>Consistent with 10(6A) of the GBRMP Act which requires at least one member of the Authority (also referred to as the Board) to be an Indigenous person with knowledge of, or experience concerning, Indigenous issues relating to the Marine Park.</li> <li>Safeguarding the Reef's heritage values, including Indigenous heritage, is one of the four goals of the Authority.</li> <li>Implementation of: <ul style="list-style-type: none"> <li><b>REC4</b> to collaborate with Traditional Owners to undertake an assessment of the Indigenous heritage values of the Region;</li> <li><b>REC5</b> to develop and implement knowledge management systems for Indigenous and historic heritage information;</li> <li><b>REC13</b> to review and update the <i>Great Barrier Reef Marine Park Heritage Strategy</i> to guide management actions to strengthen recognition and protection of heritage values</li> <li><b>REC28</b> to develop a comprehensive management framework and an Indigenous heritage strategy for Traditional Owner use and management of the Great Barrier Reef</li> </ul> </li> </ul> <p>will increase the role of Indigenous people in managing the Marine Park.</p>	<ul style="list-style-type: none"> <li>The permission system supports the objects of the GBRMP Act.</li> <li>Consistent with Regulation 88Q(a) of the GBRMP Regulation, the Authority must consider the potential impacts of the conduct proposed on the environment and on the social, cultural and heritage values of the Marine Park or a part of the Marine Park — this includes Indigenous social, cultural and heritage values.</li> <li>The commitments made under the Program Report will require the Authority to amend its policies and guidance material (that will be updated from time to time) which are used in the determination of permissions, so they explicitly require the consideration of matters of national environmental significance and relevant attributes, including Indigenous and historic heritage values.</li> </ul>
h) Promotes the use of Indigenous peoples' knowledge of biodiversity with the involvement of, and in cooperation with the owners of the knowledge.	<ul style="list-style-type: none"> <li>Consistent with 10(6A) of the GBRMP Act which requires at least one member of the Authority (also referred to as the Board) to be an Indigenous person with knowledge of, or experience concerning, Indigenous issues relating to the Marine Park.</li> <li>The Authority has strong collaborative relationships with Traditional Owners, including through the development of Traditional Use of Marine Resources Agreements (TUMRAs) and the Indigenous Reef Advisory Committee.</li> <li>Implementation of: <ul style="list-style-type: none"> <li><b>REC4</b> to collaborate with Traditional Owners to undertake an assessment of the Indigenous heritage values of the Region;</li> <li><b>REC5</b> to develop and implement knowledge management systems</li> </ul> </li> </ul>	<ul style="list-style-type: none"> <li>The permission system supports the objects of the GBRMP Act.</li> <li>Consistent with Regulation 88Q(a) of the GBRMP Regulation, the Authority must consider the potential impacts of the conduct proposed on the environment and on the social, cultural and heritage values of the Marine Park or a part of the Marine Park — this includes Indigenous social, cultural and heritage values.</li> <li>The commitments made under the Program Report will require the Authority to develop and/or strengthen knowledge and information management tools including systems for managing Indigenous and historic heritage.</li> </ul>

Endorsement Criteria	Addressing the endorsement criteria in the Authority's comprehensive management program (Part B)	Addressing the endorsement criteria in the Authority's permission system (Part C)
	<p>for Indigenous and historic heritage information;</p> <ul style="list-style-type: none"> <li>- <b>REC13</b> to review and update the <i>Great Barrier Reef Marine Park Heritage Strategy</i> to guide management actions to strengthen recognition and protection of heritage values; and</li> <li>- <b>REC28</b> to develop a comprehensive management framework and an Indigenous heritage strategy for Traditional Owner use and management of the Great Barrier Reef</li> </ul> <p>will increase the capacity to use Indigenous peoples' knowledge in managing the Marine Park.</p>	
Without limiting the matters the Minister may consider when making the decision to endorse the program, the Minister will consider the manner in which the program:		
i) Identifies direct, indirect and cumulative impacts on matters of national environmental significance.	<ul style="list-style-type: none"> <li>• Reducing cumulative impacts is one of the four goals of the Authority.</li> <li>• The Authority will use the outcomes-based management framework, cumulative impact assessment guidelines and net benefit policy to build upon and better integrate current initiatives to reduce threats, restore degraded habitats, improve water quality, and re-establish connectivity and functioning of coastal ecosystems.</li> <li>• The Authority will systematically address the cumulative impacts on the Region's biodiversity, giving priority to inshore areas and those threats that have been identified for at-risk species, species groups and habitats.</li> </ul>	<ul style="list-style-type: none"> <li>• Consistent with Regulation 88Q(b) of the GBRMP Regulations, the Authority must consider options for monitoring, managing and mitigating the potential impacts of the conduct proposed in the permit application.</li> <li>• Under the commitments made in the Program Report, where an action is likely to have a significant impact on a matter of national environmental significance protected under the EPBC Act, the Authority will require applicants to consider impacts (including direct, indirect and cumulative impacts) arising from the development action in the context of all impacts affecting matters of national environmental significance.</li> <li>• The Authority has also committed to develop and implement cumulative impact assessment guidelines.</li> </ul>
j) Avoids impacts on matters of national environmental significance.	<ul style="list-style-type: none"> <li>• Consistent with the main object of the GBRMP Act.</li> <li>• Ecologically sustainable use of the Region is allowed for only if it is consistent with the main object of the GBRMP Act.</li> </ul>	<ul style="list-style-type: none"> <li>• Consistent with Regulation 88Q(b) of the GBRMP Regulations, the Authority must consider options for monitoring, managing and mitigating the potential impacts of the conduct proposed in the permit application.</li> </ul>
k) Mitigates the impacts on matters of national environmental significance.	<ul style="list-style-type: none"> <li>• Where the Authority has a high degree of control over activities (for example, tourism, research, defence, recreation) its effectiveness to avoid, mitigate, offset, deliver net benefits and adaptively manage impacts was found to be effective, or mostly effective.</li> </ul>	<ul style="list-style-type: none"> <li>• Avoid — the Authority will give highest priority to avoiding impacts on the environment and on the social, cultural and heritage values of the Marine Park (including matters of national environmental significance and outstanding universal value). Avoidance measures must consider prudent and feasible alternatives to a proposed activity. These should include, but not be limited to, consideration of alternative sites and alternate approaches to carrying out the activity.</li> </ul>
l) Offsets the impacts on matters of national environmental significance.	<ul style="list-style-type: none"> <li>• The permission system and partnerships with users of the Marine Park was considered a strength in this area.</li> <li>• The Authority will improve its management arrangements with: <ul style="list-style-type: none"> <li>- <b>REC9 – REC17</b> focused on avoiding impacts on MNES; <b>REC18 – REC12</b> focused on mitigating impacts on MNES; <b>REC23 – 24</b> focused on offsetting impacts on MNES.</li> </ul> </li> </ul>	<ul style="list-style-type: none"> <li>• Mitigate — potential impacts that cannot be avoided must be minimised. Mitigation measures must consider direct, indirect and</li> </ul>

Endorsement Criteria	Addressing the endorsement criteria in the Authority's comprehensive management program (Part B)	Addressing the endorsement criteria in the Authority's permission system (Part C)
		<p>cumulative impacts, and account for the likely spatial and temporal scales of impacts across the duration of the proposed activity.</p> <ul style="list-style-type: none"> <li>• Offset — offsets will only be considered where impacts cannot be avoided or mitigated, and where residual impacts will not exceed standards or thresholds for ecosystem health. Requirements for offsets will apply to all residual impacts. Offsets will be required to deliver measurable conservation outcomes within the timeframes relevant to affected matters of national environmental significance and related attributes and environmental processes.</li> <li>• The commitments made under the Program Report will require the Authority to develop a Great Barrier Reef offset policy (that will be updated from time to time) to enhance the condition of values relevant to matters of national environmental significance.</li> </ul>
m) Contributes to the enhancement of the existing environment and management of existing threats.	<ul style="list-style-type: none"> <li>• Consistent with the main object of the GBRMP Act.</li> <li>• Ecologically sustainable use of the Region is allowed for only if it is consistent with the main object of the GBRMP Act.</li> <li>• Chapter 7 of the Strategic Assessment Report assesses the current condition and trend for the Region's values and processes. Section 1.3 of the Program Report outlines the management framework based on outcomes and targets.</li> <li>• The desired outcome for each value is informed by the assessment of its current condition and trend as evaluated in the strategic assessment. This approach recognises that the condition and trend of each value and process varies across the Region.</li> <li>• The outcome for each value and process is a combination of the strategic assessment findings on condition and trend and the international and national obligations to protect them.</li> <li>• For example, if a value is assessed to be good condition, the outcome is that its condition is maintained and enhanced. However, if a value is assessed to be in poor condition, the outcome is that its condition is restored to good condition. If the trend of the value is assessed to be declining, the outcome is that the decline is halted and reversed.</li> <li>• The Authority will improve its management arrangements with <b>REC25 – REC30</b> focused on enhancing MNES.</li> </ul>	<ul style="list-style-type: none"> <li>• Consistent with 88Q(a) of the GBRMP Regulations, the Authority must consider the potential impacts of the conduct proposed on the environment and on the social, cultural and heritage values of the Marine Park or a part of the Marine Park.</li> <li>• The Authority applies an avoid-mitigate-offset-net benefit-adaptive management hierarchy considered in the assessment of impacts of activities on matters of national environmental significance.</li> <li>• The commitments made under the Program Report will require the Authority to develop a Great Barrier Reef net benefit policy (that will be updated from time to time) to enhance the condition of values relevant to matters of national environmental significance.</li> </ul>
n) Demonstrates adaption to reasonable climate change scenarios.	<ul style="list-style-type: none"> <li>• The strategic assessment identified climate change as one of four categories of high risk impacts.</li> <li>• The Authority has limited jurisdictional responsibility for addressing climate</li> </ul>	<ul style="list-style-type: none"> <li>• Consistent with 88Q(a) of the GBRMP Regulations, the Authority must consider the potential impacts of the conduct proposed on the environment and on the social, cultural and heritage values of the Marine Park or a part of the Marine Park; and 88Q(f) any other</li> </ul>



Endorsement Criteria	Addressing the endorsement criteria in the Authority's comprehensive management program (Part B)	Addressing the endorsement criteria in the Authority's permission system (Part C)
	<p>change in the broad sense. However, the Authority contributes significantly to the development of international best practice for managing responses to climate change and extreme weather issues as they relate to the Reef's marine ecosystems. This is chiefly done through research and monitoring, partnerships with research institutions, government agencies and stakeholder groups and education, community awareness and stakeholder engagement programs.</p> <ul style="list-style-type: none"> <li>• The Authority has a lead role in facilitating awareness of the impacts from climate change and extreme weather and in building resilience in the Marine Park.</li> <li>• The Authority has an advisory role to other agencies in relation to mitigation and adaptation to climate change and extreme weather in the Region.</li> <li>• The <i>Great Barrier Reef Climate Change Adaptation Strategy and Action Plan 2012–2017</i> acknowledges the important role the Authority plays in informing national and international climate policy and providing knowledge to support effective management of inshore areas.</li> <li>• The Authority will improve its management arrangements with <b>REC36 – REC39</b> focused on adapting to climate change.</li> </ul>	<p>matters relevant to the orderly and proper management of the Marine Park.</p> <ul style="list-style-type: none"> <li>• The Authority recognises the increasing significance of climate change and extreme weather in the Region and as one of the four high risk impacts (climate change, catchment run-off, degradation of coastal ecosystems and direct use), it will require consideration in permit assessments.</li> <li>• The commitments made under the Program Report will require the Authority to update its assessment processes so that climate change impacts are appropriately considered in its decision-making processes.</li> </ul>
o) Commitments in the program must be adequately resourced throughout its life	<ul style="list-style-type: none"> <li>• The Authority will continue foundational management arrangements and strengthen management within the Authority by realigning its current business systems and processes. More effective use of existing resources and stronger cooperation with partner agencies will ensure the Authority is well placed to build on its existing management arrangements.</li> <li>• The Authority will embed the strengthened management measures within existing policies, plans and programs, where appropriate. It will stage implementation of the remaining measures, including its new initiatives, based on their priority and the availability of resources.</li> </ul>	<ul style="list-style-type: none"> <li>• The Authority has committed to the implementation of the measures outlined in Part C of the Program Report.</li> <li>• The speed with which the commitments will be achieved will be based on priority and availability of resources.</li> </ul>

## 13.8 Program review, modification and continuous improvement

The terms of reference for this strategic assessment require that the Authority identify and analyse likely circumstances and procedures that may result in the review or modification of the program. The arrangements described below complement adaptive management measures set out in Chapter 13.

### 13.8.1 Review and continuous improvement of Program Report Part B

Under the GBRMP Act, the Authority has an ongoing responsibility to protect and manage the Great Barrier Reef. Over the life of the proposed program and beyond, the Authority will continue to aim for best practice management of the Region, adapting to changing conditions and government priorities as required. The Authority will continue the adaptive management approach outlined above to fulfil its responsibilities and respond to change.

It is anticipated the outcomes-based management framework, described in the accompanying Program Report, will provide sufficient flexibility to deal with changing circumstances over the next 25 years, and allow for modification as required.

By virtue of the provisions of the Act, the Authority has a clear and continuing responsibility to protect the Great Barrier Reef for future generations.

### 13.8.2 Review and continuous improvement of Program Report Part C

There will be an initial review of the permission system's progress in achieving objectives and outcomes for matters of national environmental significance 24 months following any approval of a class of actions, and submitted to the Department of the Environment within six months of commencement of the review. Subsequent program evaluation reviews will be aligned with the *Great Barrier Reef Outlook Report* (five yearly) and form part of the independent review of management effectiveness conducted for each *Outlook Report*.

The framework for the review will be agreed between the Authority and the Department of the Environment within six months following endorsement.

The purpose of the review will be to assess the performance of the permission system against objectives and outcomes for matters of national environmental significance including ensuring that impacts are not unacceptable. Each review will include a detailed evaluation of a sample of all relevant decisions made by the Authority to ensure appropriate consideration has been given to matters of national environmental significance.

The review's findings will be provided to the Minister for the Environment and will be published on the Department of the Environment's and the Authority's websites. The review will enable the Authority to determine if refinements to management arrangements and standards are required, to ensure the commitments and objectives for matters of national environmental significance are being delivered by the permission system.

---

## References

1. Great Barrier Reef Marine Park Authority 2012, *Informing the outlook for Great Barrier Reef coastal ecosystems*, GBRMPA, Townsville.
2. Bennett, J., Lawrence, P., Johnstone, R. and Shaw, R. 2005, Adaptive management and its role in managing Great Barrier Reef water quality, *Marine Pollution Bulletin* 51(1-4): 70-75.
3. Day, J. 2008, The need and practice of monitoring, evaluating and adapting marine planning and management—lessons from the Great Barrier Reef, *Marine Policy* 32(5): 823-831.
4. Beeden, R., Maynard, J.A., Marshall, P.A., Heron, S.F. and Willis, B.L. 2012, A framework for responding to coral disease outbreaks that facilitates adaptive management, *Environmental Management* 49(1): 1-13.
5. Dobbs, K., Day, J., Skeat, H., Baldwin, J., Molloy, F., McCook, L.J., Johnson, M., Elliot, B., Skeat, A., Vohland, K., Wachenfeld, D. and Kenchington, R. 2011, Developing a long-term outlook for the Great Barrier Reef, Australia: a framework for adaptive management reporting underpinning an ecosystem-based management approach, *Marine Policy* 35(2): 233-240.
6. Eberhard, R., Waterhouse, J., Robinson, C., Taylor, B., Hart, B., Parslow, J. and Grayson, R. 2008, Adaptive management strategies for Great Barrier Reef water quality improvement plans, *Geographical Research* 46: 303-313.
7. Eberhard, R., Robinson, C.J., Waterhouse, J., Parslow, J., Hart, B., Grayson, R. and Taylor, B. 2009, Adaptive management for water quality planning—from theory to practice, *Marine and Freshwater Research* 60(11): 1189-1195.
8. Hughes, T.P., Gunderson, L.H., Folke, C., Baird, A.H., Bellwood, D.R., Berkes, F., Crona, B., Helfgott, A., Leslie, H., Norberg, J., Nyström, M., Olsson, P., Osterblom, H., Scheffer, M., Schuttenberg, H., Steneck, R.S., Tengö, M., Troell, M., Walker, B., Wilson, J. and Worm, B. 2007, Adaptive management of the Great Barrier Reef and the Grand Canyon World Heritage Areas, *Ambio* 36(7): 586-592.
9. McCook, L.J., Ayling, T., Cappel, M., Choat, J.H., Evans, R.D., DeFreitas, D.M., Heupel, M., Hughes, T.P., Jones, G.P., Mapstone, B., Marsh, H., Mills, M., Molloy, F.J., Pitcher, C.R., Pressey, R.L., Russ, G.R., Sutton, S., Sweatman, H., Tobin, R., Wachenfeld, D. and Williamson, D. 2010, Adaptive management of the Great Barrier Reef: A globally significant demonstration of the benefits of networks of marine reserves, *Proceedings of the National Academy of Sciences* 107(43): 18278-18285.
10. Fernbach, M. and Nairn, K. 2007, *Reef Recollections: an oral history of the Great Barrier Reef*, Great Barrier Reef Marine Park Authority, Townsville.

## Appendix 1

---

### **Agreement with the then Minister for Sustainability, Environment, Water, Population and Communities**







**ENVIRONMENT PROTECTION AND BIODIVERSITY  
CONSERVATION ACT 1999  
Part 10 Strategic Assessments  
Section 146 Agreement**

Strategic assessment of the impacts of actions on the values of the Great Barrier Reef World Heritage Area, other relevant matters of national environmental significance, and the management arrangements to deal with such impacts under the jurisdiction of the Great Barrier Reef Marine Park Authority.

between

**THE MINISTER FOR SUSTAINABILITY, ENVIRONMENT, WATER, POPULATION AND COMMUNITIES ON  
BEHALF OF THE COMMONWEALTH GOVERNMENT**

and

**THE GREAT BARRIER REEF MARINE PARK AUTHORITY**

**CONTENTS**

1 PARTIES.....	3
2 DEFINITIONS .....	3
3 PREAMBLE .....	5
4 BACKGROUND .....	6
5 TERMS OF REFERENCE FOR THE STRATEGIC ASSESSMENT REPORT .....	6
6 PREPARATION OF THE STRATEGIC ASSESSMENT REPORT AND PROGRAM REPORT .....	7
7 ENDORSEMENT OF THE PROGRAM .....	8
8 APPROVAL OF ACTIONS UNDER THE PROGRAM .....	9
9 VARIATION, CONFLICT RESOLUTION AND TERMINATION .....	9
SIGNED BY: .....	10
Attachment A .....	11
Attachment B .....	12
Attachment C .....	13

## 1 PARTIES

The Parties to this agreement are:

The Minister for Sustainability, Environment, Water, Population and Communities on behalf of the Commonwealth Government

and

The Great Barrier Reef Marine Park Authority.

## 2 DEFINITIONS

2.1 Unless stated otherwise in this Agreement, the definitions, meanings and terms in the *Environment Protection and Biodiversity Conservation Act 1999* apply to this Agreement and its attachments.

2.2 In this Agreement:

**agreement** means this strategic assessment agreement entered into between the Minister for Sustainability, Environment, Water, Population and Communities and the Great Barrier Reef Marine Park Authority on the date the last Party executes this agreement.

**day** means a business day as measured in Canberra, Australian Capital Territory.

**Department** means the Australian Government department administering the *Environment Protection and Biodiversity Conservation Act 1999* (Cth).

**endorsement criteria** means the endorsement criteria detailed in Attachment C of this Agreement.

**EPBC Act** means the *Environment Protection and Biodiversity Conservation Act 1999* (Cth).

**GBRMP** means the Great Barrier Reef Marine Park as established by the *Great Barrier Reef Marine Park Act 1975* (Cth).

**GBRMPA** means the Great Barrier Reef Marine Park Authority as established by the *Great Barrier Reef Marine Park Act 1975* (Cth).

**GBRMP Act** means the *Great Barrier Reef Marine Park Act 1975* (Cth).

**GBRMPA management arrangements** means those management arrangements under the jurisdiction of GBRMPA, and may include partnership and collaborative arrangements with other Commonwealth and Queensland Government agencies and partnerships with stakeholders and members of the Great Barrier Reef coastal community. These management arrangements include, but are not limited to, those described in Attachment B.

**GBRWhA** means the Great Barrier Reef World Heritage Area and is the same as the area described in Schedule 1 to the GBRMP Act.

**Great Barrier Reef Region** has the same meaning as described in the GBRMP Act.

**Minister** means the Minister responsible for administering the EPBC Act or the Minister's authorised delegate.

**MNES** means matters of national environmental significance protected under Part 3 of the EPBC Act.

**Queensland strategic assessment** is defined in clause 3.5 of this Agreement.

**Parties** mean the Minister and the GBRMPA together, each a **Party**.

**Program** is the 'policy, plan or program' (section 146 of the EPBC Act) of the GBRMPA management arrangements that will apply to those parts of the strategic assessment area as described in the Program Report and for which endorsement by the Minister is sought.

**Program Report** means a report prepared specifically for this strategic assessment that describes the Program.

**strategic assessment** means the strategic assessment to which this Agreement relates, namely the strategic assessment of the impacts, including cumulative impacts, of actions on the environmental values of the strategic assessment area, relevant MNES and the GBRMPA management arrangements to deal with such impacts.

**strategic assessment area** is the Great Barrier Reef Region (map at Attachment A) and areas outside the Great Barrier Reef Region, to the extent that actions in those areas may affect the Great Barrier Reef Region.

**Strategic Assessment Report** means the report prepared in accordance with the Terms of Reference assessing the likely impacts, including cumulative impacts, on MNES from implementing the Program.

**Supplementary Report** means the report submitted to the Department following public exhibition of the draft Strategic Assessment Report which addresses and responds to public comments including revisions to the Strategic Assessment Report.

**Terms of Reference** means the terms of reference for the preparation of the Strategic Assessment Report.

**Work Plan** means a plan jointly developed by the Parties for undertaking the strategic assessment, documenting a common understanding about what strategic assessment tasks will be completed by whom, by when, and what resources are required for those tasks, amongst other things.

### 3 PREAMBLE

- 3.1 Australia's Great Barrier Reef is the largest coral reef ecosystem on earth. In 1975 the Great Barrier Reef Region was established and today provides for the long term protection and conservation of the environment, biodiversity and heritage values of the Great Barrier Reef Region.
- 3.2 The Great Barrier Reef Region extends more than 2,300 km along the Queensland coastline and covers 346,000 square kilometres. In 1981 the area was listed as a World Heritage property for its outstanding universal value and in 2007 it was listed as a National Heritage property.
- 3.3 Both the Australian and Queensland Governments have legislative responsibilities within the GBRWHA. The *Great Barrier Reef Intergovernmental Agreement 2009* sets out the joint management arrangements between the two governments to ensure an integrated and collaborative approach to the management of the marine and land environments within and adjacent to the GBRWHA.
- 3.4 The strategic assessment will examine the impacts, including cumulative impacts, of actions on the GBRWHA, other relevant MNES, and the GBRMPA management arrangements to deal with such impacts.
- 3.5 The strategic assessment will complement the Queensland Government's strategic assessment of the impacts of actions on the values of the GBRWHA under the Queensland coastal management, planning and development framework (the Queensland strategic assessment). The GBRMPA strategic assessment combined with the Queensland strategic assessment will form a comprehensive strategic assessment of the entire GBRWHA property and its values and the adjacent coastal zone.
- 3.6 The Parties agree that the GBRWHA and the adjacent coastal zone contain MNES that are highly inter-related. The Parties commit to:
- (a) undertaking a strategic assessment of the impacts, including cumulative impacts, of actions on all MNES under the jurisdiction of GBRMPA, other than nuclear actions.
  - (b) developing a Work Plan for the strategic assessment describing governance arrangements, key milestones and deliverables, public exhibition requirements, communication and data sharing agreements and other responsibilities of the Parties to ensure timely completion of the strategic assessment.
  - (c) working collaboratively with the Queensland Government on the complementary Queensland strategic assessment.
- 3.7 The strategic assessment does not affect the application of the EPBC Act or GBRMPA Act in the strategic assessment area in relation to actions or activities already approved or for which approval has been sought through the assessment and approval process
- (a) established under Parts 7, 8 and 9 of the EPBC Act; or
  - (b) established under Part 2A of the *Great Barrier Reef Marine Park Regulations 1983* (Cth).

### 4 BACKGROUND

- 4.1 Subsection 146(1) of the EPBC Act allows the Minister to agree in writing with a person responsible for the adoption or implementation of a policy, plan or program to assess the impacts of actions taken under that policy, plan or program on MNES.
- 4.2 This Agreement is intended to be an agreement for the purposes of subsection 146(1) of the EPBC Act to assess the impacts of actions taken under the Program.
- 4.3 A draft Strategic Assessment Report and draft Program Report will be made available for public comment for a minimum of at least 28 days. Following the public comment period, a Supplementary Report (addressing public comments) and a revised Program Report will be submitted to the Minister. After considering these, the Minister may decide to endorse the Program if satisfied that the reports adequately address the impacts and endorsement criteria to which this Agreement relates and that recommended modifications (if any) to the Program made by the Minister, or modifications having the same effect, have been made.
- 4.4 The Parties acknowledge that the endorsement of the Program itself does not constitute any approval for the taking of actions under Part 10 of the EPBC Act.
- 4.5 If the Minister endorses the Program, the Minister may then approve the taking of an action, or a class of actions, in accordance with the Program and the EPBC Act. The effect of this approval decision is that any actions or class of actions approved under section 146B would not need further approval by the Minister under the EPBC Act if taken in accordance with the endorsed Program.
- 4.6 The Parties acknowledge that, where proponents propose to take an action in accordance with the Program that is not the subject of an approval under section 146B, they are able to seek approval for that action through the assessment and approval process established under Parts 7, 8 and 9 of the EPBC Act. Paragraphs 87(3)(b) and 136(2)(e) of the EPBC Act provide that the strategic assessment report for a policy, plan or program must be taken into account in deciding the level of assessment and approval for actions that are referred under the EPBC Act.

### 5 TERMS OF REFERENCE FOR THE STRATEGIC ASSESSMENT REPORT

- 5.1 The Parties will prepare draft Terms of Reference for the strategic assessment report that include the following (at a minimum):
- Purpose and description of the Program;
  - Description of the area in which the Program will be implemented;
  - MNES affected by the Program;
  - Identification and analysis of the potential impacts, including cumulative impacts, on MNES;
  - Consideration of existing pressures on MNES;
  - Measures to enhance the existing environment, including management of existing threats;



- Measures to avoid, mitigate and offset likely impacts, including impacts of cumulative impacts, on MNES;
- Consideration of ecologically sustainable development principles;
- Adaptive management and risk management; Auditing and reporting requirements; and
- Review, modification and abandonment of the Program.

5.2 The GBRMPA will seek public comment on the draft Terms of Reference. The GBRMPA will provide the draft Terms of Reference for public comment by notice consistent with arrangements for other major environmental impact assessment projects, including:

- (a) posting on the GBRMPA's website;
- (b) publishing in newspapers circulating in Queensland; and
- (c) circulation to key stakeholder groups agreed between the Department and the GBRMPA.

5.3 The notice must state:

- (a) that the draft Terms of Reference are available for public comment;
- (b) how copies may be obtained;
- (c) the contact details for obtaining further information; and
- (d) that public comments are invited for a period of at least 28 days as specified by the Minister.

5.4 Following the public comment period for the draft Terms of Reference in accordance with clause 5.3(d), the GBRMPA will submit to the Minister:

- (a) proposed Terms of Reference;
- (b) public responses relating to the draft Terms of Reference; and
- (c) a report on how the public responses have been taken into account.

5.5 Following receipt of the proposed Terms of Reference, the Minister will notify the GBRMPA that the proposed Terms of Reference:

- (a) are approved, or
- (b) are not approved, in which case the Minister will notify the GBRMPA of any concerns and invite the GBRMPA to provide revised Terms of Reference which take those concerns into account.

5.6 Within 15 working days of receipt of the revised Terms of Reference from the GBRMPA in accordance with clause 5.5(b), the Minister will either:

- (a) notify the GBRMPA of the Minister's approval of the revised Terms of Reference; or
- (b) provide approved Terms of Reference that meet the Minister's requirements.

## 6 PREPARATION OF THE STRATEGIC ASSESSMENT REPORT AND PROGRAM REPORT

6.1 The GBRMPA will prepare a Strategic Assessment Report and Program Report in accordance with this Agreement and the Terms of Reference approved in accordance with clause 5.5(a) or clause 5.6.

6.2 The GBRMPA will provide a draft Strategic Assessment Report and draft Program Report to the Minister for comment prior to finalising the documents for public comment.

6.3 The GBRMPA must release the draft Strategic Assessment Report and draft Program Report for public comment by notice consistent with arrangements for other major environmental impact assessment projects, including:

- (a) posting on the GBRMPA's website;
- (b) publishing in newspapers circulating in Queensland; and
- (c) circulation to key stakeholder groups agreed between the Department and the GBRMPA.

6.4 The notice must state:

- (a) that the draft Strategic Assessment Report and draft Program Report are available for public comment;
- (b) how copies may be obtained;
- (c) provide contact details for obtaining further information; and
- (d) that public comments are invited on the draft reports for a period of at least 28 days as specified by the Minister.

6.5 The GBRMPA will:

- (a) prepare a Supplementary Report and revise the Program Report, taking into account any public comments received in accordance with clause 6.4(d);
- (b) provide drafts of these reports to the Minister for comment prior to their finalisation; and
- (c) incorporate any recommendations by the Minister.

6.6 The Department agrees to assist the GBRMPA in ensuring that the reports adequately address the requirements for strategic assessments described in Part 10 of the EPBC Act and to provide comments in a timely manner.

## 7 ENDORSEMENT OF THE PROGRAM

7.1 The GBRMPA will submit to the Minister:

- (a) the Strategic Assessment Report (which was exhibited for public comment);
- (b) the Supplementary Report (explaining how relevant public responses have been taken into account and addressed in the impact assessment and revised Program);
- (c) the Program Report (incorporating any revisions in light of public comments);
- (d) public comments received during the consultation relating to the Strategic Assessment Report and Program Report;
- (e) any other documents required to support the GBRMPA's submission.

7.2 If the Minister is not satisfied that the reports adequately address the impacts of actions on MNES in the strategic assessment area, or that the Program does not provide for adequate protection of MNES, then:

- (a) the Minister can make recommendations to amend the Program;
- (b) the GBRMPA may seek clarification from the Minister on the recommendations;
- (c) the GBRMPA will then submit to the Minister for consideration the revised Program, and a summary of how the Minister's recommendations were given effect;



- (d) the Minister will consider the revised Program, and any supporting material provided, and may accept it as the final Program; and
- (e) the Minister may request further modifications if still not satisfied that the Program provides for adequate protection of MNES.

7.3 The Minister may endorse the Program if satisfied that the reports submitted under clause 7.1 adequately address the impacts to which this Agreement relates, and that any recommended modifications to the Program (clause 7.2) or modifications having the same effect have been made. In considering whether to endorse the Program the Minister will also consider the endorsement criteria at Attachment C.

## 8 APPROVAL OF ACTIONS UNDER THE PROGRAM

8.1 The Minister may approve, or approve with conditions, the taking of an action or class of actions in accordance with the endorsed Program under Section 146B of the EPBC Act. In doing so, the Minister must act in accordance with sections 146F-M of the EPBC Act. This includes considering MNES in the area affected by the Program and economic and social matters.

8.2 The Minister will seek comment from other Australian Government Ministers with administrative responsibilities relating to the actions before approving the taking of an action, or class of actions, pursuant to section 146C of the EPBC Act.

8.3 Both Parties will make publicly available electronically through their websites the endorsed Program (including the Program Report), the Strategic Assessment Report, the Supplementary Report and any approval decision and conditions.

8.4 Approvals for actions or classes of actions may occur progressively after endorsement of the Program.

## 9 VARIATION, CONFLICT RESOLUTION AND TERMINATION

9.1 This Agreement may only be varied by written agreement (including electronic communications) between the Parties and in accordance with the EPBC Act.

9.2 Where there is a dispute between the Parties to this Agreement on a particular matter, the Parties will consult in the spirit of mutual cooperation in relation to that matter and will use their best endeavours to negotiate a mutually acceptable resolution.

9.3 Either party may at any time terminate this Agreement by written correspondence to the other Party.

9.4 The Parties do not intend this Agreement to create contractual or other legal obligations, or that a breach of the Agreement will give rise to any cause of action, or right to take legal proceedings.

9.5 To avoid any doubt, nothing in clause 9.4 is intended to override the provisions of the EPBC Act.

SIGNED BY:



The Hon. Tony Burke MP  
Minister for Sustainability, Environment, Water,  
Population and Communities

16.2.12

Dated



Dr Russell Reichelt  
Chairperson, Great Barrier Reef Marine Park Authority

Dated 6/1/2012

## Attachment A

### STRATEGIC ASSESSMENT AREA

The strategic assessment area is the Great Barrier Reef Region (refer Figure 1) and areas outside the Great Barrier Reef Region, to the extent that actions in those areas may affect the Great Barrier Reef Region.

Figure 1: Great Barrier Reef Region



## Attachment B

### GBRMPA MANAGEMENT ARRANGEMENTS

Management of the GBRWHA relies upon a number of Commonwealth and Queensland agencies using a combination of management tools (including zoning plans, plans of management, permits, species recovery plans, the Reef Water Quality Protection Plan and shipping management arrangements), along with various management approaches (including education, planning, environmental impact assessment, monitoring, stewardship, enforcement) to regulate access, and to control and/or mitigate impacts associated with activities (such as tourism, fisheries, shipping) or address pressures (including climate change or declining water quality).

For the purpose of this strategic assessment, *GBRMPA's management arrangements* mean those management arrangements under the jurisdiction of GBRMPA. They include, but are not limited to, the following:

- Statutory instruments, including regulations, zoning plans, plans of management and permits;
- Non-statutory mechanisms including policies, position statements and guidelines;
- Partnership and collaborative arrangements with other Commonwealth and Queensland Government agencies;
- Partnerships with Traditional Owners in the management of marine resources;
- Partnership and stewardship programs, including education programs and engagement with local governments, communities, Indigenous persons and industry; and
- Research and monitoring programs.

## Attachment C

### ENDORSEMENT CRITERIA FOR STRATEGIC ASSESSMENT

When deciding whether to endorse the Program, the Minister must be satisfied that the Strategic Assessment Report adequately addresses the impacts to which this Agreement relates, and that any recommendations by the Minister to modify the Program have been responded to appropriately.

In determining whether or not to endorse the Program, the Minister will have regard to the extent to which the Program meets the objects of the EPBC Act. In particular, that it:

- protects the environment, especially those aspects of the environment that are MNES;
- promotes ecologically sustainable development through the conservation and ecologically sustainable use of natural resources;
- promotes the conservation of biodiversity;
- provides for the protection and conservation of heritage;
- promotes a cooperative approach to the protection and management of the environment; and
- assists in the co-operative implementation of Australia's international environmental responsibilities.

Without limiting the matters the Minister may consider when making the decision to endorse the Program, the Minister will consider the manner in which the Program:

- identifies direct, indirect and cumulative impacts on MNES;
- avoids impacts on MNES;
- mitigates the impacts on MNES;
- offsets the impacts on MNES;
- contributes to the enhancement of the existing environment and management of existing threats; and
- demonstrates adaption to reasonable climate change scenarios.

Commitments in the Program must be adequately resourced throughout its life. The Program must demonstrate an effective system of adaptive management that addresses uncertainty and contingency management as well as procedures for monitoring, independent auditing and public reporting on implementation.

The Program must address all of the above matters for it to be considered for endorsement by the Minister in accordance with the EPBC Act.






## Appendix 2

---

# Great Barrier Reef Region Strategic Assessment Terms of Reference





© Commonwealth of Australia  
Published by the Great Barrier Reef Marine Park Authority August 2012

ISBN 978-1-921682-89-6

This work is copyright. Apart from any use as permitted under the *Copyright Act 1968*, no part may be reproduced by any process without the prior written permission of the Great Barrier Reef Marine Park Authority.

Requests and inquiries concerning reproduction and rights should be addressed to:

General Manager, Communication and Policy Coordination  
Great Barrier Reef Marine Park Authority,  
PO Box 1379, Townsville Qld 4810.

Further information is available from:



**Australian Government**  
**Great Barrier Reef**  
**Marine Park Authority**

2-68 Flinders Street East (PO Box 1379)  
Townsville Queensland 4810  
Telephone +617 4750 0700  
Fax +617 4772 6093  
Web site [www.gbrmpa.gov.au](http://www.gbrmpa.gov.au)

## TABLE OF CONTENTS

### INTRODUCTION ..... 1

Background .....	1
Purpose .....	1
Scope .....	1
Context .....	2
Great Barrier Reef and Coastal Zone values .....	2
Reports to be prepared .....	3
Timeframes .....	3
Development of the Terms of Reference .....	4
Strategic Assessment process .....	4
Use of the term 'Program' .....	4

### TERMS OF REFERENCE ..... 5

1. Purpose and Description of the Program .....	5
2. Matters of National Environmental Significance .....	5
3. Assessment of Impacts on Matters of National Environmental Significance .....	6
4. Measures to Address Impacts .....	6
5. Projected Condition of Matters of National Environmental Significance .....	8
6. Proposed Program .....	8
7. Strategic Assessment Process .....	9
8. Endorsement Criteria .....	10

TERMS OF REFERENCE



## INTRODUCTION

### Background

The Australian and Queensland governments are working together to undertake a comprehensive strategic assessment of the Great Barrier Reef World Heritage Area and adjacent coastal zone. The comprehensive strategic assessment will help identify, plan for and manage existing and emerging risks so that the unique values of the Great Barrier Reef are protected and managed.

There are two components to the comprehensive strategic assessment - a marine component and a coastal component. The Great Barrier Reef Marine Park Authority (the Authority) will lead the marine component involving a strategic assessment of the Great Barrier Reef Region (the Strategic Assessment). The Queensland Government will lead the coastal component involving a strategic assessment of the Great Barrier Reef Coastal Zone.

### Purpose

The purpose of these Terms of Reference is to set out the requirements for the Authority in preparing the Strategic Assessment Report for the Great Barrier Reef Region Strategic Assessment.

The Strategic Assessment will assess the likely impacts of actions on relevant matters of national environmental significance as defined in the *Environment Protection and Biodiversity Conservation Act 1999*, including the Outstanding Universal Value of the Great Barrier Reef World Heritage Area, and the Authority's management arrangements to deal with such impacts.



Map 1: Great Barrier Reef Region

### Scope

The scope of the Strategic Assessment is defined by both geographic area and the management arrangements within the Authority's jurisdiction.

The area to be considered by the Authority in its Strategic Assessment is the Great Barrier Reef Region, as defined in the *Great Barrier Reef Marine Park Act 1975*, and areas outside that Region to the extent that actions in those areas may affect that Region (the Strategic Assessment area). The Great Barrier Reef Region covers an area from the tip of Cape York in the north to past Lady Elliot Island in the south, with mean low water as its western boundary and extending eastwards a distance of between 70 and 250 km (Map 1).

All management arrangements under the jurisdiction of the Authority will be considered together with partnerships and collaborative arrangements with other Queensland and Australian government agencies and partnerships with stakeholders and members of the Great Barrier Reef coastal community.

TERMS OF REFERENCE

1

The Authority and the Queensland Government will work closely together on those aspects of the two strategic assessments where they have joint management responsibilities, for example shipping, water quality and island management.

Due to the size and complexity of the Great Barrier Reef, the Authority's Strategic Assessment will include both a broad examination of the Great Barrier Reef Region and more focussed studies of specific locations and issues.

### Context

The Australian and Queensland governments share responsibility for managing the Great Barrier Reef. The adaptive, ecosystem-based management approach for the Great Barrier Reef is constantly reviewed and updated in response to new and emerging issues.

In 2009, the Great Barrier Reef Outlook Report identified that the Great Barrier Reef ecosystem is at a crossroad and decisions made now are likely to determine its long-term future. Climate change, declining water quality from catchment runoff, coastal development and remaining impacts from fishing were identified as the biggest risks to the future of the Reef. Since this 2009 report, some of the identified risks to the Reef have increased. These include increases in shipping activity as a result of port expansions; population growth as a result of expanding urban and industrial activities along the Great Barrier Reef coast; intensification and changes in land use within the Great Barrier Reef catchment; and extreme weather events including flooding and cyclones.

The comprehensive strategic assessment is part of the Australian and Queensland governments' adaptive approach to managing the Great Barrier Reef. Once complete the comprehensive strategic assessment will strengthen the protection of the Great Barrier Reef and guide its management by providing greater certainty on where sustainable uses can occur, the type of activities that will be allowed and the conditions under which activities may proceed.

The comprehensive strategic assessment is being carried out under Part 10 of the *Environment Protection and Biodiversity Conservation Act 1999*. Agreements between the Minister for Sustainability, Environment, Water, Population and Communities (the Minister) and both the Authority and the Queensland Government to undertake the complementary strategic assessments are available at: <http://www.environment.gov.au/epbc/notices/assessments/great-barrier-reef.html>

The comprehensive strategic assessment also forms part of the Australian Government's response to the World Heritage Committee's concerns regarding the impacts of development on the Great Barrier Reef World Heritage Area.

### Great Barrier Reef and Coastal Zone values

In accordance with the *Environment Protection and Biodiversity Conservation Act 1999*, the assessment will address impacts on **matters of national environmental significance** as defined in that Act, including world heritage properties, national heritage properties, nationally threatened species and ecological communities, nationally listed migratory species, wetlands of international importance, Commonwealth marine areas and the Great Barrier Reef Marine Park. The Strategic Assessment will describe the extent to which these matters apply to the Great Barrier Reef Region, the values which underpin these matters and ecosystem processes critical to their functioning.

As part of the strategic assessment, there will also be explicit consideration of matters relating to the Outstanding Universal Value of the Great Barrier Reef World Heritage Area.

Outstanding Universal Value is a concept central to world heritage properties. It relates to the exceptional qualities of global significance that make an area worthy of special protection. The Great Barrier Reef was listed as a world heritage property in 1981 on the basis that it met all four of the natural criteria which contribute to its Outstanding Universal Value. These criteria relate to the property's biodiversity, its significant ongoing biological and geological processes, its superlative natural phenomena and exceptional natural beauty, and its evolutionary history. Outstanding Universal Value also includes the concept of 'integrity' which is a measure of wholeness or intactness of the property's natural heritage and its attributes.

2

Great Barrier Reef Region Strategic Assessment

## Reports to be prepared

The Great Barrier Reef Region Strategic Assessment will deliver two reports:

- **Program Report** containing a detailed description of the Authority's management arrangements, including future commitments, to protect and manage matters of national environmental significance, including the Outstanding Universal Value of the Great Barrier Reef World Heritage Area.
- **Strategic Assessment Report** prepared in accordance with these Terms of Reference.

The Great Barrier Reef Coastal Zone component of the comprehensive strategic assessment will be made up of two similar reports, as shown in Figure 1. Where there are joint management responsibilities these aspects will be addressed in both the Authority's and Queensland Government's reports.




Figure 1: Reports of the Comprehensive Strategic Assessment for the Great Barrier Reef World Heritage Area and Adjacent Coastal Zone

Drafts of the Program Report and the Strategic Assessment Report will be released together for public comment prior to their finalisation and submission to the Minister for Sustainability, Environment, Water, Population and Communities.

## Timeframes

It is anticipated that a draft Strategic Assessment Report and Program Report for the Great Barrier Reef Region Strategic Assessment will be made available for public comment early to mid 2013.

It is intended that the comprehensive strategic assessment for the Great Barrier Reef World Heritage Area and adjacent coastal zone will be finalised by mid to late 2013.

Prior to finalisation of the comprehensive strategic assessment, the Australian and Queensland governments will continue to apply high environmental standards when considering development proposals that may affect the Great Barrier Reef World Heritage Area.

The life of the Program described in the Program Report will be 25 years, during which time there will be monitoring, evaluation and periodic review of the Program.

## Development of the Terms of Reference

To ensure a consistent approach to developing the two strategic assessments, the Terms of Reference have been developed in close cooperation with the Queensland Government and the Australian Government's Department of Sustainability, Environment, Water, Population and Communities.

Draft Terms of Reference were made available for public comment between 18 February and 30 April 2012. A report on submissions received and how public comments were considered in finalising the Terms of Reference is available on the Authority's website at [www.gbmpa.gov.au](http://www.gbmpa.gov.au).

## Strategic Assessment process

In preparing the Strategic Assessment, the Authority will:

- Work in close consultation and collaboration with the Queensland Government, the Australian Government's Department of Sustainability, Environment, Water, Population and Communities and other relevant Australian Government agencies.
- Engage stakeholders and the community throughout the Strategic Assessment process using the Authority's network of advisory committees together with regional and issue-based meetings.
- Communicate with the wider community throughout the Strategic Assessment process using the Authority's website, fact sheets, media releases, advertisements and newsletters.
- Use methods consistent with, and build upon approaches employed in the Great Barrier Reef Outlook Report 2009, where relevant.
- Engage independent expertise to assess the effectiveness of the Authority's Program and commission a peer review of the Strategic Assessment Report.
- Use the best available information to undertake the Strategic Assessment, including scientific data, expert opinion, and Traditional Owner and community knowledge.

## Use of the term 'Program'

The word 'Program' is used throughout the Terms of Reference. In simple terms it means those management arrangements ('policy, plan or program') under the jurisdiction of the Authority. Management arrangements include, but are not limited to, those described in section 4.1.2 of the Terms of Reference.



## TERMS OF REFERENCE

In preparing the Great Barrier Reef Region Strategic Assessment Report, the Authority must comply with the requirements set out below.

### 1. Purpose and Description of the Program

- 1.1 Provide an overview of the current Program. For the purposes of the Strategic Assessment the life of the Program is 25 years. The overview is to include a description of:
- The purpose of the Program, including Program objectives.
  - The area to which the Program applies (the Strategic Assessment area).
  - Legislation, plans, policies and other mechanisms that make up the Program, including Program commitments.
  - Relevant activities within the scope of the Program.
  - International, national, state and regional context (environmental, social, cultural and economic) in which the Program operates, including activities outside the Strategic Assessment area that may influence the Program.
  - Relevant national, state, regional and local planning or management frameworks that affect the Program and contribute to protection and management the matters of national environmental significance.
  - Identification of how long the Program will be in effect and the process for review of the Program, including adaptive management.
  - Identification of the relevant authorities responsible for the implementation of the Program.

### 2. Matters of National Environmental Significance

#### 2.1 Description of matters of national environmental significance

- 2.1.1 Describe the extent to which the following relevant matters of national environmental significance, as defined in the *Environment Protection and Biodiversity Conservation Act 1999* apply to the Strategic Assessment area.
- world heritage properties, including consideration of the Outstanding Universal Value of the Great Barrier Reef World Heritage Area
  - national heritage places
  - wetlands of international importance
  - listed threatened species and ecological communities
  - listed migratory species
  - Commonwealth marine areas
  - the Great Barrier Reef Marine Park.

The description must:

- Identify key terrestrial, coastal and marine environmental, biodiversity and heritage values and/or attributes which underpin the relevant matters of national environmental significance, including the Outstanding Universal Value of the Great Barrier Reef World Heritage Area.

TERMS OF REFERENCE

5

- Describe ecosystem processes considered critical to the functioning of the relevant matters of national environmental significance, including the Outstanding Universal Value of the Great Barrier Reef World Heritage Area.
- Provide sufficient information to allow an understanding the connectivity between the relevant matters of national environmental significance, including the Outstanding Universal Value of the Great Barrier Reef World Heritage Area.

#### 2.2 Condition and trend of matters of national environmental significance

- Describe the current condition and trend of key indicators of the relevant matters of national environmental significance, including the Outstanding Universal Value of the Great Barrier Reef World Heritage Area.
- For world heritage values, benchmark the current condition of key indicators of Outstanding Universal Value against the retrospective statement of Outstanding Universal Value which describes the state of the Great Barrier Reef World Heritage Area at the time of listing in 1981.
- Identify key information gaps and processes to address critical information needs.

### 3. Assessment of Impacts on Matters of National Environmental Significance

#### 3.1 Actual and potential impacts

- Describe the environmental, social, cultural and economic drivers affecting the relevant matters of national environmental significance, including the Outstanding Universal Value of the Great Barrier Reef World Heritage Area.
- Describe and analyse the actual and potential impacts on the relevant matters of national environmental significance, including the Outstanding Universal Value of the Great Barrier Reef World Heritage Area, in the Strategic Assessment area, including:
  - impacts from past, present and future activities
  - direct, indirect, consequential and cumulative impacts
  - the likely impacts of climate change.
- Describe the spatial and temporal scale at which impacts and their effects on the relevant matters of national environmental significance, including the Outstanding Universal Value of the Great Barrier Reef World Heritage Area, are occurring or are likely to occur.
- Identify key information gaps and processes to address critical information needs.

### 4. Measures to Address Impacts

#### 4.1 Current Program

4.1.1 Describe and assess the effectiveness of the Program to:

- Identify the relevant matters of national environmental significance, including the Outstanding Universal Value of the Great Barrier Reef World Heritage Area, and determine their current condition and trend, including spatial and non-spatial approaches.
- Identify and analyse direct, indirect, consequential and cumulative impacts, including the methods used to determine these types of impacts.
- Consider environmental, social, cultural and economic issues.
- Avoid, mitigate, offset and adaptively manage impacts.
- Address uncertainty and risk.

6

Great Barrier Reef Region Strategic Assessment

- f) Provide certainty regarding where uses may occur, the type of activities allowed, conditions under which activities may proceed and circumstances where impacts are likely to be unacceptable.
- g) Halt and reverse any declines and enhance the condition of the relevant matters of national environmental significance, including mechanisms to deliver a 'net benefit' to the condition of the relevant matters of national environmental significance, including the Outstanding Universal Value of the Great Barrier Reef World Heritage Area,
- h) Adapt to reasonable climate change scenarios.
- i) Integrate with related local, Queensland and Australian government programs to protect and manage the relevant matters of national environmental significance, including the Outstanding Universal Value of the Great Barrier Reef World Heritage Area.
- j) Meet Australia's international responsibilities in relation to the environment and protection of world heritage.
- k) Monitor, evaluate and report on the:
  - i. Condition and trends of the relevant matters of national environmental significance, including the Outstanding Universal Value of the Great Barrier Reef World Heritage Area.
  - ii. Impacts of activities, including the setting of targets to benchmark management performance.

4.1.2 In assessing the effectiveness of the Program, consideration must be given to:

- a) Statutory instruments, including regulations, zoning plans, plans of management and permits.
- b) Non-statutory mechanisms including policies, position statements and guidelines.
- c) Partnership and collaborative arrangements with Queensland and other Australian government agencies.
- d) Partnerships with Traditional Owners in the management of marine resources.
- e) Partnership and stewardship programs, including education programs and engagement, with local governments, communities, Indigenous persons, business and industry.
- f) Research and monitoring programs.
- g) Compliance and enforcement programs.
- h) Resourcing of the Program.

#### 4.2 Demonstration cases

- 4.2.1 Develop demonstration cases to assess in finer detail the effectiveness of the Program to protect and manage the relevant matters of national environmental significance, including the Outstanding Universal Value of the Great Barrier Reef World Heritage Area, and to guide improvements to the Program.
- 4.2.2 Demonstration cases are to be chosen by the Australian and Queensland governments. Criteria that will be used to guide this selection process include, but are not limited to:
  - a) Where multiple impacts are acting or predicted to act upon a region, locality or value.
  - b) To examine a specific management approach or method to identify a set of values/attributes or to assess a range of impacts/pressures.
  - c) To demonstrate connectivity across coastal and marine systems.
  - d) To demonstrate the integration of environmental, social, cultural and economic benefits in decision-making.
  - e) To improve understanding of factors affecting Great Barrier Reef ecosystem resilience.
  - f) Where lessons or outcomes could transfer to other areas.

- g) Opportunities to build capacity for future management.
- h) To examine the effectiveness of management across local, Queensland and Australian government jurisdictions.

## 5. Projected Condition of Matters of National Environmental Significance

- 5.1 Describe the projected condition of the relevant matters of national environmental significance, including the Outstanding Universal Value of the Great Barrier Reef World Heritage Area, based on an evaluation of their:
  - a) Current status and trends.
  - b) Actual and potential impacts.
  - c) The effectiveness of the Program to protect the relevant matters of national environmental significance, including the Outstanding Universal Value of the Great Barrier Reef World Heritage Area, and manage impacts.
  - d) An understanding of ecosystem resilience.
  - e) An assessment of overall risks to the relevant matters of national environmental significance, including the Outstanding Universal Value of the Great Barrier Reef World Heritage Area.

## 6. Proposed Program

### 6.1 Recommendations for changes to the Program

- 6.1.1 Recommend changes to the current Program to improve its effectiveness to deliver its objectives, including outcomes that protect the relevant matters of national environmental significance, including the Outstanding Universal Value of the Great Barrier Reef World Heritage Area. Recommendations for improvements should specifically address the matters listed in section 4.1.1.

- 6.1.2 Consistent with section 6.1.1 above, recommend improvements to related local, state and national government programs.

### 6.2 Principles of Ecologically Sustainable Development

Describe how the principles of ecologically sustainable development have been applied in the proposed Program. The principles of ecologically sustainable development as described in section 3A of the *Environment Protection and Biodiversity Conservation Act 1999* are:

- a) Decision-making processes should effectively integrate both long-term and short-term economic, environmental, social and equitable considerations.
- b) If there are threats of serious or irreversible environmental damage, lack of full scientific certainty should not be used as a reason for postponing measures to prevent environmental degradation.
- c) The principle of inter-generational equity—that the present generation should ensure that the health, diversity and productivity of the environment is maintained or enhanced for the benefit of future generations.
- d) The conservation of biological diversity and ecological integrity should be a fundamental consideration in decision-making.
- e) Improved valuation, pricing and incentive mechanisms should be promoted.

### 6.3 Adaptive management

- a) Identify the key adaptive management measures in the proposed Program that address uncertainties and risks inherent in the decision making process.



- b) Describe how the adaptive management measures will be implemented to ensure the relevant matters of national environmental significance, including the Outstanding Universal Value of the Great Barrier Reef World Heritage Area, are effectively protected and managed over the life of the Program.

#### 6.4 Monitoring and reporting

- a) Describe the monitoring, review and public reporting process that will be used to examine whether the Program adequately protects and manages the relevant matters of national environmental significance, including the Outstanding Universal Value of the Great Barrier Reef World Heritage Area, and the framework for measuring success.
- b) Describe the processes for adapting the Program in response to new information.
- c) Nominate the parties responsible for undertaking the monitoring, review and reporting and for implementing any actions arising.

#### 6.5 Review, modification or abandonment

- a) Identify and analyse likely circumstances and procedures that may result in the review, modification or abandonment of the Program. This is to include a discussion of how any commitments under the Program would continue to be met under these situations.

## 7. Strategic Assessment Process

#### 7.1 Collaboration with the Queensland Government and other Australian Government agencies

- a) Undertake the Strategic Assessment in consultation and collaboration with the Queensland Government, the Australian Government's Department of Sustainability, Environment, Water, Population and Communities and other relevant Australian Government agencies.

#### 7.2 Community engagement

- a) Document how the community and stakeholders were engaged in the Strategic Assessment process and how views and comments were taken into account in the preparation of the Strategic Assessment Report and the Program Report.
- b) Along with the draft Program Report, make the draft Strategic Assessment Report available for public comment for a period of no less than 28 days.
- c) Provide the Minister with a report on the public submissions received on the draft Reports, together with proposed final drafts of the Strategic Assessment Report and Program Report, incorporating any revisions made in response to public comments.

#### 7.3 Independent review

- a) Engage independent expertise to undertake an assessment of the effectiveness of the current Program to protect and manage the relevant matters of national environmental significance, including the Outstanding Universal Value of the Great Barrier Reef World Heritage Area.
- b) Arrange for the content of the draft Strategic Assessment Report to be peer reviewed by at least three appropriately qualified persons.
- c) Provide the Minister with the independent assessment of management effectiveness, the peer review comments and a report identifying how the findings of the independent assessment and peer review have been considered in the Strategic Assessment Report and the Program Report.

#### 7.4 Information and assessments

- a) Use the best available information to undertake the Strategic Assessment, including scientific data, expert opinion, and Traditional Owner and community knowledge.
- b) Document the methods used to undertake the Strategic Assessment.

- c) For information used in the Strategic Assessment, indicate where possible:
  - i. the source of the information
  - ii. how recent the information is
  - iii. the reliability and limitations of the assessment.

## 8. Endorsement Criteria

Describe how the Strategic Assessment Report meets the Endorsement Criteria set out below:

#### Endorsement Criteria

When deciding whether to endorse the Program, the Minister must be satisfied that the *Strategic Assessment Report* adequately addresses the impacts to which this Agreement relates, and that any recommendations by the Minister to modify the Program have been responded to appropriately.

In determining whether or not to endorse the Program, the Minister will have regard to the extent to which the Program meets the objects of the *Environment Protection and Biodiversity Conservation Act 1999*. In particular, that it:

- a) Protects the environment, especially those aspects of the environment that are matters of national environmental significance.
- b) Promotes ecologically sustainable development through the conservation and ecologically sustainable use of natural resources.
- c) Promotes the conservation of biodiversity.
- d) Provides for the protection and conservation of heritage.
- e) Promotes a cooperative approach to the protection and management of the environment.
- f) Assists in the co-operative implementation of Australia's international environmental responsibilities.
- g) Recognises the role of Indigenous people in the conservation and ecologically sustainable use of Australia's biodiversity.
- h) Promotes the use of Indigenous peoples' knowledge of biodiversity with the involvement of, and in co-operation with, the owners of the knowledge.

Without limiting the matters the Minister may consider when making the decision to endorse the Program, the Minister will consider the manner in which the Program:

- i) Identifies direct, indirect and cumulative impacts on matters of national environmental significance.
- j) Avoids impacts on matters of national environmental significance.
- k) Mitigates the impacts on matters of national environmental significance.
- l) Offsets the impacts on matters of national environmental significance.
- m) Contributes to the enhancement of the existing environment and management of existing threats.
- n) Demonstrates adaption to reasonable climate change scenarios.

Commitments in the Program must be adequately resourced throughout its life. The Program must demonstrate an effective system of adaptive management that addresses uncertainty and contingency management as well as procedures for monitoring, independent auditing and public reporting on implementation.

The Program must address all of the above matters for it to be considered for endorsement by the Minister in accordance with the *Environment Protection and Biodiversity Conservation Act 1999*.

END



## Appendix 3

---

### **Statement of Outstanding Universal Value Great Barrier Reef – Property ID 154**







**STATEMENT OF OUTSTANDING UNIVERSAL VALUE**  
**GREAT BARRIER REEF - PROPERTY ID 154**

**Brief synthesis:** As the world's most extensive coral reef ecosystem, the Great Barrier Reef is a globally outstanding and significant entity. Practically the entire ecosystem was inscribed as World Heritage in 1981, covering an area of 348,000 square kilometres and extending across a contiguous latitudinal range of 14° (10°S to 24°S). The Great Barrier Reef (hereafter referred to as GBR) includes extensive cross-shelf diversity, stretching from the low water mark along the mainland coast up to 250 kilometres offshore. This wide depth range includes vast shallow inshore areas, mid-shelf and outer reefs, and beyond the continental shelf to oceanic waters over 2,000 metres deep.

Within the GBR there are some 2,500 individual reefs of varying sizes and shapes, and over 900 islands, ranging from small sandy cays and larger vegetated cays, to large rugged continental islands rising, in one instance, over 1,100 metres above sea level. Collectively these landscapes and seascapes provide some of the most spectacular maritime scenery in the world.

The latitudinal and cross-shelf diversity, combined with diversity through the depths of the water column, encompasses a globally unique array of ecological communities, habitats and species. This diversity of species and habitats, and their interconnectivity, make the GBR one of the richest and most complex natural ecosystems on earth. There are over 1,500 species of fish, about 400 species of coral, 4,000 species of mollusk, and some 240 species of birds, plus a great diversity of sponges, anemones, marine worms, crustaceans, and other species. No other World Heritage property contains such biodiversity. This diversity, especially the endemic species, means the GBR is of enormous scientific and intrinsic importance, and it also contains a significant number of threatened species. At time of inscription, the IUCN evaluation stated "... if only one coral reef site in the world were to be chosen for the World Heritage List, the Great Barrier Reef is the site to be chosen".

**Natural beauty and phenomena (Criterion iii now Criterion VII):** The GBR is of superlative natural beauty above and below the water, and provides some of the most spectacular scenery on earth. It is one of a few living structures visible from space, appearing as a complex string of reefal structures along Australia's northeast coast.

From the air, the vast mosaic patterns of reefs, islands and coral cays produce an unparalleled aerial panorama of seascapes comprising diverse shapes and sizes. The Whitsunday Islands provide a magnificent vista of green vegetated islands and spectacular sandy beaches spread over azure waters. This contrasts with the vast mangrove forests in Hinchinbrook Channel, and the rugged vegetated mountains and lush rainforest gullies that are periodically cloud-covered on Hinchinbrook Island.

On many of the cays there are spectacular and globally important breeding colonies of seabirds and marine turtles, and Raine Island is the world's largest green turtle breeding area. On some continental islands, large aggregations of over-wintering butterflies periodically occur.

Beneath the ocean surface, there is an abundance and diversity of shapes, sizes and colours; for example, spectacular coral assemblages of hard and soft corals, and thousands of species of reef fish provide a myriad of brilliant colours, shapes and sizes. The internationally renowned Cod Hole near Lizard Island is one of many significant tourist attractions. Other superlative natural phenomena include the annual coral spawning, migrating whales, nesting turtles, and significant spawning aggregations of many fish species.

**Major stages of the Earth's evolutionary history (Criterion i now Criterion VIII):** The GBR, extending 2,000 kilometres along Queensland's coast, is a globally outstanding example of an ecosystem that has evolved over millennia. The area has been exposed and flooded by at least four glacial and interglacial cycles, and over the past 15,000 years reefs have grown on the continental shelf.

During glacial periods, sea levels dropped, exposing the reefs as flat-topped hills of eroded limestone. Large rivers meandered between these hills and the coastline extended further east. During interglacial periods, rising sea levels caused the formation of continental islands, coral cays and new phases of coral growth. This environmental history can be seen in cores of old massive corals.

Today the GBR forms the world's largest coral reef ecosystem, ranging from inshore fringing reefs to mid-shelf reefs, and exposed outer reefs, including examples of all stages of reef development. The processes of geological and geomorphological evolution are well represented, linking continental islands, coral cays and reefs. The varied seascapes and landscapes that occur today have been moulded by changing climates and sea levels, and the erosive power of wind and water, over long time periods.

One-third of the GBR lies beyond the seaward edge of the shallower reefs; this area comprises continental slope and deep oceanic waters and abyssal plains.

**Ecological and biological processes (Criterion ii now Criterion IX):** The globally significant diversity of reef and island morphologies reflects ongoing geomorphic, oceanographic and environmental processes. The complex cross-shelf, longshore and vertical connectivity is influenced by dynamic oceanic currents and ongoing ecological processes such as upwellings, larval dispersal and migration.

Ongoing erosion and accretion of coral reefs, sand banks and coral cays combine with similar processes along the coast and around continental islands. Extensive beds of *Halimeda* algae represent active calcification and accretion over thousands of years.

Biologically the unique diversity of the GBR reflects the maturity of an ecosystem that has evolved over millennia; evidence exists for the evolution of hard corals and other fauna. Globally significant marine faunal groups include over 4,000 species of molluscs, over 1,500 species of fish, plus a great diversity of sponges, anemones, marine worms, crustaceans, and many others. The establishment of vegetation on the cays and continental islands exemplifies the important role of birds, such as the Pied Imperial Pigeon, in processes such as seed dispersal and plant colonisation.

Human interaction with the natural environment is illustrated by strong ongoing links between Aboriginal and Torres Strait Islanders and their sea-country, and includes numerous shell deposits (middens) and fish traps, plus the application of story places and marine totems.

**Habitats for conservation of biodiversity (Criterion iv now Criterion X):** The enormous size and diversity of the GBR means it is one of the richest and most complex natural ecosystems on earth, and one of the most significant for biodiversity conservation. The amazing diversity supports tens of thousands of marine and terrestrial species, many of which are of global conservation significance.

As the world's most complex expanse of coral reefs, the reefs contain some 400 species of corals in 60 genera. There are also large ecologically important inter-reefal areas. The shallower marine areas support half the world's diversity of mangroves and many seagrass species. The waters also provide major feeding grounds for one of the world's largest populations of the threatened dugong. At least 30 species of whales and dolphins occur here, and it is a significant area for humpback whale calving.

Six of the world's seven species of marine turtle occur in the GBR. As well as the world's largest green turtle breeding site at Raine Island, the GBR also includes many regionally important marine turtle rookeries.

Some 242 species of birds have been recorded in the GBR. Twenty-two seabird species breed on cays and some continental islands, and some of these breeding sites are globally significant; other seabird species also utilize the area. The continental islands support thousands of plant species, while the coral cays also have their own distinct flora and fauna.

**Integrity:** The ecological integrity of the GBR is enhanced by the unparalleled size and current good state of conservation across the property. At the time of inscription it was felt that to include virtually the entire Great Barrier Reef within the property was the only way to ensure the integrity of the coral reef ecosystems in all their diversity.

A number of natural pressures occur, including cyclones, crown-of-thorns starfish outbreaks, and sudden large influxes of freshwater from extreme weather events. As well there is a range of human uses such as tourism, shipping and coastal developments including ports. There are also some disturbances facing the GBR that are legacies of past actions prior to the inscription of the property on the World Heritage list.

At the scale of the GBR ecosystem, most habitats or species groups have the capacity to recover from disturbance or withstand ongoing pressures. The property is largely intact and includes the fullest possible representation of marine ecological, physical and chemical processes from the coast to the deep abyssal waters enabling the key interdependent elements to exist in their natural relationships.

Some of the key ecological, physical and chemical processes that are essential for the long-term conservation of the marine and island ecosystems and their associated biodiversity occur outside the boundaries of the property and thus effective conservation programs are essential across the adjoining catchments, marine and coastal zones.

**Protection and management requirements:** The GBR covers approximately 348,000 square kilometres. Most of the property lies within the GBR Marine Park: at 344,400 square kilometres, this Federal Marine Park comprises approximately 99% of the property. The GBR Marine Park's legal jurisdiction ends at low water mark along the mainland (with the exception of port areas) and around islands (with the exception of 70 Commonwealth managed islands which are part of the Marine Park). In addition the GBR also includes over 900 islands within the jurisdiction of Queensland, about half of which are declared as 'national parks', and the internal waters of Queensland that occur within the World Heritage boundary (including a number of long-established port areas).

The World Heritage property is and has always been managed as a multiple-use area. Uses include a range of commercial and recreational activities. The management of such a large and iconic world heritage property is made more complex due to the overlapping State and Federal jurisdictions. The Great Barrier Reef Marine Park Authority, an independent Australian Government agency, is responsible for protection and management of the GBR Marine Park. The Great Barrier Reef Marine Park Act 1975 was amended in 2007 and 2008, and now provides for "the long term protection and conservation of the Great Barrier Reef Region" with specific mention of meeting "... Australia's responsibilities under the World Heritage Convention."

Queensland is responsible for management of the Great Barrier Reef Coast Marine Park, established under the Marine Parks Act 2004 (Qld). This is contiguous with the GBR Marine Park and covers the area between low and high water marks and many of the waters within the jurisdictional limits of Queensland. Queensland is also responsible for management of most of the islands.

The overlapping jurisdictional arrangements mean that the importance of complementary legislation and complementary management of islands and the surrounding waters is well recognised by both governments. Strong cooperative partnerships and formal agreements exist between the Australian Government and the Queensland Government. In addition, strong relationships have been built between governments and commercial and recreational industries, research institutions and universities. Collectively this provides a comprehensive management influence over a much wider context than just the marine areas and islands.

Development and land use activities in coastal and water catchments adjacent to the property also have a fundamental and critical influence on the values within the property. The Queensland Government is responsible for natural resource management and land use planning for the islands, coast and hinterland adjacent to the GBR. Other Queensland and Federal legislation also protects the property's Outstanding Universal Value addressing such matters as water quality, shipping management, sea dumping, fisheries management and environmental protection.

The Federal Environment Protection and Biodiversity Conservation Act 1999 (EPBC Act) provides an overarching mechanism for protecting the World Heritage values from inappropriate development, including actions taken inside or outside which could impact on its heritage values. This requires any development proposals to undergo rigorous environmental impact assessment processes, often including public consultation, after which the Federal Minister may decide, to approve, reject or approve under conditions designed to mitigate any significant impacts. A recent amendment to the EPBC Act makes the GBR Marine Park an additional 'trigger' for a matter of National Environmental Significance which provides additional protection for the values within the GBR.

The GBR Marine Park and the adjoining GBR Coast Marine Park are zoned to allow for a wide range of reasonable uses while ensuring overall protection, with conservation being the primary aim. The zoning spectrum provides for increasing levels of protection for the 'core conservation areas' which comprise the 115,000 square kilometres of 'no-take' and 'no-entry' zones within the GBR.

While the Zoning Plan is the 'cornerstone' of management and provides a spatial basis for determining where many activities can occur, zoning is only one of many spatial management tools

and policies applied to collectively protect the GBR. Some activities are better managed using other spatial and temporal management tools like Plans of Management, Special Management Areas, Agreements with Traditional Owners and permits (often tied to specific zones or smaller areas within zones, but providing a detailed level of management not possible by zoning alone). These statutory instruments also protect the Outstanding Universal Value of the property.

Many Aboriginal and Torres Strait Island peoples undertake traditional use of marine resource activities to provide traditional food, practice their living maritime culture, and to educate younger generations about traditional and cultural rules and protocols. In the GBR these activities are managed under both Federal and Queensland legislation and policies including Traditional Use of Marine Resource Agreements (TUMRAs) and Indigenous Land Use Agreements (ILUAs). These currently cover some 30 per cent of the GBR inshore area, and support Traditional Owners to maintain cultural connections with their sea country.

Similarly non-statutory tools like site management and Industry Codes of Practice contribute to the protection of World Heritage values. Some spatial management tools are not permanently in place nor appear as part of the zoning, yet achieve effective protection for elements of biodiversity (e.g. the temporal closures that are legislated across the GBR prohibit all reef fishing during specific moon phases when reef fish are spawning).

Other key initiatives providing increased protection for the GBR include the comprehensive Great Barrier Reef Outlook Report, (and its resulting 5-yearly reporting process); the Reef Water Quality Protection Plan; the GBR Climate Change Action Plan; and the Reef Guardians Stewardship Programs which involve building relationships and working closely with those who use and rely on the GBR or its catchment for their recreation or their business.

The 2009 Outlook Report identified the long-term challenges facing the GBR; these are dominated by climate change over the next few decades. The extent and persistence of damage to the GBR ecosystem will depend to a large degree on the amount of change in the world's climate and on the resilience of the GBR ecosystem to such change. This report also identified continued declining water quality from land-based sources, loss of coastal habitats from coastal development, and some impacts from fishing, illegal fishing and poaching as the other priority issues requiring management attention for the long-term protection of the GBR.

Emerging issues since the 2009 Outlook Report include proposed port expansions, increases in shipping activity, coastal development and intensification and changes in land use within the GBR catchment; population growth; the impacts from marine debris; illegal activities; and extreme weather events including floods and cyclones.

Further building the resilience of the GBR by improving water quality, reducing the loss of coastal habitats and increasing knowledge about fishing and its effects and encouraging modified practices, will give the GBR its best chance of adapting to and recovering from the threats ahead, including the impacts of a changing climate.

## Appendix 4

---

# Assessment of Management Effectiveness Terms of Reference







## TERMS OF REFERENCE

### ASSESSMENT OF MANAGEMENT EFFECTIVENESS BY INDEPENDENT EXPERTS FOR THE GREAT BARRIER REEF WORLD HERITAGE AREA STRATEGIC ASSESSMENT

#### 1. BACKGROUND

The Australian and Queensland governments are working together to undertake a comprehensive strategic assessment of the Great Barrier Reef World Heritage Area and adjacent coastal zone. The comprehensive strategic assessment will help identify, plan for and manage existing and emerging risks so that the unique values of the Great Barrier Reef are protected and managed.

There are two components to the comprehensive strategic assessment - a marine component and a coastal component. The Great Barrier Reef Marine Park Authority (the Authority) will lead the marine component involving a strategic assessment of the Great Barrier Reef Region (the Strategic Assessment). The Queensland Government will lead the coastal component involving a strategic assessment of the Great Barrier Reef Coastal Zone.

#### 2. PURPOSE

The Strategic Assessment Terms of Reference (SA TOR) sets out the requirements for the Authority in preparing the Strategic Assessment Report and Program Report for the marine component. The GBRMPA Strategic Assessment's key objectives are:

- i. **Identifying and describing values to be protected.** The first step in the assessment is to identify values that underpin matters of national environmental significance in the Great Barrier Reef Region. Values will be described and information gaps identified.
- ii. **Identifying and analysing threats and pressures to the values.** Threats and pressures impacting on values identified above will be described. It is important to understand how different threats and pressures impact on values in order to assess the effectiveness of management. Pressures may include climate change, extreme weather, and coastal development.

iii. **Assessing management effectiveness.** The assessment will examine how the Authority's management arrangements protect values. It will look at how impacts are avoided, mitigated, offset and/or adaptively managed across the life of the activity. It will look monitoring, compliance and stewardship programs in addition to environmental impact assessment and planning tools. It is proposed that particular cases be used as examples to examine management effectiveness, such as looking closely at management of a particular region or species.

- iv. **Recommending changes to management arrangements.** Recommendations for changes to management arrangements will be made based on the above assessment.

The Strategic Assessment requires the management effectiveness of the current Program to be described and assessed (sections 4.1.1 - 4.1.2 of the SA TOR). The purpose of the present Terms of Reference (TOR) is to set out the requirements for the independent review of management effectiveness of the Authority's current Program<sup>1</sup> against the SA TOR to help inform what improvements to management arrangements are needed for the Program Report.

<sup>1</sup> The word 'Program' is used throughout the Terms of Reference. In simple terms it means those management arrangements ('policy, plan or program') under the jurisdiction of the Authority. Management arrangements include, but are not limited to, those described in section 4.1.2 of the GBRVHA Terms of Reference.

## 3. SCOPE

The strategic assessment area relevant to this TOR is the Great Barrier Reef Region which includes the Great Barrier Reef Marine Park (including 70 Commonwealth Islands) and the 13 Ports in 12 exclusion areas. It does not include Internal Waters of Queensland or the 980 Queensland Islands within the Great Barrier Reef World Heritage Area. However, the management Program of the Great Barrier Reef Marine Park Authority applies to the Great Barrier Reef Marine Park and to matters that relate to the Marine Park, namely:

- the use or management of an area which would or might affect the Marine Park; and
- the use of a place outside the Marine Park for a purpose relating to the Marine Park.

Therefore, the management arrangements of the Authority directly address activities that occur in the Marine Park itself and seek to influence activities outside the Marine Park that are affecting or may affect the Marine Park. In addition, under Section 66(2)(e) of the Regulations, the Authority may regulate or prohibit acts, whether in the Marine Park or elsewhere, that may pollute water in a manner harmful to animals and plants in the Marine Park.

Therefore the scope of this independent assessment should primarily concentrate on management activities within the Authority's jurisdiction within the Great Barrier Reef Region, joint management arrangements with Queensland (e.g. joint Marine Park permitting and joint compliance and enforcement through the Field Management Program) where they exist, and any activities that occur outside the Great Barrier Reef Region that are affecting or may affect the Marine Park.

A key component of the Strategic Assessment is stakeholder engagement. An Engagement Strategy has been developed (Attachment B) and should be taken into consideration when determining approaches for stakeholder engagement as part of the independent review of management effectiveness.

## 4. CONTEXT

The TOR for the review must be considered in the context of:

- i. background information provided above
- ii. the approved Great Barrier Reef Region Strategic Assessment Terms of Reference (Attachment A)
- iii. the methodology used to assess management effectiveness as part of the Outlook Report 2009 (refer Hockings and Gilligan (2009)<sup>2</sup>)
- iv. the Strategic Assessment – Great Barrier Reef Stakeholder Engagement Plan (Attachment B)
- v. timeframes for delivery of the Strategic Assessment and Program report.

## 5. PROJECT DETAILS

### 5.1 Project Title

"Great Barrier Reef Region Strategic Assessment - Independent Review of Management Effectiveness"

### 5.2 Duration of Project

Start Date: "Insert the Project Start Date"

End Date: 31 May 2013

<sup>2</sup> Hockings, M. & Gilligan, B. 2009 Assessment of management effectiveness for the 2009 Great Barrier Reef Outlook Report: report to the Great Barrier Reef Marine Park Authority. Great Barrier Reef Marine Park Authority, Townsville, Australia.

### 5.3 Proposed Audience


The primary audience for the Management Effectiveness Report is the Government (both Commonwealth and State), the Australian community and the international community (e.g. The World Heritage Committee).

## 6. PROJECT OUTCOMES

- 6.1 Provide a comprehensive independent review of the effectiveness of the Authority's current Program to:
- Identify the relevant matters of national environmental significance, including the Outstanding Universal Value of the Great Barrier Reef World Heritage Area, and determine their current condition and trend, including spatial and non-spatial approaches.
  - Identify and analyse direct, indirect, consequential and cumulative impacts on relevant matters of national environmental significance, including the methods used to determine these types of impacts.
  - Consider environmental, social, cultural and economic issues in its decision-making processes.
  - Avoid, mitigate, offset and adaptively manage impacts from activities.
  - Address uncertainty and risk.
  - Provide certainty regarding where uses may occur, the type of activities allowed, conditions under which activities may proceed and circumstances where impacts are likely to be unacceptable.
  - Halt and reverse any declines and enhance the condition of the relevant matters of national environmental significance, including mechanisms to deliver a 'net benefit' to the condition of the relevant matters of national environmental significance, including the Outstanding Universal Value of the Great Barrier Reef World Heritage Area.
  - Adapt to reasonable climate change scenarios.
  - Integrate with related local, Queensland and Australian government programs to protect and manage the relevant matters of national environmental significance, including the Outstanding Universal Value of the Great Barrier Reef World Heritage Area.
  - Meet Australia's international responsibilities in relation to the environment and protection of world heritage.
  - Monitor, evaluate and report on the:
    - Condition and trends of the relevant matters of national environmental significance, including the Outstanding Universal Value of the Great Barrier Reef World Heritage Area
    - Impacts of activities, including the setting of targets to benchmark management.
  - Assess in finer detail the effectiveness of the Program to protect and manage the relevant matters of national environmental significance, including the Outstanding Universal Value of the Great Barrier Reef World Heritage Area, and to guide improvements to the Program, in up to 10 Demonstration Cases.
- 6.2 In assessing the effectiveness of the Authority's Program, consideration must be given to:
- Statutory instruments, including regulations, zoning plans, plans of management and permits.
  - Non-statutory mechanisms including policies, position statements and guidelines.

- Partnership and collaborative arrangements with Queensland and other Australian government agencies.
- Partnerships with Traditional Owners in the management of marine resources.
- Partnership and stewardship programs, including education programs and engagement, with local governments, communities, Indigenous persons, business and industry.
- Research and monitoring programs.
- Compliance and enforcement programs.
- Resourcing of the Program.

- 6.1 The independent review should follow the established framework for evaluating management effectiveness as specified in the Great Barrier Reef Outlook Report 2009. Each of the items listed in 6.1. are Desired Outcomes. The framework should consider how effective the Authority's Program is at achieving these desired outcomes in accordance with the framework below.



- 6.2 The independent review of management effectiveness must include engagement with relevant stakeholders to determine their views on management effectiveness of the current Program (listed in item 2.3 (b) of this TOR). Delivery mechanisms are to be determined by consultants. At a minimum, an online survey of relevant stakeholder's views on the effectiveness of the current management arrangements must be undertaken. The GBRMPA will provide a list of stakeholders and their contact details to the independent reviewer.

- 6.3 The review must provide findings on the following (as a minimum):

- activities that are considered effectively managed
- gaps and deficiencies in management arrangements

iii. possible additional, improved or alternative arrangements that may be effective in achieving the objectives of the Strategic Assessment TOR.

**7. PROVISION OF ASSESSMENT SERVICES**

The service to be provided comprises a written report delivered as a consensus view of the Independent Experts, addressing the Project Outcomes and in accordance with the timeframes outlined in the Project Deliverables (see 8 below),

**8. PROJECT DELIVERABLES (generally, to be finalized in contracts)**

Appointment of Independent Reviewers	September 2012
Assessment methodology and stakeholder engagement strategy finalised	September 2012
Stakeholder Engagement Completed	October 2012
Draft Assessment Report delivered	December 2012
Final Assessment Report delivered	February 2013





## Appendix 5

### **Traditional Owner and Stakeholder Engagement – Report on Workshops and Surveys**





---

# **Great Barrier Reef Region Strategic Assessment Traditional Owner and Stakeholder Engagement – Report on Workshops and Surveys**

---

## Table of contents

Background.....	5
Approach to Stakeholder Engagement.....	5
Objectives .....	5
Staged engagement .....	5
Using existing advisory groups .....	6
Workshops and surveys .....	6
Stakeholder presentations and meetings .....	7
Stage one: Values and impacts .....	7
Stage one workshops .....	7
Stage one workshop analysis .....	7
Traditional Owner workshops .....	9
Follow-up surveys .....	10
Stage one outcomes .....	11
Stage two: Management effectiveness .....	11
Testing the methodology .....	12
Survey results .....	12
Demonstration case studies .....	13
Stage two workshops.....	13
Condition and trends.....	14
Management effectiveness.....	14
Future management actions .....	14
Stage two outcomes .....	15
Summary.....	15
Appendix 1: Background documents.....	16

## List of figures

Figure 1 Stakeholder and community engagement process .....	6
Figure 2 Importance of Great Barrier Reef elements according to survey respondents .....	10
Figure 3 Background documents used in the preparation of the Stakeholder Engagement Report.....	16

## List of tables

Table 1 Important natural elements and values consistently mentioned in workshops .....	9
Table 2 Survey respondents rank top three threats to the Great Barrier Reef.....	11



## Background

The Australian and Queensland governments are working together to undertake a comprehensive strategic assessment of the Great Barrier Reef World Heritage Area and the adjacent coastal zone. The goal is to create an agreed, long-term plan for sustainable development within the Great Barrier Reef Region that provides greater certainty for industry and decision making in the area, while ensuring the values of the World Heritage Area are protected into the future. The strategic assessment's key objectives are:

- **Identifying and describing values to be protected:** Values that underpin matters of national environmental significance in the Great Barrier Reef Region are identified and described, and information gaps are identified.
- **Identifying and analysing impacts on values:** It is important to understand how different activities impact on values in order to assess the effectiveness of management.
- **Assessing management effectiveness:** This relates to how management arrangements protect values by looking at how impacts are avoided, mitigated, offset and/or adaptively managed across the life of an activity. It assesses monitoring, compliance and stewardship programs as well as environmental impact assessment and planning tools.
- **Recommending changes to management arrangements:** Recommendations for changes to management arrangements will be made based on the above assessment.

The terms of reference for the strategic assessment state that in the preparation of the assessment, the Great Barrier Reef Marine Park Authority (the Authority) will:

*Engage stakeholders and the community throughout the strategic assessment process using the Authority's network of advisory committees together with regional and issue-based meetings and use the best available information to undertake the strategic assessment, including scientific data, expert opinion, and Traditional Owner and community knowledge.*

This report provides an overview of the stakeholder and community engagement undertaken and the general outcomes, including how feedback has been incorporated into the strategic assessment. It does not include the public submissions for the draft terms of reference in May 2012, which is covered by a separate report. The outcomes of the public submissions for the Draft Strategic Assessment Report will also be covered by a separate report.

## Approach to Stakeholder Engagement

Following the release of the strategic assessment's terms of reference in August 2012, a stakeholder engagement plan was developed to provide a structured approach for stakeholder and community input during development of the assessment. The approach is anchored in the following principle:

*To work directly with the public throughout the process to ensure that public issues and concerns are consistently understood, considered and included as appropriate.*

### Objectives

The stakeholder engagement plan's objectives highlight the value of incorporating stakeholder knowledge and perspectives from the beginning of the strategic assessment and throughout the process:


- to improve decision making by gathering a range of information, views and experiences from stakeholders as part of the strategic assessment process
- to provide a transparent framework for the development of a common understanding of the broad range of issues and to foster informed and engaged stakeholders and communities as part of the process
- to seek feedback from stakeholders on the potential impact of different approaches, acknowledging that there may not be consensus
- to provide a mechanism to better understand implementation issues that may need to be considered.

### Staged engagement

The stakeholder engagement plan used the key requirements in the terms of reference to construct a two-stage process for seeking stakeholder input:

- Stage one to understand stakeholders' values of the Great Barrier Reef Region and the pressures and potential impacts on those values
- Stage two to investigate stakeholders' views of current management effectiveness and their aspirations for future management actions.

The components of the staged process (Figure 1) are outlined in further detail below. A list of background documents relating to the staged engagement process can be found at Appendix 1.



**Figure 1 Stakeholder and community engagement process**

## Using existing advisory groups

The stakeholder engagement plan established a framework for targeted stakeholder engagement to ensure community views were captured, while meeting the timelines of the strategic assessment process. A key element was the use of existing expert and community advisory groups.

### Reef Advisory Committees:

The Authority's Reef Advisory Committees are competency-based committees comprising a cross-section of stakeholder interests with relevant expertise and experience. The four Reef Advisory Committees are: Indigenous, Catchment and Coastal, Tourism and Recreation and Ecosystem.

The Reef Advisory Committees provided advice on the engagement process for the strategic assessment, have been kept informed on the progress of the strategic assessment, and have provided ongoing advice in areas relevant to their expertise.

### Local Marine Advisory Committees:

Local Marine Advisory Committees are voluntary, community-based committees established by the Authority to provide advice on management issues. There are 12 committees which cover the Great Barrier Reef Region: Cape York; Douglas; Cairns; Cassowary Coast; Hinchinbrook; Townsville; Bowen–Burdekin; Whitsundays; Mackay; Capricorn Coast; Gladstone and Burnett.

The committees played an integral role in all stages of the stakeholder engagement process. The Local Marine Advisory Committee Chairs' meeting in October 2012, which was attended by two representatives from each region, provided feedback on proposed approaches as well as being a source of critical input for the strategic assessment.

## Workshops and surveys

Opportunities were provided for further stakeholder input through a series of purpose-designed workshops and follow-up surveys. The aim of the workshops and surveys was to ensure a spectrum of stakeholder views was gathered which were representative of the wider community's views and opinions.

Individuals were invited to participate in the process through the Authority's stakeholder and community networks. This targeted consultation achieved representation from a diverse range of interests including Traditional Owners; local government; ports, shipping and related development sectors; mining and resource sectors; research

organisations; tourism industry; commercial fishers; recreational users including fishers; natural resource managers; farmers; conservation groups and the broader community.

## Stakeholder presentations and meetings

Authority staff undertook a significant number of presentations and discussions with stakeholders including sectoral interests and other government agencies.

## Stage one: Values and impacts

The terms of reference (section 2.1) requires the strategic assessment to identify and describe the values that underpin matters of national environmental significance. These wide-ranging ecological, social, cultural and economic values may differ between industries, communities and individuals. Capturing these differences and mapping and describing values was an essential step early in the strategic assessment. Discussions with stakeholders were critical in order to understand the many different ways people interact with and benefit from the Great Barrier Reef. These 'community benefits' had not previously been comprehensively studied and recorded.

The terms of reference (section 3.1) also requires the assessment to describe and analyse the actual and potential impacts on those values. Stakeholder involvement is fundamental to understanding the potential impacts on their social, cultural and economic values.

### Stage one workshops

A series of workshops were held to gather stakeholders' views about values and impacts. A total of 290 people attended the workshops between August and October 2012:

- Three purpose-designed stakeholder workshops were held in Townsville, Cairns and Rockhampton in August 2012.
- The same process was used in 12 mini-workshops at the regular Local Marine Advisory Committee meetings in September 2012.
- Input from Traditional Owners was gathered at two specific workshops in Cairns and Rockhampton in September 2012.

The workshops consisted of overview presentations about the strategic assessment and opportunities for stakeholders to provide input through roundtable discussions. Participants from diverse backgrounds were seated at tables, together with Authority facilitators promoting the discussions, which led to a broad range of thinking and a shared awareness among participants of the complexity of the issues and wide range of community views.

Participants were asked three prescribed questions to obtain responses that would inform the strategic assessment:

- What are the natural elements of the Great Barrier Reef which are important to you, and why are they important?
- Based on the natural elements you identified, what are the factors that might impact on these?
- What are some of the key elements you would like to see in the Great Barrier Reef in 25 years?

This last aspirational question related to a requirement in the terms of reference (section 1.1) for an overview of management for the next 25 years. Participants were provided with a copy of the vision statement from the *25-year Strategic Plan for the Great Barrier Reef World Heritage Area* developed in 1994. They were asked if the statement was still relevant today, and were invited to reword it or write their own vision statement.

When thinking about values, participants were asked to consider the categories of social, economic, Indigenous cultural, heritage and aesthetic values. Participants were also asked to assess the condition and trends of these values, and to identify impacts on them. Their views were captured on individual sheets and collective butcher's paper. The information was later entered into electronic databases which grouped the different responses into broad categories.

### Stage one workshop analysis

A report on each of the stakeholder workshops described and compared responses to the three questions. The reports highlighted some notable differences in the findings from each workshop. For example, people from far north Queensland (that is, around Hinchinbrook and further north) saw connectivity, integrity and health of the whole Great Barrier Reef ecosystem as being essential to maintain into the future. While several individuals participating in the other workshops also saw these as very important, people living further south tended to focus more on impacts at specific sites, especially where human activity was obviously affecting the natural, social, heritage and cultural values of the site. Consistent values to emerge from all of the five large workshops are presented in Table 1.

Factors impacting on values that were mentioned consistently across all workshops were:

- coastal development including urban sprawl and increased industrialisation of the coast

- port development and associated increase in shipping
- mining
- climate change
- extreme weather events
- overfishing
- declining water quality.

Key elements that workshop participants would like to see in the Great Barrier Reef in 25 years fell into nine broad categories:

- healthy Reef
- protection
- sustainable use
- effective Reef management
- regulated coastal development
- stewardship
- communication and education
- research and monitoring
- consultation and engagement.



**Table 1 Important natural elements and values consistently mentioned in workshops**

Reef attribute	Reef elements	Values mentioned consistently in workshops
Specific sites/ habitats	Islands, beaches and coastlines; estuaries, deep water, bays, inlets and coral reefs	<b>Social:</b> supports recreation, for example walking, camping, snorkelling, diving, fishing, wildlife watching, relaxing, spending time with family and friends, education, health, lifestyle, stewardship <b>Aesthetic:</b> reefs and islands can be seen from space, place of natural wonder, spectacular pristine beauty, awesome, spiritual, majestic and calming, looked upon with pride, unique habitats <b>Natural:</b> obligation to have it there for our children <b>Cultural:</b> significance for Traditional Owners, for example locations of fish traps, sites for traditional use of marine resources and ceremonies; other cultural heritage values, for example locations of ship wrecks, lighthouses, sites of Cook's landings <b>Economic:</b> supports commercial fishing, shipping (deep water), tourism and recreation
	Wetlands	<b>Natural:</b> important breeding and feeding grounds for a variety of fish species, migratory birds and other animals <b>Cultural:</b> Indigenous medicine and food sources <b>Social:</b> supports recreation, for example wildlife watching, education <b>Economic:</b> supports commercial fishing and tourism
	Seagrass meadows	<b>Natural:</b> important breeding and feeding grounds for a variety of fish species, turtles, dugong and other animals <b>Social:</b> supports recreation, for example snorkelling, stewardship, education <b>Cultural:</b> supports dugong and turtle populations <b>Economic:</b> supports commercial fishing and tourism
Biodiversity	Fishes, coral, dolphins, crocodiles, birds, whales, dugong, sharks, rays, sea snakes, turtles	<b>Economic:</b> supports commercial fishing and tourism <b>Social:</b> supports recreation, for example snorkelling, diving, wildlife watching, education <b>Cultural:</b> supports traditional hunting of some species <b>Natural:</b> having such a large protected area for species to move through without risk <b>Aesthetic:</b> 'wow' factor
Processes	Connectivity	<b>Natural:</b> linking biophysical processes, supporting biodiversity <b>Cultural:</b> connection of people to land and sea through stories, also in terms of the way people moved from inland areas to the sea seasonally in search of food. Connectivity occurs from west to east (i.e. catchment to sea) and from north to south through travel/trade routes.
	Integrity	<b>Natural:</b> linking biophysical processes, supporting biodiversity <b>Cultural:</b> maintaining sites for traditional use of marine resources, ceremonies and stories
	Spawning (Coral/fish)	<b>Economic:</b> supports commercial fishing and tourism <b>Natural:</b> contributes to biodiversity
	Water quality	<b>Social:</b> supports human health; recreation, for example swimming, snorkelling, diving <b>Economic:</b> supports commercial fishing and tourism <b>Aesthetic:</b> water clarity; colour of water

## Traditional Owner workshops

Aboriginal and Torres Strait Islander people are the Traditional Owners of the Great Barrier Reef Region. Today, there are approximately 70 Traditional Owner clan groups whose customary estates include sea country within the Great Barrier Reef Marine Park. Traditional Owners have a unique knowledge and understanding of many coastal, island and marine environments adjacent to and within the Marine Park.

Traditional Owners took a holistic approach to the questions about values and impacts, considering all elements of the land, sea and people together. A major issue for consideration mentioned by the Cairns workshop participants was the connectivity between Traditional Owners and sea country, and between land and sea. A major issue for consideration mentioned by the Rockhampton workshop participants was the capacity for Traditional Owners of the area to share traditional ecological knowledge and pass it on to the next generation.

Key impacts on values identified by Traditional Owners include:


- lack of recognition and support for Traditional Owners
- dispossession from country/ loss of access to traditional grounds
- loss of language and cultural knowledge leading to loss of cultural practices and management
- (lack of ) capacity of Aboriginal rangers to enforce infringements
- unauthorised entry to sites of cultural significance

- coastal development
- vandalism — damage and graffiti
- marine debris and litter
- ships and boats cutting across storylines
- overfishing
- water quality
- extreme weather events
- lack of cultural knowledge-sharing between agencies (for example sacred sites).

## Follow-up surveys

Workshop participants were invited to answer a short online survey to gain a deeper understanding about what is important to them and why. The survey was sent to people who had attended one of the workshops. Of the 290 workshop participants, 135 people (47 per cent) responded to the survey. Results give an indication of the relative importance that this group of people place on Great Barrier Reef values, threats and their aspirations.

The respondents were asked to rate different elements where *1 = not important* and *5 = extremely important*. Ecosystem health, coral reefs and the integrity of the whole Great Barrier Reef received the highest scores (see Figure 2). The most commonly chosen reason (that is, value) for why respondents considered elements to be important was *natural beauty*, followed closely by *scientific*; then *understanding and appreciation*; *health*; and *recreation and enjoyment*. Values were equally weighted and respondents could choose as many values as they wished for each element. So although *natural beauty* and *scientific* were the most commonly selected values, it does not necessarily mean that respondents thought they were the most important value. Rather, it means that many of the respondents attributed natural beauty and scientific values to most elements on the list.



**Figure 2 Importance of Great Barrier Reef elements according to survey respondents**

Respondents were asked to rate a list of possible threats to the Great Barrier Reef. A total of 118 out of 135 people responded to this question (see Table 2). Water quality, climate change and extreme weather were seen as the most serious threats.

Fourteen respondents thought one key threat not listed was *urban and coastal development*. It was not included as a specific item because it is a multi-faceted factor which includes a range of human activities and associated infrastructure already listed (that is, shipping, ports, agriculture, aquaculture, fishing, recreation and tourism). Twelve respondents thought another threat related to management effectiveness including the adequacy of legislation, zoning and enforcement.

The most important 25-year aspiration for the Great Barrier Reef was *ecosystem health/biodiversity protection* as identified by 40 per cent of respondents. More than two-thirds of respondents rated it as their first or second most important aspiration. The other aspiration most commonly identified was *preservation for future generations*.

**Table 2 Survey respondents rank top three threats to the Great Barrier Reef**

Great Barrier Reef threats	Threat rating (number of responses) 1 = highest threat 3 = third highest threat			Response rate (out of a total of 118)
	1	2	3	
Water quality	27	33	16	76 (64%)
Climate change	40	17	11	68 (58%)
Extreme weather (for example cyclones)	14	9	14	37 (31%)
Crown-of-thorns starfish	8	14	16	38 (32%)
Ports	8	15	19	42 (36%)
Shipping	7	8	17	32 (27%)
Agriculture	4	10	10	24 (20%)
Illegal fishing or poaching	3	5	5	13 (11%)
Commercial fishing	1	5	6	12 (10%)
Recreational fishing	2	0	2	4 (3%)
Aquaculture	1	0	1	2 (2%)
Tourism	1	2	1	4 (3%)
Recreation (excluding fishing)	2	0	0	2 (2%)

## Stage one outcomes

The values and impacts information collected from the stakeholder workshops and surveys informed a number of different components of the strategic assessment process:

- The information provided valuable source data which was incorporated into the identification and analysis of values and impacts.
- The information provided valuable insights into community benefits — a requirement for strategic assessment which had not been previously comprehensively studied.
- The workshops and surveys developed a common understanding among stakeholders of the key issues, and ensured the Authority understood issues from the viewpoints of the wider community.
- The information established a base from which to move forward in assessing management effectiveness and potential management arrangements that may be required to address some of the major risks.
- The results were provided to external consultants undertaking independent studies as part of the strategic assessment, including management effectiveness and aesthetic values studies.
- The insights gained will also inform the development of the Outlook Report 2014 and a proposed update of the Authority's Heritage Strategy.

## Stage two: Management effectiveness

The strategic assessment is being undertaken to evaluate and improve (where necessary) the effectiveness of the Authority's current management arrangements. To ensure an impartial and objective approach, three independent reviewers were engaged to independently assess the Authority's management effectiveness. Gathering the views of stakeholders was an important part of the reviewers' brief as it recognised that stakeholders have valuable knowledge and practical experience of the management issues and the tools used by the Authority.

## Testing the methodology

The reviewers undertook an assessment based on the International Union for Conservation of Nature (IUCN) framework for assessing management effectiveness, also used for the *Great Barrier Reef Outlook Report 2009*. This methodology was presented at the annual meeting of Local Marine Advisory Committee Chairs in Townsville in October 2012. The presentation included an explanation of the approach and described the opportunity for stakeholders to submit their views through an online survey. Committee Chairs piloted the survey and contributed to its final design. Survey respondents could choose from a list of management issues and indicate the Authority's strengths and weaknesses in management of that issue, and then rate the effectiveness of each management tool on a four-point scale (effective, mostly effective, partially effective, not effective). They could also indicate if they had no opinion or believed the tool was not applicable to that issue.

The representatives formed discussion groups to test the survey instrument. Each table examined three of the following management topics, looking at strengths, weaknesses and rating management effectiveness:

- Biodiversity protection
- Climate change and extreme weather
- Coastal ecosystems
- Community benefits
- Defence activities
- Commercial fishing
- Recreational fishing
- Indigenous heritage
- Non-Indigenous heritage
- Marine-based tourism
- Ports
- Shipping
- Recreation (non-extractive)
- Research activities
- Water quality.

The tools assessed were:

- Act and Regulations
- Zoning Plan
- plans of management
- permits (including environmental impact assessment)
- Traditional Use of Marine Resources Agreements
- compliance (formal and informal)
- policy documents
- site infrastructure
- partnerships
- education and community awareness
- stewardship and best practice
- research and monitoring.

The contributions of Local Marine Advisory Committee representatives were recorded and their comments were used to improve the survey's content and presentation.

The external consultants distributed the survey electronically to members of Local Marine Advisory Committees, Reef Advisory Committees and interested participants of the regional workshops. Participants without access to the internet were contacted and invited to undertake the survey over the phone or in writing. Face-to-face interviews were undertaken with Traditional Owners to ensure their views were documented.

## Survey results

A total of 95 people made 172 assessments of the 15 management topics. For some management topics, the number of people responding was quite low, making statistical analysis difficult. For instance, the management topics of non-Indigenous heritage, defence activities and climate change and extreme weather each received less than 10 assessments. Respondents may not have had an opinion on the topic or may have been unsure of the relevant management tools and arrangements. However, some general patterns of effectiveness were evident for most of the tools.



Surveys results show a perceived lack of effectiveness for most of the management tools, and the independent reviewers recommended the Authority investigate these views with its existing advisory committees. Only the Zoning Plan was considered to be effective or mostly effective by more than half of the respondents. Compliance and policy documents were considered to be the least effective tools, with fewer than one in three respondents considering them to be effective. Problems with compliance and enforcement of Regulations were also commonly cited as problematic in the qualitative comments of respondents. The other commonly cited issue limiting the effectiveness of management tools was the Authority's limited jurisdictional responsibility. Respondents thought many issues fell partly or largely under the responsibility of state and local governments, with a perceived lack of commitment at those levels to the protection of the Great Barrier Reef.

Traditional Owners expressed wide-ranging views, including the desire for the development of stronger partnerships and greater involvement in decision making and management. The need for education and communication with the wider community was also raised.

## **Demonstration case studies**

Demonstration case studies were developed as part of the strategic assessment to provide more detailed snapshots of management effectiveness and provide recommendations on possible improvements.

The Local Marine Advisory Committee Chairs' meeting provided an opportunity for stakeholder input across a number of demonstration case studies, in particular the regionally-specific studies. Authority staff gave a presentation on the demonstration case studies at the meeting and explained the interactions of drivers, activities, values and management. For the purposes of the feedback sessions, representatives were grouped in regionally-specific tables together with Authority staff who facilitated and recorded dialogue. Committee representatives were requested to:

- provide advice on where management topics were of concern locally
- provide advice on how management arrangements could be improved locally
- exchange ideas on a range of proposed approaches to overcome challenges.

Tables were assigned four key management topics relevant to their region. Participants were asked to focus their discussion on these management topics, especially from a socio-economic perspective, and to address two key questions:

- What are the three or four key local pressures in your area and where are they occurring? Are there any other issues in your area and where are they happening?
- For the three or four key local pressures identified, what management arrangements are working in your area and what are not? Moving forward — what arrangements are needed? If there are no problems, is this because management arrangements are working or is it just luck?

At the end of the session, each group presented feedback which was tape-recorded and transcribed. The participants contributed detailed local knowledge including conflicts of use, new pressures and insights into management undertaken by other government agencies. Issues raised included the impacts from a growing number of fly-in fly-out miners who are moving to regional centres and becoming new recreational users on the Reef.

## **Stage two workshops**

Stakeholders and community members who participated in the workshops held between August and October 2012 were invited to reconvene to provide further advice on elements of the strategic assessment. The workshops would also provide an opportunity to update participants on the progress of the strategic assessment.

The objectives of the stage two workshops were:

- to test key findings to date and gain insights on issues of community concern
- to identify any discrepancies between expert opinion and on-ground community knowledge which may require further review
- to ensure the final reports are accessible and easy to understand for stakeholders and the wider community.

A total of 105 stakeholders attended workshops in Cairns, Townsville and Rockhampton in December 2012. The workshops included representatives of Traditional Owner groups, with strong attendances in Cairns and Rockhampton in particular.

The workshops began with presentations by Authority staff that provided updates on the progress of the strategic assessment, including outcomes of the previous workshops and other stakeholder engagement which had been undertaken. The workshop broke into roundtable discussions and feedback sessions which were facilitated and recorded by Authority staff.

Participants were asked for input on three key components:

- the condition and trends for values of the Great Barrier Reef
- management effectiveness of the Authority
- future management actions to protect the Great Barrier Reef.

## Condition and trends

The first feedback session focused on condition and trends. Authority staff explained the processes, data collection and mapping which had been undertaken to evaluate values and impacts. A list of 85 elements, including habitats, species, ecosystem processes, heritage and community benefits, had been developed, including using the information gathered in previous expert and stakeholder workshops. Each element had been graded in terms of its condition and trend, based on literature reviews, expert opinion and community knowledge.

Participants were asked to comment on the condition and trend in terms of:

- whether the approach was easy to understand
- whether the format was easy to understand
- their views on the gradings
- suggestions for improvements.

The information provided at the workshops had been summarised in a series of tables and for many participants it represented a large amount of information to absorb and evaluate in a short period of time. The following issues were consistently raised about the format and approach:

- improve and simplify the presentation of information
- clearly explain the methodology and categories
- make the grading parameters more consistent
- ensure the final reports can be used as management tools (for example, for Indigenous rangers).

Participants provided wide-ranging comments about the content of the tables, with many common concerns:

- the geographic scale is too broad to provide meaningful results of condition and trend
- highlight hotspots to avoid people assuming whole regions are in poor condition
- general summaries may be sensationalised by the media
- ensure transparency in the process used for scoring grades
- if confidence in the data is low, don't grade the element's condition and trend
- the positive trend of the community benefits was questioned.

## Management effectiveness

External consultants led the second component of the workshop, where they provided participants with their preliminary assessment of the Authority's management effectiveness and asked for feedback. Each table was provided with up to three management topics which had been graded from 'ineffective' to 'very effective'. Participants were asked to re-evaluate the grades to provide a sense of whether the community agreed with the outcomes.

Participants generally supported the grades, however discussions consistently returned to the difficulty of grading the Authority on management topics which were outside its jurisdiction. Workshop participants raised the issue of confusing federal, state and local government jurisdictions as a barrier to effective management, and the need to overcome these differences in order to effectively manage the Great Barrier Reef.

Traditional Owners at the Cairns workshop raised concerns that the management tools focused on the restriction of take of turtle and dugong rather than high-level, cooperative management approaches which could be achieved through mechanisms such as Indigenous Land Use Agreements. Traditional Owners at the Rockhampton workshop said management should be more holistic and consider the connections of land and water, and that the impacts of development should be controlled.

## Future management actions

The final session provided participants with an opportunity to consider the status and trends of values and current management practices, and to recommend future management actions which would help to protect those values. The responses were wide-ranging and showed the wealth of knowledge and insights of stakeholders and the community.

Many of the recommendations reflected earlier comments about management effectiveness:

- a whole of government approach to strategically plan for development, rather than react to proposals
- a tri-lateral agreement between all levels of government
- a seamless regulatory framework which recognises land and sea connections
- stronger partnerships and greater involvement in decision making and management by Traditional Owners
- streamlined application processes and quicker responses by the Authority
- increased compliance efforts

- continued focus on education and stewardship.

## Stage two outcomes

The management effectiveness information collected during stage two from the Local Marine Advisory Committee Chair's meeting, demonstration cases, surveys and workshops fed directly into the strategic assessment report:

- stakeholder feedback informed and improved the design and presentation of the values and trends information, and led to a re-evaluation of some of the grades
- the survey results generally confirmed the independent consultants' findings on the Authority's management effectiveness and provided insights into management gaps and overlaps
- the workshops also provided an opportunity to update stakeholders on the progress of the assessment, including how their previous input had informed the process
- the workshops also gave insights into the community's aspirations for the future of the Reef and its management.

## Summary


The stakeholder and community engagement undertaken during the strategic assessment provided the Authority with an opportunity to better understand how people value the Great Barrier Reef and their views on threats to those values. It provided valuable insights into how the community views management of the Great Barrier Reef, and how management could be improved.

One of the key successes has been involving stakeholders from the beginning of the process. They were invited to join the Authority in an inclusive process of learning more about the values, pressures and management issues facing the Great Barrier Reef Region. Their views were treated with respect and in return they shared a wealth of local information and knowledge for the strategic assessment.

The timeframes for such a complex and substantial assessment were tight, with less than four months to undertake consultation. Stakeholders were often inundated with multiple requests for information and feedback which had to be accommodated around their everyday work and life commitments. The Authority wishes to acknowledge the considerable contributions made by stakeholders and community members. Those contributions have enriched the process and significantly improved the outcome of the strategic assessment.

## Appendix 1: Background documents

**Figure 3 Background documents used in the preparation of the Stakeholder Engagement Report**





## Appendix 6

---

**Matters of national environmental significance  
protected under the EPBC Act that could be impacted**



## Matters of national environmental significance protected under the EPBC Act that could be impacted by activities under the Program

Each matter is referenced to the relevant section of the Strategic Assessment Report (SAR).

References in normal text refer specifically to the matter, references in *italics* refer to the category into which the matter fits. Some matters occur in the Coastal Zone and are referenced accordingly.

Also note that Sections 4.9 – Environmental Processes; and 4.10 – Connections between matters of national environmental significance; outline the overlap, connections and importance of connectivity for protected matters within and adjacent to the Great Barrier Reef Region.

Species/Location	GBRMPA SAR
<b>World Heritage Places that could be impacted by activities under the program</b>	
<i>Outcomes: The outstanding universal value of World Heritage properties are identified, protected, conserved, presented and transmitted to future generations. Australia's World Heritage properties are managed to ensure that Australia is not subject to international sanction, none of the properties are placed on the 'in danger' list and none are delisted. (Standards for Accreditation of Environmental Approvals under the EPBC Act 1999 – 2014)</i>	
Great Barrier Reef	4.2
Wet Tropics of Queensland	4.2
<b>Great Barrier Reef Marine Park</b>	
<i>Outcomes: The outstanding universal value of the Great Barrier Reef Marine Park, as a World Heritage property, is identified, protected, conserved, presented and transmitted to future generations. The environmental, biodiversity and heritage values of the Great Barrier Reef Marine Park are protected and conserved for the long term, consistent with the objects of the Great Barrier Reef Marine Park Act 1975. (Standards for Accreditation of Environmental Approvals under the EPBC Act 1999 – 2014)</i>	
Great Barrier Reef	4.3
<b>National Heritage Places that could be impacted by activities under the program</b>	
<i>Outcome: The outstanding value to the nation of National Heritage properties is identified, protected, conserved, presented and transmitted to future generations of Australians. (Standards for Accreditation of Environmental Approvals under the EPBC Act 1999 – 2014)</i>	
Great Barrier Reef	4.4
Wet Tropics of Queensland	4.3.1 - Biodiversity
<b>Commonwealth Marine Environment</b>	
<i>Outcome: The ecosystem functioning and integrity of Commonwealth marine waters are maintained or enhanced in full conformity with relevant marine bioregional plans. (Standards for Accreditation of Environmental Approvals under the EPBC Act 1999 – 2014)</i>	
Commonwealth marine area in the Great Barrier Reef Region	4.5
<b>Listed Migratory species that could be impacted by activities under the program</b>	
<i>Outcome: The survival and conservation status of listed migratory species and their critical habitat will be promoted and enhanced, consistent with Australia's international obligations. (Standards for Accreditation of Environmental Approvals under the EPBC Act 1999 – 2014)</i>	
<b>Birds</b>	
Barn Swallow ( <i>Hirundo rustica</i> )	4.6.7 - Seabirds
Bar-tailed Godwit ( <i>Limosa lapponica</i> )	Table 4.4
Black-browed Albatross ( <i>Thalassarche melanophris</i> )	4.6.7 - Seabirds
Black-faced Monarch ( <i>Monarcha melanopsis</i> )	4.6.8 - Shorebirds
Black-naped Tern ( <i>Sterna sumatrana</i> )	Table 4.4
Black-tailed Godwit ( <i>Limosa limosa</i> )	Table 4.4
Black-winged Monarch ( <i>Monarcha frater</i> )	4.6.8 - Shorebirds

Species/Location	GBRMPA SAR
Bridled Tern ( <i>Sterna anaethetus</i> )	Table 4.4
Broad-billed Sandpiper ( <i>Limicola falcinellus</i> )	4.6.7 - Seabirds
Brown Booby ( <i>Sula leucogaster</i> )	Table 4.4
Campbell Albatross ( <i>Thalassarche impavida</i> )	4.6.7 - Seabirds
Caspian Plover ( <i>Charadrius asiaticus</i> )	Table 4.4
Caspian Tern ( <i>Sterna caspia</i> )	Table 4.4
Cattle Egret ( <i>Ardea ibis</i> )	Table 4.4
Chatham Albatross ( <i>Thalassarche eremita</i> )	4.6.7 - Seabirds
Common Noddy ( <i>Anous stolidus</i> )	Table 4.4
Common Sandpiper ( <i>Actitis hypoleucos</i> )	Table 4.4
Common Tern ( <i>Sterna hirund</i> )	Table 4.4
Curlew Sandpiper ( <i>Calidris ferruginea</i> )	Table 4.4
Double-banded Plover ( <i>Charadrius bicinctus</i> )	4.6.8 - Shorebirds
Eastern Curlew ( <i>Numenius madagascariensis</i> )	Table 4.4
Eastern Great Egret, White Egret ( <i>Egretta alba</i> )	Table 4.4
Eastern Reef Egret ( <i>Egretta sacra</i> )	Table 4.4
Flesh-footed Shearwater, Fleshy-footed Shearwater ( <i>Puffinus carneipes</i> )	4.6.7 - Seabirds
Fork-tailed Swift ( <i>Apus pacificus</i> )	4.6.8 - Shorebirds
Glossy Ibis ( <i>Plegadis falcinellus</i> )	Table 4.4
Great Frigatebird, Greater Frigatebird ( <i>Fregata minor</i> )	Table 4.4
Great Knot ( <i>Calidris tenuirostris</i> )	Table 4.4
Greater Sand Plover, Large Sand Plover ( <i>Charadrius leschenaultii</i> )	Table 4.4
Greenshank ( <i>Tringa nebularia</i> )	Table 4.4
Grey Plover ( <i>Pluvialis squatarola</i> )	Table 4.4
Grey-headed Albatross ( <i>Thalassarche chrysostoma</i> )	Tables 4.4 and 4.7
Grey-tailed Tattler ( <i>Heteroscelus brevipes</i> )	Table 4.4
Latham's Snipe, Japanese Snipe ( <i>Gallinago hardwickii</i> )	Table 4.4
Lesser Crested Tern ( <i>Sterna bengalensis</i> )	Table 4.4
Lesser Frigatebird, Least Frigatebird ( <i>Fregata ariel</i> )	Table 4.4
Lesser Golden Plover ( <i>Pluvialis dominica</i> )	Table 4.4
Lesser Sand Plover, Mongolian Plover ( <i>Charadrius mongolus</i> )	Table 4.4
Little Curlew, Little Whimbrel ( <i>Numenius minutes</i> )	Table 4.4
Little Tern ( <i>Sterna albifrons</i> )	Table 4.4
Marsh Sandpiper, Little Greenshank ( <i>Tringa stagnatilis</i> )	Table 4.4
Masked Booby ( <i>Sula dactylatra</i> )	Table 4.4
Northern Giant Petrel ( <i>Macronectes halli</i> )	Tables 4.4 and 4.7
Oriental Plover, Oriental Dotterel ( <i>Charadrius veredus</i> )	4.6.8 - Shorebirds
Pacific Golden Plover ( <i>Pluvialis fulva</i> )	4.6.8 - Shorebirds
Painted Snipe ( <i>Rostratula benghalensis (sensu lato)</i> )	4.6.8 - Shorebirds



Species/Location	GBRMPA SAR
Pin-tailed Snipe ( <i>Gallinago stenura</i> )	4.6.8 - Shorebirds
Rainbow Bee-eater ( <i>Merops ornatus</i> )	4.6.8 - Shorebirds
Red Knot, Knot ( <i>Calidris canutus</i> )	Table 4.4
Red-footed Booby ( <i>Sula sula</i> )	Table 4.4
Red-necked Stint ( <i>Calidris ruficollis</i> )	Table 4.4
Roseate Tern ( <i>Sterna dougallii</i> )	Table 4.4
Ruddy Turnstone ( <i>Arenaria interpres</i> )	Table 4.4
Rufous Fantail ( <i>Rhipidura rufifrons</i> )	4.6.8 - Shorebirds
Salvin's Albatross ( <i>Thalassarche salvini</i> )	4.6.7 - Seabirds
Sanderling ( <i>Calidris alba</i> )	Table 4.4
Sarus Crane ( <i>Grus antigone</i> )	4.6.8 - Shorebirds
Satin Flycatcher ( <i>Myiagra cyanoleuca</i> )	4.6.8 - Shorebirds
Sharp-tailed Sandpiper ( <i>Calidris acuminata</i> )	Table 4.4
Shy Albatross, Tasmanian Shy Albatross ( <i>Thalassarche cauta cauta</i> )	4.6.7 - Seabirds
Sooty Albatross ( <i>Phoebastria fusca</i> )	Tables 4.4 and 4.7
Southern Giant-Petrel ( <i>Macronectes giganteus</i> )	Table 4.4
Spectacled Monarch ( <i>Symposiachrus trivirgatus</i> )	4.6.8 - Shorebirds
Streaked Shearwater ( <i>Calonectris leucomelas</i> )	4.6.7 - Seabirds
Swinhoe's Snipe ( <i>Gallinago megala</i> )	Table 4.4
Terek Sandpiper ( <i>Xenus cinereus</i> )	Table 4.4
Wandering Albatross ( <i>Diomedea exulans</i> )	Tables 4.4 and 4.7
Wedge-tailed Shearwater ( <i>Puffinus pacificus</i> )	Table 4.4
Whimbrel ( <i>Numenius phaeopus</i> )	Table 4.4
White-bellied Sea-Eagle ( <i>Haliaeetus leucogaster</i> )	Table 4.4
White-capped Albatross ( <i>Thalassarche steadi</i> )	4.6.7 - Seabirds
White-throated Needletail ( <i>Hirundapus caudacutus</i> )	4.6.8 - Shorebirds
White-winged Black Tern ( <i>Chlidonias leucopterus</i> )	Table 4.4
Wilson's Storm Petrel ( <i>Chlidonias leucopterus</i> )	Table 4.4
Wood Sandpiper ( <i>Tringa glareola</i> )	Table 4.4
<b>Dugong</b>	
Dugong (Dugong dugon)	Table 4.4
<b>Other Cetaceans</b>	
Australian Snubfin Dolphin ( <i>Orcaella heinsohni</i> )	Table 4.4
Indo-Pacific Humpback Dolphin ( <i>Sousa chinensis</i> )	Table 4.4
Irrawaddy Dolphin ( <i>Orcaella heinsohni</i> )	4.6.4 - Dolphins
<b>Reptiles</b>	
Flatback Turtle ( <i>Natator depressus</i> )	Table 4.4
Green Turtle ( <i>Chelonia mydas</i> )	Table 4.4
Hawksbill Turtle ( <i>Eretmochelys imbricata</i> )	Table 4.4

Species/Location	GBRMPA SAR
Leatherback Turtle, Leathery Turtle, Luth ( <i>Dermochelys coriacea</i> )	Table 4.4
Loggerhead Turtle ( <i>Caretta caretta</i> )	Table 4.4
Olive Ridley Turtle, Pacific Ridley Turtle ( <i>Lepidochelys olivacea</i> )	Table 4.4
Salt-water Crocodile, Estuarine Crocodile ( <i>Crocodylus porosus</i> )	Table 4.4
<b>Sharks</b>	
Giant Manta Ray, Chevron Manta Ray, Pacific Manta Ray, Pelagic Manta Ray, Oceanic Manta Ray ( <i>Manta birostris</i> )	4.6.6 - Sharks
Great White Shark ( <i>Carcharodon carcharias</i> )	Table 4.4
Longfin Mako ( <i>Isurus paucus</i> )	Table 4.4
Porbeagle, Mackerel Shark ( <i>Lamna nasus</i> )	Table 4.4
Shortfin Mako, Mako Shark ( <i>Isurus oxyrinchus</i> )	Table 4.4
Whale Shark ( <i>Rhincodon typus</i> )	Table 4.4
<b>Whales</b>	
Antarctic Minke Whale, Dark-shoulder Minke Whale ( <i>Balaenoptera bonaerensis</i> )	Table 4.4
Blue Whale ( <i>Balaenoptera musculus</i> )	Table 4.4
Bryde's Whale ( <i>Balaenoptera edeni</i> )	Table 4.4
Humpback Whale ( <i>Megaptera novaeangliae</i> )	Table 4.4
Killer Whale, Orca ( <i>Orcinus orca</i> )	Table 4.4
Sei Whale	Tables 4.4 and 4.7
Southern Right Whale ( <i>Eubalaena australis</i> )	4.6.3 - Whales
Sperm Whale ( <i>Physeter macrocephalus</i> )	Table 4.4
<b>Listed Threatened Species and Ecological Communities that could be impacted by activities under the program</b>	
<p><i>Outcome: The survival and conservation status of listed threatened species and ecological communities will be promoted and enhanced, including through the conservation of habitat critical to the survival of a species or community and other measures contained in any recovery plans, threat abatement plans or conservation advices.</i></p> <p><i>(Standards for Accreditation of Environmental Approvals under the EPBC Act 1999 – 2014)</i></p>	
<b>Birds</b>	
Australasian Bittern ( <i>Botaurus poeciloptilus</i> ) (E)	Coastal Zone
Australian Fairy Tern ( <i>Sternula nereis nereis</i> )	Coastal Zone
Australian Painted Snipe ( <i>Rostratula australis</i> ) (E)	Coastal Zone
Black-breasted Button-quail ( <i>Turnix melanogaster</i> )	Coastal Zone
Black-browed Albatross ( <i>Thalassarche melanophris</i> )	4.7.4 - Seabirds
Black-throated Finch (southern) ( <i>Poephila cincta cincta</i> ) (E)	Coastal Zone
Buff-breasted Button-quail ( <i>Turnix olivii</i> ) (E)	Coastal Zone
Campbell Albatross ( <i>Thalassarche melanophris impavida</i> )	4.7.4 - Seabirds
Chatham Albatross ( <i>Thalassarche eremite</i> ) (E)	4.7.4 - Seabirds
Coxen's Fig-Parrot ( <i>Cyclopsitta diophthalma coxeni</i> ) (E)	Coastal Zone
Crimson Finch (white-bellied) ( <i>Neochmia phaeton evangelinae</i> )	Coastal Zone
Golden-shouldered Parrot ( <i>Psephotus chrysopterygius</i> ) (E)	Coastal Zone
Gouldian Finch ( <i>Erythrura gouldiae</i> ) (E)	Coastal Zone
Grey-headed Albatross ( <i>Thalassarche chrysostoma</i> )	Tables 4.4 and 4.7

Species/Location	GBRMPA SAR
Herald Petrel ( <i>Pterodroma heraldica</i> ) (CE)	Table 4.7
Kermadec Petrel (western) ( <i>Pterodroma neglecta neglecta</i> )	4.7.4 - Seabirds
Masked Owl (northern) ( <i>Tyto novaehollandiae kimberli</i> )	Coastal Zone
Northern Giant Petrel ( <i>Macronectes halli</i> )	Tables 4.4 and 4.7
Red Goshawk ( <i>Erythrotriorchis radiatus</i> )	Coastal Zone
Salvin's Albatross ( <i>Thalassarche cauta salvini</i> )	4.7.4 - Seabirds
Shy Albatross, Tasmanian Shy Albatross ( <i>Thalassarche cauta cauta</i> )	4.7.4 - Seabirds
Sooty Albatross ( <i>Phoebastria fusca</i> )	Tables 4.4 and 4.7
Southern Cassowary (Australian), Southern Cassowary ( <i>Casuarius casuarius johnsonii</i> ) (E)	Coastal Zone
Southern Giant-Petrel ( <i>Macronectes giganteus</i> ) (E)	Table 4.7
Squatter Pigeon (southern) ( <i>Geophaps scripta scripta</i> )	Coastal Zone
Star Finch (eastern), Star Finch (southern) ( <i>Neochmia ruficauda ruficauda</i> ) (E)	Coastal Zone
Wandering Albatross ( <i>Diomedea exulans</i> )	Tables 4.4 and 4.7
White-bellied Storm-Petrel (Tasman Sea), White-bellied Storm-Petrel (Australasian) ( <i>Fregetta grallaria grallaria</i> )	4.7.4 - Seabirds
White-capped Albatross ( <i>Thalassarche cauta steadi</i> )	4.7.4 - Seabirds
Yellow Chat (Dawson) ( <i>Epthianura crocea macgregori</i> ) (CE)	Coastal Zone
<b>Ecological Communities</b>	
Brigalow ( <i>Acacia harpophylla</i> dominant and co-dominant)	4.3.1 - Biodiversity
Broad leaf tea-tree ( <i>Melaleuca viridiflora</i> ) woodlands in high rainfall coastal north Queensland	4.3.1 - Biodiversity
Coolibah - Black Box Woodlands of the Darling Riverine Plains and the Brigalow Belt South Bioregions	4.3.1 - Biodiversity
Littoral Rainforest and Coastal Vine Thickets of Eastern Australia	4.3.1 - Biodiversity
Lowland Subtropical Rainforest of Subtropical Australia (CE)	4.3.1 - Biodiversity
Semi-evergreen vine thickets of the Brigalow Belt (North and South) and Nandewar Bioregions	4.3.1 - Biodiversity
Weeping Myall Woodlands	4.3.1 - Biodiversity
<b>Fish</b>	
Freshwater Sawfish ( <i>Pristis pristis</i> )	Table 4.7
Honey Blue-eye ( <i>Pseudomugil mellis</i> )	Coastal Zone
Lake Eacham Rainbowfish ( <i>Melanotaenia eachamensis</i> ) (E)	Coastal Zone
Opal Cling Goby ( <i>Stiphodon semoni</i> ) (CE)	Coastal Zone
<b>Frogs</b>	
Armoured Mistfrog ( <i>Litoria lorica</i> ) (CE)	Coastal Zone
Common Mistfrog ( <i>Litoria rheocola</i> ) (E)	Coastal Zone
Kuranda Tree Frog ( <i>Litoria myola</i> ) (E)	Coastal Zone
Lace-eyed Tree Frog ( <i>Nyctimystes dayi</i> ) (E)	Coastal Zone
Mountain Mistfrog ( <i>Litoria nyakalensis</i> ) (CE)	Coastal Zone
Waterfall Frog, Torrent Tree Frog ( <i>Litoria nannotis</i> ) (E)	Coastal Zone
<b>Mammals</b>	

Species/Location	GBRMPA SAR
Bare-rumped Sheath-tail Bat ( <i>Saccolaimus saccolaimus nudiclunatus</i> ) (CE)	Coastal Zone
Brush-tailed Rabbit-rat, Brush-tailed Tree-rat, Pakooma ( <i>Conilurus penicillatus</i> )	Coastal Zone
Burrowing Bettong (Shark Bay) ( <i>Bettongia lesueur lesueur</i> )	Coastal Zone
Greater Large-eared Horseshoe Bat ( <i>Rhinolophus philippinensis (large form)</i> ) (E)	Coastal Zone
Grey-headed Flying-fox ( <i>Pteropus poliocephalus</i> )	Coastal Zone
Koala (combined populations of Queensland, New South Wales and the Australian Capital Territory) ( <i>Phascolarctos cinereus</i> (combined populations of Qld, NSW and the ACT))	Coastal Zone and some islands
Large-eared Pied Bat, Large Pied Bat ( <i>Chalinolobus dwyeri</i> )	Coastal Zone
Mahogany Glider ( <i>Petaurus gracilis</i> ) (E)	Coastal Zone
Northern Bettong ( <i>Bettongia tropica</i> ) (E)	Coastal Zone
Northern Hopping-mouse, Woorrentinta ( <i>Notomys aquilo</i> )	Coastal Zone
Northern Quoll ( <i>Dasyurus hallucatus</i> ) (E)	Coastal Zone
Proserpine Rock-wallaby ( <i>Petrogale Persephone</i> ) (E)	Coastal Zone
Semon's Leaf-nosed Bat, Greater Wart-nosed Horseshoe-bat ( <i>Hipposideros semoni</i> ) (E)	Coastal Zone
South-eastern Long-eared Bat ( <i>Nyctophilus corbeni</i> )	Coastal Zone
Spectacled Flying-fox ( <i>Pteropus conspicillatus</i> )	Coastal Zone
Spotted-tailed Quoll or Yarri (North Queensland subspecies) ( <i>Dasyurus maculatus gracilis</i> ) (E)	Coastal Zone
Subantarctic Fur Seal ( <i>Arctocephalus tropicalis</i> )	Table 4.7
Water Mouse, False Water Rat, Yirrkoo ( <i>Xeromys myoides</i> )	Coastal Zone and some islands
<b>Plants</b>	
<i>Actephila foetida</i>	4.3.1 - Biodiversity
a palm ( <i>Hydriastele costata</i> )	4.3.1 - Biodiversity
a sedge ( <i>Eleocharis retroflexa</i> )	4.3.1 - Biodiversity
a shrub ( <i>Cyclophyllum costatum</i> )	4.3.1 - Biodiversity
an aquatic herb ( <i>Aponogeton prolifer</i> ) (E)	4.3.1 - Biodiversity
an orchid ( <i>Cepobaculum carronii</i> )	4.3.1 - Biodiversity
an orchid ( <i>Durabaculum mirbelianum</i> ) (E)	4.3.1 - Biodiversity
an orchid ( <i>Durabaculum nindii</i> ) (E)	4.3.1 - Biodiversity
Ant Plant ( <i>Myrmecodia beccarii</i> )	4.3.1 - Biodiversity
Antelope Orchid ( <i>Ceratobium antennatum</i> ) (E)	4.3.1 - Biodiversity
<i>Aponogeton bullosus</i> (E)	4.3.1 - Biodiversity
<i>Asplenium wildii</i>	4.3.1 - Biodiversity
Black Ironbox ( <i>Eucalyptus raveretiana</i> )	4.3.1 - Biodiversity
Blue Tassel-fern ( <i>Phlegmariurus dalhousieanus</i> ) (E)	4.3.1 - Biodiversity
Bluegrass ( <i>Dichanthium setosum</i> )	4.3.1 - Biodiversity
Byfield Matchstick ( <i>Comesperma oblongatum</i> )	4.3.1 - Biodiversity
<i>Cajanus mareebensis</i> (E)	4.3.1 - Biodiversity
<i>Calophyllum bicolor</i>	4.3.1 - Biodiversity
<i>Canarium acutifolium</i> var. <i>Acutifolium</i>	4.3.1 - Biodiversity



Species/Location	GBRMPA SAR
Cape York Vanda ( <i>Vanda hindsii</i> )	4.3.1 - Biodiversity
Cardwell Beard Orchid ( <i>Calochilus psednus</i> ) (E)	4.3.1 - Biodiversity
Cardwell Midge Orchid ( <i>Genoplesium tectum</i> ) (E)	4.3.1 - Biodiversity
<i>Carronia pedicellata</i> (E)	4.3.1 - Biodiversity
<i>Cepobaculum johannis</i>	4.3.1 - Biodiversity
<i>Chingia australis</i> (E)	4.3.1 - Biodiversity
Cooktown Orchid ( <i>Dendrobium bigibbum</i> )	4.3.1 - Biodiversity
Cooktown Orchid ( <i>Vappodes phalaenopsis</i> )	4.3.1 - Biodiversity
Dwarf Butterfly Orchid, Cooktown Orchid ( <i>Vappodes lithocola</i> ) (E)	4.3.1 - Biodiversity
<i>Crepidium lawleri</i> (E)	4.3.1 - Biodiversity
<i>Cycas megacarpa</i> (E)	4.3.1 - Biodiversity
<i>Cycas ophiolitica</i> (E)	4.3.1 - Biodiversity
<i>Cycas silvestris</i>	4.3.1 - Biodiversity
<i>Cyperus cephalotes</i> (E)	4.3.1 - Biodiversity
<i>Diplazium cordifolium</i>	4.3.1 - Biodiversity
<i>Diplazium pallidum</i> (E)	4.3.1 - Biodiversity
<i>Dipodium pictum</i> (E)	4.3.1 - Biodiversity
<i>Drosera prolifera</i>	4.3.1 - Biodiversity
<i>Endiandra cooperana</i> (E)	4.3.1 - Biodiversity
<i>Eremochloa muricata</i> (E)	4.3.1 - Biodiversity
<i>Fimbristylis adjuncta</i> (E)	4.3.1 - Biodiversity
<i>Gardenia actinocarpa</i> (E)	4.3.1 - Biodiversity
<i>Germainia capitata</i>	4.3.1 - Biodiversity
Glen Geddes Bloodwood ( <i>Corymbia xanthope</i> )	4.3.1 - Biodiversity
Hairy-joint Grass ( <i>Arthraxon hispidus</i> )	4.3.1 - Biodiversity
Hann Gardenia ( <i>Gardenia psidioides</i> )	4.3.1 - Biodiversity
Holly-leaved Graptophyllum, Mt Blackwood Holly ( <i>Graptophyllum ilicifolium</i> )	4.3.1 - Biodiversity
Lesser Swamp-orchid ( <i>Phaius australis</i> ) (E)	4.3.1 - Biodiversity
<i>Lindsaea pulchella</i> var. <i>blanda</i>	4.3.1 - Biodiversity
<i>Marsdenia brevifolia</i>	4.3.1 - Biodiversity
<i>Medicosma obovata</i>	4.3.1 - Biodiversity
Miniature Moss-orchid, Hoop Pine Orchid ( <i>Bulbophyllum globuliforme</i> )	4.3.1 - Biodiversity
Mt Berryman Phebalium ( <i>Phebalium distans</i> ) (CE)	4.3.1 - Biodiversity
Mt Larcom Silk Pod ( <i>Parsonsia larcomensis</i> )	4.3.1 - Biodiversity
Myola Palm, Myola Archontophoenix ( <i>Archontophoenix myolensis</i> ) (E)	4.3.1 - Biodiversity
<i>Myriophyllum coronatum</i>	4.3.1 - Biodiversity
Native Moth Orchid ( <i>Phalaenopsis rosenstromii</i> ) (E)	4.3.1 - Biodiversity
<i>Neisosperma kilneri</i>	4.3.1 - Biodiversity
<i>Neoroepera buxifolia</i>	4.3.1 - Biodiversity

Species/Location	GBRMPA SAR
<i>Omphalea celata</i>	4.3.1 - Biodiversity
<i>Oreogrammitis reinwardtii</i>	4.3.1 - Biodiversity
Pale Chandelier Orchid ( <i>Acriopsis emarginata</i> )	4.3.1 - Biodiversity
<i>Phaleria biflora</i>	4.3.1 - Biodiversity
<i>Phlegmariurus lockyeri</i>	4.3.1 - Biodiversity
<i>Pimelea leptospermoides</i>	4.3.1 - Biodiversity
<i>Plectranthus gratus</i>	4.3.1 - Biodiversity
<i>Polyscias bellendenkerensis</i>	4.3.1 - Biodiversity
<i>Prostanthera clotteniana</i> (CE)	4.3.1 - Biodiversity
<i>Pultenaea setulosa</i>	4.3.1 - Biodiversity
Rat's Tail Tassel-fern ( <i>Phlegmariurus filiformis</i> ) (E)	4.3.1 - Biodiversity
<i>Rhinerrhizopsis matutina</i>	4.3.1 - Biodiversity
<i>Ristantia gouldii</i>	4.3.1 - Biodiversity
Rock Tassel-fern, Water Tassel-fern ( <i>Phlegmariurus squarrosus</i> ) (E)	4.3.1 - Biodiversity
<i>Samadera bidwillii</i>	4.3.1 - Biodiversity
Siah's Backbone, Sia's Backbone, Isaac Wood ( <i>Streblus pendulinus</i> ) (E)	4.3.1 - Biodiversity
Square Tassel Fern ( <i>Phlegmariurus tetrastichoides</i> )	4.3.1 - Biodiversity
Swamp Lily, Greater Swamp-orchid ( <i>Phaius tancarvilleae</i> ) (E)	4.3.1 - Biodiversity
<i>Syzygium velarum</i>	4.3.1 - Biodiversity
<i>Tephrosia leveillei</i>	4.3.1 - Biodiversity
Thin Feather Orchid ( <i>Tropilis callitrophilis</i> )	4.3.1 - Biodiversity
Three-leaved Bosistoa, Heart-leaved Bosistoa, Yellow Satinheart, Heart-leaved Bonewood ( <i>Bosistoa transversa</i> s. Lat.)	4.3.1 - Biodiversity
<i>Toechima pterocarpum</i> (E)	4.3.1 - Biodiversity
<i>Tomophyllum walleri</i>	4.3.1 - Biodiversity
Velvet Jewel Orchid ( <i>Zeuxine polygonoides</i> )	4.3.1 - Biodiversity
<i>Vrydagzynea grayi</i> (E)	4.3.1 - Biodiversity
Water Tassel-fern ( <i>Phlegmariurus marsupiiiformis</i> )	4.3.1 - Biodiversity
Wedge-leaf Tuckeroo ( <i>Cupaniopsis shirleyana</i> )	4.3.1 - Biodiversity
<i>Xanthostemon formosus</i> (E)	4.3.1 - Biodiversity
Yarwun Whitewood ( <i>Atalaya collina</i> ) (E)	4.3.1 - Biodiversity
<b>Reptiles</b>	
Collared Delma ( <i>Delma torquata</i> )	Coastal Zone
Dunmall's Snake ( <i>Furina dunmalli</i> )	Coastal Zone
Fitzroy River Turtle, Fitzroy Tortoise, Fitzroy Turtle, White-eyed River Diver ( <i>Rheodytes leukops</i> )	Coastal Zone
Flatback Turtle ( <i>Natator depressus</i> )	Table 4.7
Green Turtle ( <i>Chelonia mydas</i> )	Table 4.7
Hawksbill Turtle ( <i>Eretmochelys imbricata</i> )	Table 4.7
Leatherback Turtle, Leathery Turtle, Luth ( <i>Dermochelys coriacea</i> ) (E)	Table 4.7

Species/Location	GBRMPA SAR
Loggerhead Turtle ( <i>Caretta caretta</i> ) (E)	Table 4.7
Olive Ridley Turtle, Pacific Ridley Turtle ( <i>Lepidochelys olivacea</i> ) (E)	Table 4.7
Ornamental Snake ( <i>Denisonia maculata</i> )	Coastal Zone
Yakka Skink ( <i>Egernia rugosa</i> )	Coastal Zone
<b>Sharks</b>	
Dwarf Sawfish, Queensland Sawfish ( <i>Pristis clavata</i> )	Table 4.7
Great White Shark ( <i>Carcharodon carcharias</i> )	Table 4.7
Green Sawfish, Dindagubba, Narrowsnout Sawfish ( <i>Pristis zijsron</i> )	Table 4.7
Grey Nurse Shark (east coast population) ( <i>Carcharias Taurus</i> (east coast population)) (CE)	Table 4.7
Largetooth Sawfish, Freshwater Sawfish, <i>Leichhardt's Sawfish</i> , Northern Sawfish ( <i>Pristis pristis</i> )	4.6.6 - Sharks
Speartooth Shark ( <i>Glyphis glyphis</i> ) (CE)	Table 4.7
Whale Shark ( <i>Rhincodon typus</i> )	Table 4.7
<b>Whales</b>	
Blue Whale ( <i>Balaenoptera musculus</i> ) (E)	Table 4.7
Fin Whale ( <i>Balaenoptera physalus</i> )	Table 4.7
Humpback Whale ( <i>Megaptera novaeangliae</i> )	Table 4.7
Sei Whale	Tables 4.4 and 4.7
Southern Right Whale ( <i>Eubalaena australis</i> ) (E)	4.7.2 – Marine Mammals
<b>Listed cetaceans (protected by Division 3, Part 13 of the EPBC Act) that may be impacted by activities under the Program</b>	
<p><i>Outcome: The survival and conservation status of listed species and ecological communities is promoted and enhanced, including through the conservation of habitat critical to the survival of a species or community and other measures contained in any recovery plans, threat abatement plans or conservation advices.</i></p> <p><i>(Standards for Accreditation of Environmental Approvals under the EPBC Act 1999 – 2014)</i></p>	
<b>Other Cetaceans</b>	
Bottlenose Dolphin ( <i>Tursiops truncatus</i> ( <i>sensu stricto</i> ))	4.6.4 - Dolphins
Common Dolphin, Short-beaked Common Dolphin ( <i>Delphinus delphis</i> )	4.6.4 - Dolphins
Fraser's Dolphin, Sarawak Dolphin ( <i>Lagenodelphis hosei</i> )	4.6.4 - Dolphins
Indian Ocean Bottlenose Dolphin, Spotted Bottlenose Dolphin ( <i>Tursiops aduncus</i> )	4.6.4 - Dolphins
Long-snouted Spinner Dolphin ( <i>Stenella longirostris</i> )	4.6.4 - Dolphins
Risso's Dolphin, Grampus ( <i>Grampus griseus</i> )	4.6.4 - Dolphins
Rough-toothed Dolphin ( <i>Steno bredanensis</i> )	4.6.4 - Dolphins
Spotted Dolphin, Pantropical Spotted Dolphin ( <i>Stenella attenuata</i> )	4.6.4 - Dolphins
Striped Dolphin, Euphrosyne Dolphin ( <i>Stenella coeruleoalba</i> )	4.6.4 - Dolphins
<b>Whales</b>	
Blainville's Beaked Whale, Dense-beaked Whale ( <i>Mesoplodon densirostris</i> )	4.6.3 - Whales
Cuvier's Beaked Whale, Goose-beaked Whale ( <i>Ziphius cavirostris</i> )	4.6.3 - Whales
Dwarf Sperm Whale ( <i>Kogia simus</i> )	4.6.3 - Whales
False Killer Whale ( <i>Pseudorca crassidens</i> )	4.6.3 - Whales
Melon-headed Whale ( <i>Peponocephala electra</i> )	4.6.3 - Whales

Species/Location	GBRMPA SAR
Minke Whale ( <i>Balaenoptera acutorostrata</i> )	4.6.3 - Whales
Pygmy Killer Whale ( <i>Feresa attenuata</i> )	4.6.3 - Whales
Pygmy Sperm Whale ( <i>Kogia breviceps</i> )	4.6.3 - Whales
Short-finned Pilot Whale ( <i>Globicephala macrorhynchus</i> )	4.6.3 - Whales
Strap-toothed Beaked Whale, Strap-toothed Whale, Layard's Beaked Whale ( <i>Mesoplodon layardii</i> )	4.6.3 - Whales
<b>Listed marine species that may be impacted by activities under the program</b>	
<p><i>Outcome: The survival and conservation status of listed species and ecological communities is promoted and enhanced, including through the conservation of habitat critical to the survival of a species or community and other measures contained in any recovery plans, threat abatement plans or conservation advices.</i></p> <p><i>(Standards for Accreditation of Environmental Approvals under the EPBC Act 1999 – 2014)</i></p>	
<b>Birds</b>	
Australian Pratincole ( <i>Stiltia Isabella</i> )	4.6.7 - Seabirds
Barn Swallow ( <i>Hirundo rustica</i> )	4.6.7 - Seabirds
Bar-tailed Godwit ( <i>Limosa lapponica</i> )	Table 4.4
Black Noddy ( <i>Anous minutus</i> )	4.6.7 - Seabirds
Black-browed Albatross ( <i>Thalassarche melanophris</i> )	4.6.7 – Seabirds and 4.7.4 - Seabirds
Black-faced Monarch ( <i>Monarcha melanopsis</i> )	4.6.7 - Seabirds
Black-naped Tern ( <i>Sterna sumatrana</i> )	Table 4.4
Black-tailed Godwit ( <i>Limosa limosa</i> )	Table 4.4
Black-winged Monarch ( <i>Monarcha frater</i> )	4.6.7 - Seabirds
Black-winged Petrel ( <i>Pterodroma nigripennis</i> )	4.6.7 - Seabirds
Black-winged Stilt ( <i>Himantopus himantopus</i> )	4.6.7 - Seabirds
Bridled Tern ( <i>Sterna anaethetus</i> )	Table 4.4
Broad-billed Sandpiper ( <i>Limicola falcinellus</i> )	4.6.7 - Seabirds
Brown Booby ( <i>Sula leucogaster</i> )	Table 4.4
Campbell Albatross ( <i>Thalassarche impavida</i> )	4.6.7 - Seabirds
Caspian Tern ( <i>Sterna caspia</i> )	Table 4.4
Cattle Egret ( <i>Ardea ibis</i> )	Table 4.4
Chatham Albatross ( <i>Thalassarche eremite</i> )	4.6.7 – Seabirds and 4.7.4 - Seabirds
Common Noddy ( <i>Anous stolidus</i> )	Table 4.4
Common Sandpiper ( <i>Actitis hypoleucos</i> )	Table 4.4
Crested Tern ( <i>Sterna bergii</i> )	4.6.7 - Seabirds
Curlew Sandpiper ( <i>Calidris ferruginea</i> )	Table 4.4
Double-banded Plover ( <i>Charadrius bicinctus</i> )	4.6.7 - Seabirds
Eastern Curlew ( <i>Numenius madagascariensis</i> )	Table 4.4
Flesh-footed Shearwater, Fleshy-footed Shearwater ( <i>Puffinus carneipes</i> )	4.6.7 - Seabirds
Fork-tailed Swift ( <i>Apus pacificus</i> )	4.6.7 - Seabirds
Great Egret, White Egret ( <i>Ardea modesta</i> )	Table 4.4
Great Frigatebird, Greater Frigatebird ( <i>Fregata minor</i> )	Table 4.4



Species/Location	GBRMPA SAR
Great Knot ( <i>Calidris tenuirostris</i> )	Table 4.4
Greater Sand Plover, Large Sand Plover ( <i>Charadrius leschenaultii</i> )	Table 4.4
Grey Plover ( <i>Pluvialis squatarola</i> )	Table 4.4
Grey-tailed Tattler ( <i>Heteroscelus brevipes</i> )	Table 4.4
Latham's Snipe, Japanese Snipe ( <i>Gallinago hardwickii</i> )	Table 4.4
Lesser Crested Tern ( <i>Sterna bengalensis</i> )	Table 4.4
Lesser Frigatebird, Least Frigatebird ( <i>Fregata ariel</i> )	Table 4.4
Lesser Sand Plover, Mongolian Plover ( <i>Charadrius mongolus</i> )	Table 4.4
Little Curlew, Little Whimbrel ( <i>Numenius minutus</i> )	Table 4.4
Little Ringed Plover ( <i>Charadrius dubius</i> )	4.6.7 - Seabirds
Little Tern ( <i>Sterna albifrons</i> )	Table 4.4
Magpie Goose ( <i>Anseranas semipalmata</i> )	4.6.7 - Seabirds
Marsh Sandpiper, Little Greenshank ( <i>Tringa stagnatilis</i> )	Table 4.4
Masked Booby ( <i>Sula dactylatra</i> )	Table 4.4
Oriental Plover, Oriental Dotterel ( <i>Charadrius veredus</i> )	4.6.7 - Seabirds
Osprey ( <i>Pandion haliaetus</i> )	Table 4.4
Pacific Golden Plover ( <i>Pluvialis fulva</i> )	4.6.7 - Seabirds
Painted Snipe ( <i>Rostratula benghalensis</i> ( <i>sensu lato</i> ))	4.6.7 - Seabirds
Pectoral Sandpiper ( <i>Calidris melanotos</i> )	4.6.7 - Seabirds
Pin-tailed Snipe ( <i>Gallinago stenura</i> )	4.6.7 - Seabirds
Rainbow Bee-eater ( <i>Merops ornatus</i> )	4.6.7 - Seabirds
Red Knot, Knot ( <i>Calidris canutus</i> )	Table 4.4
Red-capped Plover ( <i>Charadrius ruficapillus</i> )	4.6.7 - Seabirds
Red-footed Booby ( <i>Sula sula</i> )	Table 4.4
Red-necked Avocet ( <i>Recurvirostra novaehollandiae</i> )	4.6.7 - Seabirds
Red-necked Stint ( <i>Calidris ruficollis</i> )	Table 4.4
Red-tailed Tropicbird ( <i>Phaethon rubricauda</i> )	4.6.7 - Seabirds
Roseate Tern ( <i>Sterna dougallii</i> )	Table 4.4
Ruddy Turnstone ( <i>Arenaria interpres</i> )	Table 4.4
Ruff (Reeve) ( <i>Philomachus pugnax</i> )	4.6.7 - Seabirds
Rufous Fantail ( <i>Rhipidura rufifrons</i> )	4.6.7 - Seabirds
Salvin's Albatross ( <i>Thalassarche cauta salvini</i> )	4.6.7 - Seabirds
Sanderling ( <i>Calidris alba</i> )	Table 4.4
Satin Flycatcher ( <i>Myiagra cyanoleuca</i> )	4.6.7 - Seabirds
Sharp-tailed Sandpiper ( <i>Calidris acuminata</i> )	Table 4.4
Shy Albatross, Tasmanian Shy Albatross ( <i>Thalassarche cauta cauta</i> )	4.6.7 - Seabirds
Silver Gull ( <i>Larus novaehollandiae</i> )	4.6.7 - Seabirds
Southern Giant-Petrel ( <i>Macronectes giganteus</i> )	Table 4.7
Spectacled Monarch ( <i>Symposiachrus trivirgatus</i> )	4.6.7 - Seabirds

Species/Location	GBRMPA SAR
Streaked Shearwater ( <i>Calonectris leucomelas</i> )	4.6.7 - Seabirds
Swinhoe's Snipe ( <i>Gallinago megala</i> )	Table 4.4
Terek Sandpiper ( <i>Xenus cinereus</i> )	Table 4.4
Wandering Tattler ( <i>Heteroscelus incanus</i> )	Table 4.4
Wedge-tailed Shearwater ( <i>Puffinus pacificus</i> )	Table 4.4
Whimbrel ( <i>Numenius phaeopus</i> )	Table 4.4
White-bellied Sea-Eagle ( <i>Haliaeetus leucogaster</i> )	Table 4.4
White-capped Albatross ( <i>Thalassarche cauta steadii</i> )	4.6.7 - Seabirds
White-throated Needletail ( <i>Hirundapus caudacutus</i> )	4.6.7 - Seabirds
Wood Sandpiper ( <i>Tringa glareola</i> )	Table 4.4
<b>Dugong</b>	
Dugong ( <i>Dugong dugon</i> )	Table 4.4
<b>Fish</b>	
Anderson's Pipefish, Shortnose Pipefish ( <i>Micrognathus andersonii</i> )	4.3.1 - Biodiversity
Australian Messmate Pipefish, Banded Pipefish ( <i>Corythoichthys intestinalis</i> )	4.3.1 - Biodiversity
Banded Pipefish, Ringed Pipefish ( <i>Doryrhamphus dactyliophorus</i> )	4.3.1 - Biodiversity
Barred Short-bodied Pipefish, Girdled Pipefish ( <i>Choeroichthys cinctus</i> )	4.3.1 - Biodiversity
Beady Pipefish, Steep-nosed Pipefish ( <i>Hippichthys penicillus</i> )	4.3.1 - Biodiversity
Belly-barred Pipefish, Banded Freshwater Pipefish ( <i>Hippichthys spicifer</i> )	4.3.1 - Biodiversity
Bentstick Pipefish, Bend Stick Pipefish, Short-tailed Pipefish ( <i>Trachyrhamphus bicoarctatus</i> )	4.3.1 - Biodiversity
Blue-speckled Pipefish, Blue-spotted Pipefish ( <i>Hippichthys cyanospilos</i> )	4.3.1 - Biodiversity
Bluestripe Pipefish, Indian Blue-stripe Pipefish, Pacific Blue-stripe Pipefish ( <i>Doryrhamphus excisus</i> )	4.3.1 - Biodiversity
Brock's Pipefish ( <i>Halicampus brocki</i> )	4.3.1 - Biodiversity
Cleaner Pipefish, Janss' Pipefish ( <i>Doryrhamphus janssi</i> )	4.3.1 - Biodiversity
D'Arros Pipefish ( <i>Cosmocampus darrosanus</i> )	4.3.1 - Biodiversity
Davao Pughead Pipefish ( <i>Bulbonaricus davaoensis</i> )	4.3.1 - Biodiversity
Double-end Pipehorse, Double-ended Pipehorse, Alligator Pipefish ( <i>Syngnathoides biaculeatus</i> )	4.3.1 - Biodiversity
Duncker's Pipehorse ( <i>Solegnathus dunckeri</i> )	4.3.1 - Biodiversity
Fijian Banded Pipefish, Brown-banded Pipefish ( <i>Corythoichthys amplexus</i> )	4.3.1 - Biodiversity
Flagtail Pipefish, Masthead Island Pipefish ( <i>Doryrhamphus negrosensis</i> )	4.3.1 - Biodiversity
Flat-face Seahorse ( <i>Hippocampus planifrons</i> )	4.3.1 - Biodiversity
Gibbs' Pipefish ( <i>Festucalex gibbsi</i> )	4.3.1 - Biodiversity
Girdled Pipefish ( <i>Festucalex cinctus</i> )	4.3.1 - Biodiversity
Glittering Pipefish ( <i>Halicampus nitidus</i> )	4.3.1 - Biodiversity
Hairy Pipefish ( <i>Urocampus carinirostris</i> )	4.3.1 - Biodiversity
Hedgehog Seahorse ( <i>Hippocampus spinosissimus</i> )	4.3.1 - Biodiversity
Javelin Pipefish ( <i>Lissocampus runa</i> )	4.3.1 - Biodiversity
Kellogg's Seahorse, Great Seahorse ( <i>Hippocampus kelloggi</i> )	4.3.1 - Biodiversity

Species/Location	GBRMPA SAR
Madura Pipefish, Reticulated Freshwater Pipefish ( <i>Hippichthys heptagonus</i> )	4.3.1 - Biodiversity
Manado Pipefish, Manado River Pipefish ( <i>Microphis manadensis</i> )	4.3.1 - Biodiversity
Maxweber's Pipefish ( <i>Cosmocampus maxweberi</i> )	4.3.1 - Biodiversity
Mother-of-pearl Pipefish ( <i>Vanacampus margaritifer</i> )	4.3.1 - Biodiversity
Mud Pipefish, Gray's Pipefish ( <i>Halicampus grayi</i> )	4.3.1 - Biodiversity
Offshore Pipefish ( <i>Micrognathus natans</i> )	4.3.1 - Biodiversity
Orange-spotted Pipefish, Ocellated Pipefish ( <i>Corythoichthys ocellatus</i> )	4.3.1 - Biodiversity
Ornate Ghostpipefish, Harlequin Ghost Pipefish, Ornate Ghost Pipefish ( <i>Solenostomus paradoxus</i> )	4.3.1 - Biodiversity
Pacific Short-bodied Pipefish, Short-bodied Pipefish ( <i>Choeroichthys brachysoma</i> )	4.3.1 - Biodiversity
Painted Pipefish, Reef Pipefish ( <i>Nannocampus pictus</i> )	4.3.1 - Biodiversity
Pale-blotched Pipefish, Spined Pipefish ( <i>Phoxocampus diacanthus</i> )	4.3.1 - Biodiversity
Pallid Pipehorse, Hardwick's Pipehorse ( <i>Solegnathus hardwickii</i> )	4.3.1 - Biodiversity
Paxton's Pipefish ( <i>Corythoichthys paxtoni</i> )	4.3.1 - Biodiversity
Pig-snouted Pipefish ( <i>Choeroichthys suillus</i> )	4.3.1 - Biodiversity
Pygmy Seahorse ( <i>Hippocampus bargibanti</i> )	4.3.1 - Biodiversity
Red-hair Pipefish, Duncker's Pipefish ( <i>Halicampus dunckeri</i> )	4.3.1 - Biodiversity
Reef-top Pipefish ( <i>Corythoichthys haematopterus</i> )	4.3.1 - Biodiversity
Reticulate Pipefish, Yellow-banded Pipefish, Network Pipefish ( <i>Corythoichthys flavofasciatus</i> )	4.3.1 - Biodiversity
Ribboned Pipehorse, Ribboned Seadragon ( <i>Haliichthys taeniophorus</i> )	4.3.1 - Biodiversity
Robust Ghost Pipefish, Blue-finned Ghost Pipefish ( <i>Solenostomus cyanopterus</i> )	4.3.1 - Biodiversity
Samoa Pipefish ( <i>Halicampus matafae</i> )	4.3.1 - Biodiversity
Sawtooth Pipefish ( <i>Maroubra perserrata</i> )	4.3.1 - Biodiversity
Schultz's Pipefish ( <i>Corythoichthys schultzi</i> )	4.3.1 - Biodiversity
Sculptured Pipefish ( <i>Choeroichthys sculptus</i> )	4.3.1 - Biodiversity
Shortpouch Pygmy Pipehorse ( <i>Acentronura tentaculata</i> )	4.3.1 - Biodiversity
Short-tail Pipefish, Short-tailed River Pipefish ( <i>Microphis brachyurus</i> )	4.3.1 - Biodiversity
Softcoral Pipefish, Soft-coral Pipefish ( <i>Siokunichthys breviceps</i> )	4.3.1 - Biodiversity
Spiny Pipehorse, Australian Spiny Pipehorse ( <i>Solegnathus spinosissimus</i> )	4.3.1 - Biodiversity
Spiny Seahorse, Thorny Seahorse ( <i>Hippocampus histrix</i> )	4.3.1 - Biodiversity
Spiny-snout Pipefish ( <i>Halicampus spinirostris</i> )	4.3.1 - Biodiversity
Spotted Seahorse, Yellow Seahorse ( <i>Hippocampus kuda</i> )	4.3.1 - Biodiversity
Straightstick Pipefish, Long-nosed Pipefish, Straight Stick Pipefish ( <i>Trachyrhamphus longirostris</i> )	4.3.1 - Biodiversity
Thorntail Pipefish, Thorn-tailed Pipefish ( <i>Micrognathus brevirostris</i> )	4.3.1 - Biodiversity
Three-keel Pipefish ( <i>Campichthys tricarinatus</i> )	4.3.1 - Biodiversity
Tiger Pipefish ( <i>Filicampus tigris</i> )	4.3.1 - Biodiversity
Tryon's Pipefish ( <i>Campichthys tryoni</i> )	4.3.1 - Biodiversity
Western Spiny Seahorse, Narrow-bellied Seahorse ( <i>Hippocampus angustus</i> )	4.3.1 - Biodiversity
Whiskered Pipefish, Ornate Pipefish ( <i>Halicampus macrorhynchus</i> )	4.3.1 - Biodiversity

Species/Location	GBRMPA SAR
White's Seahorse, Crowned Seahorse, Sydney Seahorse ( <i>Hippocampus whitei</i> )	4.3.1 - Biodiversity
Widebody Pipefish, Wide-bodied Pipefish, Black Pipefish ( <i>Stigmatopora nigra</i> )	4.3.1 - Biodiversity
Zebra Seahorse ( <i>Hippocampus zebra</i> )	4.3.1 - Biodiversity
<b>Reptiles</b>	
a sea krait ( <i>Laticauda colubrina</i> )	4.3.1 - Biodiversity
a sea krait ( <i>Laticauda laticaudata</i> )	4.3.1 - Biodiversity
a seasnake ( <i>Hydrophis vorisi</i> )	4.3.1 - Biodiversity
Beaked Seasnake ( <i>Enhydrina schistosa</i> )	4.3.1 - Biodiversity
Black-banded Robust Seasnake ( <i>Hydrophis melanosoma</i> )	4.3.1 - Biodiversity
Black-headed Seasnake ( <i>Hydrophis atriceps</i> )	4.3.1 - Biodiversity
Dubois' Seasnake ( <i>Aipysurus duboisii</i> )	4.3.1 - Biodiversity
Elegant Seasnake ( <i>Hydrophis elegans</i> )	4.3.1 - Biodiversity
Flatback Turtle ( <i>Natator depressus</i> )	Tables 4.4 and 4.7
Freshwater Crocodile, Johnston's Crocodile, Johnston's River Crocodile ( <i>Crocodylus johnstoni</i> )	4.6.2 - Estuarine crocodiles
Green Turtle ( <i>Chelonia mydas</i> )	Tables 4.4 and 4.7
Hawksbill Turtle ( <i>Eretmochelys imbricata</i> )	Tables 4.4 and 4.7
Horned Seasnake ( <i>Acalyptophis peronii</i> )	4.3.1 - Biodiversity
Large-headed Seasnake ( <i>Hydrophis pacificus</i> )	4.3.1 - Biodiversity
Leatherback Turtle, Leathery Turtle, Luth ( <i>Dermochelys coriacea</i> )	Tables 4.4 and 4.7
Loggerhead Turtle ( <i>Caretta caretta</i> )	Tables 4.4 and 4.7
Olive Ridley Turtle, Pacific Ridley Turtle ( <i>Lepidochelys olivacea</i> )	Tables 4.4 and 4.7
Olive Seasnake ( <i>Aipysurus laevis</i> )	4.3.1 - Biodiversity
Olive-headed Seasnake ( <i>Disteira major</i> )	4.3.1 - Biodiversity
Salt-water Crocodile, Estuarine Crocodile ( <i>Crocodylus porosus</i> )	Table 4.4
Slender Seasnake ( <i>Hydrophis gracilis</i> )	4.3.1 - Biodiversity
Small-headed Seasnake ( <i>Hydrophis mcdowelli</i> )	4.3.1 - Biodiversity
Spectacled Seasnake ( <i>Disteira kingii</i> )	4.3.1 - Biodiversity
Spine-bellied Seasnake ( <i>Lapemis hardwickii</i> )	4.3.1 - Biodiversity
Spine-tailed Seasnake ( <i>Aipysurus eydouxii</i> )	4.3.1 - Biodiversity
Spotted Seasnake, Ornate Reef Seasnake ( <i>Hydrophis ornatus</i> )	4.3.1 - Biodiversity
Stokes' Seasnake ( <i>Astrotia stokesii</i> )	4.3.1 - Biodiversity
Turtle-headed Seasnake ( <i>Emydocephalus annulatus</i> )	4.3.1 - Biodiversity
Yellow-bellied Seasnake ( <i>Pelamis platurus</i> )	4.3.1 - Biodiversity
<b>Ramsar Wetlands that could be impacted by activities under the program</b>	
<p><i>Outcome: The ecological character of each Ramsar wetland is maintained, and conservation and wise and sustainable use of the wetland is promoted for the benefit of humanity in a way that is compatible with maintenance of the natural properties of the ecosystem. This is to be achieved through the implementation of ecosystem approaches, within the context of sustainable development.</i></p> <p><i>(Standards for Accreditation of Environmental Approvals under the EPBC Act 1999 – 2014)</i></p>	
Bowling Green Bay	4.8.2
Shoalwater and Corio Bays Area	4.8.1



Species/Location	GBRMPA SAR
<b>Commonwealth Heritage Places that could be impacted by activities under the program</b>	
<i>Outcome: Commonwealth Heritage places are managed to identify, protect, conserve, present and transmit, to all generations, their Commonwealth Heritage values. (Schedule 7B – EPBC Regulations)</i>	
ABC Radio Studios	Coastal Zone
Dent Island Lightstation	4.3.3 - Indigenous and Historic heritage values
Lady Elliot Island Lightstation	4.3.3 - Indigenous and Historic heritage values
Low Island and Low Islets Lightstation	4.3.3 - Indigenous and Historic heritage values
North Reef Lightstation	4.3.3 - Indigenous and Historic heritage values
Shoalwater Bay Military Training Area	5.4.7 - Defence activities
Tully Training Area	5.4.7 - Defence activities



## Appendix 7

---

### Permission system case studies

## Table of contents

<b>1</b>	<b>How the Program provides for protection of matters of national environmental significance .....</b>	<b>3</b>	4.4	Relevant Impacts .....	45
1.1	Matters of national environmental significance.....	3	4.5	Assessment of activities under the permission system .....	45
<b>2</b>	<b>Listed migratory species, listed threatened species and ecological communities .....</b>	<b>8</b>	4.6	Safeguards under the Permission System .....	47
2.1	Description .....	8	4.7	Case study - an activity with the potential to impact the Great Barrier Reef World Heritage Property, National Heritage Places, the Great Barrier Reef Marine Park and Commonwealth Marine Area .....	49
2.1.1	Listed migratory species .....	8			
2.1.2	Listed threatened species and ecological communities.....	8			
2.1.3	Migratory species, threatened species and ecological communities within the Great Barrier Reef Marine Park .....	8			
2.2	Key environmental values .....	9			
2.3	Sensitivities .....	9			
2.4	Relevant impacts.....	11			
2.5	Assessment of activities under the permission system.....	11			
2.6	Safeguards under the Program .....	13			
2.7	Case study - an activity with the potential to impact listed migratory, listed threatened species and ecological communities .....	14			
<b>3</b>	<b>Wetlands of international importance.....</b>	<b>25</b>			
3.1	Description .....	25			
3.1.1	Wetlands of international importance .....	25			
3.1.2	Ramsar wetlands .....	25			
3.2	Key environmental values .....	25			
3.3	Sensitivities .....	25			
3.4	Relevant impacts.....	26			
3.5	Assessment of activities under the permission system.....	27			
3.6	Safeguards under the Program .....	28			
3.7	Case study - an activity with the potential to impact on a Ramsar wetland.....	29			
<b>4</b>	<b>World Heritage, National Heritage, Great Barrier Reef Marine Park, Commonwealth Marine Areas .....</b>	<b>41</b>			
4.1	Description .....	41			
4.1.1	World heritage property .....	41			
4.1.2	National heritage places .....	41			
4.1.3	The Great Barrier Reef Marine Park .....	41			
4.1.4	Commonwealth marine areas.....	41			
4.2	Key environmental values .....	42			
4.3	Sensitivities .....	43			



# 1 How the Program provides for protection of matters of national environmental significance

## 1.1 Matters of national environmental significance

Part 3 of the *Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act) provides that a person must not take an action that has, will have, or is likely to have a significant impact on matters of national environmental significance without an approval under Part 9 of the Act.

Relevant matters of national environmental significance to the strategic assessment are:

- World heritage properties, including consideration of the Outstanding Universal Value of the Great Barrier Reef World Heritage Area
- National heritage places
- Wetlands of international importance
- Listed threatened species and ecological communities
- Listed migratory species
- Commonwealth marine areas
- The Great Barrier Reef Marine Park.<sup>a</sup>

The Program applies to activities requiring approval under the Great Barrier Reef Marine Park Authority's (the Authority) permission system.<sup>b</sup>

The objectives of the permission system are that all permitted activities must be carried out in a way that ensures impacts to the Marine Park, including matters of national environmental significance and the broader environment, are carried out in a manner that is consistent with the objectives of the *Great Barrier Reef Marine Park Act 1975* (GBRMP Act) and the principles of ecologically sustainable use.

The permission system ensures protection of the Great Barrier Reef Region, which necessarily includes matters of national environmental significance.

Under reg 88Q(a) of the GBRMP Regulations, the Authority must consider the potential impacts of the conduct proposed to be permitted by the permission on the environment and on the social, cultural and heritage values of the Marine Park or a part of the Marine Park.

The 'environment'<sup>c</sup> is defined as including (among other things) ecosystems and their constituent parts, including people and communities, natural and physical resources, and the qualities and characteristics of locations, places and areas. This would relevantly include the ecological character of declared Ramsar wetlands, listed threatened species and ecological communities, listed migratory species, the environment in a Commonwealth marine area and the environment in the Great Barrier Reef Marine Park.

'Heritage values'<sup>d</sup> are defined as the natural and cultural environment having aesthetic, historic, scientific or social significance. This would relevantly include the world heritage values of declared World Heritage properties and that the national heritage values of declared National Heritage places would fall within the social, cultural and heritage values of the Marine Park.

Given the relevant matters of national environmental significance fall broadly within the mandatory considerations that the Authority must consider, they would be relevant considerations. Consistent with the recommendations outlined in Chapter 12 (Table 12.2), in particular REC8A, the Authority has made a commitment in the Program Report to update, develop and implement policies and guidelines to assist permit applicants to meet permission system requirements.

Under the permission system, the Authority has committed to amending its Permit Application Form to include a section which specifically addresses matters of national environmental significance. The permit applicant must identify the matters of national environmental significance which are likely to be impacted by the proposed activity. The applicant must ensure there is sufficient information available in the permit application to enable a comprehensive and systematic evaluation of the impacts of the proposed activity on the Marine Park (including matters of national environmental significance) by the Authority. All the matters of national environmental significance identified in the Terms of Reference for the Strategic Assessment Report are accounted for under the permission system.

In accordance with the permission system, the Authority will seek further information in relation to a permit application that does not provide sufficient information to enable the Authority to assess the likely impacts to the

<sup>a</sup> Note: the broader environment of the Great Barrier Reef Marine Park is the matter protected under the Great Barrier Reef Marine Park controlling provision of Part 3 in the EPBC Act

<sup>b</sup> Activities covered by the permission system are outlined in the Program Report Section 10.5.

<sup>c</sup> 'environment' has the same meaning in the GBRMP Act (refer section 3 (1A)) as it does in the EPBC Act.

<sup>d</sup> 'heritage value' has the same meaning in the GBRMP Act (refer section 3 (1A)) as it does in the EPBC Act.

Marine Park (including matters of national environmental significance) with a high level of confidence. Recommendations in Chapter 12 of the Strategic Assessment Report and Program Report include a commitment by the Authority to develop a suite of guidelines to assist permit applicants to meet permission system requirements. Failure to provide further information within 20 business days will result in the application being taken to have been withdrawn.

The permission system addresses and makes specific commitments to ensure protection of matters of national environmental significance (refer to Part C of the Program), including through reference to statutory instruments such as the GBRMP Act and Regulations, the *Great Barrier Reef Zoning Plan 2003* (the Zoning Plan), plans of management, policies and relevant international agreements. Additional information is provided in this Appendix, for each of the relevant matters of national environmental significance, to further demonstrate how the permission system ensures the Marine Park, including matters of national environmental significance, is protected and conserved, consistent with the objects of the GBRMP Act.

Australia's relevant international treaties and obligations are primarily given effect through the EPBC Act (for example, the Ramsar Convention and the World Heritage Convention). The commitments and undertakings in the Program to matters of national environmental significance have a direct line of sight with Australia's international obligations where relevant. Under the Program the Authority will supply to the Department of the Environment the necessary information to facilitate Australia's reporting on these obligations.

Future listings of matters of national environmental significance will automatically be covered by the permission system.

Table 1 outlines the potential impacts of activities on matters of national environmental significance. A description of relevant matters of national environmental significance, including an analysis of key environmental values or 'attributes' and supporting environmental processes, is provided in Chapter 4. A detailed description of impacts, past and present, together with the risk of impacts is provided in Chapter 6 and Chapter 10 respectively.

Subsequent sections of this Appendix address each matter of national environmental significance relevant to the strategic assessment at the controlling provision level. Each section describes:

- how the matters of national environmental significance are protected under the EPBC Act, and where relevant, the international agreements underpinning those protections
- key environmental values of each matter of national environmental significance
- sensitivities of each matter of national environmental significance
- relevant impacts to matters of national environmental significance
- assessment of activities under the Program
- safeguards for the protection of matters of national environmental significance including the Authority's commitment to develop standards for determining acceptability for each matter of national environmental significance.

Hypothetical case studies demonstrate how the permission system provides for the assessment of matters of national environmental significance within the Great Barrier Reef Marine Park. The entire extent of the Marine Park lies within the Great Barrier Reef Region (refer to table 1.2 in Chapter 1 of this report). Impacts to key environmental values are used to illustrate the systems, practices, and standards which will be applied by the Authority to achieve acceptable outcomes for each matter of national environmental significance. For each hypothetical permit application made, examples are provided for how the Authority undertakes an assessment of the nature and scale of activities and makes a determination based on the acceptability of those activities on the environment. The four key steps (scoping, assessment, decision, auditing and compliance) in the permission assessment and decision process are considered for each case study. Detailed information on this process is provided in Part C of the Program Report.

Table 1 Impact of activities in the Great Barrier Reef Region on matters of national environmental significance

	Abbreviated title	Impacts	Scale	Activities in the Region covered by the Zoning Plan								Matters protected under Part 3 WH: World Heritage MP: GBR Marine Park NH: National Heritage CMA: Marine Environment MSTS: Listed migratory & listed threatened species Ramsar: Wetlands of international importance	
				Port activities	Traditional use	Tourism	Fishing — commercial	Fishing — recreational	Recreation	Shipping	Defence activities		Research activities
Catchment run-off (6.4.2)	Outbreak of crown-of-thorns starfish	Outbreak of crown-of-thorns starfish (i.e. when the density exceeds about 30 starfish per hectare)	Regional				●	●					WH, MP, NH, CMA
	Sediments from catchment run-off	Sediments entering the Region in run-off from the catchment	Reef-wide	●									WH, MP, NH, CMA, MSTS, Ramsar
	Urban and industrial discharge	Point and diffuse-source land-based discharge of pollutants from urban, commercial, agricultural and industrial land use and mining, including polluted water, sewage, wastewater and stormwater	Local	●									WH, MP, NH, CMA, MSTS, Ramsar
Degradation of coastal ecosystems (6.4.3)	Acid sulphate soils	Exposure of potential acid sulphate soils	Local	●									WH, MP, NH, CMA, MSTS, Ramsar
	Artificial barriers to flow	Artificial barriers to riverine and estuarine flow including breakwalls, weirs, dams, gates, ponded pastures, and weeds causing changes to hydrology, groundwater and ecological connectivity	Regional	●									WH, MP, NH, CMA, MSTS, Ramsar
	Atmospheric pollution	Atmospheric pollution, including coal dust	Local	●						●			WH, MP, NH, CMA, MSTS, Ramsar
	Coastal reclamation	Coastal land reclamation, including for ports and groynes	Local	●		●	●	●	●				WH, MP, NH, CMA, MSTS, Ramsar
	Light impacts (artificial)	An increased amount of artificial light	Local	●						●			MSTS
	Modifying supporting terrestrial habitats	Clearing or modifying supporting terrestrial habitats such as wetlands, saltmarshes, mangroves and sand dunes — this also includes trampling and damage from recreational vehicle use	Regional	●		●	●	●	●				MSTS, Ramsar

	Abbreviated title	Impacts	Scale	Activities in the Region covered by the Zoning Plan								Matters protected under Part 3 WH: World Heritage MP: GBR Marine Park NH: National Heritage CMA: Marine Environment MSTS: Listed migratory & listed threatened species Ramsar: Wetlands of international importance	
				Port activities	Traditional use	Tourism	Fishing — commercial	Fishing — recreational	Recreation	Shipping	Defence activities	Research activities	
Direct use (6.4.4)	Dredging	Dredging of the seafloor, including maintenance dredging of current transportation and service corridors	Local	●		●				●	●		WH, MP, NH, CMA, MSTS, Ramsar
	Dumping and resuspension of dredge material	Sea dumping of dredge material including smothering, loss and modification of seabed habitats and resuspension	Regional	●						●			WH, MP, NH, CMA, MSTS, Ramsar
	Exotic species and diseases*	Introduction of exotic species and diseases from aquaculture operations, hull fouling, ballast release, imported bait and release of aquarium specimens to the Region, plus the introduction of weeds and feral animals to islands	Regional			●	●	●	●	●			MSTS, Ramsar
	Extraction — death of discarded species	Death of non-retained species from fishing, collecting, hunting, scientific sampling and Queensland’s Shark Control Program	Regional				●	●				●	MSTS, Ramsar
	Extraction — fishing in spawning aggregations	Fishing in unprotected fish spawning aggregations	Regional				●	●					MSTS, Ramsar
	Extraction — herbivores	Retained take (extraction) of herbivores through commercial, recreational and traditional use	Reef-wide		●		●	●				●	WH, MP, NH, CMA, MSTS, Ramsar
	Extraction — lower order predators	Retained take (extraction) of lower order predators (e.g. coral trout and snapper) through commercial, recreational and traditional fishing	Reef-wide		●		●	●				●	WH, MP, NH, CMA, MSTS, Ramsar
	Extraction — lower trophic orders	Retained take (extraction) of lower trophic orders (e.g. scallops, sea cucumbers and prawns) through commercial, recreational and traditional fishing	Reef-wide		●		●	●				●	MSTS, Ramsar
	Extraction — top order predators	Retained take (extraction) of top order predators (e.g. sharks) through commercial, recreational and traditional fishing and the Queensland Shark Control Program	Reef-wide		●		●	●				●	MSTS, Ramsar
	Illegal fishing and poaching	Illegal fishing, collecting and poaching (foreign or domestic) including of species of conservation concern	Reef-wide				●	●					WH, MP, NH, CMA, MSTS, Ramsar
	Marine debris*	Manufactured material discarded, disposed of or abandoned in the marine and coastal environment (including discarded fishing gear and plastics)	Reef-wide	●		●	●	●	●			MSTS, Ramsar	





## 2 Listed migratory species, listed threatened species and ecological communities

### 2.1 Description

#### 2.1.1 Listed migratory species

Listed migratory species protected under the EPBC Act pass through or over Australian waters during their annual migrations. Examples of migratory species are shorebirds (many of which breed in the Northern hemisphere), sea birds (for example albatrosses and petrels), mammals (e.g. whales) and reptiles (e.g. sea turtles).

The list of migratory species established under Section 209 of the EPBC Act comprises:

- migratory species which are native to Australia and are included in the appendices to the Bonn Convention (Convention on the Conservation of Migratory Species of Wild Animals Appendices I and II);
- migratory species included in annexes established under the Japan-Australia Migratory Bird Agreement (JAMBA) and the China-Australia Migratory Bird Agreement (CAMBA); and native, migratory species identified in a list established under, or an instrument made under, an international agreement approved by the Environment Minister, such as the Republic of Korea-Australia Migratory Bird Agreement (ROKAMBA).

#### 2.1.2 Listed threatened species and ecological communities

Protected listed threatened species and ecological communities are listed by the Minister for the Environment under the EPBC Act.

Listed threatened species that are considered matters of national environmental significance (under Section 18 and 18A) are listed under the categories of 'extinct in the wild', 'critically endangered', 'endangered', or 'vulnerable'. The definitions of these relevant threatened categories are provided in Section 179 of the EPBC Act as follows:

- **'Extinct in the Wild'** – a species that is not extinct but is known only to survive in cultivation, in captivity or as a naturalised population well outside its past range, or it has not been recorded in its known and or expected habitat, at appropriate seasons, anywhere in its past range, despite exhaustive surveys over a time frame appropriate to its life cycle and form.
- **'Critically endangered'** – a species that is listed as critically endangered is a species that, at a particular time, is facing an extremely high risk of extinction in the wild in the immediate future as determined in accordance with the prescribed criteria in Division 7.1 of the EPBC Regulations.
- **'Endangered'** – a species that is not critically endangered and is facing a very high risk of extinction in the wild and the near future as determined in accordance with the prescribed criteria in Division 7.1 of the EPBC Regulations.
- **'Vulnerable'** – a species that is not critically endangered or endangered and is facing a high risk of extinction in the wild in the medium-term future, as determined in accordance with the prescribed criteria in Division 7.1 of the EPBC Regulations.

Listed ecological communities are listed under the categories: critically endangered, endangered or vulnerable. Ecological communities that are considered matters of national environmental significance (under Section 18 and 18A) are those that are listed as 'critically endangered', or 'endangered'. The definitions of these threatened ecological communities categories are provided in s182 of the EPBC Act as follows:

- **'Critically endangered'** – an ecological community that is facing an extremely high risk of extinction in the wild in the immediate future, as determined in accordance with the prescribed criteria in Division 7.1 of the EPBC Regulations.
- **'Endangered'** – an ecological community that is not critically endangered and is facing a very high risk of extinction in the wild in the near future as determined in accordance with the prescribed criteria in Division 7.1 of the EPBC Regulations.

#### 2.1.3 Migratory species, threatened species and ecological communities within the Great Barrier Reef Marine Park

For a list and description of threatened species subject to this Strategic Assessment and the permission system see Section 4.7. Further information on listed threatened species under the EPBC Act can be found on the Department of the Environment's website: <http://www.environment.gov.au/topics/threatened-species-ecological-communities>

There are currently no ecological communities listed under the EPBC Act which occur in the Great Barrier Reef Marine Park, see Section 4.7. Ecological communities may be listed in the future. Impacts on these would be assessed by the permission system in accordance with the Program.

For a list of migratory species subject to this Strategic Assessment and the permission system, see Section 4.6. Further information on migratory species protected under the EPBC Act can be found on the Department of Environment's website: <http://www.environment.gov.au/topics/biodiversity/migratory-species>. For a list of listed threatened species, listed ecological communities and listed migratory species subject to this Strategic Assessment and the Program, see Appendix 6.

## 2.2 Key environmental values

Habitat critical to the survival of a species or ecological community refers to areas that are necessary:

- for activities such as foraging, breeding, roosting, or dispersal
- for the long-term maintenance of the species or ecological community (including the maintenance of species essential to the survival of the species or ecological community, such as pollinators)
- to maintain genetic diversity and long term evolutionary development, or
- for the reintroduction of populations or recovery of the species or ecological community.

Such habitat may be, but is not limited to: habitat identified in a recovery plan for the species or ecological community as habitat critical for that species or ecological community; and/or habitat listed on the Register of Critical Habitat maintained by the Minister under the EPBC Act or biologically important areas, as identified through the departments regional planning..

A 'population of a species' is defined under the EPBC Act as an occurrence of the species in a particular area. In relation to critically endangered, endangered or vulnerable threatened species, occurrences include but are not limited to:

- a geographically distinct regional population, or collection of local populations, or
- a population, or collection of local populations, that occurs within a particular bioregion.

Ecological communities are unique and naturally occurring groups of plants and animals. Their presence can be determined by factors such as soil type, position in the landscape, climate and water availability.

An area of 'important habitat' for a migratory species is:

- habitat utilised by a migratory species occasionally or periodically within a region that supports an ecologically significant proportion of the population of the species
- habitat that is of critical importance to the species at particular life-cycle stages
- habitat utilised by a migratory species which is at the limit of the species range
- habitat within an area where the species is declining.

The Authority has carried out a systematic analysis of key environmental values or 'attributes' and supporting environmental processes relevant to these matters of national environmental significance. These are summarised in Table 4.8.

## 2.3 Sensitivities

Species (listed under Section 178 of the EPBC Act) and ecological communities (Section 181 of the EPBC Act) are given different categories of protection based on their levels of decline, threats and potential for recovery. This is known as their '*conservation status*'. In undertaking an assessment under the Program, the species/community's conservation status under the EPBC Act must be taken into account in determining the acceptability of the impact. For example, an activity may have a more significant impact on species in the critically endangered category than a vulnerable species.

The Authority uses EPBC Act significance guidelines in its initial screening of whether an action is likely to have a significant impact on listed threatened species and ecological communities and therefore if it requires further assessment of the acceptability of those impacts.

Examples of the different considerations which are applied to species and ecological communities by conservation status under the EPBC Act are illustrated in the following examples from the *EPBC Act Significant Impact Guidelines 1.1* outlined below:

### Extinct in the wild species

An action is likely to have a significant impact on extinct in the wild species if there is a real chance or possibility that it will:

- adversely affect a captive or propagated population or one recently introduced/reintroduced to the wild, or
- interfere with the recovery of the species or its reintroduction into the wild.

### Critically endangered or endangered species

An action is likely to have a significant impact on a critically endangered or endangered species if there is a real chance or possibility that it will:

- lead to a long-term decrease in the size of a population
- reduce the area of occupancy of the species

- fragment an existing population into two or more populations
- adversely affect habitat critical to the survival of a species
- disrupt the breeding cycle of a population
- modify, destroy, remove, isolate or decrease the availability or quality of habitat to the extent that the species is likely to decline
- result in invasive species that are harmful to a critically endangered or endangered species becoming established in the endangered or critically endangered species' habitat
- introduce disease that may cause the species to decline, or
- interfere with the recovery of the species.

### **Vulnerable species**

An action is likely to have a significant impact on a vulnerable species if there is a real chance or possibility that it will:

- lead to a long-term decrease in the size of an important population of a species
- reduce the area of occupancy of an important population
- fragment an existing important population into two or more populations
- adversely affect habitat critical to the survival of a species
- disrupt the breeding cycle of an important population
- modify, destroy, remove or isolate or decrease the availability or quality of habitat to the extent that the species is likely to decline
- result in invasive species that are harmful to a vulnerable species becoming established in the vulnerable species' habitat
- introduce disease that may cause the species to decline, or
- interfere substantially with the recovery of the species.

### **Ecological communities (critically endangered or endangered)**

An action is likely to have a significant impact on a critically endangered or endangered ecological community if there is a real chance or possibility that it will:

- reduce the extent of an ecological community
- fragment or increase fragmentation of an ecological community
- adversely affect habitat critical to the survival of an ecological community
- modify or destroy abiotic (non-living) factors (such as water, nutrients, or soil) necessary for an ecological community's survival
- cause a substantial change in the species composition of an occurrence of an ecological community, including causing a decline or loss of functionally important species, for example through flora or fauna harvesting
- cause a substantial reduction in the quality or integrity of an occurrence of an ecological community, including, but not limited to:
  - assisting invasive species, that are harmful to the listed ecological community, to become established, or
  - causing regular mobilisation of fertilisers, herbicides or other chemicals or pollutants into the ecological community which kill or inhibit the growth of species in the ecological community, or
- interfere with the recovery of an ecological community.

### **Migratory species**

An action is likely to have a significant impact on migratory species if there is a real chance or possibility that it will:

- substantially modify (including by fragmenting, altering fire regimes, altering nutrient cycles or altering hydrological cycles), destroy or isolate an area of important habitat for a migratory species
- result in an invasive species that is harmful to the migratory species becoming established in an area of important habitat for the migratory species
- seriously disrupt the lifecycle (breeding, feeding, migration or resting behaviour) of an ecologically significant proportion of the population of a migratory species.

In assessing the sensitivity of the impacts of actions on values, consideration must be given to the current condition of values. The Authority has carried out a comprehensive assessment of the condition and trend of values or 'attributes' relevant to each matters of national environmental significance in Chapter 7 of this Report. As many values were assessed to be in poor and declining condition, the Authority acknowledges that impacts deemed not to have a significant impact in the past, may be assessed to have a likely significant impact in the future. To assist consideration of whether an action is likely to have a significant impact, the Authority has committed to the development and implementation of significant impact guidelines for the Great Barrier Reef Marine Park.



Permit applications will be considered on a case by case basis, with consideration given to the objectives of the zone in which the proposed conduct will take place, as set out in the Zoning Plan.

The Authority will make decisions consistent with the objects of the GBRMP Act which includes the main object at Section 2A(1) to 'provide for the long-term protection and conservation of the environment, biodiversity and heritage values of the Great Barrier Reef Region'.

Environment has the same meaning as in the EPBC Act and is defined as including (among other things) ecosystems and their constituent parts, including people and communities, natural and physical resources, and the qualities and characteristics of locations, places and areas. This includes listed threatened species and ecological communities, and listed migratory species.

The Authority has committed in its Program to take into account the conservation status under the EPBC Act of a species/ecological community and EPBC guidance documentation in determining whether an activity is consistent with the objects of the GBRMP Act. Decisions will be made against the commitments in Table 10 of the Program.

## **2.4 Relevant impacts**

Impacts from all activities covered by the Program (e.g. those requiring a permit under the Zoning Plan as outlined in the Program Report), including those in relation to the construction, operation and (if relevant) decommissioning of a facility, are addressed by the permission system. Permitted activities have the potential to impact listed threatened species, ecological communities and migratory species through a variety of sources depending on the location and nature of the action.

Potential impacts on listed threatened species, ecological communities and migratory species include direct, indirect (upstream, downstream, facilitated) and cumulative impacts to those species, or impacts to their habitat.

Chapters 6, 7, 9 and 10 of this Report identify the key impacts to listed threatened species, ecological communities and migratory species. These chapters identify the cumulative impacts from climate-related impacts (increased sea temperature and cyclone activity), impacts that affect nesting, feeding and breeding habitats such as clearing and modifying supporting terrestrial habitats (which may also increase light and noise impacts) and direct impacts from activities that pose a threat to survival (for example, illegal fishing and poaching, death of discarded species and extraction of top order predators). Specific impact assessments of proposals must be undertaken in the context of the general level of degradation that has been identified in each habitat type within this report (See Chapter 7).

Recent severe tropical cyclones have damaged wide areas of marine habitat on which dugong and marine turtles rely. In addition, seagrass mortality as a result of floods, especially in early 2011, is thought to be responsible for severe effects on dugong populations.

Pelagic foraging seabirds are also highly vulnerable to changes in ocean currents and sea temperature as many search for food in upwellings, eddies and other sea surface temperature gradients.

Listed shorebirds are affected by the clearing and modification of coastal habitats. Previous broad scale land clearing, principally in the southern two-thirds of the Great Barrier Reef catchment, has indirectly caused major changes to the Region's near shore environment. This has, in turn, affected the shorebirds that use these habitats. In addition, changes to catchment ecosystems, such as infilling wetlands, and constructing levee banks and drainage, have affected connectivity and the habitats for listed shorebirds. Chapter 10 describes the future risk of impacts on values and attributes relevant to each matters of national environmental significance.

## **2.5 Assessment of activities under the permission system**

When assessing the impacts of an activity on a listed threatened species, ecological communities and migratory species, the permission system requires the applicant to describe the environment, identifying any listed threatened species, ecological communities and migratory species likely to be affected by the proposed activity, and provide sufficient information to understand the habitat and environment which may be affected by the proposed activity. The applicant further needs to examine how these values or elements may be affected and how impacts can be addressed and substantially reduced.

The permission system requires the Authority to analyse the potential impacts (direct, indirect and cumulative) to listed threatened species, ecological communities and migratory species or their habitats/environments and provide adequate opportunity for consultation. The Authority has committed to the development of cumulative impact assessment guidelines to provide a comprehensive, consistent and transparent approach to the assessment of potential impacts. Relevant documents that are considered when assessing the potential impacts to listed threatened species, ecological communities and migratory species include:

- Recovery Plans
- Threat Abatement Plans
- Conservation Advice
- Wildlife Conservation Plans

Relevant international obligations that the Authority is consistent with when assessing the potential impacts to listed threatened/migratory species and ecological communities include:

- the Convention on Biological Diversity
- the Apia Convention
- Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES)
- the Bonn Convention
- CAMBA
- JAMBA
- an international agreement approved under ss209(4) of the EPBC Act.

### **Recovery Plans**

The Minister for the Environment may make or adopt and implement recovery plans for threatened fauna, threatened flora (other than conservation dependent species) and threatened ecological communities listed under the EPBC Act. Recovery plans set out the research and management actions necessary to stop the decline of, and support the recovery of, listed threatened species or threatened ecological communities. The aim of a recovery plan is to maximise the long term survival in the wild of a threatened species or ecological community.

Recovery plans should state what must be done to protect and restore important populations of threatened species and habitat, as well as how to manage and reduce threatening processes. Recovery plans achieve this aim by providing a planned and logical framework for key interest groups and responsible government agencies to coordinate their work to improve the plight of threatened species and/or ecological communities.

### **Threat Abatement Plans**

The EPBC Act provides for the identification and listing of key threatening processes. Key threatening processes threaten or may threaten the survival, abundance or evolutionary development of a native species or ecological community. For example, invasive species associated with listed key threatening processes are predation by the European red fox, feral rabbits or unmanaged goats.

The assessment of a threatening process as a key threatening process is the first step to addressing the impact of a particular threat under Commonwealth law. The Australian Government Minister for the Environment may decide whether to have a threat abatement plan for a threatening process in the list of key threatening processes established under the EPBC Act.

Threat abatement plans provide for the research, management, and any other actions necessary to reduce the impact of a listed key threatening process on native species and ecological communities. Implementing the plan should assist the long term survival in the wild of affected native species or ecological communities. Threat abatement plans contain objectives and actions that relate to mitigating or reversing the impacts of a key threatening process.

### **Conservation Advice**

When a native species or ecological community is listed as threatened under the EPBC Act, conservation advice is developed to assist its recovery. Conservation advice provides guidance on immediate recovery and threat abatement activities that can be undertaken to ensure the conservation of a newly listed species or ecological community.

Conservation advice includes practical on-ground activities that can be implemented by local communities, natural resource management groups or interested individuals, such as landholders.

Conservation advice may also include broader management actions that can be undertaken by organisations such as local councils, government agencies or non-government organisations, to protect the threatened species or ecological community on a regional level.

### Wildlife conservation plans

The Minister for the Environment may make a wildlife conservation plan for the purposes of the protection, conservation and management of the following:

- (a) a listed migratory species that occurs in Australia or an external Territory;
- (b) a listed marine species that occurs in Australia or an external Territory;
- (c) a species of cetacean that occurs in the Australian Whale Sanctuary;
- (d) a conservation dependent species.

A wildlife conservation plan must provide for the research and management actions necessary to support survival of the migratory species, marine species, species of cetacean or conservation dependent species concerned. Plans may cover one or more species.

Chapter 10 describes the future risk of impacts on values and attributes relevant to each matters of national environmental significance.

## 2.6 Safeguards under the Program

Under the EPBC Act, an action or class of actions will not be approved if it would be inconsistent with:

- Australia's obligations under:
  - The Convention on Biological Diversity
  - the Apia Convention
  - Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES)
  - the Bonn Convention
  - CAMBA
  - JAMBA
  - an international agreement approved under ss209(4) of the EPBC Act, or
- a recovery plan for the species or a threat abatement plan.

Assessment of an action must also have regard to any approved conservation advice for the species or community.

The Authority is committed to the survival and conservation status of listed threatened species, ecological communities and migratory species being promoted and enhanced through the conservation of critical habitats and other relevant measures contained in relevant plans or advices.

The Authority will make decisions consistent with the objects of the GBRMP Act which includes the main object at section 2A(1) to 'provide for the long-term protection and conservation of the environment, biodiversity and heritage values of the Great Barrier Reef Region'. Environment has the same meaning as in the EPBC Act and is defined as including (among other things) ecosystems and their constituent parts, including people and communities, natural and physical resources, and the qualities and characteristics of locations, places and areas. This necessarily includes matters of national environmental significance.

In making decisions that are consistent with the objects of the GBRMP Act, the Authority must consider the potential impacts of the conduct proposed to be permitted by the permission on the environment and on the social, cultural and heritage values of the Marine Park or a part of the Marine Park.

Consideration of any relevant recovery plan, threat abatement plan, wildlife conservation plan or conservation advice relevant to the management of impacts of an activity is a requirement of the permission system. Permit application assessments also have regard to the significant impact criteria in the *EPBC Act Significant Impact Guidelines 1.1* for listed threatened species, ecological communities and migratory species that may or are likely to be impacted by the activity in the context of demonstrating consistency with the objects of the GBRMP Act. The Authority's consideration of the acceptability of an activity will also be informed by the permit applicant's avoid-mitigate-offset-net benefit-adaptive management approach as proposed in their application and subsequent assessment report(s).

Under the permission system the Authority will make decisions consistent with the GBRMP Act:

- Consideration will be given to:
  - the preservation of a listed threatened species, ecological community a migratory species or an area of important habitat for a migratory species
  - the following conventions and agreements:
    - The Bonn Convention
    - China Australia Migratory Birds Agreement (CAMBA)
    - Japan Australia Migratory Birds Agreement (JAMBA)

- Republic of Korea Australia Migratory Birds Agreement (ROKAMBA)
- An international agreement approved under Subsection 209(4) of the EPBC Act.
- Australia's obligations under the Biodiversity Convention, the Apia convention and CITES.
- a recovery plan or threat abatement plan for a listed threatened species, ecological community or migratory species.
- The Authority maintains policy guidance that applicants should have regard to in the preparation of their Permit Applications. Policy guidance will be updated to include explicit reference to:
  - consideration of all attributes relevant to matters of national environmental significance
  - consideration of the listing category and protection of the listed threatened species, ecological community or migratory species
  - consideration of the protection of an area of important habitat for a listed migratory species
  - relevant GBRMP Act and EPBC Act guidance documents, including but not limited to, recovery plans, threat abatement plans, conservation advice and wildlife conservation plans.

In undertaking assessments, the Authority has regard to relevant policy documents, recovery plans, threat abatement plans, conservation advice, wildlife conservation plans and guidelines and plans of management on the Department of the Environment website.

The Authority's assessment process also ensures an assessment of the proponent's measures to avoid, mitigate, offset and provide for net-benefit and adaptive management of potential impacts to threatened species, ecological communities and migratory species.

All permitted activities must be carried out in a manner that is consistent with the objectives of the GBRMP Act, ecosystem-based management and the principles of ecologically sustainable use.

## 2.7 Case study - an activity with the potential to impact listed migratory, listed threatened species and ecological communities

**Scenario:** A drilling and blasting operation for the expansion of a shipping lane will be undertaken in close proximity to loggerhead and green turtles and associated foraging and nesting habitats.

A hypothetical<sup>5</sup> seagrass system ecological community, listed as endangered, is also potentially impacted.

Key impacts of the activity include:

- marine debris
- noise and light pollution
- physical damage, ship grounding, vessel strike
- waste discharges and spills
- species displacement from key ecological functions (foraging, resting, breeding)
- potential introduction of exotic species through hull fouling
- permanent and temporary loss of habitat
- underwater habitat smothering
- water quality impacts, primarily changes in sediment loads.

These impacts, their cumulative effects, and past and present effects on values, including on matters of national environmental significance, are described in detail in Chapter 6.

- The drilling and blasting activity has the potential to have a significant impact on one or more listed threatened/migratory species and the seagrass ecological community. The Authority has determined it will be subject to an Environmental Impact Statement (EIS) assessment in accordance with the Authority's permit assessment policies.
- The species at risk have the following threatened status. They are also migratory species:
  - Loggerhead turtle (*Caretta caretta*) listed as endangered under the EPBC Act, listed migratory species (Bonn Convention)
  - Green turtle (*Chelonia mydas*) listed as vulnerable under the EPBC Act, listed migratory species (Bonn Convention).
- The seagrass ecological community is listed as endangered under the EPBC Act.

<sup>5</sup> There are no EPBC listed ecological communities in the GBRMP. A hypothetical ecological community is therefore used to illustrate the efficacy of the Program to protect these.



A Public Environment Report (PER) is an alternative assessment option under the Program. Both EIS and PER assessments require an Environmental Management Plan (EMP) to be prepared and assessed which describes how control measures are implemented and managed on site.

A drilling and blasting operation is an activity classified as 'carrying out works' which requires permission (a permit) under the Great Barrier Reef Zoning Plan. As such, this activity would require a permit application to be submitted, assessed and issued to determine whether the potential impacts of the project on the broader environment, threatened and migratory species and the listed ecological community will be consistent with the objects of the GBRMP Act.

The Authority's assessment process requires the permit application to describe the environment and identify any listed threatened/migratory species and listed ecological communities that are likely to be affected by the proposed activity. The application should contain a summary of the potential impacts (direct, indirect and cumulative) of all proposed activities. It must provide sufficient information to understand the habitat which may be affected by the proposed activity.

The permission system requires the Authority to analyse the potential impacts of the proposed activity (direct, indirect and cumulative) on listed threatened/migratory species and listed ecological communities, their habitat and ensure impacts are acceptable. The types of studies and fields of investigation which would usually occur in this scenario include: investigation into underwater noise effects on the species ecology, assessment of nesting periods and the behavioural responses to noise, light pollution and habitat removal and water quality impacts on seagrass habitats.

**Table 2 How a permit application would meet EPBC Act requirements for listed migratory species, listed threatened species and listed ecological communities.**

Outcome from EPBC Act	How listed migratory species, listed threatened species and ecological communities are considered under the Authority's permission system
<p><b>SCOPING:</b> The purpose of the scoping phase is to proactively discuss the initial concepts of a proposed action in order to explain regulatory arrangements and identify potential mitigations. For actions which will likely have a significant impact on the Marine Park, including matters of national environmental significance, reference is made to relevant GBRMP Act and EPBC Act guidance material. A Permit Application Assessment Fee must be paid before the accompanying Permit Application Form is considered. The Permit Application Form must contain sufficient information to enable a clear understanding of the intended action.</p>	
<p><b>Activities with a likely significant impact must be assessed under legislation with regards to matters of national environmental significance</b></p>	<p><b>Mechanism (legislative &amp; policy):</b></p> <p>Section 268 of the EPBC Act requires that a Commonwealth agency must not take any action that contravenes a recovery plan or a threat abatement plan. As a Commonwealth statutory authority, the Authority must comply with this requirement.</p> <p>Section 286 of the EPBC Act requires that a Commonwealth agency must take all reasonable steps to act in accordance with a wildlife conservation plan. As a Commonwealth statutory authority, the Authority must comply with this requirement. GBRMP Regulations 88Q (mandatory considerations) and 88R (discretionary considerations) outline the Authority's permit assessment considerations.</p> <p>Under Regulation 88Q(a) of the GBRMP Regulations, the Authority must consider the potential impacts of the conduct proposed to be permitted by the permission on the environment and on the social, cultural and heritage values of the Marine Park or a part of the Marine Park. This relevantly includes matters of national environmental significance.</p> <p>Under Regulation 88E the Authority would seek further information for a permit application that does not provide sufficient detail to evaluate whether an activity is having a significant impact. This includes:</p> <ul style="list-style-type: none"> <li>• a description of the relevant matters of national environmental significance attributes likely to be affected by the proposed activity</li> <li>• the direct, indirect and cumulative impacts of the proposed activity</li> <li>• the risks of those impacts to matters of national environmental significance.</li> </ul> <p>Failure to provide sufficient detail would result in the application being withdrawn.</p> <p><b>Program commitment(s):</b></p> <p>The Authority will:</p> <ul style="list-style-type: none"> <li>• require consideration of relevant EPBC policy documents, including relevant significant impact guidelines and reporting tools, in the scoping phase of the permission assessment and decision process.</li> </ul>

Outcome from EPBC Act	How listed migratory species, listed threatened species and ecological communities are considered under the Authority's permission system
	<ul style="list-style-type: none"> <li>• require consideration of prudent and feasible alternatives which may have a lower impact on the environment and amend its policy documentation to include reference to this requirement.</li> <li>• consider any relevant international conventions, Commonwealth State agreements, plans of management, recovery plans, wildlife conservation plan, threat abatement plans, approved conservation advice under the EPBC Act, EPBC Act guidelines, policies and information in assessing and determining permissions under its permission system.</li> <li>• consider relevant Department of the Environment policy documents, guidelines, plans of management and other online data sources available on the Department's website in its scoping, assessment and decision-making process.</li> <li>• implement guidelines to assist applicants to determine whether an action is likely to have a significant impact on the Great Barrier Reef Marine Park matters of national environmental significance. The Authority will seek to have these guidelines approved by the Minister.</li> <li>• amend policies and guidance material used in the determination of permissions so they explicitly require the consideration of matters of national environmental significance and relevant attributes, including: <ul style="list-style-type: none"> <li>○ biodiversity and environmental processes</li> <li>○ Indigenous and historic heritage values</li> <li>○ community benefits derived from the environment, including those not easily represented or measured (such as aesthetic values).</li> </ul> </li> <li>• develop guidance material to assist in determining the acceptability of impacts. Criteria shall include consistency of assessment outcomes with regulatory objectives, outcomes for the Marine Park, including matters of national environmental significance and relevant GBRMP Act and EPBC Act guidelines and standards.</li> <li>• develop specific guidance documents for notifications under Part 5 of the Zoning Plan to make reference to consideration of matters of national environmental significance under the EPBC Act.</li> </ul> <p>The Authority will require applicants to consider:</p> <ul style="list-style-type: none"> <li>• the impacts of actions on matters of national environmental significance and relevant attributes and environmental processes as described in the Strategic Assessment Report, and as amended in accordance with the Authority's policy review procedures.</li> <li>• impacts (including direct, indirect and cumulative impacts) arising from the development action in the context of all impacts affecting the matters of national environmental significance.</li> </ul> <p><b>Case study:</b></p> <ul style="list-style-type: none"> <li>• Under the Zoning Plan a drill and blast operation is considered to be 'carrying out works'. This type of activity can only occur in certain parts of the Marine Park – namely areas where the purpose of the activity is consistent with the objective of the relevant zone and only with the written permission of the Authority.</li> <li>• The Authority determines that proposed activity is consistent with the objectives of the relevant zone and that the proponent requires a permit from the Authority for the carrying out of works.</li> <li>• For actions which may have a significant impact on the Marine Park, including matters of national environmental significance, reference is made to: <ul style="list-style-type: none"> <li>○ GBRMP Act, Regulations, Zoning Plan, plans of management and relevant policy documents including the Authority's environmental impact management, position statements and guidelines;</li> <li>○ EPBC Act policy documents, including relevant significant impact guidelines, the EPBC Act Species Profile and Threats Database (SPRAT), listing advice policy statements, recovery plans, conservation advices, threat abatement plans and wildlife conservation plans available on the Department of the Environment website.</li> </ul> </li> <li>• Information available through online data sources hosted by the Department of the Environment, such as the Environment Reporting Tool, augments local knowledge in determining whether matters of national environmental significance are likely to occur in the area of interest.</li> <li>• EPBC Act Significant Impact Guidelines 1.1, as they relate to listed species</li> </ul>

Outcome from EPBC Act	How listed migratory species, listed threatened species and ecological communities are considered under the Authority's permission system
	<p>and ecological communities include consideration of the conservation status under the EPBC Act of the species and community. The Authority, in evaluating whether the activity is consistent with the objects of the GBRMP Act, will take into account either EPBC Act species specific significance guidelines or, where none exist, the following generic considerations from the EPBC Act Significant Impact Guidelines 1.1. An action is likely to have significant impact on critically endangered, endangered or vulnerable species or ecological communities if there is a real chance or possibility that it will:</p> <ul style="list-style-type: none"> <li>○ lead to a long-term decrease in the size of a population</li> <li>○ reduce the area of occupancy for the species or extent of the ecological community</li> <li>○ fragment an existing population or ecological community into two or more populations</li> <li>○ reduce the extent of an ecological community</li> <li>○ adversely affect habitat critical to the survival of a species or ecological community</li> <li>○ disrupt the breeding cycle of a population</li> <li>○ modify, destroy, remove, isolate or decrease the availability or quality of habitat to the extent that the species or ecological community is likely to decline</li> <li>○ result in invasive species that are harmful to a critically endangered or endangered species becoming established in their habitats</li> <li>○ introduce disease that may cause species to decline</li> <li>○ interfere with the recovery of the species</li> <li>○ cause a substantial change in the species composition of an occurrence of an ecological community, including causing a decline or loss of functionally important species, for example through regular burning or flora or fauna harvesting</li> <li>○ modify or destroy abiotic (non-living) factors (such as water, nutrients, or soil) necessary for an ecological community's survival, including reduction of groundwater levels, or substantial alteration of surface water drainage patterns</li> <li>○ cause a substantial reduction in the quality or integrity of an occurrence of an ecological community, including, but not limited to: <ul style="list-style-type: none"> <li>▪ assisting invasive species, that are harmful to the listed ecological community, to become established, or</li> <li>▪ causing regular mobilisation of fertilisers, herbicides or other chemicals or pollutants into the ecological community which kill or inhibit the growth of species in the ecological community, or</li> </ul> </li> <li>○ interfere with the recovery of an ecological community.</li> </ul> <p>It is likely to have a significant impact on migratory species if there is a real chance or possibility that it will:</p> <ul style="list-style-type: none"> <li>○ substantially modify (including by fragmenting, altering nutrient cycles or altering hydrological cycles), destroy or isolate an area of important habitat for a migratory species</li> <li>○ result in an invasive species that is harmful to the migratory species becoming established in an area of important habitat for the migratory species</li> <li>○ seriously disrupt the lifecycle (breeding, feeding, migration or resting behaviour) of an ecologically significant proportion of the population of a migratory species</li> </ul> <ul style="list-style-type: none"> <li>• Information available through online data sources hosted by the Department of the Environment, such as the Environment Reporting Tool, augments local knowledge in determining whether matters of national environmental significance are likely to occur in the area of interest.</li> <li>• The Authority seeks the advice of the Queensland Parks and Wildlife Service and other relevant agencies with an administrative responsibility relating to the proposed activity, on the potential impacts of the proposed activity and level of assessment.</li> <li>• Under this scenario, the Authority determines that the proposed activity is likely to have a significant impact on the green and loggerhead turtles and the listed seagrass ecological community and requires assessment by an</li> </ul>

Outcome from EPBC Act	How listed migratory species, listed threatened species and ecological communities are considered under the Authority's permission system
	Environmental Impact Statement (EIS) which also requires an Environmental Management Plan (EMP) to be prepared.
<p><b>Applicant is required to provide adequate information to enable the decision maker to make a decision.</b></p> <ul style="list-style-type: none"> <li>- <b>The applicant describes impacts to matters of national environmental significance</b></li> </ul>	<p><b><i>Mechanism (legislative &amp; policy):</i></b></p> <p>GBRMP Regulation 88A enables the Authority to approve permit application forms. The Authority can vary the approved application form to meet legal and management responsibilities.</p> <p>GBRMP Regulation 88E enables the Authority to request further information, for the purpose of assessing an application, and for such information to be provided within 20 business days (or for a longer period, as determined by the Authority) or the application is taken to have been withdrawn.</p> <p>GBRMP Regulation 103 provides that persons shall not furnish false information in a permit application.</p> <p>Failure to provide sufficient information to describe the specific threatened/migratory species, habitats, important populations, or the ecological community potentially at risk from the activity in detail sufficient to evaluate all impacts to assess whether the activity is consistent with the objects of the GBRMP Act and to determine the level of assessment under the Program (e.g. by EIS) would result in the application being withdrawn.</p> <p><b><i>Program commitment(s):</i></b></p> <p>The Authority will:</p> <ul style="list-style-type: none"> <li>• amend its Permit Application Form to include a section which specifically addresses matters of national environmental significance.</li> <li>• seek to pursue amendments to the Great Barrier Reef Marine Park Regulations 1983 to explicitly outline the levels of assessment under the permission system.</li> <li>• update its permissions guidance material to outline the general information requirements for applicants according to the level of assessment.</li> </ul> <p><b><i>Case study:</i></b></p> <ul style="list-style-type: none"> <li>• The applicant will have access to guidance material developed by the Authority relating to the Marine Park, including matters of national environmental significance, that proponent's should have regard to in the preparation of their application.</li> <li>• An application must be submitted to the Authority. It must provide sufficient information on the drilling and blasting so assessors can clearly understand what the intended activity is, how it is to be carried out, where it will occur, timing and the likely nature and scale of environmental impacts.</li> <li>• Descriptions must identify any threatened/migratory species and ecological communities; including a sufficient description of the turtles and seagrass community which may be at risk from the activity. The application must provide sufficient information for the Authority to identify potentially impacted turtle habitats, migratory populations and the significance of the area for each species/population potentially affected. Information should be included on the context of the seagrass ecological community, such as the range, condition and extent of the community.</li> <li>• This will include a description of the occurrence of these species/populations within the project area and within any areas (zone of influence) that may potentially be affected by upstream and downstream impacts such as water quality or noise effects.</li> <li>• The description of the activity and receiving environment's characteristics must be sufficient to inform consideration of the 'nature and scale' of the impacts by the Authority in its assessment. In this case, any elements of an activity relevant to impacts on marine turtles and the seagrass community must be sufficiently described including operational details relevant to the listed turtle species and seagrass community e.g.: <ul style="list-style-type: none"> <li>○ The timing and duration of the activity in relation to the use of known habitat, including biologically important areas/populations (for example the breeding, hatching or migration periods for sea turtles). The application will require information on all impacts on the species and community including but not limited to: <ul style="list-style-type: none"> <li>▪ duration, timing and magnitude of drilling and blasting activities</li> <li>▪ details of other disturbances such as light, increased vessel movements, permanent and or temporary placement of objects in</li> </ul> </li> </ul> </li> </ul>



Outcome from EPBC Act	How listed migratory species, listed threatened species and ecological communities are considered under the Authority's permission system
	<p>turtle habitats</p> <ul style="list-style-type: none"> <li>▪ quantification of any habitat loss</li> <li>▪ quantification and duration of water quality impacts</li> <li>▪ discharges or emissions</li> <li>▪ extent and duration of disturbance to seagrass meadows</li> <li>▪ and any other impacts or activities that have the potential to adversely impact directly on: <ul style="list-style-type: none"> <li>• individuals,</li> <li>• important populations or their habitats</li> <li>• seagrass extent or composition.</li> </ul> </li> <li>• The application must demonstrate that appropriate contingency plans, including appropriate emergency response measures are in place to ensure that the applicant would be able to effectively respond to an unplanned oil spill or discharge event. These plans would form part of the Environment Management Plan required for this activity to proceed (see Monitoring Compliance and Auditing below).</li> <li>• If an application does not include sufficient information for the purposes of assessing the likely nature and scale of environmental impacts additional information or documents must be provided at the request of the Authority.</li> </ul>
<p><b>ASSESSMENT:</b> The purpose of the assessment phase is to enable an informed decision on whether or not a permit will be granted to undertake an action within the Great Barrier Reef Marine Park. During the assessment phase, the Authority considers criteria outlined in the Great Barrier Reef Marine Park Regulations 1983, and relevant GBRMP Act and EPBC Act guidance material.</p>	
<p><b>Public consultation occurs in a transparent manner and outcomes are taken into account</b></p>	<p><b>Mechanism (legislative &amp; policy):</b></p> <p>GBRMP Regulations 88Q (mandatory considerations) and 88R (discretionary considerations) outline the Authority's permit assessment considerations. Under Regulation 88E the Authority would seek further information for a permit application that does not provide sufficient detail to evaluate whether an activity is having a significant impact. This includes:</p> <ul style="list-style-type: none"> <li>• a description of the relevant matters of national environmental significance attributes likely to be affected by the proposed activity</li> <li>• the direct, indirect and cumulative impacts of the proposed activity</li> <li>• the risks of those impacts to matters of national environmental significance.</li> </ul> <p>Regulation 88D of the GBRMP Regulations requires applications to be advertised if the Authority considers that granting a permission may restrict the reasonable use by the public of part of the Great Barrier Reef Marine Park.</p> <p>GBRMP Regulations 88D provide that if a permit applicant does not publish an advertisement as required, the application is taken to have been withdrawn.</p> <p>GBRMP Regulation 88Q provides that public comments must be considered by the Authority when making decisions on whether to grant the permit.</p> <p><b>Program commitment(s):</b></p> <p>The Authority will:</p> <ul style="list-style-type: none"> <li>• publish notification of all permit applications on the internet as soon as practicable after receiving a permit application.</li> <li>• seek to pursue amendments to the GBRMP Regulations (in particular regulation 88D) to require advertising of applications where the granting of a permission may restrict the reasonable use by the public of a part of the Marine Park or is likely to have a significant impact on the Marine Park, including on matters of national environmental significance.</li> <li>• update relevant policies to explicitly state that appropriate consultation will be undertaken with persons or organisations, including Commonwealth and State government agencies, whose functions, interests or activities could be impacted by the proposed activity.</li> </ul> <p><b>Case study:</b></p> <ul style="list-style-type: none"> <li>• The application is assessed by the permit assessor/project manager who determines if species and/or seagrass specific expert advice is required. In this case it is standard practice for it to be referred to relevant parties and experts for comment. Expert advice may be sought from the Australian Government Department of the Environment if required.</li> </ul>

Outcome from EPBC Act	How listed migratory species, listed threatened species and ecological communities are considered under the Authority's permission system
	<ul style="list-style-type: none"> <li>• The Authority also undertakes appropriate consultation with persons or organisations, including Australian and State government agencies, whose functions, interests or activities could be impacted by the proposed activity.</li> <li>• The application, including the EIS, is made available for a period no less than 30 business days for public comment via the Authority's website.</li> <li>• The applicant is required to publish an advertisement for projects under assessment and for the EIS to be made available to the public during a defined public comment period. This advertisement must be published in a newspaper circulating in the part of the State of Queensland adjacent to that part of the Marine Park in which the conduct is proposed to occur. The advertisement is to include an invitation for interested persons to make comment on the application within a period of not less than 30 days. The advertisement will also be published on the Authority's website.</li> </ul>
<p><b>The detail and rigour of the assessment process is commensurate with the scale and potential impacts of the activity i.e. risk-based.</b></p>	<p><b><i>Mechanism (legislative &amp; policy):</i></b></p> <p>Section 268 of the EPBC Act requires that a Commonwealth agency must not take any action that contravenes a recovery plan or a threat abatement plan, as a Commonwealth Statutory Authority, the Authority must comply with this requirement.</p> <p>Section 286 of the EPBC Act requires that a Commonwealth agency must take all reasonable steps to act in accordance with a wildlife conservation plan. As a Commonwealth statutory authority, the Authority must comply with this requirement.</p> <p>GBRMP Regulations 88Q (mandatory considerations) and 88R (discretionary considerations) outline the Authority's permit assessment considerations.</p> <p>The Authority's Environmental Impact Assessment Policy and Environmental Assessment and Management Risk Assessment Guidelines (developed in accord with the Australian Standards Environmental Risk Management Handbook HB203:2006) provide guidance in the assessment of risk including analysis of the likelihood and consequence of identified hazards being realised to obtain a Hazard Risk Grade.</p> <p><b><i>Program commitment(s):</i></b></p> <p>The Authority will:</p> <ul style="list-style-type: none"> <li>• develop and maintain regionally-based standards for ecosystem health (including water quality) that support best management practices, and may be used to inform: <ul style="list-style-type: none"> <li>○ the assessment of point source and cumulative impacts on the Marine Park, including matters of national environmental significance</li> <li>○ actions to mitigate impacts and cumulative impacts on the Marine Park, including matters of national environmental significance</li> <li>○ outcomes-based decision-making and considerations for determining 'acceptability' of impacts on the Marine Park, including matters of national environmental significance, based the commitments in Table 10 of the Program Report.</li> </ul> </li> <li>• implement guidelines requiring consideration of regionally-based standards for ecosystem health to improve the assessment of impacts including cumulative impacts on the Region's values.</li> <li>• facilitate development of a dredging and dredge material disposal policy</li> <li>• improve the hydrodynamic modelling guidelines so they take account of longshore drift, resuspension and inter-annual current variability.</li> <li>• develop guidance material to be used in the determination of permissions to require the consideration of the impacts of underwater noise, particularly on at-risk species and inshore species.</li> <li>• integrate relevant components, as required, of the Department of the Environment's EPBC Act assessment templates and manuals into its policy documents as they are reviewed and updated.</li> </ul> <p><b><i>Case study:</i></b></p> <ul style="list-style-type: none"> <li>• The assessment must clearly demonstrate the risks of impacts arising from the activity impacting directly and indirectly and cumulatively on turtle species and the ecological community.</li> <li>• The Authority undertakes a risk assessment in permit assessments demonstrating that the impacts of the blasting and drilling activity on will be of an acceptable level at all times during the conduct of the activity. The</li> </ul>

Outcome from EPBC Act	How listed migratory species, listed threatened species and ecological communities are considered under the Authority's permission system
	<p>assessment must demonstrate that all risks are acceptable. In this case, the Authority's assessment will consider:</p> <ul style="list-style-type: none"> <li>○ habitat types, seasonality, important populations and whether there are critical life stages such as feeding, breeding, hatchling recruitment or migration that are likely to be affected by artificial lighting and any other potential impacts arising from the activity,</li> <li>○ the extent of the seagrass ecological community the potential for its fragmentation, or impacts on factors necessary for its survival, e.g. species composition) arising from the activity. Assessment will consider the potential for introduction of invasive species, pollutants, smothering and whether the activity will interfere with the seagrass community's recovery.</li> </ul> <ul style="list-style-type: none"> <li>• The Authority's assessment must consider any relevant information from recovery plans, conservation advices, wildlife conservation plans or threat abatement plans under the EPBC Act. Assessments must take into account the conservation status under the EPBC Act of each species and ecological community that may be impacted, the principal threats and priority actions as outlined in the relevant plans and analysis of the impacts of drilling and blasting against the requirements of these plans.</li> <li>• The assessment will also specifically address the likelihood of occurrence of unplanned events and demonstrate how they are proposed to be managed to acceptable levels.</li> <li>• Levels of acceptability will be based on the commitments in Table 10 of the Program Report, which draw on the EPBC Act's standards. Acceptability considerations must also take into account the different conservation status under the EPBC Act of the threatened/migratory species. For example: <ul style="list-style-type: none"> <li>○ The acceptable level of impact on a habitat for the endangered Loggerhead turtle with limited range and no alternative habitats may differ from the Green turtle which is listed as vulnerable and may not be as susceptible to impacts from the activity at the specific location described in the permit application. Consideration in this case would depend on whether there is an important population of the Green turtle in the impact area and the nature and scale of impacts on that population.</li> <li>○ The acceptable level of impact on the migratory values of the listed turtle species will depend on the extent to which the area supports an ecologically significant proportion of the population of the species and is an important habitat for the species regarding life cycle, is at the limit of the species range, or is within an area where the turtles are declining.</li> <li>○ The Authority will also have regard to relevant policies and guidelines such as the EPBC Act Significant Impact Guidelines 1.1<sup>6</sup>, offsets and cumulative impact guidelines where appropriate.</li> </ul> </li> <li>• The Authority must demonstrate that impacts have been reduced to acceptable levels including when measured against principal threats identified in any relevant recovery plan conservation advice, wildlife conservation plan or threat abatement plan.</li> <li>• The Authority's assessment must contain any relevant legislative or other controls that are necessary for managing the impacts to the threatened/migratory species and ecological communities (such as recovery actions) and their associated habitats. In establishing performance outcomes and standards the Authority must consider any outcomes, and targets and performance criteria in relevant statutory recovery plans for the marine turtles.</li> <li>• The legislative requirements of the Program and information published on the Department of the Environment website form a comprehensive suite to ensure that the Authority will not permit an activity that will have an unacceptable impact on a listed threatened species</li> <li>• The Authority would refuse a permit application for an activity that was not consistent with the objects of the GBRMP Act, for example, where there would be unacceptable impacts on the Marine Park, including on listed threatened/migratory species and ecological communities.</li> </ul>
<p><b>EIS process follows 'avoid/mitigate/offset/enhance principles'. Any residual significant impacts are offset appropriately.</b></p>	<p><b>Mechanism (legislative &amp; policy):</b></p> <p>GBRMP Regulation 88Q(b) requires that the Authority must consider options for monitoring, managing and mitigating the potential impacts of the proposed conduct.</p>

<sup>6</sup>Commonwealth of Australia (2013). Ibid.

Outcome from EPBC Act	How listed migratory species, listed threatened species and ecological communities are considered under the Authority's permission system
	<p data-bbox="571 232 1410 315">GBRMP Regulation 88X -The Authority must, after taking into account the matters that it is required or permitted to take into account under the GBRMP Act and Regulations, make a decision on the application.</p> <p data-bbox="571 342 847 371"><b>Program commitment(s):</b></p> <p data-bbox="571 376 754 405">The Authority will:</p> <ul data-bbox="592 409 1410 779" style="list-style-type: none"> <li>• implement guidelines for the application of Great Barrier Reef offsets to maintain the condition of the Marine Park, including matters of national environmental significance and relevant attributes and environmental processes, where impacts cannot be avoided or mitigated: <ul data-bbox="628 524 1369 607" style="list-style-type: none"> <li>◦ The guidelines will seek to deliver an outcome equivalent to, or better than, the outcome that would apply if the EPBC Act Environmental Offsets Policy were applied.</li> </ul> </li> <li>• implement a Great Barrier Reef net benefit policy to enhance the condition of the Marine Park, including matters of national environmental significance and relevant attributes and environmental processes.</li> <li>• implement guidelines on cumulative impact assessment to inform a transparent, consistent and systematic approach to the assessment of impacts from activities within and adjacent to the Region.</li> </ul> <p data-bbox="571 806 703 835"><b>Case study:</b></p> <ul data-bbox="592 840 1410 1749" style="list-style-type: none"> <li>• The assessment of the permit application uses an 'avoid, mitigate, offset, net-benefit and adaptive management methodology' of controls</li> <li>• The Authority uses a risk assessment to evaluate the efficacy of controls to avoid, mitigate, offset, provide net-benefit and adaptive management of potential impacts on threatened/migratory species and ecological communities. The assessment ensures that vessel transit areas avoid known aggregations of species, appropriate lookouts are maintained and ensure that blasting/drilling do not affect species and artificial light impacts do not extend to nesting beaches for marine turtles. Impacts on the extent of the seagrass ecological community and any threats to its recovery are also considered.</li> <li>• Enhancement of conditions for species and communities is also a consideration of the risk assessment process.</li> <li>• The Authority's assessment documentation must demonstrate that an assessment of the appropriateness of avoidance, mitigation, offset, net-benefit and adaptive management measures are in place to reduce impacts on species and communities in the Marine Park. The assessment must demonstrate that impacts on the listed species will be acceptable and are not inconsistent with any priority recovery actions identified in relevant recovery plans, conservation advices or wildlife conservation plans.</li> <li>• The assessment must ensure that the identified avoidance, mitigation, offset, net-benefit and adaptive management measures, such as operational monitoring/environmental site supervisors, Deeds of Agreement and bonds are appropriate for managing impacts.</li> <li>• The proponent must demonstrate in their assessment documentation that impacts have been reduced to levels where ecological character will be maintained and enhanced. Considerations must include consistency of assessment outcomes with: <ul data-bbox="628 1621 1369 1749" style="list-style-type: none"> <li>◦ Regulatory objectives</li> <li>◦ Outcomes for matters of national environmental significance including relevant attributes and environmental processes, and</li> <li>◦ Relevant GBRMP Act and EPBC Act guidelines and standards.</li> </ul> </li> </ul>
	<p data-bbox="148 1778 1410 1883"><b>DECISION:</b> The purpose of the decision phase is to grant or refuse a permission based on information provided within the assessment phase. A permission may be granted with any conditions appropriate to the attainment of the objects of the GBRMP Act, (which may include conditions relating to the principles of ecologically sustainable use and the protection of matters of national environmental significance).</p>
<p data-bbox="148 1912 555 2018"><b>EIS process ensures unacceptable impacts on matters of national environmental significance do not occur (including implementation)</b></p>	<p data-bbox="571 1912 932 1942"><b>Mechanism (legislative &amp; policy):</b></p> <p data-bbox="571 1946 1410 2051">The permission system is established by the GBRMP Act as a mechanism to achieve the objects of the Act. The permission system regulates use of the Marine Park in ways consistent with ecosystem-based management and the principles of ecologically sustainable use.</p> <p data-bbox="571 2056 1390 2085">GBRMP Regulation 88ZE A permission may be granted subject to any conditions</p>



Outcome from EPBC Act	How listed migratory species, listed threatened species and ecological communities are considered under the Authority's permission system
	<p>appropriate to the attainment of the objects of the Act.</p> <p>GBRMP Regulation 88ZP A permission may be modified, suspended and revoked.</p> <p>The Authority may grant or refuse a permission based on the assessment. A permission may be granted with any conditions appropriate to the attainment of the objects of the GBRMP Act and the protection of matters of national environmental significance, the conditions can be modified, suspended or revoked.</p> <p><b>Program commitment(s):</b></p> <p>The Authority will</p> <ul style="list-style-type: none"> <li>• update the Environmental Impact Management Policy to enable the effective integration of all the commitments in the Program Report.</li> </ul> <p><b>Case study:</b></p> <ul style="list-style-type: none"> <li>• The assessment must demonstrate that appropriate avoidance, mitigation, offset, net benefit and adaptive management measures are in place to reduce impacts on affected species and ecological communities in the Marine Park. The measures must ensure that impacts will be acceptable and are not inconsistent with management plans or management principles relevant to affected species and ecological communities.</li> <li>• The Authority will apply conditions to ensure an unacceptable impact on listed turtles and the seagrass community does not occur as a result of the proposed action. Conditions will be developed having regard to recovery plans, conservation advices, wildlife conservation plans and threat abatement plans.</li> <li>• The conditions must ensure that the identified avoidance, mitigation, offset, net benefit and adaptive management measures, such as operational monitoring/environmental site supervisors, Deeds of Agreement and bonds are appropriate for managing impacts for the life of the project.</li> <li>• The Authority will not grant a permit for an activity inconsistent with regulatory objectives set out under the GBRMP Act.</li> </ul>
<p><b>AUDIT &amp; COMPLIANCE:</b> The purpose of the audit and compliance phase is to monitor the environmental performance of permitted activities against required standards and permit conditions. Where the results of performance monitoring indicate the required standards have not been met, penalties and/or remediation will be required. Permits may also be suspended or revoked depending on the severity of the non-compliant behaviour.</p>	
Monitoring, compliance and auditing	<p><b>Mechanism (legislative &amp; policy):</b></p> <p>Section 38EA of the GBRMP Act provides penalties for the contravention of a permission condition. Activities must be carried out in accordance with a Great Barrier Reef Marine Park Permission and any conditions.</p> <p>Section 38BA of the GBRMP Act states that it is an offence to conduct an activity in the Marine Park without Permission.</p> <p><b>Program commitment(s):</b></p> <p>The Authority will:</p> <ul style="list-style-type: none"> <li>• implement guidelines to improve the collection, management, analysis and interpretation of information derived from the Authority's permission system and compliance monitoring. The information will be used to better inform adaptive management strategies.</li> <li>• develop and/or strengthen knowledge and information management tools (including spatial mapping tools, hydrological connectivity tools and systems for managing Indigenous and historic heritage information) so their use can enhance and support: <ul style="list-style-type: none"> <li>○ the effects of impacts of actions on the Marine Park, including matters of national environmental significance</li> <li>○ measures to avoid, mitigate and offset impacts on the Marine Park, including matters of national environmental significance</li> <li>○ actions to deliver net benefits and best practice adaptive management.</li> </ul> </li> <li>• strengthen its system for monitoring and managing compliance risks associated with permit conditions. It will review and make any necessary improvements to internal policies, procedures, training protocols and mechanisms to systematically enhance the effectiveness of its compliance and auditing process. This will include specific reference to matters of</li> </ul>

Outcome from EPBC Act	How listed migratory species, listed threatened species and ecological communities are considered under the Authority's permission system
	<p data-bbox="630 235 991 264">national environmental significance.</p> <p data-bbox="571 293 703 322"><b>Case study:</b></p> <ul data-bbox="592 329 1406 1792" style="list-style-type: none"> <li data-bbox="592 329 1406 591">• The Authority's Environmental Impact management policy states that an Environmental Management Plan (EMP) is required for activities that may impact on the marine turtles and the seagrass community. The EMP outlines the proposed activity, its possible impacts on the Marine Park (including listed species and communities) and specifies strategies to minimise potential impacts, such as avoidance, mitigation, offset, net-benefit and adaptive management measures. An EMP is required for all activities undergoing an EIS assessment. The EMP needs to be approved by the Authority before implementation and before the commencement of activities permitted under the permit.</li> <li data-bbox="592 598 1406 703">• The EMP must include identification of potential environmental impacts on marine turtles and the seagrass community, how on-water activities will be managed to reduce these impacts, a monitoring program, emergency response plans and any relevant issue-based plans.</li> <li data-bbox="592 710 1406 869">• Independent experts are selected and directly contracted by the Authority review EMPs to ensure the environmental management measures, set out in the EMP for the protection of marine turtles and the seagrass community, are being followed and adhered to. The results of the review are reported to the Authority. An independent expert is required for all activities undergoing an EIS assessment.</li> <li data-bbox="592 875 1406 1115">• An Environmental Site Supervisor and Monitoring Consultant may be required (in addition to an independent expert who reviews EMPs) to ensure the activity is undertaken in accordance with the permission to ensure protection of marine turtles and the seagrass community. These persons must be funded by, but independent of, the proponent and be without any conflict of interest. The Site Supervisor and Monitoring consultants have an 'on the ground role' to liaise with the applicant to ensure the conditions of the Permission are implemented and upheld. The person selected for these roles are selected by the Authority</li> <li data-bbox="592 1122 1406 1196">• The Authority will select and directly contract the Monitoring Consultant and Environmental Site Supervisor. These are required for all high risk activities undergoing an EIS assessment.</li> <li data-bbox="592 1202 1406 1308">• The Environmental Site Supervisor may be engaged for the whole project or any relevant phases. For example a marine species site supervisor may be engaged for the period of drilling, blasting, during which there are likely impacts on marine turtles.</li> <li data-bbox="592 1314 1406 1473">• An Advisory Committee may be established for the project to provide advice to the Authority. The Advisory Committee may include representatives from Government, the proponent, community, the Environmental Site Supervisor, and experts in the fields of science and management. An Advisory Committee may be used to advise on mitigation measures in place for the protection of the seagrass community.</li> <li data-bbox="592 1480 1406 1608">• The Authority requires an environmental performance audit of activities against permission conditions for developments requiring an EIS. If environmental performance or compliance does not meet required standards for the protection of marine turtles and the seagrass community then penalties and/or remediation will be required.</li> <li data-bbox="592 1615 1406 1742">• The Authority can determine the need for a cash bond or bank guarantee and a Deed of Agreement to provide ongoing indemnity and, where necessary, to ensure there is an ongoing contractual obligation to protect the environment if the permit lapses.</li> <li data-bbox="592 1749 1406 1792">• Civil penalties are available for offences under the GBRMP Act for breaches of Permission conditions or undertaking an activity without Permission.</li> </ul>

### **3 Wetlands of international importance**

#### **3.1 Description**

##### **3.1.1 Wetlands of international importance**

The Convention on Wetlands of International importance, also known as the Ramsar Convention encourages the designation of sites containing representative, rare or unique wetlands, or wetlands that are important for conserving biological diversity. Once designated, these sites are added to the Convention's List of Wetlands of International Importance and become known as Ramsar wetlands.

A declared Ramsar wetland is a wetland, or part of a wetland, designated by the Commonwealth under the Ramsar Convention for inclusion in the List of Wetlands of International Importance (Article 2 of the Ramsar Convention). The Ramsar Convention sets out the criteria for identifying Wetlands of International Importance which relate to the wetland types represented by the site and the importance of the site for conserving biological diversity.

The Minister for the Environment may declare that a specified wetland be declared Ramsar wetland by notice in the Gazette if the Minister is satisfied that the wetland is of international significance, or is likely to be of international significance because of its ecology, botany, zoology, limnology or hydrology; and the ecological character of some or all of the wetland is under threat.

In designating a wetland as a Ramsar site, countries agree to establish and oversee a management framework aimed at conserving the wetland and ensuring its wise use. Wise use under the Convention is broadly defined as maintaining the ecological character of a wetland.

##### **3.1.2 Ramsar wetlands**

There are two Ramsar wetlands within or adjacent to the Great Barrier Reef Marine Park:

- Shoalwater and Corio Bays Area Ramsar site, and
- Bowling Green Bay Area Ramsar site.

Both of these areas are described in Section 4.8 of the Strategic Assessment Report. Further information on Ramsar wetlands under the EPBC Act can be found on the Department of the Environment's website: <http://www.environment.gov.au/topics/water/water-our-environment/wetlands/ramsar-convention-wetlands>.

A list of Ramsar Wetlands subject to this Strategic Assessment and the Program, along with other relevant matters of national environmental significance, are provided in Appendix 6.

#### **3.2 Key environmental values**

Under the Ramsar Convention, a wide variety of natural and human-made habitat types ranging from rivers to coral reefs can be classified as wetlands. Wetlands include swamps, marshes, billabongs, lakes, salt marshes, mudflats, mangroves, coral reefs, fens, peat bogs, or bodies of water - whether natural or artificial, permanent or temporary. Water within these areas can be static or flowing; fresh, brackish or saline; and can include inland rivers and coastal or marine water to a depth of six meters at low tide. There are even underground wetlands (DoE website, 2014).

The Authority has carried out a systematic analysis of key environmental values or 'attributes' and supporting environmental processes relevant to Ramsar wetlands covered by the Strategic Assessment. These are based on Ramsar Convention Information Sheets and are summarised in Table 4.8.

#### **3.3 Sensitivities**

Under the Australian Ramsar Management Principles, one of the primary purposes of management of a Ramsar wetland is to maintain the ecological character of the wetland. Under the EPBC Act, any action that is likely to result in a significant impact on the ecological character of a Ramsar wetland requires approval (Sections 16 and 17B of the EPBC Act).

Ecological Character is defined by the Ramsar Convention Resolution IX.1 (Annex A) as the combination of the ecosystem components, processes, and benefits and services that characterise the wetland at a given point in time. The phrase 'at a given point in time' refers to the time of designation (Ramsar Convention Resolution VI.1, paragraph 2.1).

The Authority uses EPBC Act significance guidelines in its initial screening of whether an action is likely to have a significant impact on the Marine Park, including Ramsar wetlands, and therefore if it requires further assessment of the acceptability of those impacts. Under the *EPBC Act Significant Impact Guidelines 1.1*, an action is likely to

have a significant impact on the ecological character of the declared Ramsar wetland if there is a real chance or possibility that it will result in:

- areas of the wetland being destroyed or substantially modified
- a substantial and measurable change in the hydrological regime of the wetland, for example, substantial change to the volume, timing, duration, and frequency of ground and surface water flows to and within the wetland
- the habitat or lifecycle of native species, including invertebrate fauna and fish species, dependent upon the wetland being seriously affected
- a substantial and measurable change in the water quality of the wetland – for example a substantial change in the level of salinity, pollutants, or nutrients in the wetland, or water temperature which may adversely impact on biodiversity, ecological integrity, social amenity or human health
- an invasive species that is harmful to the ecological character of the wetland being established (or an existing invasive species being spread) in the wetland.

Permit applications will be considered on a case by case basis, with consideration given to the objectives of the zone in which the proposed conduct will take place, as set out in the Zoning Plan.

The Authority will make decisions consistent with the objects of the GBRMP Act which includes the main object at Section 2A(1) to 'provide for the long-term protection and conservation of the environment, biodiversity and heritage values of the Great Barrier Reef Region'.

Environment has the same meaning as in the EPBC Act and is defined as including (among other things) ecosystems and their constituent parts, including people and communities, natural and physical resources, and the qualities and characteristics of locations, places and areas. This includes the ecological character of declared Ramsar wetlands.

The Authority has committed in its Program to take into account Ramsar Information Sheets, ecological character descriptions and plans of management for Ramsar wetlands in determining whether an activity is consistent with the objects of the GBRMP Act. Decisions will be made against the commitments in Table 10 of the Program. The Authority will use EPBC guidance documentation in acceptability assessments. The Program's commitments will ensure that the Authority acts consistently with the Australian Ramsar management principles and Australia's obligations under the Ramsar Convention.

Activities taking place a significant distance from the Ramsar wetlands in or adjacent to the Great Barrier Reef Marine Park can still result in impacts to the ecological character of the Ramsar site. For example, dependent on ocean currents and weather conditions an activity that originates some distance from the Ramsar wetlands could impact on the Ramsar wetlands. Under the Program, the potential impacts to all Ramsar sites, including future listings, identified as being at risk from a proposed activity will be considered, including those that are a significant distance from the location of the activity, where appropriate. Joint regulatory/management arrangements are in place with the Queensland Government and Commonwealth Department of the Environment regarding these types of assessments.

In assessing the sensitivity of the impacts of actions on values, consideration must be given to the current condition of values. The Authority has carried out a comprehensive assessment of the condition and trend of values or 'attributes' relevant to each matters of national environmental significance in Chapter 7 of this Report. As many values were assessed to be in poor and declining condition, the Authority acknowledges that impacts deemed not to have a significant impact in the past, may be assessed to have a likely significant impact in the future. To assist consideration of whether an action is likely to have a significant impact, the Authority has committed to the development and implementation of significant impact guidelines for the Great Barrier Reef Marine Park.

### 3.4 Relevant impacts

All impacts from activities covered by the Program (e.g. those requiring a permit under the Zoning Plan), including those in relation to the construction, operation and (if relevant) decommissioning of a project, are addressed by the permission system. Activities have the potential to impact Ramsar wetlands through a variety of sources depending on the location and nature of the action. Potential impacts on Ramsar wetlands include direct, indirect (upstream, downstream, facilitated) and cumulative impacts<sup>7</sup>.

Chapters 6, 7, 9, and 10 of this Report identify the impacts to Ramsar wetlands. These chapters identify the cumulative impacts from climate-related impacts (increased sea temperature and cyclone activity). Chapter 6 specifically deals with the Shoalwater and Corio Bays Area Ramsar site. Specific impact assessments of proposals must be undertaken in the context of the general level of degradation which has been identified in each habitat type within this report (see Table 6.6).

The potential impacts on any Ramsar wetland are dependent on the supporting and critical components (for example flora or fauna present in the wetland), processes (for example breeding activities) and services (for example provision of a key habitat) that make up the ecological character of a wetland.

<sup>7</sup> EPBC Act Significant Impact Guidelines 1.1



### 3.5 Assessment of activities under the permission system

When assessing the impacts of an activity on a Ramsar wetland, the permission system requires a permit application to describe the environment, identifying any part of the ecological character of the wetland that is likely to be affected by the proposed activity. The permission system requires analysis of the potential impacts to the ecological character of the wetland and provision of sufficient information to inform consideration and understand the ecological character of the wetland which may be affected by the proposed activity. The applicant further needs to examine how these values or elements may be affected and how impacts can be addressed and substantially reduced.

The permission system requires the Authority to analyse the potential impacts of the proposed activity (direct, indirect and cumulative) to the ecological character of the wetland and provide adequate opportunity for consultation. The Authority has committed to the development of cumulative impact assessment guidelines to provide a comprehensive, consistent and transparent approach to the assessment of potential impacts. Relevant documents that should be considered when assessing the potential impacts to a Ramsar site include:

- Ramsar information sheets
- Ecological Character Descriptions
- Ramsar Plans of Management

#### **Ramsar information sheets**

Contracting Parties to the Ramsar Convention are required to provide a Ramsar Information Sheet for all sites designated as Wetlands of International Importance under the Ramsar Convention. Ramsar information sheets need to be provided to the Ramsar Secretariat at the time of nomination of a site to the List of Wetlands of International Importance. Furthermore, parties to the Ramsar Convention have a commitment to provide updated Ramsar Information Sheet information for all of their Ramsar sites at intervals of six years or when there are any significant changes in the sites' ecological character.

The Ramsar Information Sheet provides essential data on each designated Wetland of International Importance, in order to allow analysis of Ramsar-listed wetlands around the world, provide baseline data for measuring changes in the ecological character of wetlands listed under the Ramsar Convention, and provide material for publications which inform the public about Ramsar sites. Under the EPBC Act, the detailed written description of a designated wetland in the Ramsar Information Sheet legally defines the 'declared Ramsar wetland'.

There are Ramsar information sheets for the Bowling Green Bay, Corio Bays and Shoalwater Bay Ramsar sites available through the Australian Government Department of the Environment's website: <http://www.environment.gov.au/cgi-bin/wetlands/alphablist.pl>.

#### **Ecological Character Descriptions**

Ecological Character Descriptions (ECDs) supplement the description of the ecological character contained in the Ramsar Information Sheet submitted under the Ramsar Convention for each listed wetland and, collectively, form an official record of the ecological character of the site.

ECDs describe the ecological character of a wetland at the time of its listing as a Wetland of International Importance. The description of ecological character is a requirement under the Ramsar Convention and the Australian Ramsar Management Principles. The information below is sourced from the *National Framework and Guidance for Describing the Ecological Character of Australia's Ramsar Wetlands* (Australian Government 2008).

The Ecological Character Description for a Ramsar wetland is also used to:

- Assist in implementing Australia's obligations under the Ramsar Convention, as stated in Schedule 6 (Managing wetlands of international importance) of the Environment Protection and Biodiversity Conservation Regulations 2000, including describing and maintaining the ecological character of declared Ramsar wetlands in Australia.
- Assist any person considering a proposed activity that may impact on a declared Ramsar wetland.

ECDs for both the Bowling Green Bay and Shoalwater Bay Ramsar sites are currently being developed and will be available through the Australian Government Department of the Environment's website once finalised: <http://www.environment.gov.au/cgi-bin/wetlands/alphablist.pl>

### Ramsar Plans of Management

Plans of management are used to formulate and implement planning so as to promote the wise use and conservation of wetlands. Plans of management should be consistent with the Ramsar Convention, Schedule 6 of the EPBC Act Regulations 2000 (the Australian Ramsar Management Principles) and relevant *National Guidelines for Ramsar Wetlands – Implementing the Ramsar Convention in Australia*.

Section 328 of the EPBC Act states that the Commonwealth is required to make plans of management for Ramsar sites entirely on Commonwealth land but not within a Commonwealth Reserve. For all other Ramsar wetlands best endeavours are being used to ensure that there is a management plan in place that is consistent with the Australian Ramsar Management Principles (Schedule 6 of the EPBC Regulations).

According to the Australian Ramsar Management Principles, the primary purpose of management of a Ramsar wetland is to describe and maintain the ecological character of the wetland. Additionally, the regulations note that before an action is taken, the likely impact on the wetlands ecological character should be assessed.

There is no Commonwealth plan of management for the Bowling Green Bay Area Ramsar site however a management plan has been prepared for the Queensland state Bowling Green Bay National Park. A plan of management is currently being developed for the Shoalwater and Corio Bays Ramsar site (DoE Website, 2014). Plans of management are available through the Australian Government Department of the Environment's website: <http://www.environment.gov.au/cgi-bin/wetlands/alphablist.pl>

Chapter 10 describes the future risk of impacts on values and attributes relevant to each matters of national environmental significance.

## 3.6 Safeguards under the Program

Under the EPBC Act, an action should not be approved if it would be inconsistent with:

- maintaining the ecological character of the wetland, or
- providing for the conservation and sustainable use of the wetland.

The Authority's is committed to maintaining the ecological character of Ramsar wetlands through the conservation and sustainable use of Ramsar wetlands.

The Authority will make decisions consistent with the objects of the GBRMP Act which includes the main object at section 2A(1) to 'provide for the long-term protection and conservation of the environment, biodiversity and heritage values of the Great Barrier Reef Region'. Environment has the same meaning as in the EPBC Act and is defined as including (among other things) ecosystems and their constituent parts, including people and communities, natural and physical resources, and the qualities and characteristics of locations, places and areas. This necessarily includes matters of national environmental significance.

In making decisions that are consistent with the objects of the GBRMP Act, the Authority must consider the potential impacts of the conduct proposed to be permitted by the permission on the environment and on the social, cultural and heritage values of the Marine Park or a part of the Marine Park.

The Authority requires permit applications and assessments under the permission system to be consistent with Australia's obligations under the Ramsar convention. Consideration of any Ramsar Information Sheet, Ecological Character Description, or plan of management relevant to the management of impacts of an activity is a requirement of the permission system. Permit application assessments also have regard to the significant impact criteria in the *EPBC Act Significant Impact Guidelines 1.1* for Ramsar wetlands that may or are likely to be impacted by the activity in the context of demonstrating acceptable levels of impacts. Assessment of the level of acceptability will also be informed by the permit applicant's measures to address the Authority's avoid-mitigate-offset-net benefit-adaptive management hierarchy for the assessment of impacts as outlined in their application and subsequent assessment report(s).

Under the permission system the Authority will make decisions consistent with the GBRMP Act:

- Consideration will be given to:
  - plans of management for a Ramsar wetland to preserve the ecological character of a Ramsar wetland.
  - the Australian Ramsar management principles.
  - relevant policy documents, guidelines, Ramsar Information Sheets, Ecological Character Descriptions and plans of management on the Department of the Environment website or reference on that website.
- The Authority maintains policy guidance that stakeholders should have regard to in the preparation of their Permit Applications. Policy guidance will be updated to include explicit reference to:

- consideration of all attributes relevant to matters of national environmental significance
- consideration of the protection of the ecological character of Ramsar wetlands
- relevant GBRMP Act and EPBC Act guidance documents to be considered by Permit Applicants in preparing Permit Applications including, but not limited to, Ramsar Information Sheets and Ecological Character Descriptions.

In undertaking assessments, the Authority has regard to relevant policy documents, guidelines, online data sources, Ramsar information sheets, Ecological Character Descriptions and plans of management on the DoE website.

The Authority's assessment process also ensures an assessment of the proponent's measures to avoid, mitigate, offset and provide for net-benefit and adaptive management of potential impacts to the ecological character of Ramsar wetlands.

All permitted activities must be carried out in a manner that is consistent with the objectives of the GBRMP Act, ecosystem-based management and the principles of ecologically sustainable use.

### 3.7 Case study - an activity with the potential to impact on a Ramsar wetland

**Scenario:** An eco-lodge for bird watching is proposed to be developed on land adjacent to the Bowling Green Bay Ramsar Wetland, with a jetty and a pipeline infrastructure passing through the wetland into the General Use Zone of Bowling Green Bay within the Great Barrier Reef Marine Park boundary. The proposal potentially includes bird viewing platforms in the wetland and small vessel docking facilities off the jetty (does not include dredging). Key aspects of the proposal include;

- a land based eco lodge accommodation facility which will include removal of ecosystems to be replaced with infrastructure, hardened surfaces and artificial landscapes.
- site access and service supply (electricity and water)
- a jetty and pipeline structure from the facility, through the Ramsar wetland and into the Great Barrier Reef Marine Park
- facility operations for 24 hours per day
- boardwalks and viewing platforms in the Ramsar wetland areas, passenger
- ferry loading from the jetty, (does not require access channel or berth dredging)

Key impacts of the activity may include:

- water quality impacts including altered hydrodynamics and changes in nutrient, sediment and other pollutant loads (e.g. from catchment run-off, urban and industrial discharge)
- impacts to marine habitats including disturbances to coral reef, seagrass and other benthic habitats
- impacts to supporting terrestrial habitats including modifying/changing/permanent and temporary loss of wetland habitat
- compromising habitat and amenity values
- interactions with protected species (e.g. marine debris, modifying supporting terrestrial habitats) marine debris
- noise pollution
- physical damage (e.g. increased visitation)
- vessel strike on wildlife
- waste discharges and small spills
- wildlife disturbance from key ecological functions (foraging, resting, breeding)
- light pollution.

These impacts, their cumulative effects, and past and present effects on values, including on matters of national environmental significance, are described in detail in Chapter 6.

There is no ecological character description for the Bowling Green Bay Ramsar wetland. The Ramsar information sheet lists the criterion for which the site was listed and the justifications for listing. For the purposes of this scenario, references to the ecological character of the Ramsar wetland include consideration of the site's Ramsar listing criteria. The following criteria are potentially impacted by the activity:

**Criterion 1: A wetland should be considered internationally important if it contains a representative, rare, or unique example of a natural or near-natural wetland type found within the appropriate biogeographic region.**

- The site is representative of many coastal and seasonal wetlands in the area, but is particularly significant for its diversity and extent of wetland types.
- The Bowling Green Bay site is representative of the richest coastal wetland habitats typical of the coastal wet-dry tropics of north-eastern Australia.

- The site plays a substantial role in protection against erosion by cyclones. The site provides an extremely important bait fish, finfish and crustacean nursery, providing critical habitat at particular stages in their lifecycle.

**Criterion 2: A wetland should be considered internationally important if it supports vulnerable, endangered, or critically endangered species or threatened ecological communities.**

- The site provides feeding grounds for vulnerable species such as dugongs, estuarine crocodiles and green turtles, and several bird species.

**Criterion 3: A wetland should be considered internationally important if it supports populations of plant and/or animal species important for maintaining the biological diversity of a particular biogeographic region.**

- The site is particularly important for the abundance and diversity of avifauna, which depend on maintenance of the site for continued use.
- The site regularly supports substantial numbers of all Australian waterbird groups, including post breeding populations of brolgas (4000), magpie geese (10,000), and various species of Anatidae (4000-5000).

**Criterion 4: A wetland should be considered internationally important if it supports plant and/or animal species at a critical stage in their life cycles, or provides refuge during adverse conditions.**

- The site is of special significance as breeding and feeding habitat for brolgas and magpie geese.

**Criterion 5: A wetland should be considered internationally important if it regularly supports 20,000 or more waterbirds.**

- The site seasonally supports a wide range of waterbird species well in excess of 20,000 individuals.

**Criterion 6: A wetland should be considered internationally important if it regularly supports 1% of the individuals in a population of one species or subspecies of waterbird.**

- Although population sizes are poorly known in northern Australia, it is likely that, seasonally, populations of brolgas exceed 1% of the total population.

Section 330 of the EPBC Act requires that a Commonwealth agency must not authorise any person to do anything that may contravene a plan made under Section 328 for management of a Ramsar wetland. If no plan made under Section 328 exists, a Commonwealth agency must take all reasonable steps to ensure that acts relating to a Ramsar wetland are not inconsistent with the Ramsar Management Principles (Schedule 6, EPBC Regulations). As a Commonwealth statutory authority, the Authority must comply with this requirement and ensure that all reasonable steps are taken to act consistently with the Ramsar Management Principles.

There is no Commonwealth Plan of Management for the Ramsar site however a management plan has been prepared for the Queensland state Bowling Green Bay National Park. The purpose of the plan is to maintain the naturally occurring diversity of flora and fauna, habitat types and other natural features of the area, with particular attention to the site's wetlands. The park management plan is also focussed on the aboriginal custodians of the area, with the maintenance of cultural resources and engagement in decision making a management priority. The plan of management will be consulted during the assessment regarding the proposal, as context to the wetlands values and character.

Under the GBRMP Act, a proposal for a facility requires permission (a permit) in the General Use Zone of the Great Barrier Reef Marine Park (GBRMP). As such, this activity would require a permit application to be submitted, assessed and issued to determine whether the potential impacts of the project on the Ramsar wetland and the broader environment will be acceptable. Although the activity is outside the boundary of the Ramsar wetland, there is potential for the activity to have 'downstream and facilitated' impacts on the wetland, such as water quality impacts, impacts on bird species, crocodiles, dugongs and turtles.

The activity has the potential to have a significant impact on the criteria in the Ramsar information sheet and would be subject to an Environmental Impact Statement (EIS) assessment in accordance with the Authority's environmental impact assessment and management policies including: the Environmental Impact Management Policy, Environmental Assessment and Management Risk Assessment Guidelines, Structures Policy, Water Quality Guidelines for the Great Barrier Reef Marine Park.

The Authority's assessment process requires the permit application to provide a description of the environment, impacts and identify the ecological character of the Ramsar wetland that is likely to be affected by the proposed activity identify the matters of national environmental significance that may be affected by the proposed activity, and provide sufficient information to inform decision making.

The permission system requires the Authority to analyse the potential impacts of the proposed activity (direct, indirect and cumulative) on matters of national environmental significance (such as Ramsar wetlands) and ensure impacts are identified and are not unacceptable. The types of studies and fields of investigation which would be required to support a permit application under such a scenario would include:

- Connecting processes, such as ocean hydrology, coastal processes, ground and surface water linkages and the effect of natural and long-term events. Information would be required to show the spatial and temporal scales in which the site is connected to its adjacent environments.



- Ecological functions and processes that are connected to and support ecosystems at the proposed site. This would include cataloguing the ecosystems, habitat and species within the site, and describing the sites relationship to ecosystems, habitats and species within the landscape.
- The current use and amenity adjacent to and supported by the site.
- The environments assimilative capacity and its ability to retain ecological health under the proposed land use change. This would include information on current ecological condition adjacent to and within the site.

**Table 3 How a permit application would meet EPBC Act requirements for Ramsar Wetlands**

Outcome from EPBC Act	How Ramsar Wetlands are considered under the Authority's permission system
	<p><b>SCOPING:</b> The purpose of the scoping phase is to proactively discuss the initial concepts of a proposed action in order to explain regulatory arrangements and identify potential mitigations. For actions which will likely have a significant impact on the Marine Park, including matters of national environmental significance, reference is made to relevant GBRMP Act and EPBC Act guidance material. A Permit Application Assessment Fee must be paid before the accompanying Permit Application Form is considered. The Permit Application Form must contain sufficient information to enable a clear understanding of the intended action.</p>
<p><b>Activities with a likely significant impact must be assessed under legislation with regards to matters of national environmental significance</b></p>	<p><b>Mechanism (legislative &amp; policy):</b></p> <p>Section 330 of the EPBC Act requires that a Commonwealth agency must not contravene a plan for listed wetlands in Commonwealth areas. As a Commonwealth statutory authority, the Authority must comply with this requirement.</p> <p>Section 334 of the EPBC Act requires that a Commonwealth agency must take all reasonable steps to ensure it exercises its powers and functions in relation to a wetland in a State in a way that is not inconsistent with:</p> <ul style="list-style-type: none"> <li>• the Ramsar Convention</li> <li>• the Australian Ramsar management principles</li> <li>• a plan prepared for managing the property with the State as describe in Section 333 of the EPBC Act.</li> </ul> <p>As a Commonwealth statutory authority, the Authority must comply with this requirement.</p> <p>GBRMP Regulations 88Q (mandatory considerations) and 88R (discretionary considerations) outline the Authority's permit assessment considerations.</p> <p>Under regulation 88Q(a) of the GBRMP Regulations, the Authority must consider the potential impacts of the conduct proposed to be permitted by the permission on the environment and on the social, cultural and heritage values of the Marine Park or a part of the Marine Park. This relevantly includes matters of national environmental significance.</p> <p>Under Regulation 88E the Authority would seek further information for a permit application that does not provide sufficient detail to evaluate whether an activity is having a significant impact. This includes:</p> <ul style="list-style-type: none"> <li>• a description of the relevant matters of national environmental significance attributes likely to be affected by the proposed activity</li> <li>• the direct, indirect and cumulative impacts of the proposed activity</li> <li>• the risks of those impacts to matters of national environmental significance.</li> </ul> <p>Failure to provide sufficient detail would result in the application being withdrawn.</p> <p><b>Program Commitment(s):</b></p> <p>The Authority will:</p> <ul style="list-style-type: none"> <li>• require consideration of relevant EPBC policy documents, including relevant significant impact guidelines and reporting tools, in the scoping phase of the permission assessment and decision process.</li> <li>• require consideration of prudent and feasible alternatives which may have a lower impact on the environment and amend its policy documentation to include reference to this requirement.</li> <li>• consider any relevant international conventions, Commonwealth State agreements, plans of management, recovery plans, wildlife conservation plans, threat abatement plans, approved conservation advice under the EPBC Act, EPBC Act guidelines, policies and information in assessing and determining permissions under its permission system.</li> <li>• consider relevant Department of the Environment policy documents, guidelines, plans of management and other online data sources available on the Department's website in its scoping, assessment and decision-making</li> </ul>

Outcome from EPBC Act	How Ramsar Wetlands are considered under the Authority's permission system
	<p>process.</p> <ul style="list-style-type: none"> <li>• implement guidelines to assist applicants to determine whether an action is likely to have a significant impact on the Great Barrier Reef Marine Park matters of national environmental significance. The Authority will seek to have these guidelines approved by the Minister.</li> <li>• amend policies and guidance material used in the determination of permissions so they explicitly require the consideration of matters of national environmental significance and relevant attributes, including: <ul style="list-style-type: none"> <li>○ biodiversity and environmental processes</li> <li>○ Indigenous and historic heritage values</li> <li>○ community benefits derived from the environment, including those not easily represented or measured (such as aesthetic values).</li> </ul> </li> <li>• develop guidance material to assist in determining the acceptability of impacts. Criteria shall include consistency of assessment outcomes with regulatory objectives, outcomes for the Marine Park, including matters of national environmental significance and relevant GBRMP Act and EPBC Act guidelines and standards.</li> <li>• develop specific guidance documents for notifications under Part 5 of the Zoning Plan to make reference to consideration of matters of national environmental significance under the EPBC Act.</li> </ul> <p>The Authority will require applicants to consider:</p> <ul style="list-style-type: none"> <li>• the impacts of actions on matters of national environmental significance and relevant attributes and environmental processes as described in the Strategic Assessment Report, and as amended in accordance with the Authority's policy review procedures.</li> <li>• impacts (including direct, indirect and cumulative impacts) arising from the development action in the context of all impacts affecting the matters of national environmental significance.</li> </ul> <p><b>Case study:</b></p> <ul style="list-style-type: none"> <li>• Under the Zoning Plan the proposed activity is considered to be 'carrying out works' and 'operating a facility' also within the Marine Park (jetty/pipeline structure, eco lodge). This type of activity can only occur in certain parts of the Marine Park – namely areas where the purpose of the activity is consistent with the objective of the relevant zone and only with the written permission of the Authority.</li> <li>• The Authority determines that proposed activity is consistent with the objectives of the relevant zone (General Use) and that the proponent requires a permit from the Authority for the carrying out of works/operating a facility within the Marine Park.</li> <li>• For actions which may have a significant impact on the Marine Park, including matters of national environmental significance, reference is made to: <ul style="list-style-type: none"> <li>○ GBRMP Act, Regulations, Zoning Plan, plans of management and relevant policy documents including the Authority's environmental impact management, position statements and guidelines;</li> <li>○ EPBC Act policy documents, including relevant significant impact guidelines, the EPBC Act Species Profile and Threats Database (SPRAT), listing advice policy statements, recovery plans, conservation advices, threat abatement plans and wildlife conservation plans available on the Department of the Environment website.</li> </ul> </li> <li>• Information available through online data sources hosted by the Department of the Environment, such as the Environment Reporting Tool, augments local knowledge in determining whether matters of national environmental significance are likely to occur in the area of interest.</li> <li>• EPBC Act Significant Impact Guidelines 1.1, as they relate to Ramsar Wetlands, includes consideration of the conservation status under the EPBC Act of the species and community. The Authority, in evaluating whether the activity is likely to have significant impacts, will take into, the following generic considerations from the EPBC Act Significant Impact Guidelines 1.1, which state that an action is likely to have a significant impact on the ecological character of a declared Ramsar wetland if there is a real change or possibility that it will result in: <ul style="list-style-type: none"> <li>○ Areas of the wetland being destroyed or substantially modified</li> <li>○ A substantial and measurable change in the hydrological regime of the</li> </ul> </li> </ul>

Outcome from EPBC Act	How Ramsar Wetlands are considered under the Authority's permission system
	<p>wetland, for example, a substantial change to the volume, timing, duration and frequency of ground and surface water flows to and within the wetland</p> <ul style="list-style-type: none"> <li>○ The habitat or lifecycle of native species, including invertebrate fauna and fish species, dependent upon the wetland being seriously affected</li> <li>○ A substantial and measurable change in the water quality of the wetland – for example, a substantial change in the level of salinity, pollutants, or nutrients in the wetland, or water temperature which may adversely impact on biodiversity, ecological integrity, social amenity or human health, or</li> <li>○ An invasive species that is harmful to the ecological character of the wetland being established (or an existing invasive species being spread) in the wetland.</li> </ul> <ul style="list-style-type: none"> <li>• The Authority seeks the advice of the Queensland Parks and Wildlife Service and other relevant agencies with an administrative responsibility relating to the proposed activity, on the potential impacts of the proposed activity and level of assessment.</li> <li>• Under this scenario the Authority determines the proposed activity is likely to have a significant impact on the ecological character of the Ramsar wetland (considering its Ramsar listing criteria and Ramsar Management Principles and requires assessment, as enabled by the GBRMP Regulations, by an Environmental Impact Statement (EIS) which also requires an Environmental Management Plan (EMP) to be prepared.</li> </ul>
<p><b>Applicant is required to provide adequate information to enable the decision maker to make a decision.</b></p> <p><b>The applicant describes impacts to matters of national environmental significance</b></p>	<p><b><i>Mechanism (legislative &amp; policy):</i></b></p> <p>GBRMP Regulation 88A enables the Authority to approve permit application forms. The Authority can vary the approved application form to meet legal and management responsibilities.</p> <p>GBRMP Regulation 88E enables the Authority to request further information, for the purpose of assessing an application, and for such information to be provided within 20 business days (or for a longer period, as determined by the Authority) or the application is taken to have been withdrawn.</p> <p>GBRMP Regulation 103 provides that persons shall not furnish false information in a permit application.</p> <p>Failure to provide the information that did not adequately describe the underpinning ecosystem functions and processes and the ecological character of the Ramsar wetland potentially at risk from the activity in detail sufficient to evaluate all impacts to assess whether the activity is having a significant impact and determine the level of assessment under the Program (e.g. by EIS) would result in the application being withdrawn.</p> <p><b><i>Program commitment(s):</i></b></p> <p>The Authority will:</p> <ul style="list-style-type: none"> <li>• amend its Permit Application Form to include a section which specifically addresses matters of national environmental significance.</li> <li>• seek to pursue amendments to the Great Barrier Reef Marine Park Regulations 1983 to explicitly outline the levels of assessment under the permission system.</li> <li>• update its permissions guidance material to outline the general information requirements for applicants according to the level of assessment.</li> </ul> <p><b><i>Case study:</i></b></p> <ul style="list-style-type: none"> <li>• The applicant will have access to guidance material developed by the Authority relating to the Marine Park, including matters of national environmental significance, that proponent's should have regard to in the preparation of their application.</li> <li>• An application must be submitted to the Authority. It must provide sufficient information on the proponents and the proposed facility so assessors can clearly understand what the intended activity is. With regard to the Ramsar wetland, the proponent will be required to provide information on the following in the application: <ul style="list-style-type: none"> <li>○ Describe the nature, scale, timing and duration of the proposed development, including associated activities and actions that occur in relation to the proposal.</li> <li>○ Include any potential expansion of the site's infrastructure in the future, and detail the scale of the final design and operation of the facility at the</li> </ul> </li> </ul>

Outcome from EPBC Act	How Ramsar Wetlands are considered under the Authority's permission system
	<p>site.</p> <ul style="list-style-type: none"> <li>○ Identify the Ramsar wetland's attributes likely to be affected potentially at risk from the proposal.</li> <li>○ Identify and describe the potential temporal and spatial impacts on the ecosystem functions, processes and species in the landscape.</li> <li>○ From this landscape assessment, describe the specific values and critical components, functions and processes that make up the ecological character of the Ramsar wetland, and their connection to the surrounding landscape.</li> <li>○ Specifically, with regard to the site's Ramsar listing criteria and the Ramsar Management Principles, identify areas that are significant to supporting water birds and habitats that support rare or threatened species (e.g. dugongs, turtles, and brolgas). Descriptions must identify any threatened species, include a sufficient description of the threatened species which may be at risk from the activity, and provide sufficient information to inform consideration by the Authority to identify potentially impacted habitats, including the significance of the area for each species potentially affected. This will include a description of the occurrence of these species within the project area and within any areas that may potentially be affected by indirect impacts such as water quality or noise effects.</li> <li>○ Identify and describe safeguards that maintain the ecological character of the wetland (acceptability of the impact). The proponent must demonstrate that appropriate contingency plans, including appropriate emergency response measures are in place to ensure that the proponent would be able to effectively respond to unplanned events. These plans would form part of the Environment Management Plan required for this activity to proceed.</li> <li>○ Provide specific information on potential impacts of a jetty/pipeline structure and associated activities on: protected, vulnerable or iconic species; coastal processes; water quality; hydrodynamics; noise; light; air quality; social values; cultural values and heritage values.</li> </ul> <ul style="list-style-type: none"> <li>• Descriptions must identify any direct, indirect or cumulative impacts and risks to the Ramsar Wetland from the proposed facility.</li> <li>• The information provided should be sufficient to inform the Authority's risk assessment processes and demonstrate that there will be no unacceptable residual impacts and risk to the Ramsar wetland.</li> <li>• If an application does not include sufficient information for the purposes of assessing the likely nature and scale of environmental impacts additional information or documents must be provided at the request of the Authority.</li> </ul>
<p><b>ASSESSMENT:</b> The purpose of the assessment phase is to enable an informed decision on whether or not a permit will be granted to undertake an action within the Great Barrier Reef Marine Park. During the assessment phase, the Authority considers criteria outlined in the Great Barrier Reef Marine Park Regulations 1983, and relevant GBRMP Act and EPBC Act guidance material.</p>	
<p><b>Public consultation occurs in a transparent manner and outcomes are taken into account</b></p>	<p><b>Mechanism (legislative &amp; policy):</b></p> <p>GBRMP Regulations 88Q (mandatory considerations) and 88R (discretionary considerations) outline the Authority's permit assessment considerations.</p> <p>Under Regulation 88E the Authority would seek further information for a permit application that does not provide sufficient detail to evaluate whether an activity is having a significant impact. This includes:</p> <ul style="list-style-type: none"> <li>• a description of the relevant matters of national environmental significance attributes likely to be affected by the proposed activity</li> <li>• the direct, indirect and cumulative impacts of the proposed activity</li> <li>• the risks of those impacts to matters of national environmental significance.</li> </ul> <p>Regulation 88D of the GBRMP Regulations requires applications to be advertised if the Authority considers that granting a permission may restrict the reasonable use by the public of part of the Great Barrier Reef Marine Park.</p> <p>GBRMP Regulations 88D provide that if a permit applicant does not publish an advertisement as required, the application is taken to have been withdrawn.</p> <p>GBRMP Regulation 88Q provides that public comments must be considered by the Authority when making decisions on whether to grant the permit.</p> <p><b>Program commitment(s):</b></p>



Outcome from EPBC Act	How Ramsar Wetlands are considered under the Authority's permission system
	<p>The Authority will:</p> <ul style="list-style-type: none"> <li>publish notification of all permit applications on the internet as soon as practicable after receiving a permit application.</li> <li>seek to pursue amendments to the GBRMP Regulations (in particular regulation 88D) to require advertising of applications where the granting of a permission may restrict the reasonable use by the public of a part of the Marine Park or is likely to have a significant impact on the Marine Park, including on matters of national environmental significance.</li> <li>update relevant policies to explicitly state that appropriate consultation will be undertaken with persons or organisations, including Commonwealth and State government agencies, whose functions, interests or activities could be impacted by the proposed activity.</li> </ul> <p><b>Case study:</b></p> <ul style="list-style-type: none"> <li>The Authority has a set of criteria that it must consider in deciding whether or not to grant a permission or impose conditions on an application for permission to construct the jetty/pipeline structures in the Marine Park. The Authority also has discretionary considerations which it may choose to consider in relation to the application.</li> <li>The grant of the permit for the facility/pipeline that passes through the Ramsar wetland into the Great Barrier Reef (Coastal) and Great Barrier Reef (Commonwealth) Marine Park is likely to restrict reasonable use of part of the Marine Park by the public. Under this scenario, the Great Barrier Reef Marine Park Regulation requires that the proponent publicly advertises the proposal. With reference to potential impacts on a Ramsar wetland, information in the public advertising must include: <ul style="list-style-type: none"> <li>an introduction about the company applying for the permit</li> <li>an overview of works or program to be undertaken</li> <li>a description of values and attributes relevant to matters of national environmental significance that are likely to be affected</li> <li>maps of the proposed location, showing relevant site management arrangements, a drawing or plan of the proposal, location of the Ramsar wetland and important and sensitive ecological features (such as nesting sites or locally important feeding areas for threatened species)</li> <li>a description of potential impacts, including those already described that potentially impact on elements that underpin the Ramsar wetlands ecological character</li> <li>a description of how environmental impacts will be managed, using the avoid, mitigate, offset, net-benefit and adaptive management hierarchy, and outlining the acceptability of the potential impact</li> <li>benchmark the impacts of activities on affected values/attributes against appropriate standards and relevant outcomes</li> <li>how the proposal will impact access by other users</li> <li>frequently asked questions, relevant policies and contact details.</li> </ul> </li> <li>This advertisement must be published in a newspaper circulating in the part of the State of Queensland adjacent to that part of the Marine Park in which the conduct is proposed to occur. The advertisement is to include an invitation for interested persons to make comment on the application within a period of not less than 30 days. The advertisement will also be published on the Authority's website.</li> <li>Following the public comment period, the Authority may issue a news release and/or schedule a meeting to discuss the outcome of the public comment phase. The Authority will prepare a Summary of Submissions Report which will be published on its website and circulated to the proponent and all those who sent in submissions.</li> <li>The Authority also undertakes appropriate consultation with persons or organisations, including Australian and State government agencies, whose functions, interests or activities could be impacted by the proposed activity.</li> </ul>
<p>The detail and rigour of the assessment process is commensurate with the scale and potential impacts of the activity i.e. risk-based.</p>	<p><b>Mechanism (legislative &amp; policy):</b></p> <p>Section 330 of the EPBC Act requires that a Commonwealth agency must not authorise any person to do anything that may contravene a plan made under Section 328 for management of a Ramsar wetland. If no plan made under Section 328 exists, a Commonwealth agency must take all reasonable steps to ensure that acts relating to a Ramsar wetland are not inconsistent with the Ramsar</p>

Outcome from EPBC Act	How Ramsar Wetlands are considered under the Authority's permission system
	<p>Management Principles (Schedule 6, EPBC Regulations). As a Commonwealth statutory authority, the Authority must comply with this requirement.</p> <p>GBRMP Regulations 88Q (mandatory considerations) and may consider 88R (discretionary considerations) outline the Authority's permit assessment considerations. Note: where discretionary considerations (88R) are required to implement the Program's commitments these become mandatory policy considerations in the Authority's permit assessments.</p> <p>The Authority's Environmental Impact Assessment Policy and Environmental Assessment and Management Risk Assessment Guidelines (developed in accord with the Australian Standards Environmental Risk Management Handbook HB203:2006) provide guidance in the assessment of risk including analysis of the likelihood and consequence of identified hazards being realised to obtain a Hazard Risk Grade.</p> <p><b>Program Commitment(s):</b></p> <p>The Authority will:</p> <ul style="list-style-type: none"> <li>• develop and maintain regionally-based standards for ecosystem health (including water quality) that support best management practices, and may be used to inform: <ul style="list-style-type: none"> <li>○ the assessment of point source and cumulative impacts on the Marine Park, including matters of national environmental significance</li> <li>○ actions to mitigate impacts and cumulative impacts on the Marine Park, including matters of national environmental significance</li> <li>○ outcomes-based decision-making and considerations for determining 'acceptability' of impacts on the Marine Park, including matters of national environmental significance, based the commitments in Table 10 of the Program Report</li> </ul> </li> <li>• implement guidelines requiring consideration of regionally-based standards for ecosystem health to improve the assessment of impacts including cumulative impacts on the Region's values.</li> <li>• facilitate development of a dredging and dredge material disposal policy</li> <li>• improve the hydrodynamic modelling guidelines so they take account of longshore drift, resuspension and inter-annual current variability.</li> <li>• develop guidance material to be used in the determination of permissions to require the consideration of the impacts of underwater noise, particularly on at-risk species and inshore species.</li> <li>• integrate relevant components, as required, of the Department of the Environment's EPBC Act assessment templates and manuals into its policy documents as they are reviewed and updated.</li> </ul> <p><b>Case study:</b></p> <ul style="list-style-type: none"> <li>• The level of assessment required for the jetty /pipeline structure by the Authority is determined through the Authority's Environmental Impact Management Policy. The assessment levels are from (low risk) to (high risk). The levels for Authority's environmental impact management closely reflect the Australian and Queensland Government development assessment process.</li> <li>• The Authority conducts an initial risk assessment on the application and further information from the proponent for the jetty / pipeline structure. The Authority uses a risk assessment to evaluate the impacts of the proposed action.</li> <li>• The Authority considers: <ul style="list-style-type: none"> <li>○ The scale of the project and potential impacts, including the long term future for the tourism facility, and potential impacts to the environment during the construction and operation phases of the proposal.</li> <li>○ The potential sites values, and its relationship of the site to the surrounding or connected environment and environmental processes.</li> <li>○ Other factors that influence the sites values, such as seasonality and current condition</li> <li>○ Conservation planning, advice, legislation, policies and guidelines (including EPBC Act statutory and guidance documents) that is relevant to the site, adjacent environment, and ecological communities and species.</li> <li>○ The potential direct, indirect and consequential impacts and risk of the proposed facility construction and operation. Considerations include</li> </ul> </li> </ul>

Outcome from EPBC Act	How Ramsar Wetlands are considered under the Authority's permission system
	<p>such things as, individual, cumulative and synergistic effects; temporal and spatial scale; likelihood; and magnitude and consequence</p> <ul style="list-style-type: none"> <li>○ Potential measures to avoid, mitigate, offset and provide for net-benefit and adaptive management of the potential impacts from the proposal.</li> <li>• The proposed jetty/pipeline structure at the Ramsar wetland site would require the highest level of assessment as set out in the Authority's guidelines, due to likely public interest and potential impacts on ecological character of the wetland. This means that the proposal will be assessed through an Environmental Impact Assessment process.</li> </ul>
<p>EIS process follows 'avoid/mitigate/offset/enhance principles'. Any residual significant impacts are offset appropriately.</p>	<p><b>Mechanism (legislative &amp; policy):</b></p> <p>GBRMP Regulation 88Q(b) requires that the Authority must consider options for monitoring, managing and mitigating the potential impacts of the proposed conduct.</p> <p>GBRMP Regulation 88X -The Authority must, after taking into account the matters that it is required or permitted to take into account under the GBRMP Act and Regulations, make a decision on the application</p> <p><b>Policy Commitment(s):</b></p> <p>The Authority will:</p> <ul style="list-style-type: none"> <li>• implement guidelines for the application of Great Barrier Reef offsets to maintain the condition of the Marine Park, including matters of national environmental significance and relevant attributes and environmental processes, where impacts cannot be avoided or mitigated: <ul style="list-style-type: none"> <li>○ The guidelines will seek to deliver an outcome equivalent to, or better than, the outcome that would apply if the EPBC Act Environmental Offsets Policy were applied.</li> </ul> </li> <li>• implement a Great Barrier Reef net benefit policy to enhance the condition of the Marine Park, including matters of national environmental significance and relevant attributes and environmental processes.</li> <li>• implement guidelines on cumulative impact assessment to inform a transparent, consistent and systematic approach to the assessment of impacts from activities within and adjacent to the Region.</li> </ul> <p><b>Case study:</b></p> <ul style="list-style-type: none"> <li>• The assessment of the permit application uses an 'avoid, mitigate, offset, net-benefit and adaptive management methodology' of controls in the impact assessment.</li> <li>• The Authority develop a terms of reference for the Environmental Impact Assessment that requires the proponent to identify the site and adjacent values, potential impacts and risk, and measures to avoid, mitigate, offset and provide for net-benefit and adaptive management of impact on ecological character of the Ramsar wetland. The proponent will generally be requested to address the matters that include the following:</li> <li>• Detailed analysis of project specifications, including such things, but not limited to, <ul style="list-style-type: none"> <li>○ the long term future footprint for the tourism facility</li> <li>○ its production capacity</li> <li>○ emission quantity and constituents</li> <li>○ discharge quantity and constituents</li> <li>○ maximum, minimum, mean and median rates of contaminant release</li> <li>○ daily, weekly, monthly and annual release rates</li> <li>○ day to day operation of the tourism facility</li> <li>○ decommissioning.</li> </ul> </li> <li>• Detailed analysis of the sites values, and its relationship to the surrounding or connected environment and environmental processes. This may include such things as, but not limited to <ul style="list-style-type: none"> <li>○ Ecological character of the Ramsar wetland</li> <li>○ Population of species</li> <li>○ Habitat types</li> <li>○ Seasonal influence on habitats and species</li> <li>○ Critical life stages such as feeding, recruitment, breeding or migration</li> </ul> </li> </ul>

Outcome from EPBC Act	How Ramsar Wetlands are considered under the Authority's permission system
	<p>that are likely to be affected</p> <ul style="list-style-type: none"> <li>○ Marine Parks values</li> <li>○ Ecosystem functions and processes that underpin the health and resilience of ecological character and the values of the Marine Park</li> <li>○ Current health, condition, resilience and vulnerability of ecosystem types and species potentially impacted by the action.</li> </ul> <ul style="list-style-type: none"> <li>• A detailed consideration of conservation planning that may apply to the site. This may include such things as, but not limited to: <ul style="list-style-type: none"> <li>○ Relevant legislation</li> <li>○ Zoning Plans</li> <li>○ Plans of Management</li> <li>○ Policies and guidelines</li> <li>○ Recovery plans</li> <li>○ Conservation advices</li> <li>○ Threat abatement plans</li> <li>○ Conservation status of each species or ecological community that may be impacted</li> </ul> </li> <li>• A detailed analysis of direct, indirect, consequential and facilitated impacts and risk of the proposed facility construction and operation. Considerations include such things as, but not limited to: <ul style="list-style-type: none"> <li>○ Individual, cumulative and synergistic effects</li> <li>○ Temporal and spatial scale</li> <li>○ Likelihood</li> <li>○ Magnitude and consequence</li> </ul> </li> <li>• A detailed analysis, description and effectiveness of measures such as: <ul style="list-style-type: none"> <li>○ Consideration of feasible alternatives to the proposal to avoid impacts</li> <li>○ Mitigation measures</li> <li>○ Offsetting of residual impacts</li> <li>○ Net benefits</li> <li>○ Adaptive management for the life of the project</li> <li>○ Outcomes and objectives for protected values</li> </ul> </li> <li>• The proponent must demonstrate in their assessment documentation that impacts on the ecological character of the Ramsar wetland is acceptable, and that impacts have been reduced to levels where ecological character will be maintained and enhanced. Considerations must include consistency of assessment outcomes with: <ul style="list-style-type: none"> <li>○ Regulatory objectives</li> <li>○ Outcomes for matters of national environmental significance including relevant attributes and environmental processes, and</li> <li>○ Relevant GBRMP Act and EPBC Act guidelines and standards.</li> </ul> </li> </ul>
<p><b>DECISION:</b> The purpose of the decision phase is to grant or refuse a permission based on information provided within the assessment phase. A permission may be granted with any conditions appropriate to the attainment of the objects of the GBRMP Act, (which may include conditions relating to the principles of ecologically sustainable use and the protection of matters of national environmental significance).</p>	
<p><b>EIS process ensures unacceptable impacts on matters of national environmental significance do not occur (including implementation)</b></p>	<p><b>Mechanism (legislative &amp; policy):</b></p> <p>The permission system is established by the GBRMP Act as a mechanism to achieve the objects of the Act. The permission system regulates use of the Marine Park in ways consistent with ecosystem-based management and the principles of ecologically sustainable use.</p> <p>GBRMP Regulation 88ZE A permission may be granted subject to any conditions appropriate to the attainment of the Act.</p> <p>GBRMP Regulation 88ZP A permission may be modified, suspended and revoked.</p> <p>The Authority may grant or refuse a permission based on the assessment. A permission may be granted with any conditions appropriate to the attainment of the objects of the GBRMP Act and the protection of matters of national environmental significance, the conditions can be modified, suspended or revoked.</p>



Outcome from EPBC Act	How Ramsar Wetlands are considered under the Authority's permission system
	<p><b>Program commitment(s):</b> The Authority will</p> <ul style="list-style-type: none"> <li>• update the Environmental Impact Management Policy to enable the effective integration of all the commitments in the Program Report.</li> </ul> <p><b>Case study:</b></p> <ul style="list-style-type: none"> <li>• The assessment must demonstrate that appropriate avoidance, mitigation, offset, net benefit and adaptive management measures are in place to reduce impacts on the ecological character of the Ramsar wetland. The measures must ensure that impacts will be acceptable and are not inconsistent with management plans or management principles relevant to the Ramsar wetland.</li> <li>• The Authority will apply conditions to ensure an unacceptable impact on the ecological character of the Ramsar wetland, or the ecosystem functions and processes that underpin the maintenance of ecological character, do not occur as a result of the proposed action. Conditions will be developed having regard to management plans, management principles, the site's Ramsar listing criteria and the Australian Ramsar Management Principle relevant to the Ramsar wetland.</li> <li>• The conditions must ensure that the identified avoidance, mitigation, offset, net benefit and adaptive management measures, such as operational monitoring/environmental site supervisors, Deeds of Agreement and bonds are appropriate for managing impacts for the life of the project.</li> <li>• The Authority will not grant a permit for an activity inconsistent with regulatory objectives set out under the GBRMP Act.</li> </ul>
<p><b>AUDIT and COMPLIANCE:</b> The purpose of the audit and compliance phase is to monitor the environmental performance of permitted activities against required standards and permit conditions. Where the results of performance monitoring indicate the required standards have not been met, penalties and/or remediation will be required. Permits may also be suspended or revoked depending on the severity of the non-compliant behaviour.</p>	
Monitoring, compliance and auditing	<p><b>Mechanism (legislative &amp; policy):</b> Section 38EA of the GBRMP Act provides penalties for the contravention of a permission condition. Activities must be carried out in accordance with a Great Barrier Reef Marine Park Permission and any conditions. Section 38BA of the GBRMP Act states that it is an offence to conduct an activity in the Marine Park without Permission.</p> <p><b>Program commitment(s):</b> The Authority will:</p> <ul style="list-style-type: none"> <li>• implement guidelines to improve the collection, management, analysis and interpretation of information derived from the Authority's permission system and compliance monitoring. The information will be used to better inform adaptive management strategies.</li> <li>• develop and/or strengthen knowledge and information management tools (including spatial mapping tools, hydrological connectivity tools and systems for managing Indigenous and historic heritage information) so their use can enhance and support: <ul style="list-style-type: none"> <li>○ the effects of impacts of actions on the Marine Park, including matters of national environmental significance</li> <li>○ measures to avoid, mitigate and offset impacts on the Marine Park, including matters of national environmental significance</li> <li>○ actions to deliver net benefits and best practice adaptive management.</li> </ul> </li> <li>• strengthen its system for monitoring and managing compliance risks associated with permit conditions. It will review and make any necessary improvements to internal policies, procedures, training protocols and mechanisms to systematically enhance the effectiveness of its compliance and auditing process. This will include specific reference to matters of national environmental significance.</li> </ul> <p><b>Case study:</b></p> <ul style="list-style-type: none"> <li>• An Environmental Management Plan (EMP) is required for activities that may impact on the ecological character of the Ramsar wetland. The EMP outlines the proposed activity, its possible impacts on the Marine Park and the Ramsar wetland, and specifies strategies to minimise potential impacts.</li> </ul>

Outcome from EPBC Act	How Ramsar Wetlands are considered under the Authority's permission system
	<p>An EMP is required for all activities undergoing an EIS assessment. The EMP needs to be approved by the Authority before implementation and before the commencement of activities permitted under the permit.</p> <ul style="list-style-type: none"> <li>• The EMP must include identification of potential residual environmental impacts on ecological character, or on ecosystem functions and processes that underpin the maintenance of ecological character. The EMP will include how residual environmental impacts will be managed, a monitoring program, emergency response plans and any relevant issue-based plans.</li> <li>• Independent experts who are selected and directly contracted by the Authority review EMPs to ensure the environmental management measures set out in the EMP are being followed and adhered to. The results of the review are reported to the Authority. An independent expert is required for all activities undergoing an EIS assessment.</li> <li>• An Environmental Site Supervisor and Monitoring Consultant may be required (in addition to an independent expert who reviews EMPs) to ensure the activity is undertaken in accordance with the permission for the trestle structure/pipeline. These persons must be funded by, but independent of, the proponent and be without any conflict of interest. The Site Supervisor and Monitoring consultants have an 'on the ground role' to liaise with the applicant to ensure the conditions of the Permission are implemented and upheld. The person selected for these roles are selected by the Authority</li> <li>• The Authority will select and directly contract the Monitoring Consultant and Environmental Site Supervisor. These are required for all high risk activities undergoing an EIS assessment.</li> <li>• The Environmental Site Supervisor may be engaged for the whole project or any relevant phases. For example an avifauna species site supervisor may be engaged for the period of construction in the Ramsar wetland during, where impacts on bird fauna are most likely.</li> <li>• An Advisory Committee may be established for the project to provide advice to the Authority. The Advisory Committee may include representatives from Government, the proponent, community, the Environmental Site Supervisor, and experts in the fields of science and management. An Advisory Committee may be used to advise on mitigation measures in place for the protection of the Ramsar wetland.</li> <li>• The Authority requires an environmental performance audit of activities against permission conditions for developments requiring an EIS. If environmental performance or compliance does not meet required standards for the protection of the Ramsar wetland then penalties and/or remediation will be required.</li> <li>• The Authority can determine the need for a cash bond or bank guarantee and a Deed of Agreement to provide ongoing indemnity and, where necessary, to ensure there is an ongoing contractual obligation to protect the environment if the permit lapses.</li> <li>• Civil penalties are available for offences under the GBRMP Act for breaches of Permission conditions or undertaking an activity without Permission.</li> </ul>

## **4 World Heritage, National Heritage, Great Barrier Reef Marine Park, Commonwealth Marine Areas**

### **4.1 Description**

#### **4.1.1 World heritage property**

The World Heritage Convention (the Convention for the Protection of the World Cultural and Natural Heritage) (the Convention) was adopted by the United Nations Education, Scientific and Cultural Organisation (UNESCO) in 1972. Australia became a signatory (State Party) to the Convention in 1974. The World Heritage List, established by the Convention, comprises those parts of the world's cultural and natural heritage which are so important that they are considered to be of outstanding value to humanity as a whole. This level of importance is known as 'outstanding universal value'. State Parties to the Convention undertake to identify, protect, preserve and present this outstanding universal value. Australia has 19 properties on the Committee's World Heritage List.

Places on the World Heritage List each have a Statement of Outstanding Universal Value, determined by the Committee in consultation with the relevant State Party. The World Heritage values of a property listed on the World Heritage List are protected under Part 3 of the EPBC Act as matters of national environmental significance. For the purposes of the EPBC Act, a property's world heritage values are essentially the same as the world heritage attributes identified in the property's Statement of Outstanding Universal Value. This is the case for the Great Barrier Reef World Heritage Area.

The Great Barrier Reef is a world heritage property encompassing both Commonwealth and Queensland areas seaward of low water mark (see Figure 1.1, Chapter 1). Another world heritage property, the Wet Tropics of Queensland, is adjacent to the Great Barrier Reef World Heritage Area. There are close natural and cultural connections between the two properties. Further information on Great Barrier Reef World Heritage Area is provided in Section 4.2 of this report. A list of world heritage properties subject to this Strategic Assessment and the Program along with other relevant matters of national environmental significance is provided in Appendix 6.

#### **4.1.2 National heritage places**

The National Heritage List is a list of places of outstanding heritage significance to Australia. It comprises places with natural, historic and/or Indigenous values. Each place in the List has been assessed by an independent body, the Australian Heritage Council, to determine whether the place has national heritage values. The Environment Minister makes the final decision on whether a place is listed. Under the EPBC Act, a place is included on the National Heritage list if the Minister is satisfied that the place meets one or more of the National Heritage criteria prescribed in the EPBC Regulations. The listed values are then gazetted.

The Great Barrier Reef and Wet Tropics of Queensland are national heritage places relevant to the strategic assessment (both places are listed as World Heritage Properties). Further information on national heritage places is provided in Section 4.4 of this report. For a list of National Heritage places subject to this Strategic Assessment and the Program, see Appendix 6.

#### **4.1.3 The Great Barrier Reef Marine Park**

The Great Barrier Reef Marine Park is established under the GBRMP Act. The Marine Park covers 344,400 square kilometres and includes the subsoil beneath the seabed extending to a depth of 1000 metres and the airspace above extending to a height of 915 metres. It does not include Queensland islands and internal waters.

In accordance with the GBRMP Act, the Marine Park is managed to protect and conserve the environment biodiversity and heritage values of the Region. The term 'environment' is defined broadly and, as well as ecosystems, natural resources and heritage values, it encompasses physical resources — the area's geomorphological and geological aspects; and people and communities — social, economic and cultural aspects.

The Marine Park is a multiple-use area allowing ecologically sustainable use, consistent with the protection and conservation of the Region's environment, biodiversity and heritage values.

Further information on the Great Barrier Reef Marine Park is provided in Section 4.3 of this report.

#### **4.1.4 Commonwealth marine areas**

The marine environment is a matter of national environmental significance. It includes marine waters, airspace above those waters, seabed features, and all marine biota within those areas. The marine environment also includes social and cultural values, including recreational opportunities, amenity, cultural heritage, conservation and scientific significance.

The Commonwealth marine area extends beyond the outer edge of state/territory waters, generally some three nautical miles (or 5.5 km) from the coast, to the boundary of the Australia's exclusive economic zone, generally around 200 nautical miles from shore. The Coral Sea Commonwealth marine Reserve is adjacent to the Great

Barrier Reef Marine Park. Further information on Commonwealth marine areas is provided in Section 4.5 of this report.

## 4.2 Key environmental values

The strategic assessment concluded that there is substantial overlap and connections between the matters of national environmental significance and the values that support them. In addition, the matters and the values that relate to them do not occur in isolation but are embedded within complex and dynamic systems.

**World heritage values** are the matter protected under the EPBC Act. A Statement of the outstanding universal value of the Great Barrier Reef World Heritage Area is the official statement adopted by the World Heritage Committee outlining how the property met the criteria for outstanding universal value, integrity and protection and management at the time of listing. The statement was prepared retrospectively based on information that was available in 1981 and addresses the criteria in place at the time of inscription. It does, however, adopt the contemporary numbering for the relevant criteria.

The statement provides guidance on the world heritage attributes of the property. A description of the outstanding universal value of the Great Barrier Reef World Heritage Area is provided in Section 4.2 of this report. Since the Reef was inscribed on the World Heritage List, the criteria have been amended and renumbered. It is reproduced in full at Appendix 3 and is summarised below:

### Statement of Outstanding Universal Value

#### **Criteria (i) now (viii) — major stages of the Earth's evolutionary history**

The Great Barrier Reef, extending 2300 kilometres along Queensland's coast, is a globally outstanding representation of the major stages of the Earth's evolutionary history. The area has been exposed and flooded by at least four glacial and interglacial cycles, and over the past 18,000 years reefs have grown on the continental shelf. During glacial periods, sea levels dropped, extending the coastline further east and exposing the reefs as flat-topped hills of eroded limestone with large rivers meandering between them. During interglacial periods, rising sea levels caused the formation of continental islands and coral cays, and new phases of coral growth. Today's varied seascapes and landscapes have been moulded by changing climates and sea levels, and the erosive power of wind and water, over long time periods. A recent report *Geological and geomorphological features of outstanding universal value in the Great Barrier Reef World Heritage Area* identifies the geological and geomorphological features of outstanding universal value and provides an overview of the pressures affecting values. The outcomes are summarised in Section 4.3.2.

#### **Criteria (ii) now (ix) — ecological and biological processes**

The ecological and biological processes of the Region reflect the maturity of an ecosystem that has evolved over millennia. Globally significant marine fauna groups include more than 4000 species of molluscs, more than 1625 species of fish, as well as a great diversity of sponges, anemones, marine worms, crustaceans, and many other taxonomic groups. The establishment of vegetation on the cays and continental islands exemplifies the important role of birds in processes such as seed dispersal and plant colonisation. Biodiversity attributes and environmental processes which support the Great Barrier Reef are described in more detail in Section 4.3.1 and Section 4.10. Human interaction with the natural environment is illustrated by strong ongoing links between Aboriginal and Torres Strait Islanders and their land and sea country, and includes numerous shell deposits (middens), fish traps, story places and marine totems. A description of Indigenous heritage values is provided in Section 4.3.3.

#### **Criteria (iii) now (vii) — natural beauty and natural phenomena**

The Great Barrier Reef demonstrates superlative natural beauty above and below the water, providing spectacular scenery. It is one of a few living structures visible from space, appearing as a complex string of reefs along Australia's north-east coast. From the air, the vast mosaic patterns of reefs, islands and coral cays produce an unparalleled aerial panorama. The Whitsunday Islands provide a magnificent vista of green vegetated islands and spectacular sandy beaches. Hinchinbrook Island supports vast mangrove forests, rugged vegetated mountains and lush rainforest gullies. The Reef's natural phenomena include annual coral spawning, migrating whales, nesting turtles, and significant spawning aggregations of many fish species.

#### **Criteria (iv) now (x) — habitats for conservation of biodiversity**

The enormous size and diversity of the Region means it is one of the richest and most complex natural ecosystems on Earth, and one of the most significant for biodiversity conservation. The diversity of habitats supports tens of thousands of marine and terrestrial species, many of which are of global conservation significance. For example, the waters of the Region provide major feeding grounds for one of the world's largest populations of the threatened dugong and it is an important area for humpback whale calving. Furthermore, six of the world's seven species of marine turtle occur in the Region, with internationally important breeding grounds for green, loggerhead and hawksbill turtles. A more detailed description of Great Barrier Reef species and habitats is provided in Sections 4.3.1, 4.6, 4.7 and 4.8.



At the time of inscription it was considered that to include virtually the entire Great Barrier Reef within the property was the only way to ensure **the integrity** of the coral reefs in all their diversity. The Statement of Outstanding Universal Value also notes that some of the key ecological, physical and chemical processes that are essential for the long-term conservation of the marine and island ecosystems and their associated biodiversity occur outside the boundaries of the property.

The Great Barrier Reef World Heritage Area was one of 15 world heritage properties included as a **national heritage place** on the National Heritage List in 2007. While there are specific criteria that apply to the listing of national heritage places, the national heritage listing of the world heritage properties was done on the basis of those values identified by the World Heritage Committee. Therefore, for the purposes of the Strategic Assessment Report and Program, the values of the Great Barrier Reef national heritage place are taken to correspond to the world heritage criteria (see Section 4.4). The **Great Barrier Reef Marine Park** has a wide range of values, including those related to the environment, biodiversity and heritage values (see Section 4.3 of this report). They may be grouped into four broad categories:

- biodiversity, including habitats and species
- geomorphological features
- Indigenous and historic heritage
- community benefits derived from the environment, comprising the cultural, social and economic benefits derived from the Region's environment.

The **Commonwealth marine area** is a subset of the Marine Park and protection of the environment is a common aspect of both. Because of these commonalities, the Strategic Assessment Report and the Program considers the values of the Commonwealth marine area to fall into the same categories as those described for the Great Barrier Reef Marine Park (see Section 4.5). Listed marine species are considered as part of the impact assessment process under the EPBC Act and cetaceans are protected in the Commonwealth marine area under Part 13 of the EPBC Act. For a list of marine species and cetaceans subject to this Strategic Assessment, see Appendix 6.

Another important heritage feature of the Commonwealth marine area is underwater cultural values. Shipwrecks and associated relics are protected under the *Historic Shipwrecks Act 1976* (the Shipwreck Act). There are more than 1400 historic shipwrecks within the Great Barrier Reef. The Shipwrecks Act protects all shipwrecks and associated relics that are 75 years or older, regardless of whether their physical location is known. Shipwrecks younger than 75 years old can be individually declared protected. More information about shipwrecks, including the location of known protected sites, can be found at: [www.environment.gov.au/aggregation/historic-shipwrecks](http://www.environment.gov.au/aggregation/historic-shipwrecks)

The Authority has carried out a systematic analysis of key environmental values or 'attributes' and relevant supporting each of these matters of national environmental significance. These are summarised in Table 4.8. A description of environmental processes which support and connect the Great Barrier Reef environment is provided in Section 4.9.

### 4.3 Sensitivities

A range of management arrangements are in place for the protection of the Great Barrier Reef World Heritage Area, national heritage place, Marine Park and Commonwealth marine area. The Marine Park covers around 98 per cent of the World and National Heritage Area, the other two per cent being Queensland islands and internal waters.

The GBRMP Act establishes the Great Barrier Reef Marine Park Authority to manage and advise government in relation to care and development of the Marine Park. In doing so, the Authority must have regard to, and act consistently with, the protection of World Heritage values. Zoning plans are the key planning instrument through which the Marine Park is managed.

The Authority must, in preparing zoning plans, have regard to the Australian World Heritage Management Principles, National Heritage Management Principles and National Heritage values of the Great Barrier Reef, as specified under and pursuant to, the EPBC Act.

Whether the action/s would have an unacceptable impact is a function of the significance of its potential impacts and other factors such as proposed mitigation and/or offset measures. Under the EPBC Act Significant Impact Guidelines 1.1, an activity is likely to have a significant impact on the World Heritage values of a declared World Heritage property if there is a real chance or possibility that it will cause:

- one or more of the World Heritage values to be lost
- one or more of the World Heritage values to be degraded or damaged
- one or more of the World Heritage values to be notably altered, modified, obscured or diminished.

In addition to the above three impacts, the referral guidelines for the Outstanding universal value of the Great Barrier Reef World Heritage area states that an action is likely to have a significant impact on the Outstanding Universal Value of a World Heritage property if there is a real chance or possibility that it (at the site of the action or elsewhere as a result of the action) will:

- impact on the integrity of the property

Similarly, under the EPBC Act Significant Impact Guidelines 1.1, an activity is likely to have a significant impact on National Heritage values if there is a real chance or possibility that it will cause:

- one or more of the National Heritage values to be lost
- one or more of the National Heritage values to be degraded or damaged
- one or more of the National Heritage values to be notably altered, modified, obscured or diminished.

An action is likely to have a significant impact on the environment of the Great Barrier Reef Marine Park if there is a real chance or possibility that the action will:

- modify, destroy, fragment, isolate or disturb an important, substantial, sensitive or vulnerable area of habitat or ecosystem component such that an adverse impact on marine ecosystem health, functioning or integrity in the Great Barrier Reef Marine Park results
- have a substantial adverse effect on a population of a species or cetacean including its life cycle (for example, breeding, feeding, migration behaviour, life expectancy) and spatial distribution
- result in a substantial change in air quality or water quality (including temperature) which may adversely impact on biodiversity, ecological health or integrity or social amenity or human health
- result in a known or potential pest species being introduced or becoming established in the Great Barrier Reef Marine Park
- result in persistent organic chemicals, heavy metals, or other potentially harmful chemicals accumulating in the marine environment such that biodiversity, ecological integrity, or social amenity or human health may be adversely affected, or
- have a substantial adverse impact on heritage values of the Great Barrier Reef Marine Park, including damage or destruction of an historic shipwreck.

Under the EPBC Act, a person must not take an action in a Commonwealth marine area or outside a Commonwealth marine area that has, will have or is likely to have a significant impact on the environment without an approval. An action is likely to have a significant impact on the environment in a Commonwealth marine area if there is a real chance or possibility that the action will:

- result in a known or potential pest species becoming established in the Commonwealth marine area
- modify, destroy, fragment, isolate or disturb an important or substantial area of habitat such that an adverse impact on marine ecosystem functioning or integrity in a Commonwealth marine area results
- have a substantial adverse effect on a population of a marine species or cetacean including its life cycle (for example, breeding, feeding, migration behaviour, life expectancy) and spatial distribution
- result in a substantial change in air quality or water quality (including temperature) which may adversely impact on biodiversity, ecological integrity; social amenity or human health
- result in persistent organic chemicals, heavy metals, or other potentially harmful chemicals accumulating in the marine environment such that biodiversity, ecological integrity, social amenity or human health may be adversely affected, or
- have a substantial adverse impact on heritage values of the Commonwealth marine area, including damage or destruction of an historic shipwreck.

Under the EPBC Act when assessing impacts to the Great Barrier Reef Marine Park or a Commonwealth marine area all elements of the 'environment' must be considered to the extent that they apply. There are tools available to assist in identifying the values or sensitivities. For example, the Department of Environment maintains a suite of interactive tools as part of the Conservation Values Atlas that allows users to search, find and generate reports on information and data. The Conservation Values Atlas is designed to provide a visual representation of the conservation values in each marine region, including location and spatial extent where sufficient information exists. More information is available at [www.environment.gov.au/cva](http://www.environment.gov.au/cva).

Permit applications will be considered on a case by case basis, with consideration given to the objectives of the zone in which the proposed conduct will take place, as set out in the Zoning Plan.

The Authority will make decisions consistent with the objects of the GBRMP Act which includes the main object at Section 2A(1) to 'provide for the long-term protection and conservation of the environment, biodiversity and heritage values of the Great Barrier Reef Region'.

Environment has the same meaning as in the EPBC Act and is defined as including (among other things) ecosystems and their constituent parts, including people and communities, natural and physical resources, and the qualities and characteristics of locations, places and areas. This includes the environment in a Commonwealth marine area and the environment in the Marine Park.

Heritage values have the same meaning as in the EPBC Act and are defined as 'the natural and cultural environment having aesthetic, historic, scientific or social significance'. The world heritage values of declared World Heritage Places and the national heritage values of declared National Heritage places fall within the social, cultural and heritage values of the Marine Park.

In assessing the sensitivity of the impacts of actions on values, consideration must be given to the current condition of values. The Authority has carried out a comprehensive assessment of the condition and trend of

values or 'attributes' relevant to each matters of national environmental significance in Chapter 7 of this Report. As many values were assessed to be in poor and declining condition, the Authority acknowledges that impacts deemed to not to have a significant impact in the past, may be assessed to have a likely significant impact in the future. To assist consideration of whether an action consistent with the objects of the GBRMP Act, the Authority has committed to the development and implementation of significant impact guidelines for the Great Barrier Reef Marine Park. Decisions will be made against the commitments in Table 10 of the Program.

## 4.4 Relevant Impacts

Under the EPBC Act an activity would require approval if:

- the action is taken in the Great Barrier Reef Marine Park or Commonwealth marine area and the action has, will have, or is likely to have a significant impact on the environment, or
- the action is taken outside the Great Barrier Reef Marine Park or Commonwealth marine area and the action has, will have, or is likely to have a significant impact on the environment in the Great Barrier Reef Marine Park.

Impacts from all activities requiring a permit under the Zoning Plan, including those in relation to the construction, operation and (if relevant) decommissioning of a facility and the carrying out of works, are addressed by the permission system. Permitted activities have the potential to impact World Heritage values through a variety of sources depending on the location and nature of the action. An activity taken outside the boundary of a World Heritage property can have the potential to impact the property's values.

There are both direct and indirect potential impacts on the World Heritage values of the Property, which are detailed in Chapters 6, 7, 9 and 10 in this report. Chapters 6, 7, 9 and 10 of this Report identify the key impacts to the Property. A summary of the key impacts on the Property's values is found in Table 4.

**Table 4 Summary of impacts affecting OUV**

Short title of World Heritage criteria	Summary of key impacts
<b>Major stages of the Earth's evolutionary history</b>	The strategic assessment of impacts on geomorphological features shows the main impacts affecting them are cyclones and increased sediments. Cyclones have the ability to alter and weaken the structure of the geomorphological features, while sediments affect the features that are reliant upon photosynthesis (such as coral reefs and seagrass meadows) and can infill palaeochannels.
<b>Ecological and biological processes</b>	Modifying supporting terrestrial habitats has changed hydrological processes in many parts of the catchment. This has had downstream impacts on ecological and biological processes, reducing connectivity and the health of the World Heritage Area. The strong ongoing links between Aboriginal and Torres Strait Islanders and their land and sea country have been and continue to be affected by those impacts which are diminishing the natural environment of the Region.
<b>Natural beauty and phenomena</b>	Combined effects from cyclones, declining water quality and increasing sea temperature are affecting on the natural beauty and phenomena of the Great Barrier Reef ecosystem. Coastal habitat degradation and highly visible activities, such as coastal development, dredging and shipping, may have also diminished natural scenic values, especially in coastal areas. Transient impacts to the aesthetic experience of a place can include large numbers of tourists, passing ships or turbidity due to weather. Natural phenomena such as nesting marine turtles have been affected by previous commercial harvesting which has increased the species' vulnerability to subsequent impacts. The most severe current impacts affecting nesting marine turtles include death of discarded species and illegal fishing and poaching.
<b>Habitats for conservation of biodiversity</b>	Cyclones, increasing sea temperature, crown-of-thorns starfish outbreaks and increased freshwater inflow are threatening habitats and species. Almost all areas of the Region have been exposed to one or more of these threats over the last decade. The assessment shows the greatest impacts to seagrass and coral habitats arise from the adjacent catchment (specifically increased freshwater inflow, nutrients, pesticides and sediments) and from extreme weather.

## 4.5 Assessment of activities under the permission system

When assessing the impacts of an activity on the World Heritage Area, the environment of the Marine Park, National Heritage places or Commonwealth marine areas the permission system requires a permit application to describe the environment, identifying any gazetted value or element that are likely to be affected by the proposed activity, and provide sufficient information to understand the extent and risks of the impacts from the proposed activity. The applicant further needs to examine how these values or elements may be affected and how impacts can be addressed and substantially reduced.

The permission system requires the Authority to analyse the potential impacts (direct and indirect) to the environment and provide adequate opportunity for consultation. The Authority has committed to the development of cumulative impact assessment guidelines to provide a consistent and transparent approach to the assessment of potential impacts. Relevant EPBC Act documents will be considered when assessing whether a project is likely to have an impact on the environment, including but not limited to:

- Plans of management (Commonwealth Heritage)
- Gazettal instruments
- Bioregional plans
- Recovery plans, threat abatement plans, & conservation advices
- Wildlife conservation plans

### **Plans of management**

#### **World heritage properties**

Plans of management are used to formulate and implement planning so as to promote the wise use and conservation of World Heritage properties. Plans of management should be consistent with the World Heritage Convention, Schedule 5 of the EPBC Act Regulations 2000 (the Australian World Heritage Management Principles).

Section 316 of the EPBC Act states that the Commonwealth is required to make plans of management for World Heritage properties entirely within one or more Commonwealth areas, but not within a Commonwealth Reserve. Section 321 of the EPBC Act, which covers the management of World Heritage properties in States and self-governing Territories, notes that the requirement for Queensland to prepare and implement a plan does not apply for the Great Barrier Reef Marine Park, as a zoning plan must be prepared under the GBRMP Act, where regard must be had to the Australian World Heritage management principles.

For all other World Heritage properties best endeavours are being used to ensure that there is a management plan in place that is consistent with the Australian World Heritage Management Principles (Schedule 5 of the EPBC Regulations).

According to the Australian World Heritage Management Principles, the primary purpose of management of a World Heritage property is to identify, protect, conserve, present, transmit to future generations and, if appropriate, rehabilitate the World Heritage values of the property.

The *Great Barrier Reef Marine Park Act 1975* (GBRMP Act) and its Regulations govern protection and management of the Great Barrier Reef Region. The Act provides the Authority's head of power to regulate certain activities in the Region.

#### **Commonwealth Heritage places**

To protect the heritage values of Commonwealth Heritage Places they own or lease, Australian Government agencies are required to develop heritage strategies, a heritage register, and plans of management for places on the Commonwealth Heritage List in accordance with the Commonwealth Heritage management principles.

#### **Commonwealth reserves\***

Management plans provide for the protection and conservation of Commonwealth reserves. Management plans have a maximum life of 10 years and must set out how the reserves are to be managed. The plans provide certainty about the activities that will be allowed in the reserves and must be consistent with the relevant Australian IUCN Reserve Management Principles which define how the reserves should be managed.

*\*The Governor-General must not make a Proclamation under Section 344 of the EPBC Act declaring an area in the Great Barrier Reef Region to be a Commonwealth reserve under that Act.*



## **Gazettal instrument – The Zoning plan**

### **Gazettal instrument**

The Commonwealth Heritage List is a list of the natural, cultural and historic heritage places on Commonwealth land or in Commonwealth waters, or owned or managed by the Commonwealth Government. The Commonwealth Heritage values of a Commonwealth Heritage place are the Commonwealth Heritage values of the place included in the Commonwealth Heritage List for the place.

Heritage values of a place include the place's natural and cultural environment, having aesthetic, historic, scientific or social significance, or other significance for current and future generations of Australians. To be listed as National Heritage values it must be able to be shown that they reach the level of significance of 'outstanding value to the nation' against listed criteria and that this must be able to be established through a comparative analysis. If a place is determined to be included in the National Heritage list then the Environment Minister must by instrument published in the Gazette:

- the assessed place or part of the assessed place; and
- the National Heritage values of the assessed place, or that part of the assessed place, that are specified in the instrument

### **The Zoning Plan**

The primary regulatory instrument under the GBRMP Act through which protection of matters protected by Part 3 of the EPBC Act are given effect is the *Great Barrier Reef Marine Park Zoning Plan, 2003* (Zoning Plan). The Zoning Plan divides the Great Barrier Reef into eight zones setting out which activities can occur in each zone. Each zone has specific conservation and management objectives for the protection of the environmental values of that zone, including matters of national environmental significance.

The GBRMP Regulations provide for a system of permits, which are required for specified activities likely to impact on the environmental values of each zone. For each permit application the Authority undertakes an assessment of the nature and scale of activities and makes a determination based on the acceptability of those activities on the environment.

## **Bioregional plans under the EPBC Act**

Bioregional plans describe the environment and conservation values of the various regions, identifies and characterises the pressures affecting these conservation values, identifies regional priorities and outlines strategies to address them, and provides advice to decision-makers and people planning to undertake activities in the various regions in relation to their conservation values. Under s176(5) of the EPBC Act, the Minister for the Environment must have regard to a bioregional plan in making any decision under the EPBC Act to which the plan is relevant.

Note: There are no current Bioregional Plans that are applicable to the Marine Park or Great Barrier Reef World Heritage Area.

Where a National Heritage place is in a state or territory, the Australian Government must endeavor to ensure that a management plan is prepared and implemented in cooperation with the relevant state or territory government. The Minister for the Environment is responsible for preparing plans of management for National heritage places in Commonwealth marine areas.

Chapter 10 describes the future risk of impacts on values and attributes relevant to each matters of national environmental significance.

## **4.6 Safeguards under the permission system**

Under the EPBC Act, an action or class of actions should not be approved if the action/s is inconsistent with:

- Australia's obligations under the World Heritage Convention
- National Heritage Management Principles (Schedule 5B of the EPBC Regulations)
- an agreement to which the Commonwealth is party in relation to the National Heritage place
- the Australian World Heritage management principles (Schedule 5 of the EPBC Regulations)
- a plan that has been prepared for the management of the declared World Heritage property under s316 or as described in s321 of the EPBC Act.
- a plan that has been prepared for the management of the National Heritage place under s324S or as described in s324 of the EPBC Act. (EPBC Act, 146H)

An action or class of actions also cannot be approved if the action/s would have an unacceptable impact on a World Heritage property, the environment of the Great Barrier Reef Marine Park or a National Heritage place.

The Authority is committed to:

- the outstanding universal value of the Great Barrier Reef World Heritage property and national heritage places being identified, protected, conserved, presented and transmitted to future generations.
- the environmental, biodiversity and heritage values of the Great Barrier Reef Marine Park being protected and conserved; its use is ecologically sustainable; and the community is engaged in its protection and management
- maintaining and enhancing the environment of a Commonwealth marine area.

The Authority will make decisions consistent with the objects of the GBRMP Act which includes the main object at section 2A(1) to 'provide for the long-term protection and conservation of the environment, biodiversity and heritage values of the Great Barrier Reef Region'. Environment has the same meaning as in the EPBC Act and is defined as including (among other things) ecosystems and their constituent parts, including people and communities, natural and physical resources, and the qualities and characteristics of locations, places and areas. This necessarily includes matters of national environmental significance.

In making decisions that are consistent with the objects of the GBRMP Act, the Authority must consider the potential impacts of the conduct proposed to be permitted by the permission on the environment and on the social, cultural and heritage values of the Marine Park or a part of the Marine Park.

Permit application assessments have regard to the significant impact criteria in the EPBC Act Significant Impact Guidelines 1.1. Assessment of the level of acceptability will also be informed by the permit applicant's measures to address the Authority's avoid-mitigate-offset-net benefit-adaptive management hierarchy for the assessment of impacts as outlined in their application and subsequent assessment report(s).

Under the permission system the Authority will make decisions consistent with the GBRMP Act:

- Consideration will be given to:
  - Australia's obligations under the World Heritage Convention.
  - Australia's World Heritage management principles.
  - a plan that has been prepared for the management of the Great Barrier Reef declared World Heritage property under Section 316 or as described in section 321<sup>8</sup> of the EPBC Act.
  - World Heritage values of the Great Barrier Reef World Heritage property.
  - maintaining and enhancing the environment of a Commonwealth marine area
  - a plan of management for a Commonwealth reserve, and the IUCN reserve management principles.
  - a plan of management for a Commonwealth Heritage place, and the Commonwealth Heritage management principles.
  - any relevant bioregional plans.
  - preservation of the national heritage values of a national heritage place.
  - Australia's national heritage management principles.
  - a plan that has been prepared for the management of the Great Barrier Reef national heritage place under Section 324S of the EPBC Act.
- The Authority maintains policy guidance that applicants should have regard to in the preparation of their Permit Applications. Policy guidance will be updated to include explicit reference to:
  - consideration of all attributes relevant to matters of national environmental significance
  - consideration of statements of outstanding universal value, world heritage and national heritage criteria
  - relevant GBRMP Act and EPBC Act guidance documents, including but not limited to significant impact guidelines for outstanding universal value.

In undertaking assessments, the Authority will have regard to relevant policy documents, guidelines, online data sources, Statement of Outstanding Universal Value (OUV) and plans of management on the Department of the Environment website.

The Authority's assessment process also ensures an assessment of the proponent's measures to avoid, mitigate, offset and provide for net-benefit and adaptive management of potential impacts to matters of national environmental significance.

All permitted activities must be carried out in a manner that is consistent with the objectives of the GBRMP Act, ecosystem-based management and the principles of ecologically sustainable use.

---

<sup>8</sup>The GBRMP Zoning Plan 2003 is the primary planning instrument for the conservation and management of the Marine Park. Noting the existence of the Zoning Plan, Section 321 of the EPBC Act does not require the preparation and implementation of a separate plan for managing the GBR World Heritage property in relation to so much of the property as is in the GBR Marine Park.

## 4.7 Case study - an activity with the potential to impact the Great Barrier Reef World Heritage Property, National Heritage Places, the Great Barrier Reef Marine Park and Commonwealth Marine Area

### Scenario:

Dredging of a shipping channel is proposed for a hypothetical jetty at Flinders Island (within the Flinders Group) in the Far Northern Management Area of the Great Barrier Reef Marine Park. The dredging itself will take place within a Conservation Park Zone in Fly Channel and the preferred disposal site is proposed for a General Use Zone between Stanley Island and Clack Island (within the Designated Shipping Area) or 20km north of Stanley Island within the Commonwealth Marine Area. The proposal includes the construction of a 100 metre jetty including pile driving, dredging of 250,000 cubic metres and disposal at a designated dredge material disposal area. The dredging and disposal campaign will run for three weeks and operate 24 hours a day. A shipwreck registered on the National Heritage property database lies within the potential impact area. The shipwreck has cultural and indigenous cultural values. Furthermore the Great Barrier Reef is a listed National Heritage Property and the Cape York Peninsula is currently a nominated place. The proposed project triggers the controlling provisions of the Great Barrier Reef World Heritage Area, the Great Barrier Reef Marine Park, Commonwealth Marine Area and the National Heritage, Listed Threatened Species and Ecological Communities and Listed Migratory Species

Potential impacts of the activity include:

- Sediments from catchment run-off (e.g. water quality and impacts on the benthos)
- Exotic species and diseases
- Wildlife disturbance (e.g. Interactions with protected, threatened and listed migratory species, species displacement from key ecological functions such as foraging, resting, breeding)
- Amenity and aesthetic impacts (e.g. from coastal reclamation)
- Urban and industrial discharge (e.g. change in nutrient, sediment and other pollutant loads)
- Marine debris
- Noise pollution
- Light impacts - artificial (e.g. impacts on light dependent species such as corals and seagrasses)
- Vessel strike on wildlife
- Waste discharge from vessels
- Spills (e.g. oil, chemical)
- Permanent and temporary loss of benthic habitats (e.g. from dumping and resuspension of dredge material)
- Indigenous heritage impacts
- Cultural heritage impacts
- Impact on commercial fisheries
- Impact on recreational fisheries
- Impact on other users
- Impact on Princess Charlotte Bay Special Management Area.

These impacts, their cumulative effects, and past and present effects on values, including on matters of national environmental significance, are described in detail in Chapter 6.

The permission system requires the Authority to analyse the potential impacts of the proposed activity (direct, indirect and cumulative) on matters of national environmental significance (in this case the Great Barrier Reef World Heritage Area, the Great Barrier Reef Marine Park, a Commonwealth Marine Area and a properties listed on the National Heritage register) and ensure impacts are identified, minimised, mitigated and acceptable.

The types of studies and fields of investigation which would usually occur in this scenario include: Benthic surveys, hydrodynamic modelling, EPBC Act Protected Matters Report, Dredge Sediment Analysis Plan and Dredge Sediment Characterisation Report, Infauna Monitoring Report, Dredge Plume Dispersion Study and Impact Assessment, Dredge Disposal Options Assessment Report, Marine Ecology Baseline Report, Underwater Construction and Operational Noise Impact Assessment, Megafauna Report and Seagrass Report, and a Cultural and Heritage Survey Report and Construction Drawings and Environmental Management Plans for proposed activities. These reports and studies would usually be contained within an Environmental Impact Statement or Public Environment Report, however at times some of the preliminary information is provided with the initial application.

**Table 5 How a permit application would meet EPBC Act requirements for World Heritage Properties the Great Barrier Reef Marine Park, National Heritage Places, and Commonwealth Marine Areas**

Outcome from EPBC Act	How World Heritage Properties, National Heritage Places, the Great Barrier Reef Marine Park and Commonwealth Marine Areas are considered under the Authority's permission system
<p><b>SCOPING:</b> The purpose of the scoping phase is to proactively discuss the initial concepts of a proposed action in order to explain regulatory arrangements and identify potential mitigations. For actions which will likely have a significant impact on the Marine Park, including matters of national environmental significance, reference is made to relevant GBRMP Act and EPBC Act guidance material. A Permit Application Assessment Fee must be paid before the accompanying Permit Application Form is considered. The Permit Application Form must contain sufficient information to enable a clear understanding of the intended action</p>	
<p><b>Activities with a likely significant impact must be assessed under legislation with regards to matters of national environmental significance.</b></p>	<p><b>Mechanism (legislative &amp; policy):</b></p> <p>Section 318 of the EPBC Act requires that a Commonwealth agency must not authorise any person to do anything that may contravene a plan made under Section 316 for management of a World Heritage property. As a Commonwealth statutory authority, the Authority must comply with this requirement.</p> <p>Section 324U of the EPBC Act requires that a Commonwealth agency must not authorise any person to do anything that may contravene a plan made under Section 324S for management of a National Heritage place. If no plan made under Section 324S exists, a Commonwealth agency must take all reasonable steps to ensure that acts relating to a property are not inconsistent with the National Heritage management principles (Schedule 5B, EPBC Regulations). As a Commonwealth statutory authority, the Authority must comply with this requirement.</p> <p>Section 341V of the EPBC Act requires that a Commonwealth agency must not authorise any person to do anything that may contravene a plan made under Section 341S for management of a Commonwealth Heritage place. If there is no Section 341S plan, a Commonwealth agency must take all reasonable steps to ensure that acts relating to the property are not inconsistent with the Commonwealth Heritage management principles (Schedule 7B, EPBC Regulations).</p> <p>When conducting a whole of environment assessment other protected matters potentially relevant to the Great Barrier Reef are also assessed (e.g. Ramsar, listed threatened species). In undertaking a whole of the environment assessment, the Authority will ensure levels of acceptability for the Marine Park, including matters of national environmental significance, will be based on the Program's commitments in Table 10.</p> <p>GBRMP Regulations 88Q (mandatory considerations) and 88R (discretionary considerations) outline the Authority's permit assessment considerations.</p> <p>Under reg 88Q(a) of the GBRMP Regulations, the Authority must consider the potential impacts of the conduct proposed to be permitted by the permission on the environment and on the social, cultural and heritage values of the Marine Park or a part of the Marine Park. This relevantly includes matters of national environmental significance.</p> <p>Under Regulation 88E the Authority would seek further information for a permit application that does not provide sufficient detail to evaluate whether an activity is having a significant impact. This includes:</p> <ul style="list-style-type: none"> <li>• a description of the relevant matters of national environmental significance attributes likely to be affected by the proposed activity</li> <li>• the direct, indirect and cumulative impacts of the proposed activity</li> <li>• the risks of those impacts to matters of national environmental significance.</li> </ul> <p>Failure to provide sufficient detail would result in the application being withdrawn.</p>
<p><b>Program commitment(s):</b></p> <p>The Authority will:</p> <ul style="list-style-type: none"> <li>• require consideration of relevant EPBC policy documents, including relevant significant impact guidelines and reporting tools, in the scoping phase of the permission assessment and decision process.</li> <li>• require consideration of prudent and feasible alternatives which may have a lower impact on the environment and amend its policy documentation to include reference to this requirement.</li> <li>• consider any relevant international conventions, Commonwealth State agreements, plans of management, recovery plans, wildlife conservation plans, threat abatement plans, approved conservation advice under the EPBC Act, EPBC Act guidelines, policies and information in assessing and determining permissions under its permission system.</li> </ul>	



Outcome from EPBC Act	How World Heritage Properties, National Heritage Places, the Great Barrier Reef Marine Park and Commonwealth Marine Areas are considered under the Authority's permission system
	<ul style="list-style-type: none"> <li>consider relevant Department of the Environment policy documents, guidelines, plans of management and other online data sources available on the Department's website in its scoping, assessment and decision-making process.</li> <li>implement guidelines to assist applicants to determine whether an action is likely to have a significant impact on the Great Barrier Reef Marine Park matters of national environmental significance. The Authority will seek to have these guidelines approved by the Minister.</li> <li>amend policies and guidance material used in the determination of permissions so they explicitly require the consideration of matters of national environmental significance and relevant attributes, including: <ul style="list-style-type: none"> <li>biodiversity and environmental processes</li> <li>Indigenous and historic heritage values</li> <li>community benefits derived from the environment, including those not easily represented or measured (such as aesthetic values).</li> </ul> </li> <li>develop guidance material to assist in determining the acceptability of impacts. Criteria shall include consistency of assessment outcomes with regulatory objectives, outcomes for the Marine Park, including matters of national environmental significance and relevant GBRMP Act and EPBC Act guidelines and standards.</li> <li>develop specific guidance documents for notifications under Part 5 of the Zoning Plan to make reference to consideration of matters of national environmental significance under the EPBC Act.</li> </ul> <p>The Authority will require applicants to consider:</p> <ul style="list-style-type: none"> <li>the impacts of actions on matters of national environmental significance and relevant attributes and environmental processes as described in the Strategic Assessment Report, and as amended in accordance with the Authority's policy review procedures.</li> <li>impacts (including direct, indirect and cumulative impacts) arising from the development action in the context of all impacts affecting the matters of national environmental significance.</li> </ul> <p><b>Case study:</b></p> <ul style="list-style-type: none"> <li>Under the Zoning Plan the dredging and dumping of spoil is considered to be 'carrying out works' and the construction of a jetty to be the 'operation of a facility'. These types of activity can only occur in certain parts of the Marine Park – namely areas where the purpose of the activity is consistent with the objective of the relevant zone and only with the written permission of the Authority.</li> <li>The Authority determines that proposed activity is consistent with the objectives of the relevant zone and that the proponent requires a permit from the Authority for the carrying out of works.</li> <li>For actions which may have a significant impact on the Marine Park, including matters of national environmental significance, reference is made to: <ul style="list-style-type: none"> <li>GBRMP Act, Regulations, Zoning Plan, plans of management and relevant policy documents including the Authority's environmental impact management, position statements and guidelines;</li> <li>EPBC Act policy documents, including relevant significant impact guidelines, the EPBC Act Species Profile and Threats Database (SPRAT), listing advice policy statements, recovery plans, conservation advices, threat abatement plans and wildlife conservation plans available on the Department of the Environment website.</li> </ul> </li> <li>Information available through online data sources hosted by the Department of the Environment, such as the Environment Reporting Tool, augments local knowledge in determining whether matters of national environmental significance are likely to occur in the area of interest.</li> <li>EPBC Act Significant Impact Guidelines 1.1, and OUV guidelines as they relate to World Heritage Properties, the Great Barrier Reef Marine Park, National Heritage Places and Commonwealth Marine Environment, includes consideration of the conservation status under the EPBC Act. The Authority, in evaluating whether the activity is consistent with the objects of the GBRMP Act, will take into account either EPBC Act species specific significance guidelines or, where none exist, the following generic</li> </ul>

Outcome from EPBC Act	How World Heritage Properties, National Heritage Places, the Great Barrier Reef Marine Park and Commonwealth Marine Areas are considered under the Authority's permission system
	<p>considerations from the EPBC Act Significant Impact Guidelines 1.1.</p> <ul style="list-style-type: none"> <li>An action is likely to have a significant impact on the World Heritage Values if there is a real chance or possibility that it will: <ul style="list-style-type: none"> <li>cause one or more of the World Heritage values/attributes to be lost</li> <li>cause one or more of the World Heritage values/attributes to be degraded or damaged, or</li> <li>cause one or more of the World Heritage values/attributes to be notably altered, modified, obscured or diminished.</li> <li>Impact on the integrity of the property</li> </ul> <p>An action is likely to have a significant impact the National Heritage values of a National Heritage place if there is a real chance or possibility that it will cause:</p> <ul style="list-style-type: none"> <li>one or more of the National Heritage values to be lost</li> <li>one or more of the National Heritage values to be degraded or damaged, or</li> <li>one or more of the National Heritage values to be notably altered, modified, obscured or diminished.</li> </ul> </li> <li>An action is likely to have a significant impact on the Great Barrier Reef Marine Park if there is a real chance or possibility that it will cause: <ul style="list-style-type: none"> <li>modify, destroy, fragment, isolate or disturb an important, substantial, sensitive or vulnerable area of habitat or ecosystem component such that an adverse impact on marine ecosystem health, functioning or integrity in the Great Barrier Reef Marine Park results</li> <li>have a substantial adverse effect on a population of a species or cetacean including its life cycle (for example, breeding, feeding, migration behaviour, life expectancy) and spatial distribution</li> <li>result in a substantial change in air quality or water quality (including temperature) which may adversely impact on biodiversity, ecological health or integrity or social amenity or human health</li> <li>result in a known or potential pest species being introduced or becoming established in the Great Barrier Reef Marine Park</li> <li>result in persistent organic chemicals, heavy metals, or other potentially harmful chemicals accumulating in the marine environment such that biodiversity, ecological integrity, or social amenity or human health may be adversely affected, or</li> <li>have a substantial adverse impact on heritage values of the Great Barrier Reef Marine Park, including damage or destruction of an historic shipwreck.</li> </ul> </li> <li>An action is likely to have a significant impact on the environment in a Commonwealth marine area if there is a real chance or possibility that the action will: <ul style="list-style-type: none"> <li>result in a known or potential pest species becoming established in the Commonwealth marine area</li> <li>modify, destroy, fragment, isolate or disturb an important or substantial area of habitat such that an adverse impact on marine ecosystem functioning or integrity in a Commonwealth marine area results</li> <li>have a substantial adverse effect on a population of a marine species or cetacean including its life cycle (for example, breeding, feeding, migration behaviour, life expectancy) and spatial distribution</li> <li>result in a substantial change in air quality<sup>4</sup> or water quality (including temperature) which may adversely impact on biodiversity, ecological integrity; social amenity or human health</li> <li>result in persistent organic chemicals, heavy metals, or other potentially harmful chemicals accumulating in the marine environment such that biodiversity, ecological integrity, social amenity or human health may be adversely affected, or</li> <li>have a substantial adverse impact on heritage values of the Commonwealth marine area, including damage or destruction of an historic shipwreck.</li> </ul> </li> <li>Information available through online data sources hosted by the Department of the Environment, such as the Environment Reporting Tool, augments local knowledge in determining whether matters of national environmental significance are likely to occur in the area of interest.</li> </ul>

Outcome from EPBC Act	How World Heritage Properties, National Heritage Places, the Great Barrier Reef Marine Park and Commonwealth Marine Areas are considered under the Authority's permission system
	<ul style="list-style-type: none"> <li>• The Authority has delegations for other Commonwealth legislation that may be triggered by this proposal such as the <i>Environment Protection (Sea Dumping) Act 1981</i>, and the <i>Sea Installations Act 1987</i>.</li> <li>• The Authority and the Queensland Parks and Wildlife consider the significant impacts to the environment and the outstanding universal value of the Great Barrier Reef Marine Park Authority, as per the Department of the Environment's significant impact guidelines. <ul style="list-style-type: none"> <li>○ a) ecosystems and their constituent parts, including people and communities; and</li> <li>○ (b) natural and physical resources; and</li> <li>○ (c) the qualities and characteristics of locations, places and areas; and</li> <li>○ (d) heritage values of places; and</li> <li>○ (e) the social, economic and cultural aspects of a thing mentioned in paragraph (a), (b), (c) or (d).</li> </ul> </li> <li>• Examples of direct impacts that are considered may include (but are not limited to) disturbance to coral, sea grasses, benthic habitats, cultural and heritage sites, water quality, altered hydrodynamics, impacts to species and their habitats.</li> <li>• Examples of indirect impacts that are considered may include (but are not limited to) downstream impacts on water quality, increased boating, increased recreational activity, impacts on commercial fisheries, impacts to migratory species through removal of habitat, noise and light pollution and increase in shipping activity.</li> <li>• The Authority and Queensland Parks and Wildlife Service determine the level of risk associated with the action by undertaking a risk assessment of the proposed action against the Authority's policy and procedures. This particular scenario would be undertaken using the Authority's highest levels of assessment due to the sensitive receiving environment and the possibility of attracting public interest.</li> <li>• The Authority uses all the above mentioned information to determine whether the proposed action is likely to be consistent with the objects of the GBRMP Act.</li> <li>• The Authority determines, in relation to the potential impacts: <ul style="list-style-type: none"> <li>○ Whether the proposal should be permitted in the Marine Park, World Heritage Area, Commonwealth Marine Area</li> <li>○ If further information is required to assess potential impacts</li> <li>○ avoidance, mitigation, offset, net-benefit and adaptive management measures</li> <li>○ If the action can be permitted in the Marine Park, what level of assessment is required and what the controlling provisions are.</li> </ul> </li> <li>• Under this scenario, the Authority determines that the proposed activity is likely to have a significant impact on values/attributes that contribute to the Outstanding Universal Value of the Great Barrier Reef and elements of the Great Barrier Reef Marine Park and requires assessment by an Environmental Impact statement (EIS), which also requires an Environmental Management Plan (EMP) to be prepared.</li> </ul>
<p><b>Applicant is required to provide adequate information to enable the decision maker to make a decision.</b></p> <ul style="list-style-type: none"> <li>- <b>The applicant describes impacts to matters of national environmental significance</b></li> </ul>	<p><b>Mechanism (legislative &amp; policy):</b></p> <p>GBRMP Regulation 88A enables the Authority to approve permit application forms. The Authority can vary the approved application form to meet legal and management responsibilities.</p> <p>GBRMP Regulation 88E enables the Authority to request further information, for the purpose of assessing an application, and for such information to be provided within 20 business days (or for a longer period, as determined by the Authority) or the application is taken to have been withdrawn.</p> <p>GBRMP Regulation 103 provides that persons shall not furnish false information in a permit application.</p> <p>Failure to provide the information that did not adequately describe the risk from the activity to the environment and OUV in detail sufficient to evaluate all impacts to assess whether the activity is having a significant impact and determine the level of assessment under the Program (e.g. by PER or EIS) would result in the application being withdrawn.</p>

Outcome from EPBC Act	How World Heritage Properties, National Heritage Places, the Great Barrier Reef Marine Park and Commonwealth Marine Areas are considered under the Authority's permission system
	<p><b>Program commitment(s):</b></p> <p>The Authority will:</p> <ul style="list-style-type: none"> <li>• amend its Permit Application Form to include a section which specifically addresses matters of national environmental significance.</li> <li>• seek to pursue amendments to the Great Barrier Reef Marine Park Regulations 1983 to explicitly outline the levels of assessment under the permission system.</li> <li>• update its permissions guidance material to outline the general information requirements for applicants according to the level of assessment.</li> </ul> <p><b>Case study:</b></p> <ul style="list-style-type: none"> <li>• The applicant will have access to guidance material developed by the Authority relating to the Marine Park, including matters of national environmental significance, that proponent's should have regard to in the preparation of their application.</li> <li>• An application must be submitted to the Authority. It must provide sufficient information regarding the proposed activity so assessors can clearly understand what the intended activity is, where it is to be carried out, timing and the likely nature and scale of potential environmental impacts.</li> <li>• Descriptions must identify any impacts and risk to the environment and OUV from the proposed activity including: <ul style="list-style-type: none"> <li>○ a) ecosystems and their constituent parts, including people and communities; and</li> <li>○ (b) natural and physical resources; and</li> <li>○ (c) the qualities and characteristics of locations, places and areas; and</li> <li>○ (d) heritage values of places; and</li> <li>○ (e) the social, economic and cultural aspects of a thing mentioned in paragraph (a), (b), (c) or (d).</li> </ul> </li> <li>• Descriptions must identify any other matters of national environmental significance, which form part of the environment in the impacted area and include information consistent with the Program's commitments.</li> <li>• The description of the activity and receiving environment's characteristics must be sufficient to inform consideration of the 'nature and scale' of the impacts by the Authority in its assessment.</li> <li>• Cumulative impact considerations of the area and adjacent areas.</li> <li>• If the information is not initially provided in the permit application, the proponent will be requested to provide additional information (Regulation 88E) to inform the Authority's assessment against the mandatory and discretionary considerations. The proponent will be required to provide information on the following in the application or further information request: Any elements of the proposed activity relevant to impacts on matters of national environmental significance must be sufficiently described including operational details relevant to the dredging and disposal activity: <ul style="list-style-type: none"> <li>○ The timing and duration of the dredging and disposal activity in relation to the use of known habitat, including biologically important areas/populations (for example the breeding, hatching or migration periods for sea turtles). The application will require information on all impacts including but not limited to: <ul style="list-style-type: none"> <li>▪ Describe the nature, scale, timing and duration of the proposed dredging, disposal and pile driving activity, including associated activities and actions that occur in relation to the proposal.</li> <li>▪ Provide specific information on potential impacts of the jetty structure, the dredging and the disposal activity on protected, vulnerable or iconic species, coastal processes, water quality, hydrodynamics, noise, light, air quality, social values, cultural values and heritage values.</li> <li>▪ That all alternatives to avoid, mitigate, offset and provide for net-benefit and adaptive management of impacts have been considered.</li> <li>▪ details of other disturbances such as light, increased vessel movements, permanent and or temporary placement of objects related to the project.</li> <li>▪ Include any potential expansion of the site's infrastructure in the</li> </ul> </li> </ul> </li> </ul>



Outcome from EPBC Act	How World Heritage Properties, National Heritage Places, the Great Barrier Reef Marine Park and Commonwealth Marine Areas are considered under the Authority's permission system
	<p>future, and show the potential scale of the final design and operation of the facility at the site.</p> <ul style="list-style-type: none"> <li>▪ Identify the matters of national environmental significance potentially at risk from the proposal.</li> <li>▪ Identify areas that are significant to supporting rare or threatened species. Descriptions must include any threatened species, include a sufficient description of the threatened species which may be at risk from the activity, and provide sufficient information for the Authority to identify potentially impacted habitats, including the significance of the area for each species potentially affected. This will include a description of the occurrence of these species within the project area and within any areas that may potentially be affected by upstream and downstream impacts such as water quality, noise or light.</li> <li>▪ quantification of any benthic habitat loss associated with the project,</li> <li>▪ quantification (i.e. potential dredge plume footprint) and duration of water quality impacts,</li> <li>▪ discharges or emissions,</li> <li>▪ and any other impacts or activities that have the potential to adversely impact directly on individuals, important populations or their habitats.</li> </ul> <ul style="list-style-type: none"> <li>• The application would need to demonstrate that appropriate contingency plans, including appropriate emergency response measures are in place to ensure that the applicant would be able to effectively respond to an unplanned oil spill or discharge event. These plans would form part of the Environment Management Plan required for this activity to proceed (see Monitoring Compliance and Auditing below). The Environmental Management Plan will evolve throughout the assessment process and is not final at this stage of the process.</li> <li>• If an application does not include sufficient information for the purposes of assessing the likely nature and scale of environmental impacts additional information or documents must be provided at the request of the Authority.</li> </ul>
<p><b>ASSESSMENT:</b> The purpose of the assessment phase is to enable an informed decision on whether or not a permit will be granted to undertake an action within the Great Barrier Reef Marine Park. During the assessment phase, the Authority considers criteria outlined in the <i>Great Barrier Reef Marine Park Regulations 1983</i>, and relevant GBRMP Act and EPBC Act guidance material</p>	
<p><b>Public consultation occurs in a transparent manner and outcomes are taken into account.</b></p>	<p><b>Mechanism (legislative &amp; policy):</b></p> <p>GBRMP Regulations 88Q (mandatory considerations) and 88R (discretionary considerations) outline the Authority's permit assessment considerations.</p> <p>Under Regulation 88E the Authority would seek further information for a permit application that does not provide sufficient detail to evaluate whether an activity is having a significant impact. This includes:</p> <ul style="list-style-type: none"> <li>• a description of the relevant matters of national environmental significance attributes likely to be affected by the proposed activity</li> <li>• the direct, indirect and cumulative impacts of the proposed activity</li> <li>• the risks of those impacts to matters of national environmental significance.</li> </ul> <p>Regulation 88D of the GBRMP Regulations requires applications to be advertised if the Authority considers that granting a permission may restrict the reasonable use by the public of part of the Great Barrier Reef Marine Park.</p> <p>GBRMP Regulations 88D provide that if a permit applicant does not publish an advertisement as required, the application is taken to have been withdrawn.</p> <p>GBRMP Regulation 88Q provides that public comments must be considered by the Authority when making decisions on whether to grant the permit.</p> <p><b>Program commitment(s):</b></p> <p>The Authority will:</p> <ul style="list-style-type: none"> <li>• publish notification of all permit applications on the internet as soon as practicable after receiving a permit application.</li> <li>• seek to pursue amendments to the GBRMP Regulations (in particular regulation 88D) to require advertising of applications where the granting of a permission may restrict the reasonable use by the public of a part of the Marine Park or is likely to have a significant impact on the Marine Park,</li> </ul>

Outcome from EPBC Act	<p><b>How World Heritage Properties, National Heritage Places, the Great Barrier Reef Marine Park and Commonwealth Marine Areas are considered under the Authority's permission system</b></p>
	<p>including on matters of national environmental significance.</p> <ul style="list-style-type: none"> <li>• update relevant policies to explicitly state that appropriate consultation will be undertaken with persons or organisations, including Commonwealth and State government agencies, whose functions, interests or activities could be impacted by the proposed activity.</li> </ul> <p><b>Case study:</b></p> <ul style="list-style-type: none"> <li>• The application is assessed against policy and guidance documents by the permit assessor/project manager who determines whether subject matter expert advice is required.</li> <li>• In this case the proposal may be referred to relevant parties and subject matter experts for further advice e.g. scientific experts within the Authority, James Cook University, Australian Institute of Marine Science Queensland and Commonwealth environment departments.</li> <li>• The Authority also undertakes appropriate consultation with persons or organisations, including Australian and State government agencies, whose functions, interests or activities could be impacted by the proposed activity.</li> <li>• The applicant is required to publish an advertisement for projects under assessment and for the PER/EIS to be made available to the public during a defined public comment period This advertisement must be published in a newspaper circulating in the part of the State of Queensland adjacent to that part of the Marine Park in which the conduct is proposed to occur. The advertisement is to include an invitation for interested persons to make comment on the application within a period of not less than 30 days. The advertisement will also be published on the Authority's website.</li> </ul>
<p><b>The detail and rigour of the assessment process is commensurate with the scale and potential impacts of the activity i.e. risk-based.</b></p>	<p><b>Mechanism (legislative &amp; policy):</b></p> <p>Section 318 of the EPBC Act requires that a Commonwealth agency must not authorise any person to do anything that may contravene a plan made under Section 316 for management of a World Heritage property. As a Commonwealth statutory authority, the Authority must comply with this requirement.</p> <p>Section 324U of the EPBC Act requires that a Commonwealth agency must not authorise any person to do anything that may contravene a plan made under Section 324S for management of a National Heritage place. If no plan made under Section 324S exists, a Commonwealth agency must take all reasonable steps to ensure that acts relating to a property are not inconsistent with the National Heritage management principles (Schedule 5B, EPBC Regulations). As a Commonwealth statutory authority, the Authority must comply with this requirement.</p> <p>Section 341V of the EPBC Act requires that a Commonwealth agency must not authorise any person to do anything that may contravene a plan made under Section 341S for management of a Commonwealth Heritage place. If there is no Section 341S plan, a Commonwealth agency must take all reasonable steps to ensure that acts relating to the property are not inconsistent with the Commonwealth Heritage management principles (Schedule 7B, EPBC Regulations).</p> <p>When conducting a whole of environment assessment other protected matters potentially relevant to the Great Barrier Reef are also assessed (e.g. Ramsar, listed threatened species). In undertaking a whole of the environment assessment, the Authority will ensure levels of acceptability on the Marine Park, including matters of national environmental significance will be based on the Program's commitments in Table 10.</p> <p>GBRMP Regulations 88Q (mandatory considerations) and 88R (discretionary considerations) outline the Authority's permit assessment considerations.</p> <p>The Authority's Environmental Impact Assessment Policy and Environmental Assessment and Management Risk Assessment Guidelines (developed in accord with the Australian Standards Environmental Risk Management Handbook HB203:2006) provide guidance in the assessment of risk including analysis of the likelihood and consequence of identified hazards being realised to obtain a Hazard Risk Grade.</p> <p><b>Program commitment(s):</b></p> <p>The Authority will:</p> <ul style="list-style-type: none"> <li>• develop and maintain regionally-based standards for ecosystem health (including water quality) that support best management practices, and may be used to inform:</li> </ul>

Outcome from EPBC Act	How World Heritage Properties, National Heritage Places, the Great Barrier Reef Marine Park and Commonwealth Marine Areas are considered under the Authority's permission system
	<ul style="list-style-type: none"> <li>○ the assessment of point source and cumulative impacts on the Marine Park, including matters of national environmental significance</li> <li>○ actions to mitigate impacts and cumulative impacts on the Marine Park, including matters of national environmental significance</li> <li>○ outcomes-based decision-making and considerations for determining 'acceptability' of impacts on the Marine Park, including matters of national environmental significance, based the commitments in Table 10 of the Program Report</li> <li>• implement guidelines requiring consideration of regionally-based standards for ecosystem health to improve the assessment of impacts including cumulative impacts on the Region's values.</li> <li>• facilitate development of a dredging and dredge material disposal policy</li> <li>• improve the hydrodynamic modelling guidelines so they take account of longshore drift, resuspension and inter-annual current variability.</li> <li>• develop guidance material to be used in the determination of permissions to require the consideration of the impacts of underwater noise, particularly on at-risk species and inshore species.</li> <li>• integrate relevant components, as required, of the Department of the Environment's EPBC Act assessment templates and manuals into its policy documents as they are reviewed and updated.</li> </ul> <p><b>Case study:</b></p> <ul style="list-style-type: none"> <li>• The Authority's assessment will consider habitat types, seasonality, important populations and whether there are critical life stages such as feeding, recruitment, breeding or migration that are likely to be affected.</li> <li>• The Authority's assessment must consider any relevant information from all relevant plans. The Authority will have regard to relevant policy documents, guidelines, Statement of Outstanding Universal Value (OUV) and plans of management on the Department of the Environment website.</li> <li>• Assessments must take into account and assess the impacts on any other matters of national environmental significance potentially affected (e.g. listed migratory species, as these form part of the 'environment').</li> <li>• The assessment must identify the specific impacts of the activity on the environment and OUV. This will include an evaluation of impacts to determine whether the activity will result in unacceptable impacts, for example, whether it will: <ul style="list-style-type: none"> <li>○ modify, destroy, fragment, isolate or disturb an important, substantial, sensitive or vulnerable area of habitat or ecosystem component such that an adverse impact on marine ecosystem health, functioning or integrity in the Great Barrier Reef Marine Park results</li> <li>○ have a substantial adverse effect on a population of a species or cetacean including its life cycle (for example, breeding, feeding, migration behaviour, life expectancy) and spatial distribution</li> <li>○ result in a substantial change in air quality or water quality (including temperature) which may adversely impact on biodiversity, ecological health or integrity or social amenity or human health</li> <li>○ result in a known or potential pest species being introduced or becoming established in the Great Barrier Reef Marine Park</li> <li>○ result in persistent organic chemicals, heavy metals, or other potentially harmful chemicals accumulating in the marine environment such that biodiversity, ecological integrity, or social amenity or human health may be adversely affected</li> <li>○ have a substantial adverse impact on heritage values of the Great Barrier Reef Marine Park, including damage or destruction of an historic shipwreck</li> <li>○ have, or is likely to have a significant impact on the environment of the Commonwealth Marine Area</li> <li>○ one or more of the World/National Heritage values to be lost</li> <li>○ one or more of the World/National Heritage values to be degraded or damaged</li> <li>○ one or more of the World/National Heritage values to be notably altered, modified, obscured or diminished.</li> </ul> </li> <li>• The assessment must clearly demonstrate the risks of impacts arising from</li> </ul>

Outcome from EPBC Act	How World Heritage Properties, National Heritage Places, the Great Barrier Reef Marine Park and Commonwealth Marine Areas are considered under the Authority's permission system
	<p>the activity impacting directly and indirectly on the environment and OUV. The assessment must demonstrate that all risks are acceptable. This will require information on the extent of potential impacts of the dredging and disposal campaign and the extent, severity and duration of the plume footprint upon sensitive receptors. Once the plume footprint has been established the proponent must demonstrate that the reduction in water quality will not have unacceptable impacts on sensitive receptors such as coral, seagrass and the shipwreck. Any impacts on these sensitive receptors could have long-term impacts on those threatened and migratory species that depend on them. Artificial lighting from the jetty construction and the noise from pile driving would also need to be considered, as would all other potential impacts outlined and/or arising from the types of impacts described in the permit application.</p> <ul style="list-style-type: none"> <li>• The assessment would also specifically address the likelihood of occurrence of accidents and incidents and would need to demonstrate that they would be low risk.</li> <li>• Permit assessors will have regard to, in preparing the permit assessment, relevant policy documents, guidelines, Position Statements, Statement of Outstanding Universal Value (OUV) and plans of management on the Department of the Environment website. Levels of acceptability will be based on the Program's commitments in Table 10, which draw on the EPBC Act's standards. <ul style="list-style-type: none"> <li>◦ The Authority will also have regard to relevant policies and guidelines such as the Significant Impact Guidelines 1.1<sup>9</sup>, offsets and cumulative impact policies where appropriate.</li> </ul> </li> <li>• The Authority will document all potential direct and indirect impacts on the environment and OUV in their assessment report. Specifically the assessment addresses the impacts: <ul style="list-style-type: none"> <li>◦ from the activities such as pile driving, dredging and disposal, the impacts on habitat, populations and water quality, discharges, the use of artificial lighting, noise and vessel movements resulting in collision risk will be acceptable at all times, and</li> <li>◦ that feasible alternative options to avoid, mitigate, offset and provide for net-benefit and adaptive management of potential risks have been presented and assessed. This includes alternatives to the project itself, or to the layout of the project.</li> </ul> </li> <li>• The Authority's assessment must contain any relevant legislative or other controls that are necessary for managing the impacts to the environment and OUV. In establishing performance outcomes and standards the Authority must consider any outcomes, and targets and performance criteria in relevant statutory plans.</li> </ul>
<p>EIS process follows 'avoid/mitigate/offset/enhance principles'. Any residual significant impacts are offset appropriately.</p>	<p><b>Mechanism (legislative &amp; policy):</b></p> <p>GBRMP Regulation 88Q(b) requires that the Authority must consider options for monitoring, managing and mitigating the potential impacts of the proposed conduct.</p> <p>GBRMP Regulation 88X -The Authority must, after taking into account the matters that it is required or permitted to take into account under the GBRMP Act and Regulations, make a decision on the application.</p> <p><b>Program commitment(s):</b></p> <p>The Authority will:</p> <ul style="list-style-type: none"> <li>• implement guidelines for the application of Great Barrier Reef offsets to maintain the condition of the Marine Park, including matters of national environmental significance and relevant attributes and environmental processes, where impacts cannot be avoided or mitigated: <ul style="list-style-type: none"> <li>◦ The guidelines will seek to deliver an outcome equivalent to, or better than, the outcome that would apply if the EPBC Act Environmental Offsets Policy were applied.</li> </ul> </li> <li>• implement a Great Barrier Reef net benefit policy to enhance the condition of the Marine Park, including matters of national environmental significance and relevant attributes and environmental processes.</li> <li>• implement guidelines on cumulative impact assessment to inform a</li> </ul>

<sup>9</sup>Commonwealth of Australia (2013). Ibid.



Outcome from EPBC Act	How World Heritage Properties, National Heritage Places, the Great Barrier Reef Marine Park and Commonwealth Marine Areas are considered under the Authority's permission system
	<p data-bbox="679 259 1390 315">transparent, consistent and systematic approach to the assessment of impacts from activities within and adjacent to the Region.</p> <p data-bbox="624 342 751 371"><b>Case study:</b></p> <ul data-bbox="639 378 1437 1402" style="list-style-type: none"> <li data-bbox="639 378 1414 434">• The assessment of the permit application uses an 'avoid, mitigate, offset, net-benefit and adaptive management methodology' of controls.</li> <li data-bbox="639 441 1437 909">• The Authority uses a risk assessment to evaluate the environment and OUV. The risk assessment must analyse the efficacy of controls to avoid, mitigate, offset, provide net-benefit and adaptive management of potential impacts on the environment and OUV, ensuring that the dredging is absolutely necessary (avoid), that the amount of dredging is reduced as much as possible (mitigate) and that any risks associated with the dredging are offset in order to deliver a net environmental benefit. With regards to disposal at sea the proponent must demonstrate that all alternatives to at sea disposal have been considered and that the amount to be disposed of has been reduced as much as possible. The site selection of a disposal site will be critically assessed to ensure that the site with the least potential impacts is utilised whilst still considering the impacts on other users (in this case including commercial and recreational fisheries). With regards to pile driving the proponent must demonstrate that the noise associated with the activity and the artificial light impacts do not impact on threatened and migratory species. Enhancement of conditions for these species is also a consideration of the risk assessment process as is the implementation of net environmental benefits to the region.</li> <li data-bbox="639 916 1430 1128">• The assessment must demonstrate that appropriate avoidance, mitigation, offset, net-benefit and adaptive management measures are in place to reduce impacts on the environment and OUV. These measures must ensure that impacts on the environment and OUV will be acceptable and are not inconsistent with any statutory guidance or plans. The assessment must ensure that identified control measures, such as operational monitoring/environmental site supervisors, Deeds of Agreement and bonds are appropriate for managing impacts.</li> <li data-bbox="639 1135 1422 1402">• The proponent must demonstrate in their assessment documentation that impacts on matters of national environmental significance are acceptable, and that impacts have been reduced to levels where matters of national environmental significance will be maintained and enhanced. Considerations must include consistency of assessment outcomes with: <ul data-bbox="679 1274 1422 1402" style="list-style-type: none"> <li data-bbox="679 1274 935 1303">○ Regulatory objectives</li> <li data-bbox="679 1310 1422 1364">○ Outcomes for matters of national environmental significance including relevant attributes and environmental processes, and</li> <li data-bbox="679 1370 1350 1402">○ Relevant GBRMP Act and EPBC Act guidelines and standards.</li> </ul> </li> </ul>
<p data-bbox="188 1424 1422 1532"><b>DECISION:</b> The purpose of the decision phase is to grant or refuse a permission based on information provided within the assessment phase. A permission may be granted with any conditions appropriate to the attainment of the objects of the GBRMP Act, (which may include conditions relating to the principles of ecologically sustainable use and the protection of matters of national environmental significance).</p>	
<p data-bbox="188 1559 560 1666"><b>EIS process ensures unacceptable impacts on matters of national environmental significance do not occur (including implementation)</b></p>	<p data-bbox="624 1559 983 1588"><b>Mechanism (legislative &amp; policy):</b></p> <p data-bbox="624 1594 1398 1700">The permission system is established by the GBRMP Act as a mechanism to achieve the objects of the Act. The permission system regulates use of the Marine Park in ways consistent with ecosystem-based management and the principles of ecologically sustainable use.</p> <p data-bbox="624 1706 1437 1762">GBRMP Regulation 88ZE A permission may be granted subject to any conditions appropriate to the attainment of the objectives of the Act.</p> <p data-bbox="624 1769 1358 1825">GBRMP Regulation 88ZP A permission may be modified, suspended and revoked.</p> <p data-bbox="624 1832 1422 1960">The Authority may grant or refuse a permission based on the assessment. A permission may be granted with any conditions appropriate to the attainment of the objects of the GBRMP Act and the protection of matters of national environmental significance, the conditions can be modified, suspended or revoked.</p> <p data-bbox="624 1989 895 2018"><b>Program commitment(s):</b></p> <p data-bbox="624 2024 799 2054">The Authority will</p> <ul data-bbox="639 2060 1358 2087" style="list-style-type: none"> <li data-bbox="639 2060 1358 2087">• update the Environmental Impact Management Policy to enable the</li> </ul>

Outcome from EPBC Act	<b>How World Heritage Properties, National Heritage Places, the Great Barrier Reef Marine Park and Commonwealth Marine Areas are considered under the Authority's permission system</b>
	<p>effective integration of all the commitments in the Program Report.</p> <p><b>Case study:</b></p> <ul style="list-style-type: none"> <li>The Authority will apply conditions to ensure an unacceptable impact on the environment and OUV does not occur as a result of the proposed action. Conditions will be developed having regard to statutory plans. The conditions applied to projects must be specific, measurable, achievable, realistic, time related. Ultimately projects must deliver a net environmental benefit to the environment.</li> <li>The conditions must ensure that the identified avoidance, mitigation, offset, net benefit and adaptive management measures, such as operational monitoring/environmental site supervisors, Deeds of Agreement and bonds are appropriate for managing impacts for the life of the project.</li> <li>The Authority will not grant a permit for an activity inconsistent with regulatory objectives set out under the GBRMP Act.</li> </ul>
<p><b>AUDIT AND COMPLIANCE:</b> The purpose of the audit and compliance phase is to monitor the environmental performance of permitted activities against required standards and permit conditions. Where the results of performance monitoring indicate the required standards have not been met, penalties and/or remediation will be required. Permits may also be suspended or revoked depending on the severity of the non-compliant behaviour.</p>	
Monitoring, compliance and auditing	<p><b>Mechanism (legislative &amp; policy):</b></p> <p>Section 38EA of the GBRMP Act provides penalties for the contravention of a permission condition. Activities must be carried out in accordance with a Great Barrier Reef Marine Park Permission and any conditions.</p> <p>Section 38BA of the GBRMP Act states that it is an offence to conduct an activity in the Marine Park without Permission.</p> <p><b>Program commitment(s):</b></p> <p>The Authority will:</p> <ul style="list-style-type: none"> <li>implement guidelines to improve the collection, management, analysis and interpretation of information derived from the Authority's permission system and compliance monitoring. The information will be used to better inform adaptive management strategies.</li> <li>develop and/or strengthen knowledge and information management tools (including spatial mapping tools, hydrological connectivity tools and systems for managing Indigenous and historic heritage information) so their use can enhance and support: <ul style="list-style-type: none"> <li>the effects of impacts of actions on the Marine Park, including matters of national environmental significance</li> <li>measures to avoid, mitigate and offset impacts on the Marine Park, including matters of national environmental significance</li> <li>actions to deliver net benefits and best practice adaptive management.</li> </ul> </li> <li>strengthen its system for monitoring and managing compliance risks associated with permit conditions. It will review and make any necessary improvements to internal policies, procedures, training protocols and mechanisms to systematically enhance the effectiveness of its compliance and auditing process. This will include specific reference to matters of national environmental significance.</li> </ul> <p><b>Case study:</b></p> <ul style="list-style-type: none"> <li>An Environmental Management Plan (EMP) is required for activities that may impact on the environment and OUV. The EMP outlines the proposed activity, its possible impacts on the Marine Park and specifies strategies to minimise potential impacts. An EMP is required for all activities undergoing an EIS/PER assessment. The EMP needs to be approved by the Authority before implementation and before the commencement of activities permitted under the permit.</li> <li>The EMP must include identification of potential impacts on the environment and OUV, how on-water activities will be managed to reduce these impacts, a monitoring program, emergency response plans and any relevant issue-based plans.</li> <li>Independent experts who are selected and directly contracted by the Authority review EMPs to ensure the environmental management</li> </ul>

Outcome from EPBC Act	How World Heritage Properties, National Heritage Places, the Great Barrier Reef Marine Park and Commonwealth Marine Areas are considered under the Authority's permission system
	<p>measures set out in the EMP are being followed and adhered to. The results of the review are reported to the Authority. An independent expert is required for all activities undergoing an EIS assessment.</p> <ul style="list-style-type: none"> <li>• An Environmental Site Supervisor and/or a Monitoring Consultant may be required (in addition to an independent expert who reviews EMPs) to ensure the activity is undertaken in accordance with the permission to ensure protection of the environment and OUV. These persons must be funded by, but independent of, the proponent and be without any conflict of interest. The Site Supervisor and Monitoring consultants have an 'on the ground role' to liaise with the applicant to ensure the conditions of the Permission are implemented and upheld. The person selected for these roles are selected by the Authority.</li> <li>• The Authority will select and directly contract the Monitoring Consultant and Environmental Site Supervisor. These are required for all high risk activities undergoing an EIS assessment</li> <li>• The Environmental Site Supervisor may be engaged for the whole project or any relevant phases. For example a site supervisor with heritage expertise may be engaged for the period of pile driving, dredging and or disposal where impacts on OUV or other heritage sites are likely.</li> <li>• An Advisory Committee may be established for the project to provide advice to the Authority. The Advisory Committee may include representatives from Government, the proponent, community, the Environmental Site Supervisor, and experts in the fields of science, heritage and management.</li> <li>• The Authority requires an environmental performance audit of activities against permission conditions for developments requiring an EIS. If environmental performance or compliance does not meet required standards for the protection of world heritage property, marine park values, national heritage places or commonwealth marine areas then penalties and/or remediation will be required.</li> <li>• The Authority can determine the need for a cash bond or bank guarantee and a Deed of Agreement to provide ongoing indemnity and, where necessary, to ensure there is an ongoing contractual obligation to protect the environment if the permit lapses, unexpected harm occurs or the proponent is financially unable to take action.</li> <li>• Civil penalties are available for offences under the GBRMP Act for breaches of Permission conditions or undertaking an activity without Permission.</li> </ul>





## Appendix 8

---

### Terms of Reference – concordance table



TERMS OF REFERENCE	Strategic Assessment Report	Program Report
<b>1. Purpose and description of the program</b>		
1.1 Provide an overview of the current Program. For the purposes of the Strategic Assessment the life of the Program is 25 years. The overview is to include a description of:	Chapter 3 Chapter 13	5
a) The purpose of the Program, including Program objectives.	3.2, 13.2	1.2, 1.4
b) The area to which the Program applies (the Strategic Assessment area).	1.2.4, 3.3	1.6
c) Legislation, plans, policies and other mechanisms that make up the Program, including Program commitments.	Chapter 3, 13.2	2.5, 4.1, 10.3
d) Relevant activities within the scope of the Program.	3.6, 3.7	10.4
e) International, national, state and regional context (environmental, social, cultural and economic) in which the Program operates, including activities outside the Strategic Assessment area that may influence the Program.	1.1.1, 1.1.2, 1.1.3, 3.15, 5.3	2
f) Relevant national, state, regional and local planning or management frameworks that affect the Program and contribute to protection and management of matters of national environmental significance.	3.1, 3.4., 3.5, 3.6, 3.7, 3.8, 3.11, 3.12	2.4
g) Identification of how long the Program will be in effect and the process for review of the Program, including adaptive management.	1.2.7, 13.5, 13.8	1.7, 9.1, 9.2, 13.1
h) Identification of the relevant authorities responsible for the implementation of the Program.	1.1.4, 3.8	
<b>2. Matters of national environmental significance</b>		
2.1 Description of matters of national environmental significance		
2.1.1 Describe the extent to which the following relevant matters of national environmental significance, as defined in the Environment Protection and Biodiversity Conservation Act 1999 apply to the Strategic Assessment area.	Chapter 4	
a) World heritage properties, including consideration of the outstanding universal value of the Great Barrier Reef World Heritage Area	4.2	
b) national heritage places	4.4	
c) wetlands of international importance	4.8	
d) listed threatened species and ecological communities	4.7	
e) listed migratory species	4.6	
f) Commonwealth marine areas	4.5	
g) the Great Barrier Reef Marine Park.	4.3	
The description must:		
h) Identify key terrestrial, coastal and marine environmental, biodiversity and heritage values and/or attributes which underpin the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area.	4.1 to 4.8	
i) Describe ecosystem processes considered critical to the functioning of the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area.	4.3, 4.9	
j) Provide sufficient information to allow an understanding the connectivity between the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area.	4.9, 4.10	
2.2 Condition and trend of matters of national environmental significance		
a) Describe the current condition and trend of key indicators of the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area.	Chapter 7	

TERMS OF REFERENCE	Strategic Assessment Report	Program Report
b) For world heritage values, benchmark the current condition of key indicators of outstanding universal value against the retrospective statement of outstanding universal value which describes the state of the Great Barrier Reef World Heritage Area at the time of listing in 1981.	7.6.1	
c) Identify key information gaps and processes to address critical information needs.	7.7	
<b>3. Assessment of impacts on matters of national environmental significance</b>		
3.1 Actual and potential impacts		
a) Describe the environmental, social, cultural and economic drivers affecting the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area.	Chapter 5	
b) Describe and analyse the actual and potential impacts on the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area, in the Strategic Assessment area, including:	6.9, 10.11	
i. impacts from past, present and future activities	5.3, 5.4, 6.2.1, 6.3, 6.4.5, 6.11,	
ii. direct, indirect, consequential and cumulative impacts	6.3, 6.4.4, 6.8	
iii. the likely impacts of climate change.	5.2.2, 6.4.1	
c) Describe the spatial and temporal scale at which impacts and their effects on the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area, are occurring or are likely to occur.	Chapter 5, Chapter 6	
d) Identify key information gaps and processes to address critical information needs.	6.10, 7.7	
<b>4. Measures to address impacts</b>		
4.1 Current Program		
4.1.1 Describe and assess the effectiveness of the Program to:	Chapter 8	
a) Identify the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area, and determine their current condition and trend, including spatial and non-spatial approaches.	8.7.1, 8.8	
b) Identify and analyse direct, indirect, consequential and cumulative impacts, including the methods used to determine these types of impacts.	8.6.1	
c) Consider environmental, social, cultural and economic issues.	8.6.1	
d) Avoid, mitigate, offset and adaptively manage impacts.	8.6.2	
e) Address uncertainty and risk.	8.6.3	
f) Provide certainty regarding where uses may occur, the type of activities allowed, conditions under which activities may proceed and circumstances where impacts are likely to be unacceptable.	8.6.3	
g) Halt and reverse any declines and enhance the condition of the relevant matters of national environmental significance, including mechanisms to deliver a 'net benefit' to the condition of the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area,	8.6.4	
h) Adapt to reasonable climate change scenarios.	8.6.5	
i) Integrate with related local, Queensland and Australian government programs to protect and manage the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area.	8.6.6	
j) Meet Australia's international responsibilities in relation to the environment and protection of world heritage.	8.6.7	



TERMS OF REFERENCE	Strategic Assessment Report	Program Report
k) Monitor, evaluate and report on the:		
i. Condition and trends of the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area.	8.7	
ii. Impacts of activities, including the setting of targets to benchmark management performance.	8.6.8, 8.7.2	
4.1.2 In assessing the effectiveness of the Program, consideration must be given to:		
a) Statutory instruments, including regulations, zoning plans, plans of management and permits.	8.2	
b) Non-statutory mechanisms including policies, position statements and guidelines.	8.2	
c) Partnership and collaborative arrangements with Queensland and other Australian government agencies.	8.2	
d) Partnerships with Traditional Owners in the management of marine resources.	8.2.3, 8.3.2	
e) Partnership and stewardship programs, including education programs and engagement, with local governments, communities, Indigenous persons, business and industry.	8.3.3	
f) Research and monitoring programs.	8.5.8, 8.6.8, 8.7.2	
g) Compliance and enforcement programs.	8.2	
h) Resourcing of the Program.	Chapter 8	
4.2 Demonstration cases		
4.2.1 Develop demonstration cases to assess in finer detail the effectiveness of the Program to protect and manage the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area, and to guide improvements to the Program.	Chapter 9 Appendix 7	
4.2.2 Demonstration cases are to be chosen by the Australian and Queensland governments. Criteria that will be used to guide this selection process include, but are not limited to:	Chapter 9	
a) Where multiple impacts are acting or predicted to act upon a region, locality or value.	9.6, 9.7, 9.9 Appendix 7	
b) To examine a specific management approach or method to identify a set of values/attributes or to assess a range of impacts/pressures.	Chapter 9 Appendix 7	
c) To demonstrate connectivity across coastal and marine systems.	Chapter 9	
d) To demonstrate the integration of environmental, social, cultural and economic benefits in decision-making.	9.10 Appendix 7	
e) To improve understanding of factors affecting Great Barrier Reef ecosystem resilience.	9.3, 9.4	
f) Where lessons or outcomes could transfer to other areas.	9.11	
g) Opportunities to build capacity for future management.	9.11 11.2.2	
h) To examine the effectiveness of management across local, Queensland and Australian government jurisdictions.	Chapter 9 11.2.2	
<b>5. Projected condition of matters of national environmental significance</b>		
5.1 Describe the projected condition of the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area, based on an evaluation of their:		
a) Current status and trends.	11.1, 11.5	
b) Actual and potential impacts.	11.4, 11.5	

TERMS OF REFERENCE	Strategic Assessment Report	Program Report
c) The effectiveness of the Program to protect the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area, and manage impacts.	11.2	
d) An understanding of ecosystem resilience.	10.1, 10.2, 10.3, 10.4, 10.5, 10.12, 11.3	
e) An assessment of overall risks to the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area.	10.6, 10.7, 10.8, 10.9, 10.10, 10.11, 11.4, 11.6	
<b>6. Proposed Program</b>		
6.1 Recommendations for changes to the Program	Chapter 12 Chapter 13	
6.1.1 Recommend changes to the current Program to improve its effectiveness to deliver its objectives, including outcomes that protect the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area. Recommendations for improvements should specifically address the matters listed in section 4.1.1.	12.2, 12.3	5, 6.3, 7.3, 8.3, Part C
6.1.2 Consistent with section 6.1.1 above, recommend improvements to related local, state and national government programs.	12.4	
6.2 Principles of Ecologically Sustainable Development	13.4	14.2.2
a) Decision-making processes should effectively integrate both long-term and short-term economic, environmental, social and equitable considerations.	13.4	14.2.2
b) If there are threats of serious or irreversible environmental damage, lack of full scientific certainty should not be used as a reason for postponing measures to prevent environmental degradation.	13.4	14.2.2
c) The principle of inter-generational equity—that the present generation should ensure that the health, diversity and productivity of the environment is maintained or enhanced for the benefit of future generations.	13.4	14.2.2
d) The conservation of biological diversity and ecological integrity should be a fundamental consideration in decision-making.	13.4	14.2.2
e) Improved valuation, pricing and incentive mechanisms should be promoted.	13.4	14.2.2
6.3 Adaptive management	13.5	9
a) Describe how the adaptive management measures will be implemented to ensure the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area, are effectively protected and managed over the life of the Program.	13.5	9.5
6.4 Monitoring and reporting	13.6	9, 13
a) Describe the processes for adapting the Program in response to new information.	13.6.3	9, 13
b) Nominate the parties responsible for undertaking the monitoring, review and reporting and for implementing any actions arising.	13.6.4	9, 13
6.5 Review, modification or abandonment		
a) Identify and analyse likely circumstances and procedures that may result in the review, modification or abandonment of the Program. This is to include a discussion of how any commitments under the Program would continue to be met under these situations.	13.6, 13.8	
<b>7. Strategic assessment process</b>		
7.1 Collaboration with the Queensland Government and other Australian Government agencies	3.4	

TERMS OF REFERENCE	Strategic Assessment Report	Program Report
a) Undertake the Strategic Assessment in consultation and collaboration with the Queensland Government, the Australian Government's Department of Sustainability, Environment, Water, Population and Communities and other relevant Australian Government agencies.	1.2.2, 1.2.3, 1.2.6	
7.2 Community engagement		
a) Document how the community and stakeholders were engaged in the Strategic Assessment process and how views and comments were taken into account in the preparation of the Strategic Assessment Report and the Program Report.	2.11	
b) Along with the draft Program Report, make the draft Strategic Assessment Report available for public comment for a period of no less than 28 days.	Public comment period from 1 November 2013 to 31 January 2014	
c) Provide the Minister with a report on the public submissions received on the draft Reports, together with proposed final drafts of the Strategic Assessment Report and Program Report, incorporating any revisions made in response to public comments.	Supplementary Report will be provided to the Minister together with final drafts.	
7.3 Independent review		
a) Engage independent expertise to undertake an assessment of the effectiveness of the current Program to protect and manage the relevant matters of national environmental significance, including the outstanding universal value of the Great Barrier Reef World Heritage Area.	Chapter 8	
b) Arrange for the content of the draft Strategic Assessment Report to be peer reviewed by at least three appropriately qualified persons.	2.12, SKM report	
c) Provide the Minister with the independent assessment of management effectiveness, the peer review comments and a report identifying how the findings of the independent assessment and peer review have been considered in the Strategic Assessment Report and the Program Report.	2.12, Supplementary report	
7.4 Information and assessments		
a) Use the best available information to undertake the Strategic Assessment, including scientific data, expert opinion, and Traditional Owner and community knowledge.	2.10, 2.11, references for each chapter	
b) Document the methods used to undertake the Strategic Assessment.	2.7	
c) For information used in the Strategic Assessment, indicate where possible:		
i. the source of the information	2.10, references for each chapter	
ii. how recent the information is	2.10, references for each chapter	
iii. the reliability and limitations of the assessment.	2.13, 6.10, 7.7, references for each chapter	
<b>8. Endorsement criteria</b>		
Describe how the Strategic Assessment Report meets the Endorsement Criteria set out below:	13.7	14.2.3
a) Protects the environment, especially those aspects of the environment that are matters of national environmental significance.	13.7	14.2.3
b) Promotes ecologically sustainable development through the conservation and ecologically sustainable use of natural resources.	13.7	14.2.3
c) Promotes the conservation of biodiversity.	13.7	14.2.3
d) Provides for the protection and conservation of heritage.	13.7	14.2.3
e) Promotes a cooperative approach to the protection and management of the environment.	13.7	14.2.3

TERMS OF REFERENCE	Strategic Assessment Report	Program Report
f) Assists in the co-operative implementation of Australia's international environmental responsibilities.	13.7	14.2.3
g) Recognises the role of Indigenous people in the conservation and ecologically sustainable use of Australia's biodiversity.	13.7	14.2.3
h) Promotes the use of Indigenous peoples' knowledge of biodiversity with the involvement of, and in co-operation with, with the owners of the knowledge.	13.7	14.2.3
Without limiting the matters the Minister may consider when making the decision to endorse the Program, the Minister will consider the manner in which the Program:		
i) Identifies direct, indirect and cumulative impacts on matters of national environmental significance.	Chapter 6, 13.7	
j) Avoids impacts on matters of national environmental significance.	13.7, 12.3 (REC9-17)	11.6.5
k) Mitigates the impacts on matters of national environmental significance.	13.7, 12.3 (REC18-22)	11.6.5
l) Offsets the impacts on matters of national environmental significance.	13.7, 12.3 (REC23-24)	11.6.5
m) Contributes to the enhancement of the existing environment and management of existing threats.	13.7, 12.3 (REC25-30)	11.6.6
n) Demonstrates adaption to reasonable climate change scenarios.	13.7, 12.3 (REC36-38)	3.3, 6.3.5, 7.3.2
Commitments in the Program must be adequately resourced throughout its life. The Program must demonstrate an effective system of adaptive management that addresses uncertainty and contingency management as well as procedures for monitoring, independent auditing and public reporting on implementation.	13.7,	9.3
The Program must address all of the above matters for it to be considered for endorsement by the Minister in accordance with the <i>Environment Protection and Biodiversity Conservation Act 1999</i> .	13.7	14.2.3







**Australian Government**

**Great Barrier Reef  
Marine Park Authority**

[www.gbrmpa.gov.au](http://www.gbrmpa.gov.au)