

# GREAT BARRIER REEF MARINE MONITORING PROGRAM

Inshore water quality monitoring  
Annual Report 2023–24



© Commonwealth of Australia (Australian Institute of Marine Science) and James Cook University (TropWATER) 2025

Published by the Great Barrier Reef Marine Park Authority

ISSN: 2208-4096

**A catalogue record for this publication is available from the National Library of Australia**

This document is licensed for use under a Creative Commons By Attribution 4.0 International licence with the exception of the Coat of Arms of the Commonwealth of Australia, the logos of the Great Barrier Reef Marine Park Authority, Australian Institute of Marine Science, James Cook University and TropWATER, any other material protected by a trademark, content supplied by third parties and any photographs. For licence conditions see: <http://creativecommons.org/licenses/by/4.0>



**This publication should be cited as:**

Moran, D., Waterhouse, J., Petus, C., Howley, C., Lewis, S., Gruber, R., James, C., Logan, M., Bove, U., Brady, B., Choukroun, S., Connellan, K., Davidson, J., Mellors, J., O'Callaghan, M., O'Dea, C., Shellberg, J., Dick, E., Polglase, L., Tracey, D., Molinari, B., Zagorskis, I., 2025. *Great Barrier Reef Marine Monitoring Program Inshore Water Quality Monitoring: Annual Report 2023–24*. Great Barrier Reef Marine Park Authority, Townsville. 322 pp.

Front cover image: Water sample collection at Fitzroy Island, conducted by technician from the Australian Institute of Marine Science © AIMS.

The Great Barrier Reef Marine Park Authority acknowledges the continuing Sea Country management and custodianship of the Great Barrier Reef by Aboriginal and Torres Strait Island Traditional Owners whose rich cultures, heritage values, enduring connections and shared efforts protect the Reef for future generations.

**DISCLAIMER**

While reasonable efforts have been made to ensure that the contents of this document are factually correct, AIMS and JCU do not make any representation or give any warranty regarding the accuracy, completeness, currency or suitability for any particular purpose of the information or statements contained in this document. To the extent permitted by law AIMS and JCU shall not be liable for any loss, damage, cost or expense that may be occasioned directly or indirectly through the use of or reliance on the contents of this document.

Comments and questions regarding this document are welcome and should be addressed to:

Australian Institute of Marine Science  
PMB No 3  
Townsville MC, Qld 4810

TropWATER- Centre for Tropical Water and  
Aquatic Ecosystem Research  
James Cook University  
Townsville, Qld 4811  
[Tropwater@jcu.edu.au](mailto:Tropwater@jcu.edu.au)

*This project is supported by the Great Barrier Reef Marine Park Authority through funding from the Great Barrier Reef Marine Monitoring Program, the Australian Institute of Marine Science, and James Cook University.*

## CONTENTS

<b>Contents</b>	<b>ii</b>
<b>List of figures</b>	<b>iv</b>
<b>List of tables</b>	<b>ix</b>
<b>Appendices: List of figures</b>	<b>x</b>
<b>Appendices: List of tables</b>	<b>xi</b>
<b>Commonly used abbreviations, acronyms, definitions and units</b>	<b>xii</b>
<b>Acknowledgements</b>	<b>xiii</b>
<b>Executive summary</b>	<b>1</b>
Trends in key inshore water quality indicators	1
Drivers and pressures	2
Conclusions	3
<b>1 Introduction</b>	<b>5</b>
1.1 The Great Barrier Reef	5
1.2 Water quality monitoring in the Reef	5
1.3 Structure of the summary report	5
<b>2 Methods</b>	<b>7</b>
2.1 Sampling design	7
2.2 Water quality sampling	9
2.3 <i>In situ</i> loggers	11
2.4 Data analyses – Summary statistics and trends	11
2.5 Data analyses – Water Quality Index	12
2.6 Data analyses – Remote sensing monitoring products	13
2.6.1 Mapping Reef water types	13
2.6.2 Characterising composition of Reef water types	17
2.7 River discharge and catchment loads	20
<b>3 Drivers and pressures influencing water quality in 2023–24</b>	<b>25</b>
3.1 Coastal development including agriculture	25
3.2 Climate and cyclone activity	26
3.2.1 Rainfall for the Reef, NRM regions, and basins	28
3.2.2 Freshwater discharge for the Reef, NRM regions, and basins	29
<b>4 Satellite remote sensing of marine water quality</b>	<b>32</b>
4.1 Satellite remote sensing of Reef water types	32
4.1.1 Areas affected	32
4.1.2 Composition of Reef water types	36
4.1.3 Potential exposure risk to Reef ecosystems	38
4.2 Regional exposure of coastal waters and ecosystems to wet season discharge	42
4.2.1 Cape York region	42
4.2.2 Wet Tropics region	46
4.2.3 Burdekin region	50
4.2.4 Mackay-Whitsunday region	54
4.2.5 Fitzroy and Burnett-Mary regions	58
4.3 Satellite remote sensing summary and discussion	63
<b>5 Focus region water quality and Water Quality Index</b>	<b>68</b>
5.1 Cape York region	68
5.1.1 Pascoe	71
5.1.2 Stewart	77
5.1.3 Normanby	82
5.1.4 Annan-Endeavour	89
5.2 Wet Tropics region	98
5.2.1 Barron-Daintree	98
5.2.2 Russell-Mulgrave	112
5.2.3 Tully	126
5.3 Burdekin region	141
5.4 Mackay-Whitsunday region	148

<b>6</b>	<b>Conclusions.....</b>	<b>156</b>
6.1	Cape York .....	162
6.2	Wet Tropics .....	165
6.3	Burdekin .....	167
6.4	Mackay-Whitsunday .....	169
<b>7</b>	<b>References.....</b>	<b>171</b>
	<b>Appendix A: Water quality site locations and frequency of monitoring.....</b>	<b>178</b>
	<b>Appendix B: Water quality monitoring methods.....</b>	<b>183</b>
B-1	Comparison with Reef Water Quality guideline values.....	183
B-2	Calculation of the Water Quality Index .....	183
B-3	Monitoring of Reef water quality trends using remote sensing data .....	187
B-4	References.....	198
	<b>Appendix C: Additional information .....</b>	<b>200</b>
C-1	Time-series of turbidity and chlorophyll.....	200
C-2	Time-series of temperature and salinity .....	207
C-3	Summary statistics for all sites .....	208
C-4	Data used to generate remote sensing maps .....	260
C-5	Site-specific guideline values for MMP sites .....	266
C-6	Regional exposure assessments for waterbodies .....	274
C-7	References.....	277
	<b>Appendix D: Water Quality Monitoring in the Fitzroy NRM Region 2023–24.....</b>	<b>278</b>
D-1	Acknowledgements .....	278
D-2	Introduction and Background .....	278
D-3	Methods .....	279
D-4	Drivers and pressures influencing water quality in 2023–24 .....	283
D-5	Focus region water quality and Water Quality Index .....	288
D-6	Discussion and Conclusions .....	296
D-7	References.....	298
	<b>Appendix E. Scientific publications and presentations associated with the program, 2023–24 .....</b>	<b>305</b>
E-1	Publications.....	305
E-2	Presentations .....	306

## LIST OF FIGURES

Figure i: Water Quality Index scores from 2008 to 2024.....	1
Figure 1-1: Driver Pressure State Impact Response (DPSIR) framework used to guide the structure of the MMP .....	6
Figure 2-1: Site locations for water quality monitoring sampled from 2015 onwards.....	8
Figure 2-2: Methods for Reef water type, frequency, and exposure maps using Sentinel-3-OLCI true colour imagery .....	16
Figure 3-1: Trajectories of tropical cyclones affecting the Reef in the 2023–24 water year (1 October to 30 September) and in previous years. ....	27
Figure 3-2: Average daily wet season rainfall (mm d <sup>-1</sup> ) in the Reef catchment.....	28
Figure 3-3: Difference between daily average wet season rainfall (December 2023–April 2024) and the long-term wet season rainfall average (from 1961–1990).....	29
Figure 3-4: Long-term total annual discharge in ML (water year: 1 October to 30 September) for the 35 main Reef basins. ....	30
Figure 3-5: Corrected annual water year (1 October to 30 September) discharge from each NRM region (using the correction factors in Table 2-3) for 2003–04 to 2023–24 in (ML per year).....	30
Figure 4-1: Map showing the frequency of the Reef WT1–2 combined.....	33
Figure 4-2: Map showing the frequency of the Reef WT1 (FU10–21), WT2 (FU6–9), and WT3 (FU4–5) in the 2023–24 wet season (22 weeks).....	34
Figure 4-3: Areas (km <sup>2</sup> ) and percentages (%) of the Reef lagoon (total 348,839 km <sup>2</sup> ) affected by the different Reef water types.....	35
Figure 4-4: Long-term (2004–2023) concentrations of water quality parameters and Secchi disk depth boxplots for each Reef Water Type.....	37
Figure 4-5: Mean long-term concentrations of water quality parameters (top) and magnitude scores across the three Reef water types (bottom) .....	38
Figure 4-6: Map showing the reclassified surface exposure .....	40
Figure 4-7: Percentage of the a) Reef waterbodies, b) Reef regions, c) regional Reef waterbodies, d) seagrass and e) coral habitats affected by different risk categories of exposure during the 2023–24 wet season. ....	41
Figure 4-8: Long-term and current year remote sensing results for the Cape York region .....	43
Figure 4-9: Percentage of the Cape York region a) coral reef and b) surveyed seagrass habitats affected by different risk categories.....	46
Figure 4-10: Long-term and 2023–24 remote sensing results for the Wet Tropics region .....	47
Figure 4-11: Percentage of the Wet Tropics region a) coral reef and b) surveyed seagrass habitats affected by different risk categories.....	50
Figure 4-12: Long-term and current year remote sensing results for the Burdekin region .....	51
Figure 4-13: Percentage of the Burdekin region a) coral reef and b) surveyed seagrass habitats affected by different risk categories.....	54
Figure 4-14: Long-term and current year remote sensing results for the Mackay-Whitsunday region .	55
Figure 4-15: Percentage of the Mackay-Whitsunday region a) coral reef and b) surveyed seagrass habitats affected by different risk categories .....	57
Figure 4-16: Percentage of the Fitzroy region a) coral reef and b) surveyed seagrass habitats affected by different risk categories.....	60
Figure 4-17: Percentage of the Burnett-Mary region a) coral reef and b) surveyed seagrass habitats affected by different risk categories.....	62

Figure 4-18: Areas (km <sup>2</sup> ) and percentages (%) of the Reef and Reef regions, coral reefs, and surveyed seagrass affected by different risk categories of exposure during the 2023–24 wet season and the long-term (2003–2022) .....	67
Figure 5-1: Water quality sampling sites in the Cape York region shown with water body boundaries	69
Figure 5-2: Combined Sentinel satellite images showing the extent of flood plumes and maximum turbidity (FU units) observed over the week of 18–24 December 2023, following TC Jasper. ....	70
Figure 5-3: Water quality sampling sites in the Pascoe River transect with water body boundaries. ...	71
Figure 5-4: Daily discharge and sampling dates for the Pascoe River (gauge 102102A) for the 2023–24 water year. ....	72
Figure 5-5: Long-term discharge for the Olive and Pascoe Rivers. ....	72
Figure 5-6: Modelled loads of (A) TSS, DIN, and PN and (B) discharge for the Pascoe catchment (note Pascoe catchment only, does not include the Olive catchment) from 2006 to 2024. ....	73
Figure 5-7: Water quality parameters (surface and subsurface samples) and Secchi depth over distance (km) from the river mouth for the Pascoe River focus region.....	74
Figure 5-8: The annual condition WQ Index for the Pascoe focus region for 2023–24 .....	75
Figure 5-9: Sentinel satellite images showing average magnitude Pascoe River flood event and sampling locations on 25 January 2024. ....	76
Figure 5-10: Water quality sampling sites in the Stewart River transect with water body boundaries..	77
Figure 5-11: Daily discharge and sampling dates for the Stewart River (gauge 104001A) for the 2023–24 wet season. ....	78
Figure 5-12: Long-term discharge for the Stewart River (gauge 104001A – Telegraph Road). ....	78
Figure 5-13: Loads of (A) TSS, DIN and PN, and (B) discharge for the Stewart Basin from 2006 to 2024. ....	79
Figure 5-14: Water quality parameters (all surface and subsurface samples for the 2023–24 season) and Secchi depth over distance (km) from river mouth for the Stewart River focus region .....	80
Figure 5-15: Annual condition version of the WQ Index for the Stewart focus region. ....	81
Figure 5-16: Water quality sampling sites in the Normanby Basin focus region with water body boundaries. ....	82
Figure 5-17: Daily discharge and sampling dates for the Normanby River (gauge 105107A) for the 2023–24 wet season. ....	83
Figure 5-18: Long-term discharge for the Normanby River (gauge 105107A – Kalpower Crossing). ...	83
Figure 5-19: Modelled loads of (A) total suspended solids, dissolved inorganic (DIN) and particulate nitrogen (PN) and (B) discharge for the Normanby Basin.....	84
Figure 5-20: Water quality parameters (surface and subsurface) and Secchi depth over distance (km) from river mouth for the Normanby focus region .....	85
Figure 5-21: Annual condition version of the WQ Index for the Normanby focus region. ....	86
Figure 5-22: Water quality data under the influence of flood plumes from the Normanby and Kennedy rivers on 31 December 2023 and 26 March 2024 .....	87
Figure 5-23: Satellite true colour image of Kennedy and Normanby River flood plume and sample locations on 31 December 2023 (left). ....	88
Figure 5-24: Satellite true colour image of Kennedy and Normanby River flood plume and sample locations on 26 March 2024 (left). The corresponding processed imagery on the right panel shows the Forel Ule classes. ....	89
Figure 5-25: Water quality sampling sites in the Annan-Endeavour region shown with water body boundaries. ....	90
Figure 5-26: Loads of (A) total suspended solids, dissolved inorganic (DIN) and particulate nitrogen (PN) and (B) discharge for the Endeavour Basin from 2006 to 2024. ....	91

Figure 5-27: Daily discharge and sampling dates for the Endeavour Basin .....	92
Figure 5-28: Long-term discharge for the Endeavour Basin .....	92
Figure 5-29: Water quality parameters (surface and subsurface samples) and Secchi depth over distance from river mouth (km) for the Endeavour Basin focus region during ambient conditions (2023–24 water year). .....	93
Figure 5-30: Annual condition version of the WQ Index for the Annan-Endeavour focus region. ....	94
Figure 5-31: Satellite true colour image of flood plumes associated with Annan and Endeavour rivers and Tropical Cyclone Jasper .....	96
Figure 5-32: Flood event water quality data from the Annan-Endeavour transect on 20 December 2023 (blue circles) and 24 March 2024 (red circles) .....	97
Figure 5-33: Sampling sites in the Barron-Daintree focus region shown with water body boundaries. ....	98
Figure 5-34: Combined discharge from rivers in the Barron-Daintree region .....	99
Figure 5-35: Loads of (A) TSS, DIN and PN and (B) discharge for the Barron, Daintree, and Mossman Basins from 2006–2024.....	100
Figure 5-36: eReefs hydrodynamic model showing coastal hydrodynamics and drivers for the Wet Tropics region on 9 December 2023 .....	101
Figure 5-37: eReefs hydrodynamic model showing coastal hydrodynamics and drivers for the Wet Tropics region on 13 December 2023 .....	101
Figure 5-38: eReefs hydrodynamic model showing coastal hydrodynamics and drivers for the Wet Tropics region on 15 December 2023 .....	102
Figure 5-39: eReefs hydrodynamic model showing coastal hydrodynamics and drivers for the Wet Tropics region on 23 December 2023 .....	102
Figure 5-40: Temporal trends in water quality variables for the Barron-Daintree focus region .....	105
Figure 5-41: The Water Quality Index (WQ Index) for the Barron-Daintree focus region .....	106
Figure 5-42: Barron River at Myola flow gauge for the 2023–24 wet season. ....	107
Figure 5-43: Satellite true colour image on 28 December 2023 (left panel) showing the flood plume sampling sites from the Barron River. ....	108
Figure 5-44: True colour satellite images of the Barron flood plume from 24 December 2023 (left panel) and from 28 December 2023 (right panel). ....	109
Figure 5-45: Water quality data from the Cairns Transect region under the influence of flood plumes from the Barron River on 19 and 28 December 2023 including A) total suspended solids (TSS), B) oxidised nitrogen ( $\text{NO}_x$ ), and C) chlorophyll <i>a</i> .....	110
Figure 5-46: Water quality data from the Cairns Transect region under the influence of flood plumes from the Barron River on 19 and 28 December 2023 including A) dissolved inorganic phosphorus (DIP), B) particulate nitrogen (PN), and C) particulate phosphorus (PP).....	111
Figure 5-47: Water quality data from the Cairns Transect region under the influence of flood plumes on 28 December 2023 including the herbicides diuron (A) and metolachlor (B) and the insecticide imidacloprid (C) .....	112
Figure 5-48: Sampling sites in the Russell-Mulgrave focus region, shown with the water body boundaries. ....	113
Figure 5-49: Combined discharge for rivers in the Russell-Mulgrave region .....	114
Figure 5-50: Loads of (A) TSS, DIN and PN and (B) discharge for the Russell, Mulgrave and Johnstone Basins from 2006 to 2024.....	115
Figure 5-51: Water quality variables measured during ambient wet and dry season sampling in 2023–24 along the Russell-Mulgrave focus region transect. ....	116
Figure 5-52: Temporal trends in water quality variables for the Russell-Mulgrave focus region .....	119
Figure 5-53: The Water Quality Index (WQ Index) for the Russell-Mulgrave focus region.....	120

Figure 5-54: Russell-Mulgrave River (Mulgrave River at Peets Bridge + Russell River at Bucklands flow gauges) for the 2023–24 wet season. ....	121
Figure 5-55: Satellite true colour image on 19 December 2023 (left panel) from the Wet Tropics NRM region.....	122
Figure 5-56: Map of the offshore Russell-Mulgrave standard event sampling sites (left); and plots of the water column profiles (right) .....	122
Figure 5-57: Water quality data from the Russell-Mulgrave region under the influence of flood plumes on 19 December 2023 including A) total suspended solids (TSS), B) Oxidised nitrogen (NO <sub>x</sub> ), and C) chlorophyll a.....	124
Figure 5-58: Water quality data from the Russell-Mulgrave region under the influence of flood plumes on 19 December 2023 including A) dissolved inorganic phosphorus (DIP), B) particulate nitrogen (PN), and C) particulate phosphorus (PP).....	125
Figure 5-59: Water quality data from the Russell-Mulgrave region under the influence of flood plumes on 19 December 2023 including the herbicides A) diuron and B) metolachlor and C) the insecticide imidacloprid.....	126
Figure 5-60: Sampling sites in the Tully focus region, shown with the water body boundaries. ....	127
Figure 5-61: Combined discharge from rivers in the Tully region .....	128
Figure 5-62: Loads of (A) TSS, DIN and PN and (B) discharge for the Tully, Murray, and Herbert Basins from 2006 to 2024. ....	129
Figure 5-63: Water quality variables measured during ambient and event sampling in 2023–24 along the Tully focus region transect.....	130
Figure 5-64: Temporal trends in water quality variables for the Tully focus region.....	133
Figure 5-65: The Water Quality Index (WQ Index) for the Tully focus region .....	134
Figure 5-66: Tully River at Euramo flow gauge for the 2023–24 wet season. The dates of event sampling by JCU are marked as orange dots.....	135
Figure 5-67: Satellite true colour image on 13 March 2024 (left panel) showing the flood plume sampling sites from the Tully focus region.....	136
Figure 5-68: Water quality data from the Tully region under the influence of flood plumes on 16 December 2023 and 21 March 2024 including A) total suspended solids (TSS), B) Oxidised nitrogen (NO <sub>x</sub> ), and C) chlorophyll a.....	138
Figure 5-69: Water quality data from the Tully region under the influence of flood plumes on 16 December 2023 and 21 March 2024 including A) dissolved inorganic phosphorus (DIP), B) particulate nitrogen (PN), and C) particulate phosphorus (PP) .....	139
Figure 5-70: Water quality data from the Tully region under the influence of flood plumes on 16 December 2023 and the 21 March 2024 including the herbicides A) diuron and B) metolachlor and C) the insecticide imidacloprid.....	140
Figure 5-71: Sampling sites in the Burdekin focus region, shown with the water body boundaries. ...	141
Figure 5-72: Total discharge from rivers in the Burdekin region .....	142
Figure 5-73: Loads of (A) TSS, DIN and PN and (B) discharge for the Burdekin and Haughton Basins from 2006 to 2024. ....	143
Figure 5-74: Water quality variables measured during ambient and event sampling in 2023–24 along the Burdekin focus region transect. ....	144
Figure 5-75: Temporal trends in water quality variables for the Burdekin focus region .....	145
Figure 5-76: The Water Quality Index (WQ Index) for the Burdekin focus region .....	147
Figure 5-77: Sampling sites in the Mackay-Whitsunday focus region, shown with the water body boundaries.....	148
Figure 5-78: Combined discharge from rivers in the Mackay-Whitsunday region .....	149

Figure 5-79: Loads of (A) TSS, DIN and PN and (B) discharge for the Proserpine, O'Connell, Pioneer, and Plane Basins from 2006 to 2024. ....	149
Figure 5-80: Water quality variables measured during ambient and event sampling in 2023–24 along the Mackay-Whitsunday focus region transect.....	151
Figure 5-81: Temporal trends in water quality variables for the Mackay-Whitsunday focus region....	152
Figure 5-82: The Water Quality Index (WQ Index) for the Mackay-Whitsunday focus region .....	154

## LIST OF TABLES

Table 2-1: List of parameters measured during ambient and event-based water quality monitoring. ....	9
Table 2-2: Description of the Sentinel-3 Reef water types (WT) and corresponding Forel-Ule (FU) colour classes .....	15
Table 2-3: The 35 basins of the Reef catchment, the gauges for each Basin, and the correction factors used to upscale flows to provide annual discharge estimates. ....	21
Table 3-1: Annual water year discharge (ML) of the 35 main Reef basins .....	31
Table 4-1: Areas (km <sup>2</sup> ) and percentages (%) of the Reef, Reef-wide coral reefs, and Reef-wide surveyed seagrass affected by different risk categories of exposure during the 2023–24 wet season and the long-term (2003–2022). ....	39
Table 4-2: Areas (km <sup>2</sup> ) and percentages (%) of the Cape York region, Cape York coral reefs, and Cape York surveyed seagrass affected by different categories of exposure during the 2023–24 wet season and the long-term (2003–2022). ....	45
Table 4-3: Areas (km <sup>2</sup> ) and percentages (%) of the Wet Tropics region, Wet Tropics coral reefs, and Wet Tropics surveyed seagrass affected by different risk categories of exposure during the 2023–24 wet season and the long-term (2003–2022). ....	49
Table 4-4: Areas (km <sup>2</sup> ) and percentages (%) of the Burdekin region, Burdekin coral reefs, and Burdekin surveyed seagrass affected by different risk categories of exposure during the 2023–24 wet season and the long-term (2003–2022). ....	53
Table 4-5: Areas (km <sup>2</sup> ) and percentages (%) of the Mackay-Whitsunday region, Mackay-Whitsunday coral reefs, and Mackay-Whitsunday surveyed seagrass affected by different risk categories of exposure during the 2023–24 wet season and the long-term (2003–2022). ....	56
Table 4-6: Areas (km <sup>2</sup> ) and percentages (%) of the Fitzroy region, Fitzroy coral reefs, and Fitzroy surveyed seagrass affected by different risk categories of exposure during the 2023–24 wet season and the long-term (2003–2022). ....	60
Table 4-7: Areas (km <sup>2</sup> ) and percentages (%) of the Burnett-Mary region, Burnett-Mary coral reefs, and Burnett-Mary surveyed seagrass affected by different risk categories of exposure during the 2023–24 wet season and the long-term (2003–2022). ....	62
Table 5-1: Median wet season turbidity and chlorophyll concentrations from a Wetlabs FLNTU datalogger at Dawson Reef .....	94
Table 5-2: Median wet season turbidity and chlorophyll concentrations from a Wetlabs FLNTU datalogger at Forrester Reef .....	95
Table 6-1: Summary of results for some of the primary indicators measured in the MMP Inshore Water Quality program, 2023–24. ....	158
Table 6-2: Water quality indicator summary for Cape York. ....	163
Table 6-3: Water quality indicator summary for the three focus regions of the Wet Tropics. ....	165
Table 6-4: Water quality indicator summary for the Burdekin region. ....	167
Table 6-5: Water quality indicator summary for Mackay-Whitsunday. ....	169

## APPENDICES: LIST OF FIGURES

Figure B-1: Long-term water quality (WQ) concentrations and Secchi depth boxplots for each wet season colour class. ....	191
Figure B-2: Mean long-term (2004–2023) concentrations of water quality parameters across the three wet season water types in all focus regions .....	192
Figure B-3: Boundaries used for the Marine Park, each NRM region, and the coral reefs and seagrass ecosystems.....	197
Figure C-1: Time-series of daily means of chlorophyll fluorescence and turbidity measured by moored ECO FLNTUSB instruments.....	206
Figure C-2: Time-series of daily means of temperature and salinity derived from moored loggers ..	207
Figure C-3: Areas (in km <sup>2</sup> and represented as horizontal bars) of seagrass (left) and coral reefs (right) .....	274
Figure D-1: Sampling sites in the Fitzroy focus region, shown with the water body boundaries .....	288
Figure D-2: Loads of (A) TSS, DIN and PN and (B) discharge for the Fitzroy Basin from 2006 to 2024. ....	289
Figure D-3: Total discharge from rivers in the Fitzroy region .....	290
Figure D-4: Water quality variables measured during ambient sampling in 2023–24 along the Fitzroy focus region transect. ....	291
Figure D-5: Temporal trends in water quality variables for the Fitzroy focus region .....	293
Figure D-6: The Water Quality Index (WQ Index) for the Fitzroy focus region. ....	295

## APPENDICES: LIST OF TABLES

Table A-1: Description of the water quality sites sampled by AIMS, JCU and CYWP during 2023–24.	178
Table B-1: Guidelines values for four cross-shelf water bodies, provided by the Reef Authority. Values come from the Water Quality Guidelines for the Great Barrier Reef Marine Park	183
Table B-2: Wettest and driest years used to compute the typical wet and typical dry composite frequency maps in each NRM region	189
Table B-3: Reef-wide wet season guideline values used to calculate the exposure score for satellite exposure maps.	194
Table B-4: Number of collected <i>in situ</i> samples used in exposure scoring by region and water type.	195
Table C-1: Summary statistics for water quality parameters at individual monitoring sites from 1 October 2023 to 30 September 2024.	208
Table C-2: Summary of turbidity measurements from moored loggers (site locations in Section 5 and Figure D-1 for the past three water years.	258
Table C-3: Summary of water quality data collected across the Sentinel-3 Reef water types (WT) as part of the wet season event sampling of the MMP	260
Table C-4: Summary of water quality data collected in the Cape York region across the Sentinel-3 Reef water types (WT) as part of the wet season event sampling of the MMP	261
Table C-5: Summary of water quality data collected in the Wet Tropics region across the Sentinel-3 Reef water types (WT) as part of the wet season event sampling of the MMP	262
Table C-6: Summary of water quality data collected in the Burdekin region across the Sentinel-3 Reef water types (WT) as part of the wet season event sampling of the MMP	263
Table C-7: Summary of water quality data collected in the Mackay-Whitsunday region across the Sentinel-3 Reef water types (WT) as part of the wet season event sampling of the MMP	264
Table C-8: Summary of water quality data collected in the Fitzroy region across the Sentinel-3 Reef water types (WT) as part of the wet season event sampling of the MMP	265
Table C-9: Site-specific guideline values (GVs) used for comparison with water quality monitoring data.	266
Table C-10: Areas (km <sup>2</sup> ) (and percentages, %) of the Reef lagoon (total 348,839 km <sup>2</sup> ) and division by water bodies (WB)	275
Table D-1: Description of the Fitzroy water quality sites monitored during 2023–24.	280
Table D-2: Water quality indicator summary for Fitzroy region.	297

## COMMONLY USED ABBREVIATIONS, ACRONYMS, DEFINITIONS AND UNITS

### Abbreviations, acronyms, and definitions

AIMS	Australian Institute of Marine Science
BoM	Bureau of Meteorology
CDOM	colour dissolved organic matter
Chl-a	chlorophyll a
CTD	Conductivity Temperature Depth profiler
CYWP	Cape York Water Partnership
DIN	dissolved inorganic nitrogen
DOC	dissolved organic carbon
DON	dissolved organic nitrogen
DOP	dissolved organic phosphorus
ENSO	El Nino – Southern Oscillation cycle
GAMM	generalised additive mixed effect model
GV	guideline value
JCU	James Cook University
$K_D$	light attenuation coefficient
LOD	limit of detection
MMP	Marine Monitoring Program
Marine Park	Great Barrier Reef Marine Park
MODIS	Moderate Resolution Imaging Spectroradiometer
$\text{NH}_3$	ammonia
$\text{NO}_x$	nitrogen oxides
NRM	natural resource management
PN	particulate nitrogen
$\text{PO}_4$	phosphate (dissolved inorganic phosphorus)
POC	particulate organic carbon
PP	particulate phosphorus
PSII herbicide	photosystem II inhibiting herbicide
QA/QC	quality assurance/quality control
QLUMP	Queensland Land Use Mapping Program
Reef	Great Barrier Reef
Reef Authority	Great Barrier Reef Marine Park Authority
Reef 2050 WQIP	<i>Reef 2050 Water Quality Improvement Plan</i>
Reef Plan	Reef Water Quality Protection Plan
Reef 2050 Plan	<i>Reef 2050 Long-Term Sustainability Plan</i>
SDD	Secchi disk depth
TSS	total suspended solids
water year	1 October to 30 September (e.g., 2010–11)
WS colour scale	wet season colour scale
WQ Index	Water Quality Index

### Units

GL	gigalitre
m	metre
$\text{mm d}^{-1}$	millimetres per day
$\text{mg L}^{-1}$	milligram per litre
ML	megalitre
km	kilometre
$\text{km h}^{-1}$	kilometres per hour
kt	kilotonne
t	tonne
$\mu\text{g L}^{-1}$	microgram per litre

## ACKNOWLEDGEMENTS

We acknowledge the Australian Government funding through the Great Barrier Reef Marine Park Authority (the Reef Authority) for financial and technical support of this program. Thank you to Kaye Walker, Martina Prazeres, and Bronte Claussen from the Reef Authority for their overall project management and program guidance.

We are grateful to all the people involved in the field work. We acknowledge Andrew Mead from Barra Charters for the Burdekin region sampling in partnership with TropWATER James Cook University. Thanks to the Australian Institute of Marine Science (AIMS) Oceanographic Technician team, Marine Operations team, Data and Digital Engineering team, and the crews of the R.V. Cape Ferguson and R.V. Solander, who have been a critical part of this program for many years, as well as Keegan Vickers and Madi McLatchie. From the Cape York Water Partnership, we thank Eric Dick, Jeff Shellberg, Sarah Herkess, Sienna Thomason, the Lama Lama Rangers (Yintinga Aboriginal Corporation) and the Yuku Baja Muliku Land and Sea Rangers. We thank the Pascoe sampling crew: Lana Polglase, Steve Rehn (Albatross Hire) and Tilly Rehn, and acknowledge the Pascoe family group for their support for our work on their Traditional Sea Country. We are grateful to the dedicated staff of the TropWATER laboratory for their assistance with sample preparation and processing and the AIMS Analytical Technology Laboratory who have analysed a large volume of samples for this program.

## EXECUTIVE SUMMARY

The water quality component of the Great Barrier Reef Marine Monitoring Program (MMP) reports on the annual and long-term condition in inshore water quality of the Great Barrier Reef (the Reef) with reference to 19 years of monitoring data. This report presents the results for the 2023–24 water year (1 October 2023 to 30 September 2024). The program design includes the collection of samples along transects in the Cape York, Wet Tropics, Burdekin, and Mackay-Whitsunday regions year-round, with higher frequency sampling during the wet season to better characterise this period of episodic river discharge. Monitoring also occurs in the Fitzroy region through the Fitzroy Basin Marine Monitoring Program for Inshore Water Quality, and results are presented within this report (Appendix D). Satellite imagery and remote sensing are linked with *in situ* monitoring data to estimate the exposure of inshore areas to end-of-catchment loads from rivers for all Reef catchment regions.

### Trends in key inshore water quality indicators

Key water quality indicators were used to derive a Water Quality Index which communicates the long-term trend (insensitive to year-to-year variability) and annual condition (sensitive to year-to-year variability) of water quality relative to guideline values (GVs) (Figure i).

The long-term Index showed that inshore water quality was:

- **stable** this year in the Wet Tropics region after improvement between 2018 and 2022 following a trend of deterioration from 2008 to 2018,
- **stable** in the Burdekin region this year after improvement between 2021 and 2023 and a trend of deterioration from 2010 to 2018,
- **stable** in the Mackay-Whitsunday region this year after improvement over the previous five years following a trend of deterioration from 2007 to 2018, and
- **stable** in the Fitzroy region over the last three years (although this result is preliminary and more data are needed to confirm this) after a trend of improvement from 2010 to 2015.

The annual condition Index showed that inshore water quality in 2023–24 was:

- **‘moderate’** in the Cape York region, representing deterioration compared with the previous year’s ‘good’ score;
- **‘good’** in the Wet Tropics and Burdekin regions, which was an improvement in comparison to the previous several years;
- **‘moderate’** in the Mackay-Whitsunday region and continuing to improve since 2018; and
- **‘good’** in the Fitzroy region, similar to the previous three years.




Figure i: Water Quality Index scores from 2008 to 2024 for the Cape York, Wet Tropics, Burdekin, Mackay-Whitsunday, and Fitzroy regions. The Index is calculated to show the long-term trend since the start of monitoring (circles), where seasonal and short-term variability signals are removed. An updated Index version communicating annual condition is calculated from 2015 onwards (squares) that includes increased temporal and spatial sampling and relates water quality values to wet and dry season Reef water quality guidelines. The Index includes five indicators: water clarity, nitrate/nitrite, particulate nitrogen, particulate phosphorus, and chlorophyll *a*. Cape York long-term trends are not yet assessed due to lack of long-term data.

Individual water quality indicators were monitored for trends and compared against water quality GVs. This water year, GVs were:

- **met in all focus regions** for total suspended solids;
- **met in most focus regions** for particulate nitrogen, particulate phosphorus and chlorophyll *a*;
- **not met in most focus regions** for phosphate; and
- **not met in any focus regions** for Secchi depth and nitrate/nitrite.

Trend analysis shows that since 2015, most water quality indicators have shown **improvement** or **stability** in most focus regions with the exception of particulate nitrogen in the Burdekin focus region and phosphate in the Wet Tropics region, which have shown trends of deterioration. Nitrate/nitrite has shown a trend of improvement in most focus regions, although concentrations remain above (exceeding) GVs. This is a promising finding for nitrate/nitrite and if current trends continue, concentrations in some focus regions could approach GVs in the next few years. Improvement in many water quality indicators has been occurring in the last two to five years, following a period of deterioration from ~2007–2018 that occurred in many regions.

Changes in nutrient concentrations are related to changes in nutrient sources (i.e., inputs) and sinks (i.e., outputs) in the Reef lagoon and potentially changes in the rates of key ecological processes such as primary production or remineralisation. The spatial and temporal variability in the *in situ* water quality discussed in this report highlights the combination of complex factors that control water quality in the Reef lagoon including river discharge, biogeochemical processes, physical forcing, and the variability of these drivers between focus regions.

### Drivers and pressures

Environmental conditions over the 2023–24 wet season were variable across the Reef. River discharge was 1.7 times above the long-term median for the total Reef discharge but varied among Natural Resource Management (NRM) regions. The Cape York and Wet Tropics regions had discharge well above their long-term medians (2.6 and 2.1 times above, respectively). Discharge from the Cape York region for the 2023–24 water year was the highest on record (earliest records extending back to the 1968–69 water year) and the discharge from the Wet Tropics region was the highest since the 2010–11 water year. Discharge from the Burdekin region was slightly over (1.2 times) the long-term median, and discharge from the Fitzroy region was close to the long-term median. Discharge from the Mackay Whitsunday region was less than the long-term median (0.8 times). Discharge from the Burnett-Mary region was 1.7 times the long-term median.

Two cyclones affected the Reef in 2023–24: Tropical Cyclones Jasper and Kirrily. Tropical Cyclone Jasper crossed the Queensland coast as a Category 2 system near Wujal Wujal (between Mossman and Cooktown) on 13 December 2023; it had previously peaked as a Category 5 system in the Coral Sea. Following its crossing, the cyclone quickly weakened to a tropical depression but stalled over Cape York for several days bringing major flooding for the rivers of southern Cape York and northern Wet Tropics. Tropical Cyclone Kirrily crossed the coast near Townsville as a Category 1 system on 25 January 2024; it peaked in the Coral Sea as a Category 3 system but weakened quickly on its approach to the coast.

The impacts of severe flooding to rivers in the Cape York region were evident in the water quality conditions for several Cape York regions including the Annan-Endeavour, Normanby and (to a lesser extent) the Stewart, which did not experience major flooding itself but was influenced by floodwaters that impacted Princess Charlotte Bay and coastal areas to the north. Tropical Cyclone Jasper also caused flooding in the Barron and Daintree Rivers; these floodwaters were influenced by transport patterns that tended to move fresh water across the shelf north of Cape Grafton.

End-of-catchment sediment and nutrient load estimates showed distinct variations between the focus regions, which were consistent with previous years. In Cape York, the modelled loads from the Endeavour Basin for the 2023–24 water year were the highest on record and were among the highest on record for the Normanby basin. The Pascoe and Stewart basin loads were also above median levels but were less extreme than the southern Cape York basins. Of the three focus regions within the Wet Tropics NRM region, the Barron-Daintree-Mossman Basin commonly contributes the

lowest discharge and consistent loads compared to the two focus regions to the south (i.e., the Russell-Mulgrave-Johnstone and the Tully-Murray-Herbert Basins). However, in the 2023–24 water year modelled loads for the Daintree-Mossman-Barron were the highest seen over the monitoring period and contributed similar discharge and loads to the Russell-Mulgrave-Johnstone and Tully-Murray-Herbert focus regions. The combined discharge and loads calculated for the 2023–24 water year from the Burdekin-Haughton and Mackay-Whitsunday Basins were similar to the average over the past ten years.

Sentinel-3 satellite images were used to map Reef optical water types (WT). Updated satellite reference maps (long-term, wet, and dry frequency maps) were used for this report, as well as updated mean long-term concentrations of water quality parameters (Gruber *et al.*, 2024). Water types were classified depending on water colour and linked to water quality characteristics; Reef WT1 are brownish waters (enriched in sediment and dissolved organic matter), Reef WT2 are greenish waters (enriched in algae and dissolved organic matter), and Reef WT3 waters have low risk of detrimental ecological effects. There was a high frequency of exposure to Reef WT1 in inshore areas, with mid-shelf to offshore areas usually exposed to Reef WT3 only. In the mid-shelf and offshore water bodies, presence of Reef WT3 waters is often the result of oceanographic processes such as upwelling rather than direct influence of catchment discharge. The area exposed to any water quality potential risk in 2023–24 was spatially limited relative to the scale of the Reef; even so, the area of the Reef in the highest risk categories (risk categories III and IV) was about 10,000 km<sup>2</sup>. Ninety percent of the total Reef area was exposed to no or very low potential risk. Regionally, the total area of exposure to a potential risk (combined risk categories II–IV) was largely similar to the long-term means in the central/southern regions (Burdekin to Burnett-Mary regions), but a 4% increase in area of exposure was observed in both the Cape York and Wet Tropics regions. The area of marine habitat exposed to a potential risk exceeded the long-term average exposure in the Cape York region (+7% coral reef and +10% seagrass) but was similar to the long-term average exposure in the Wet Tropics, despite a small increase to the total area exposed. This is likely related to coastal oceanographic processes in the Wet Tropics region that rapidly dispersed floodwaters from the Barron and Daintree Rivers in a south-easterly cross-shelf direction. In the other regions, the coral reef and seagrass areas exposed to combined potential risk categories II–IV in 2023–24 were either similar to the long-term average or had a slight shift in exposure to lower risk categories. However, there was an increase in the seagrass area exposed to potential risk category III (+10%) from the potential risk category II (-8%) in the Burdekin region.

## Conclusions

This report presents some positive results for inshore water quality in the Reef lagoon for the 2023–24 water year. Although acute impacts to water quality conditions from Tropical Cyclone Jasper were observed within the Cape York NRM, long-term trends of stability or improvement in overall water quality were observed in all other monitored focus regions. Nevertheless, some indicators (especially nitrate/nitrite) remain above (not meeting) water quality guideline values in all focus regions and require continued improvement before guideline values may be met. The relationship between river discharge and water quality is complex and is confounded by large inter-annual and decadal rainfall variability in tropical coastal waters. Progress of catchment management practice adoption is incremental and slow response timeframes are expected between land-based changes and marine water quality. There is also a need to understand the time lags between disturbance events and responses in water quality conditions, which will vary spatially. It is therefore important to interpret the trends of improvement in this report cautiously; further work is needed to distinguish oceanographic and climatic drivers of the observed trends. Associated trends in catchment load reductions from catchment monitoring programs are an important line of evidence that need to be established before trends in inshore water quality could be related to catchment practices. Additionally, further monitoring is needed to determine if trends in inshore water quality are broad-reaching and sustained over time.

The inshore water quality component of the Marine Monitoring Program continues to provide a high value, long-term dataset to inform management of The Reef. The multiple lines of evidence utilised in the Marine Monitoring Program include *in situ* water chemistry sample collection, *in situ* water

quality time-series' from dataloggers, remote sensing analysis, and marine modelling, which all assist in providing a high degree of confidence in the interpretation of the results. The Marine Monitoring Program's outputs have a wide application across scientific, management, government, and public audiences. Findings from the Marine Monitoring Program were used extensively in the recent update of the 2022 Scientific Consensus Statement (Waterhouse *et al.*, 2024), the Great Barrier Reef Outlook Report 2024 (Great Barrier Reef Marine Park Authority, 2024), and the current review of the Reef water quality targets. Continuation of the Marine Monitoring Program and strengthening of linkages with other Paddock to Reef components will further enhance the value of the Marine Monitoring Program.

## 1 INTRODUCTION

### 1.1 The Great Barrier Reef

The Great Barrier Reef (the Reef) is the most extensive reef system in the world, comprising over 2,900 km<sup>2</sup> of coral reefs. It also includes large areas of seagrass meadows, estimated to be over 43,000 km<sup>2</sup> or ~12.5% of the total area of the Great Barrier Reef Marine Park (the Marine Park) (Great Barrier Reef Marine Park Authority, 2019). The Reef catchment is divided into six natural resource management (NRM) regions, each with differing land use, biophysical, and socio-economic characteristics.

### 1.2 Water quality monitoring in the Reef

The management of water quality remains a priority for the Great Barrier Reef Marine Park Authority (the Reef Authority) and the Australian and Queensland governments because good water quality supports the health and resilience of coastal and inshore ecosystems of the Reef (Great Barrier Reef Marine Park Authority, 2019).

In response to concerns about the impact of land-based runoff on Reef water quality, the Reef Water Quality Protection Plan was established by the Australian and Queensland governments in 2003 (Australian and Queensland governments, 2003). The Plan is updated periodically and was last updated by the Australian and Queensland governments in 2017 and was released as the *Reef 2050 Water Quality Improvement Plan* (Reef 2050 WQIP; Australian and Queensland governments, 2018a). It is a major component of the Reef 2050 Long-Term Sustainability Plan (Commonwealth of Australia, 2015, 2018, 2021, <http://www.environment.gov.au/marine/gbr/reef2050>), which provides a framework for the integrated management of the Great Barrier Reef World Heritage Area.

A key deliverable of the Reef 2050 WQIP is the Paddock to Reef Integrated Monitoring, Modelling and Reporting Program (Australian and Queensland governments, 2018b), which is used to evaluate the efficiency and effectiveness of the implementation of the Reef 2050 WQIP and report progress towards goals and targets. The Great Barrier Reef Marine Monitoring Program (MMP) forms an integral part of the Paddock to Reef Integrated Monitoring, Modelling and Reporting Program. The MMP has three components: inshore water quality, coral, and seagrass. Ecological components of the MMP (seagrass and coral health) are published in separate annual reports detailing the condition and trend of these ecosystems in relation to multiple stressors, including water quality presented in this report (for example, McKenzie *et al.*, 2024; Thompson *et al.*, 2024). In previous years, inshore pesticide monitoring had been presented in a separate report (for example, Thai *et al.*, 2020) or as part of the MMP water quality report. Pesticide sampling recommenced in the 2022–23 wet season and is now reported separately (Kaserzon *et al.*, 2024). Loads of sediments and nutrients, and concentrations of pesticides within rivers are monitored and reported by the Great Barrier Reef Catchment Loads Monitoring Program (Water Quality & Investigations, 2024).

The overarching objective of the inshore water quality monitoring program is to ‘Assess temporal and spatial trends in inshore marine water quality and link pollutant concentrations to end-of-catchment loads’ (Australian and Queensland governments, 2018b). Water quality monitoring has been delivered by the Australian Institute of Marine Science (AIMS), James Cook University (JCU) and the Reef Authority since 2005; the Cape York Water Partnership (CYWP) was added as a collaborator in 2017.

### 1.3 Structure of the summary report

This year’s annual report is presented in a shorter, summary report format. It provides a comprehensive summary of the 2023–24 monitoring efforts in a format that focuses more briefly on the observations this year and the emerging trends in water quality conditions over the long-term.

[Section 2](#) presents a summary of the program’s methods. [Section 3](#) describes the factors influencing marine water quality, referred to as drivers and pressures in the Driver-Pressure-State-Impact-

Response (DPSIR) framework (Figure 1-1). Water quality results from satellite imagery are presented in [Section 4](#) at Reef and regional scales. Detailed results from focus regions are presented in [Section 5](#), including river discharge, catchment loading, *in situ* water quality monitoring results, water quality condition relative to guideline values (GVs), and observed trends since 2005. More information on monitoring sites and methods is given in Appendix A and Appendix B. Detailed results tables and figures are included in Appendix C. Monitoring results from the Fitzroy NRM region are presented in a stand-alone chapter as Appendix D. The program's major publications and presentations from the year are reported in Appendix E.




Figure 1-1: Driver Pressure State Impact Response (DPSIR) framework used to guide the structure of the MMP derived from the Great Barrier Reef Strategic Assessment (Great Barrier Reef Marine Park Authority, 2014). The aspects highlighted in yellow are included in this report.

## 2 METHODS

This Section provides an overview of the sampling design and indicators that are monitored as part of the MMP. More details are presented in Appendices and in a separate quality assurance and quality control (QA/QC) report (Great Barrier Reef Marine Park Authority, 2024).

### 2.1 Sampling design

The MMP inshore water quality monitoring component is designed to measure the annual condition and long-term trends in coastal water quality of the Reef. Tropical waters are characterised by high seasonal variability in river discharge, as rainfall from low pressure systems causes river flood plumes to extend into the coastal ocean, while river discharge becomes negligible during low rainfall periods. Water quality monitoring by the MMP is thus conducted during ambient conditions as well as river discharge events.

Ambient monitoring refers to routine sampling during the wet (1 November–30 April) and dry (1 May–31 October) seasons outside of major flood events. It has been conducted since 2005 under the MMP, although the program design (site location, site number, and monitoring frequency) has changed over time.

Event-based monitoring occurs in response to major flood events to capture conditions within flood plumes; event-based monitoring occurs at the ambient site locations, plus additional ‘event’ sites, which may vary in location depending on the characteristics of a particular flood event. The monitoring frequency depends on the number of flood events each year.

The program covers four NRM regions including Cape York, Wet Tropics, Burdekin, and Mackay-Whitsunday regions, initially chosen on the basis of water quality risk assessments (Brodie *et al.*, 2013). Monitoring site locations were selected along expected water quality gradients related to exposure to land-based runoff. This was largely determined by increasing distance from a river mouth in a northerly direction to reflect the predominantly northward flow of surface water driven by the prevailing south-easterly winds (Brinkman *et al.*, 2011).

From 2005 to 2014, monitoring occurred 3 times per year at sites in the regions listed above (with the exception of Cape York) and additionally in the Fitzroy region (discontinued in 2015). An independent statistical review of the MMP in 2014 (Kuhnert *et al.*, 2015) showed that additional sites and higher sampling frequency would provide necessary additional statistical power for the program. The current program design was implemented in February 2015 and includes most of the sampling sites in the pre-2015 design, allowing for the continuation of the long-term time-series, and inclusion of additional sites. This program re-design was reviewed in 2021 and the increase in power to detect change in the Reef’s inshore water quality was verified (Lloyd-Jones *et al.*, 2022).

The program currently includes nine ‘focus regions’, each with 5 to 6 sites measured routinely: Pascoe, Normanby, Annan-Endeavour, and Stewart (in the Cape York NRM, all added in 2017); Barron-Daintree, Russell-Mulgrave, and Tully (all in the Wet Tropics NRM); Burdekin; and Mackay-Whitsunday (Figure 2-1). The frequency of ambient monitoring was increased in 2015, and sites are now visited 3–10 times annually, depending on the focus region.




Figure 2-1: Site locations for water quality monitoring sampled from 2015 onwards. Note that the Cape York transects were added in 2017. Monitoring sites in the Fitzroy NRM region are shown in the stand-alone report for this region (Appendix D).

The list of parameters sampled by the MMP is provided in Table 2-1 and includes:

- continuous measurement of salinity and temperature at seven sites
- continuous measurement of chlorophyll and turbidity at 16 sites
- 49 ambient sites with more frequent sampling during the wet season
- 26 event-based sites identified for sampling during flood conditions.

Table 2-1: List of parameters measured during ambient and event-based water quality monitoring. Note that +/- signs identifying the charge of the nutrient ions were omitted for brevity.

Condition	Parameter	Abbreviation	Units of Measure
Physico-chemical	Salinity	Salinity	(unitless)
	Temperature	Temperature	Celsius degree
	Light attenuation coefficient <sup>1</sup>	$K_D$	$m^{-1}$
	Secchi depth	Secchi	m
	Total suspended solids	TSS	$mg\ L^{-1}$
	Coloured dissolved organic matter	CDOM	$m^{-1}$
	Turbidity	Turb	NTU
Nutrients	Ammonia	$NH_3$	$\mu g\ N\ L^{-1}$
	Nitrite <sup>2</sup>	$NO_2$	$\mu g\ N\ L^{-1}$
	Nitrate <sup>2</sup>	$NO_3$	$\mu g\ N\ L^{-1}$
	Dissolved inorganic phosphorus	$PO_4$	$\mu g\ P\ L^{-1}$
	Silica	Si	$\mu g\ Si\ L^{-1}$
	Particulate nitrogen	PN	$\mu g\ N\ L^{-1}$
	Particulate phosphorus	PP	$\mu g\ P\ L^{-1}$
	Total dissolved nitrogen	TDN	$\mu g\ N\ L^{-1}$
	Total dissolved phosphorus	TDP	$\mu g\ P\ L^{-1}$
	Particulate organic carbon	POC	$\mu g\ C\ L^{-1}$
	Dissolved organic carbon	DOC	$\mu g\ C\ L^{-1}$
Biological	Chlorophyll <i>a</i>	Chl- <i>a</i>	$\mu g\ L^{-1}$
<sup>1</sup> Derived from vertical profiles of photosynthetically active radiation and not sampled at all sites			
<sup>2</sup> $NO_x$ is the sum of $NO_2$ and $NO_3$			

## 2.2 Water quality sampling

At each sampling location (Figure 2-1, Appendix A), vertical profiles of water salinity and temperature were measured with a Conductivity Temperature Depth (CTD) profiler (Sea-Bird Electronics SBE19plus). CTD profiles are used to characterise the water column and to identify its state of vertical mixing. Some CTD profiles included measurements of photosynthetically active radiation

(PAR), which were used to derive the light attenuation coefficient ( $K_D$ ). See the QA/QC report for a detailed description of CTD data processing (Great Barrier Reef Marine Park Authority, 2024).

Immediately following the CTD cast, discrete water samples were collected using Niskin bottles. Surface (~0.5 m below water surface) and bottom (~1 m above the seabed) samples were collected during ambient monitoring, whereas for some event-based monitoring only surface water samples were collected. Samples from the Niskin bottles were taken in duplicate and were analysed for a suite of water quality parameters (Table 2-1). Detailed descriptions of analytical chemistry techniques can be found in the QA/QC report (Great Barrier Reef Marine Park Authority, 2024). Values of water quality variables presented in this report are depth-weighted means calculated using surface and bottom samples.

Below is a brief description of each of the main water quality variables measured as part of the MMP relevant to this report. These definitions are not all-encompassing but are meant to provide a short description of what aspects of water quality they measure and what processes influence the variables:

- **Total Suspended Solids (TSS)** is a measure of the suspended particulate material in the water column. This includes suspended sediments (sand, silt, and clay), living plankton, and detrital (non-living organic) material. TSS concentrations are affected by oceanographic processes including primary production and wind and tide-driven resuspension, as well as inputs from other sources such as dredging and runoff from land.
- **Secchi depth** is a visual measure of water clarity and proxy for light penetration, which is measured using a high-contrast black and white patterned disc called a Secchi disc. The Secchi depth is the average of the vertical disappearance and reappearance depths of the disc, where clarity increases with increasing Secchi depth. Secchi depth is a simple method that has been used for over 150 years, so is excellent for assessing long-term change and for cross-system comparisons.
- **Turbidity** is a measure of light scattering caused by fine suspended particles, such as sediment, detritus, and plankton. Turbidity is affected by a wide range of factors including oceanographic processes such as resuspension of bottom sediments by wind and tides; river discharge; and anthropogenic factors such as dredging.
- **Chlorophyll a (Chl-a)** concentration is a measure of phytoplankton biomass in a water body. Phytoplankton grow quickly in response to nutrient availability, so elevated values of Chl-a can indicate increased nutrient loading.
- **Dissolved inorganic nutrients ( $\text{NH}_3$ ,  $\text{NO}_x$ ,  $\text{PO}_4$  and  $\text{Si}$ )** measure the amount of readily available nutrients for plankton growth in water samples. Inorganic nitrogen ( $\text{NH}_3$ ,  $\text{NO}_x$ ) and phosphate ( $\text{PO}_4$ ) represent around 1% of the nutrient pools in the Reef. The inorganic nutrient pools are affected by a complex range of biogeochemical processes including both natural (for example, plankton uptake, upwelling, nitrogen fixation, and remineralisation) and anthropogenic (for example, dredging and nutrient inputs from changed land use) processes.
- **Particulate nutrients (POC, PN, and PP)** are a measure of the suspended material retained on a filter with a pore size of approximately 0.7  $\mu\text{m}$ . This material consists of a minor fraction of living biomass (e.g., bacteria, phytoplankton) and a major fraction of detritus (e.g., dead cells, faecal pellets). Particulate nutrient concentrations are affected by oceanographic processes (primary production, bacterial production, resuspension, and remineralisation) as well as sources such as dredging and land-based runoff.
- **Dissolved organic carbon (DOC)** is a measure of organic carbon concentrations passing through a filter with a pore size of 0.45  $\mu\text{m}$ . DOC has a complex chemical composition and is used by bacteria as a source of energy. The DOC pool is affected by a range of production and degradation pathways. The sources include primary production by phytoplankton, zooplankton grazing, resuspension events, river runoff, and abiotic breakdown of POC. DOC can be degraded by sunlight.

## 2.3 *In situ* loggers

Continuous *in situ* chlorophyll fluorescence and turbidity were measured using WET Labs ECO FLNTUSB Combination Fluorometer and Turbidity Sensors located at 16 sites (Appendix A), which were deployed at 3 m (Cape York region) or 5 m (all other regions) below the surface and sampled at 10 min intervals. Water samples for analyses of Chl-*a* and TSS were collected three times per year to calibrate logger fluorescence and turbidity to *in situ* conditions. Diver-operated Niskin bottles were used to sample close to the moored loggers and samples were preserved and analysed in the same manner as ship-based water samples.

Daily averages of the chlorophyll and turbidity collected by the ECO FLNTUSB instruments are presented as time-series graphs in Appendix C Figure C-1. Annual means and medians of turbidity were also calculated for each site based on the ‘water year’ (1 October to 30 September the following year) and compared with the guideline value (GV) (Appendix C Table C-2).

Salinity and temperature loggers (Sea-Bird Electronics SBE37) were deployed at seven locations, with four of these being placed on fixed moorings near the O’Connell, Russell-Mulgrave, Tully, and Burdekin River mouths (Figure 2-1; Appendix A). See the QA/QC report (Great Barrier Reef Marine Park Authority, 2024) for detailed descriptions of logger pre- and post-deployment procedures. Site-specific time-series from these loggers can be found in Appendix C Figure C-2.

## 2.4 Data analyses – Summary statistics and trends

Concentrations of water quality parameters at each sampling occasion were calculated as depth-weighted means by trapezoidal integration of the data from all sampling depths. At most sites, only two vertical points are sampled (i.e., surface and bottom samples), and this method averages these values to derive the depth-weighted mean. Measurements falling below the analytical detection limit were replaced with values of half the detection limit. Summary statistics for all water quality variables are presented for all monitoring sites in Appendix C Table C-1. Concentrations were compared to site-specific GVs (Appendix C Table C-9), which are defined for Chl-*a*, PN, PP, TSS, Secchi depth, NO<sub>x</sub>, and PO<sub>4</sub> at most sites. Concentrations of water quality parameters are presented along the sampling transects for each focus region with distance from major river mouth. Trends in water quality with distance are represented with generalised additive models, fitted with a maximum of five knots and modelled with a gamma-distributed response and log-link function.

Temporal trends in key water quality variables (Chl-*a*, TSS, Secchi depth, turbidity, NO<sub>x</sub>, PO<sub>4</sub>, PN, PP, DOC, and POC) since 2005 are reported for all focus regions except Cape York. Only open coastal and mid-shelf sites are used for these analyses because GVs for enclosed coastal waters are derived differently and are not available for all variables, creating statistical imbalance.

Generalised additive mixed effects models (GAMMs) were used to decompose each irregularly spaced time-series into its trend cycles (long-term) and periodic (seasonal) components (Wood, 2006). GAMMs are an extension of additive models (which allow flexible modelling of non-linear relationships by incorporating penalised regression spline types of smoothing functions into the estimation process), where the degree of smoothing of each smooth term (and by extension, the estimated degrees of freedom of each smoother) is treated as a random effect and thus can be estimated via its variance as with other effects in a mixed modelling structure (Wood, 2006).

For each water quality variable within each focus region, the variable was modelled against a thin-plate smoother for date and a cyclical cubic regression spline (maximum of 5 knots) over months within the year. Spatial and temporal autocorrelation in the residuals was addressed by including sampling locations as a random effect and imposing a first-order continuous-time auto-regressive correlation structure (Pinheiro and Bates, 2000). All GAMMs were fitted using the *mgcv* (Wood 2006, 2011) package in R 3.6.1 (R Core Team, 2022).

GAMMs represent long-term trends in water quality variables measured during ambient periods (i.e., not peak flood events) of the dry and wet seasons and are presented in [Section 5](#). It is important to note that this trend analysis removes variability associated with wind, tides, and seasons. Thus, individual data points will have different values from raw data. This analysis is designed to detect **long-term** and **regional-scale** trends in water quality by removing the effect of short-term changes associated with weather and seasonal differences.

In order to provide a more quantitative assessment of trend, linear change in values of GAMMs was measured from the present sampling year to 2015. This period was chosen as it incorporates the MMP re-design, which was introduced in 2015; using earlier data would unbalance this analysis as the amount of sampling greatly increased in 2015. As GAMMs are de-trended to remove the effects of seasons, tides, and wind, this analysis aims to quantify long-term trends occurring outside of these cycles. Trend analysis results are presented for each focus region in [Section 5](#).

## 2.5 Data analyses – Water Quality Index

The Water Quality Index (WQ Index) is an interpretation tool developed by AIMS to visualise trends in the suite of water quality variables measured and to compare monitored water quality to existing Water Quality Guidelines (Department of Environment and Resource Management, 2009; Great Barrier Reef Marine Park Authority, 2010). The WQ Index uses a set of five key indicators:

- Water clarity,
- Chl-a concentrations,
- PN concentrations,
- PP concentrations, and
- NO<sub>x</sub> concentrations.

These five indicators are a subset of the comprehensive suite of water quality variables measured in the MMP inshore water quality program. They have been selected because GVs are available for these measures, and they can be considered as relatively robust indicators that integrate a number of bio-physical processes in the coastal ocean.

For each monitoring site, these indicators are scored based on performance relative to GVs and averaged to give an overall site-specific score. Sites are then averaged over a region or focus region to give a regional score (see Appendix B for details of Index calculation). Results are presented in [Section 5](#).

The WQ Index is calculated using two different methods due to the objectives of the program needing to report both the long-term trend in water quality condition and the annual conditions that ecosystems experience. Changes in the MMP design that occurred in 2015 (increased number of sites, increased sampling frequency, and higher sampling frequency during the wet season [1 November to 30 April]) also needed to be incorporated into WQ Index calculation. The two versions of the WQ Index have different purposes:

1. **Long-term trend:** This version is based on the pre-2015 MMP sampling design and uses only the original sites (open coastal water body) and three sampling dates per year. This sampling design had low temporal and spatial resolution and was aimed at detecting long-term trends in inshore water quality. Key aspects of this version are:
  - annual water quality GVs are used for scoring monitoring data (Table B-1),
  - only AIMS monitoring data are used,
  - a four-year running mean is applied to data to reduce the effect of sampling time on the Index, and
  - the Index is an average of five equally-weighted indicator scores: water clarity (the average of TSS and turbidity from loggers, where available), Chl-a, NO<sub>x</sub>, PN, and PP weighted equally.

In 2024, there was a minor update to the long-term Index calculation, which involved the definition of a ‘water year’. Historically, (pre-2015) the ‘water year’ had been defined as 1 June–31 May due to time constraints around laboratory sample processing and the MMP report deadline. Improvements to workflows in recent years have meant that laboratory sample analysis is now completed more quickly, and samples collected in July can be included in the report (due December of the same year). Therefore starting in 2024, the ‘water year’ was defined as 1 October–30 September to bring the long-term Index in line with all other MMP reporting products. Although all data used to calculate the long-term Index remain the same, this re-definition of the ‘water year’ has caused some minor changes in individual Index scores compared to what has been reported in previous years. These minor changes do not affect the overall trends of the long-term Index, which remain consistent with previous published reports.

**2. Annual condition:** This version is based on the post-2015 MMP sampling design and uses all sites (except enclosed coastal sites) and sampling dates per year. Key aspects of this version are:

- seasonal site-specific water quality GVs are used (i.e., wet season data are compared to a wet season GV and dry season data are compared to a dry season GV) (Appendix C Table C-9)
- both AIMS and JCU monitoring data are used,
- a running mean is not applied, and
- the Index is a hierarchical combination of scores for five indicators: water clarity (the average of TSS, Secchi depth, and turbidity from loggers, where available), productivity (combined score of Chl-a and NO<sub>x</sub>), and particulate nutrients (combined score of PN and PP), which are weighted equally.

## 2.6 Data analyses – Remote sensing monitoring products

### 2.6.1 Mapping Reef water types

The current Program utilises optical information available from medium resolution optical satellite images combined with modelling and field water quality data to monitor the Reef water quality (e.g., Petus *et al.*, 2019; Waterhouse *et al.*, 2021). Until 2020, trends in Reef marine water composition during the wet season were monitored using a combination of Moderate-Resolution Imaging Spectroradiometer (MODIS) satellite imagery and a “wet season” colour scale specifically developed for the Reef (Alvarez-Romero *et al.*, 2013). MODIS satellite pixels were reclassified into six colour classes, then into four broad groups of water type characteristics: the primary (corresponding to colour classes 1 to 4), secondary (colour class 5), and tertiary (colour class 6) wet season water types and the marine water type (Figure 2-2a and Table 2-2).

These water types represented typical colour and water quality gradients encountered in the Reef during the wet season (December to April), including river plumes. Catchment runoff in sediment-laden river discharge appears in satellite images as brownish flood plumes, while productive chlorophyll-rich waters appear with a greenish colour, and ambient (clear) marine waters are a blueish colour. Brownish-green waters also appear when sediments are re-suspended by wind or tide, and it is impossible to fully separate the direct influence of riverine plume from wind- and wave-driven sediment resuspension (some of which may have been originally derived from river discharge in previous events) in optical satellite images. Therefore, the term “wet season waters” referred collectively to flood river plumes, associated resuspension, and oceanographic processes occurring in the Reef during the wet season.

However, MODIS sensors were ageing, and the quality of the MODIS imagery was declining (MODIS-Aqua was launched in 2002). A transition to Sentinel-3 Ocean Land Colour Instrument (OLCI) satellite imagery and another colour scale (the Forel-Ule [FU] colour scale) was proposed for the continuous mapping of Reef waters in 2019 and adopted in 2020 (Petus *et al.*, 2019) (Table 2-2). The FU colour scale is an historical colour scale standard to determine water colour and classifies

waterbodies worldwide (Novoa *et al.*, 2013). It is composed of 22 colours going from indigo blue to 'cola' brown, and is applicable for all natural waters (inland, estuarine, inshore and offshore) and all environmental conditions, including wet and dry season conditions (Wernand *et al.*, 2012, 2013; Van der Woerd *et al.*, 2016; Van der Woerd and Wernand, 2018). In a case study focusing on Wet Tropics and Burdekin regions of the Reef over the 2017–18 wet season, the MODIS-Aqua WS and Sentinel-3 FU colour class maps showed very similar patterns (Petus *et al.*, 2019; Figure 2-2 and Table 2-2).

Equivalent FU water types were defined by grouping:

- FU colour classes 1–3 (equivalent to marine waters in the wet season scale used before 2020–21),
- FU colour classes 4–5 (equivalent to tertiary water type in the wet season scale),
- FU colour classes 6–9 (equivalent to the secondary water type), and
- FU colour classes  $\geq 10$  (equivalent to the primary water type).

Table 2-2: Description of the Sentinel-3 Reef water types (WT) and corresponding Forel-Ule (FU) colour classes (and comparison with MODIS wet season (WS) water types). Mean long-term (2004–2023) concentrations of water quality parameters ( $\pm$  standard deviation) across the Reef water types were updated in 2023 and are indicated in the right column (modified from Petus *et al.*, 2019 and Waterhouse *et al.*, 2018).

Reef water types	FU colour classes (and WS colour classes)	Description	Mean long-term (2004–23) concentrations of water quality parameters*
WT1 (previously primary)	FU $\geq 10$ (WS1–4)	Brownish to brownish-green turbid waters typical of inshore regions of the Reef that receive land-based discharge and/or have high concentrations of resuspended sediments during the wet season.  In flood conditions, this water type typically contains high sediment and dissolved organic matter concentrations resulting in reduced light levels. It is also enriched in coloured dissolved organic matter and phytoplankton concentrations and has elevated nutrient levels.	SDD: $2.1 \pm 2.1$ m TSS: $17.4 \pm 44.1$ mg L <sup>-1</sup> Chl-a: $1.5 \pm 2.2$ $\mu$ g L <sup>-1</sup>
WT2 (previously secondary)	FU6–9 (WS5)	Greenish to greenish-blue turbid water typical of coastal waters with colour dominated by phytoplankton (Chl-a), but also containing dissolved organic matter and fine sediment. This water body is often found in open coastal waters of the Reef as well as in the mid-shelf where relatively high nutrient availability and increased light levels due to sedimentation favour coastal productivity (Bainbridge <i>et al.</i> , 2012).	SDD: $4.6 \pm 2.8$ m TSS: $4.6 \pm 7.0$ mg L <sup>-1</sup> Chl-a: $0.6 \pm 0.7$ $\mu$ g L <sup>-1</sup>
WT3 (previously tertiary)	FU4–5 (WS6)	Greenish-blue waters corresponding to waters with slightly above-ambient suspended sediment concentrations and high light penetration typical of areas towards the open ocean. This water type includes the outer areas of river flood plumes, fine sediment resuspension around reefs and islands and oceanographic processes such as upwelling. Reef WT3 waters are associated with low land-based contaminant concentrations and the ecological relevance of these waters is likely to be minimal although not well researched. The Type III areas have a low magnitude score in the Reef exposure assessment.	SDD: $8.2 \pm 4.1$ m TSS: $2.3 \pm 3.9$ mg L <sup>-1</sup> Chl-a: $0.4 \pm 0.4$ $\mu$ g L <sup>-1</sup>
WT4 (previously marine)	FU1–3 No number	Blueish marine waters with high light penetration.	SDD**: $11.0 \pm 4.8$ m TSS**: $1.8 \pm 3.5$ mg L <sup>-1</sup> Chl-a**: $0.5 \pm 0.9$ $\mu$ g L <sup>-1</sup>

\*SDD = Secchi disk depth, TSS = total suspended solids, and Chl-a = chlorophyll-a.

\*\*Please note that the number of data points collected in the Reef WT4/Marine water type is limited in comparison to the data available in the other water types (Appendix B). Long-term concentrations of water quality parameters in the Reef WT4 are thus just given as an indication and are not used in the monitoring products presented in this report. A pilot study funded by Reef Trust Partnership collected water quality data in mid-shelf and offshore locations via the Crown of Thorns Starfish Control Program (Waterhouse *et al.*, 2023). Water quality data collected as part of this study are included in the long-term dataset to help in progressing the characterisation of the Reef WT4, but more data are still needed to improve accuracy of the characterisation.




Figure 2-2: Methods for Reef water type, frequency, and exposure maps using Sentinel-3-OLCI true colour imagery: a) Summary of the process to produce the Reef water type maps (1) downloading of the true colour imagery, (2) processing of colour class map using the Forel-Ule (FU) colour scale Toolbox implemented in the Sentinel Application Platform, and (3) reclassification into Reef water types. Images are from the Eumetsat data centre and were captured on 26 April 2022 (source: Sentinel Hub EO browser). b) Burdekin River plume (14 March 2018) (the Wet Season [WS] scale ranges from 1–6, plus marine [M]; the catchment [C] is shown in grey). This panel illustrates the similar colour patterns between the (right) Sentinel-3 Forel-Ule colour class maps and (left) the MODIS wet season

water type maps. Panel b was mapped using a supervised classification of MODIS true colour data developed by Álvarez-Romero *et al.* (2013) (modified from Devlin *et al.*, 2015).

In the 2021–22 report (Moran *et al.*, 2023), the water type (WT) terminology was modified to: Reef WT1, WT2, WT3, and WT4 instead of primary, secondary, tertiary, and marine wet season water types. This change was made in response to recognition that the previous terminology may be misleading and systematically implied the presence of flood plume waters, while the Reef WT1 (primary waters) may also represent sediment resuspension in shallower parts of the Reef lagoon, and the Reef WT3 (tertiary waters) may represent oceanographic processes such as upwelling or sediment resuspension around reefs and islands (Table 2-2). Importantly, while names of the water types changed, the definition of the water types in Table 2-2 essentially remained the same.

Several monitoring products are derived from the Sentinel-3 FU water type maps to report on water quality trends. These products map water quality gradients during the wet season and are used to:

- Map the extent of river flood plumes during high flow conditions;
- Characterise the composition of the Reef water types (mean long-term total suspended solids [TSS], chlorophyll *a* [Chl-*a*], coloured dissolved organic matter [CDOM], dissolved inorganic nitrogen [DIN], dissolved inorganic phosphorus [DIP], particulate phosphorus [PP], and particulate nitrogen [PN] concentrations and Secchi disk depth [SDD] values) and identify where mean long-term concentrations of TSS, Chl-*a*, PP, and PN are likely to be above wet season GVs. Wet season GVs for the whole of the Reef (hereafter Reef-wide GVs) are derived from De'ath and Fabricius (2008) (Appendix B Table B-4); and
- Assess the exposure of coral reefs and seagrass ecosystems to potential risk from land-based pollutants.

These products are used to illustrate wet season conditions for every wet season and to compare seasonal trends with baseline reference trends in water composition including long-term conditions, typical wet year and dry year conditions and conditions over a documented recovery period for coral reefs. Available satellite data are biased toward clear, non-cloudy days, and may underrepresent poor water quality in regions of higher rainfall and cloudiness like the Wet Tropics and Cape York. However, they provide a unique large-scale and long-term view of the Reef that is not available using water quality data only.

### 2.6.2 Characterising composition of Reef water types

The classification of four Reef water types allows mapping of a broad grouping of water type characteristics with different colours, concentrations of optically active components (TSS, CDOM, and Chl-*a*), water quality indicators (e.g., nutrient concentrations; Devlin *et al.*, 2015; Petus *et al.*, 2019), and light attenuation levels (Petus *et al.*, 2018) typically found in the Reef during the wet season (Table 2-2 and Figure 2-2). These characteristics vary the potential impact on the underlying ecological systems. In summary:

- **The brownish Reef WT1 (FU ≥ 10)** represents turbid waters from river flood plumes, and also sediment resuspension in the shallower parts of the Reef;
- **The greenish Reef WT2 (FU6–9)** represents the less turbid parts of flood plumes enriched in Chl-*a* and fine sediment. It is usually found in the inshore to mid-shelf regions of the Reef;
- **The greenish-blue Reef WT3 (FU4–5)** represents waters with suspended sediment concentrations slightly above ambient conditions and high light penetration typically found in the outer areas of river flood plumes. It can also represent oceanographic processes such as upwelling or the fine sediment resuspension around reefs and islands; and
- **The blueish Reef WT4 (FU1–3)** represents ambient waters with high light penetration and negligible concentrations of optically active water quality constituents.

Match-ups of *in situ* concentrations of water quality parameters and the four Reef water types are regularly performed to validate this concept and quantify the range and average of concentrations of water quality parameters found in each Reef water type. The previous update was in 2022–23 (Gruber *et al.*, 2024). As part of this update, all mean concentrations of water quality parameters were reviewed to ensure that the water type characterisation remains appropriate, and to improve its accuracy building on the field data that are collected every wet season. The colour class category and water type corresponding to the location and week of acquisition of each water quality sample were extracted from the archive of MODIS-Aqua (wet seasons 2003–2020) and Sentinel-3 (wet seasons 2020–2023) weekly colour class maps (see method in Appendix B). Weekly composites were used rather than daily colour class/water type data to minimise data loss due to the periodic dense cloud cover in the Reef. This approach maximises the incorporation of water quality parameters measured during each wet season since 2003–04 that can be associated with a Reef water type (and colour class) category.

Ideally, match-ups between satellite and *in situ* water quality information should be performed using field data collected  $\pm 2$  hours from the satellite overpass. This is very complicated to achieve in the MMP, which is in part focused on responsive monitoring of flood events and in areas of the Reef where the cloud cover has a major influence during the wet season. The methodology above was thus selected to maximise the number of data points used to assess the water quality characteristics of each Reef water type. The limitations are considered acceptable as the mean concentrations of water quality parameters are used as a relative measure to assign a potential risk grading for each Reef water type (see below). However, the long-term average concentration values should not be used as an exact value *per se*.

The long-term concentrations of water quality parameters were calculated using all surface data ( $<0.2$  m) available. This included data collected between December and April by JCU since the 2003–04 water year, and up to April 2023. It included data collected by AIMS and the CYWP since the 2016–17 water year and covered all regions and waterbodies of the Reef, and all Reef water types. TSS and Chl-*a* data collected in mid-shelf and offshore areas as part of a Reef Trust Partnership project with the locations via the Crown of Thorns Starfish Control Program between December 2021 and April 2023 (Waterhouse *et al.*, 2023) were also included. In previous assessments, long-term mean DIN, PP, and PN concentrations were calculated as: DIN = nitrite + nitrate + ammonia, PP = Total Phosphorus – Total Dissolved Phosphorus, and PN = Total Nitrogen – DIN, respectively. In the current assessment, mean long-term direct measurements of PP and PN were used (rather than indirectly estimated values as above), and NO<sub>x</sub> was used instead of DIN and calculated as NO<sub>x</sub> = nitrite + nitrate due to its greater robustness than ammonia as an indicator of N availability in marine waters.

Boxplots of water quality concentration and Secchi disk depth were plotted against their water type and colour class categories. The mean long-term TSS, Chl-*a*, PP, and PN concentrations were then assessed against wet season GVs as a relative measure to assign potential risk grading for each Reef water type. Reef-wide wet season GVs were derived from De'ath and Fabricius (2008) (Appendix B Table B-4).

### **Reef water type, frequency, and exposure maps**

Several summary maps are produced every wet season including frequency maps of occurrence of wet season water types and exposure maps. The area (km<sup>2</sup>) and percentage (%) of coral reefs and seagrass meadows affected by different relative categories of exposure (or potential risk) was summarised. Details are in Appendix B. For brevity, we have not included weekly panel maps of environmental and marine wet season conditions in this Summary Report.

**Reef water type maps of the 2023–24 wet season** were produced using daily Sentinel-3 OLCI Level 2 (hereafter, Sentinel-3 or S3) imagery (Figure 2-2a, Step 1) reclassified to 21 distinct colour classes defined by their colour properties and using the FU colour classification scale (Figure 2-2a, Step 2). Sentinel-3 imagery of the study area was downloaded on the EUMETSAT Data centre (<https://www.eumetsat.int/eumetsat-data-centre>). Sentinel-3 images are atmospherically corrected and were processed with the FU Satellite Toolbox implemented in the Sentinel Application Platform

(<https://step.esa.int/main/toolboxes/snap/>) and using automated tools (python scripts and ArcGIS toolboxes) developed through MMP funding.

**Frequency maps** were produced to predict the areas affected by the Reef WT1, WT2 and WT3 individually (i.e., of the brownish, greenish and greenish-blue waters, respectively) and by the Reef WT1–2 combined (previously a combination of WT1–3; this has been modified to recognise that the ecological relevance of the water quality variables at concentrations in WT3 waters is not well understood, but expected to be relatively minor).

Average frequency maps were produced for this reporting wet season (2023–24) and compared to average frequency maps representing relevant reference periods (Gruber *et al.*, 2024), including:

- (i) the long-term reference composite (2002–03 to 2021–22: 20 wet seasons),
- (ii) a documented recovery period for coral reefs (2012–2017; Thompson *et al.*, 2024) intended to represent a favourable exposure scenario, and
- (iii) composite frequency maps produced to represent typical wet year and dry year conditions, considering the wettest and driest years for each NRM region over the previous 20 years (Gruber *et al.*, 2024). This is explained further in Appendix B.

Except for the ‘coral recovery period’, reference maps (long-term, representative Wet and Dry frequency maps) were all updated in the 2022–23 report (Gruber *et al.*, 2024) (20 years: 2002–03 to 2021–22) to ensure they remain appropriate and to improve their accuracy as more satellite data are available. The previous update was in the 2018–19 reporting (Gruber *et al.*, 2020).

The presence and spatial extent of each Reef water type is the result of the complex physico-chemical transformations occurring within river plumes, but also of resuspension, transport, and other oceanographic processes. As a result, the extent of the Reef WT2 and WT3 is rarely attributed to an individual river and is usually merged into one heterogeneous area.

**Exposure maps** were produced for this reporting wet season (2023–24) for the whole of the Reef, for all focus regions and compared to reference exposure maps produced for the same timeframe as those reported for the frequency maps above. The maps were produced using an exposure assessment framework developed through a collaborative effort between the MMP monitoring providers (JCU water quality and seagrass monitoring teams and the AIMS coral monitoring team) and modified from Petus *et al.* (2016). Reference exposure composites were also reviewed in the 2022–23 report (Gruber *et al.*, 2024) to produce 20-year composite maps.

In this *magnitude × likelihood* framework, the ‘potential risk’ corresponds to an exposure to above Reef-wide wet season GV concentrations of land-based pollutants during the wet season and focuses on TSS, Chl-*a*, PP, and PN concentrations. The ‘*magnitude of the exposure*’ corresponds to the mean long-term wet season concentration of pollutants (the proportional exceedance of the Reef-wide wet season GV) mapped through the Reef WT1, WT2 and WT3. The ‘*likelihood of the exposure*’ is estimated by calculating the frequency of occurrence of each Reef water type mapped through the frequency maps (see above). The exposure for each of the water quality parameters defined is the proportional exceedance of the GV multiplied by the likelihood of exposure in each of the Reef water types.

1. **Calculation of the exposure (magnitude) scores:** The long-term mean concentrations of water quality parameters (Reef-wide) measured across the Reef water types (Section 2.6.2) were assessed against Reef-wide wet season GVs to calculate magnitude scores for TSS, Chl-*a*, PP, and PN. The GVs were calculated based on annual GVs (Great Barrier Reef Marine Park Authority, 2010) that were seasonally adjusted as described in De’ath and Fabricius (2008) (see Appendix B Table B-4). Mean long-term concentrations of water quality parameters include samples collected from the enclosed coastal zone, where high TSS, Chl-*a*, PN, and PP concentrations are likely to contribute to exceedances of the Reef-wide GVs (see Appendix B Table B-1). The only GV currently available for Secchi depth is an annual mean, and thus comparison with wet season Secchi depth data was not possible.
2. **Production of the exposure maps:** The magnitude scores were used in combination with the seasonal, long-term, coral recovery, wet-year and dry-year frequency maps (described above)

to derive seasonal, long-term, coral recovery, wet-year, and dry-year exposure maps, respectively. Exposure from each map produced was then grouped into potential risk categories (I to IV) based on a “Natural Break (or Jenks)” classification<sup>1</sup> (Appendix B-3). The exposure classes were defined by applying the Jenks classification to the mean long-term (2003–2022) exposure map, because this map presented the highest number of observations (20 wet seasons). Category I and areas not exposed were re-grouped into a unique category corresponding to no or very low exposure to a potential risk. Magnitude scores have no designated ecological significance but are used in the risk framework as a relative measure to assign relative potential risk grading for each Reef water type.

3. *Exposure assessment:* Exposure maps were overlaid with information on the spatial distribution of coral reefs and surveyed seagrass meadows to identify areas and percentages of these ecosystems that may experience exposure to pollutants during the wet season. The area (km<sup>2</sup>) and percentage (%) of coral reefs and seagrass meadows affected by the different categories of exposure (I to IV) was calculated in the Reef and marine NRM regions. Exposure maps are presented in the context of the long-term reference period (average of 20 wet seasons), the representative coral recovery period (2012–2017), and typical wet-year and dry-year composites. Areas and percentages of exposure are presented in the context of the long-term reference period.

The methods are described in further detail in Appendix B. The ‘potential risk’ is influenced by the available satellite data on cloud-free days, with the likelihood of exposure likely to be underestimated in higher rainfall and areas with high cloud cover like the Wet Tropics and Cape York regions.

## 2.7 River discharge and catchment loads

River flow is reported annually and can be derived from several sources. In many cases, river flow gauges that measure discharge (and used to calculate constituent loads) are located well upstream of the river mouth and only capture a certain proportion of the catchment/basin area. Such disparities mean that river gauge data should not be directly compared across basins and NRM regions. For example, the Daintree and Barron Basins within the Wet Tropics region contain a similar area (2,100–2,200 km<sup>2</sup>); however, the Daintree River at Bairds and the Bloomfield River at China Camp gauges collectively only measure 56% of the Daintree Basin whereas the Barron River at Myola gauge captures 89% of the Barron Basin. If gauge data are used to compare discharge between these basins, the gauge on the Barron Basin is covering a much larger proportion of the area compared to the gauges on the Daintree Basin. A scaling factor is used on these data so that discharge (and constituent loads) can be directly compared across basins and NRM regions.

To account for these differences, the relevant discharge data for each basin were compiled, where available (Table 2-3; Department of Regional Development, Manufacturing and Water [DRMW], 2024). The total annual discharge for each gauge was then up-scaled using the recommended scaling factors outlined in Puignou Lopez *et al.* (in review). This work examined different upscale factors based on basin area to total gauged area, mean annual flow for the gauged basin and the mean annual basin flow from either the Bureau of Meteorology’s G2G model (BoM, 2017; Wells *et al.*, 2018), or the Source Catchments model (McCloskey *et al.*, 2021), or the linear relationships between river gauge data and the two models. The most appropriate upscale factor was then recommended for each basin (Puignou Lopez *et al.*, in review). Where a flow gauge did not exist in a basin (i.e. Jacky Jacky Creek, Lockhart River, Jeannie River, Proserpine River, Styx River, Shoalwater Creek and Boyne River—marked with an asterisk in Table 2-3), the gauge from the nearest neighbouring basin was used. The calculation of the long-term medians for each basin has been anchored to cover the 30-year period from 1990–91 to 2019–20 water years.

---

<sup>1</sup> Jenks is a statistical procedure, embedded in ArcGIS that analyses the distribution of values in the data and finds the most evident breaks in it (i.e., the steep or marked breaks; Jenks and Caspall 1971).

There were three flow gauges that were impacted by the extreme flows generated from Tropical Cyclone Jasper over the 2023–24 water year that needed to be addressed:

- The Annan River at Beesbike gauge had no flow data between 18 December 2023 and 14 March 2024 (inclusive). To fill this gap in the record, additional data were sourced (J. Shellberg CYWP, pers. comm.) that covered the stage height/discharge of the main flood event (18–20 December). For the remaining period, a linear correlation was established with the upstream flow gauge (Annan River at Collingwood) supplied by CYWP.
- Both flow gauges in the Daintree Basin (Daintree River at Bairds and Bloomfield River at China Camp gauges) also failed with the Tropical Cyclone Jasper event. The Daintree River at Bairds gauge had no flow data for the period 17 December 2023 to 16 August 2024 (inclusive) and so could not be used in the analysis. The Bloomfield River at China Camp gauge had missing data between 23 November 2023 and 27 January 2024, and a linear relationship was established between flow data from this gauge with the neighbouring gauges from the Annan River at Beesbike and the Mossman River at Mossman to infill this missing record. The resulting Bloomfield at China Camp record was then used with an area upscale factor of 8.0 to calculate discharge for the Daintree Basin.
- The Ross River at Aplins Weir data were unavailable for the 2003–24 water year, so the flow was estimated using an upscale from the Alligator Creek at Allendale gauge.

Table 2-3. The 35 basins of the Reef catchment, the gauges for each Basin, and the correction factors used to upscale flows to provide annual discharge estimates. (\*Australian Water Resources Council).

NRM Region	Basin	AWRC No.*	Basin area (km <sup>2</sup> )	Relevant gauges	% of Basin covered by key gauges	Correction factor
Cape York	Jacky Jacky Creek	101	2,963	Jardine River at Monument*	0	1.1x + 560,000
	Olive Pascoe River	102	4,180	Pascoe River at Garraway Creek	31	3.1
	Lockhart River	103	2,883	Pascoe River at Garraway Creek*	0	1.5
	Stewart River	104	2,743	Stewart River at Telegraph Road	17	5.6
	Normanby River	105	24,399	Normanby River at Kalpowar Crossing + Hann River at Sandy Creek (from 2005/06). Previous upscale period uses Normanby at Battle Camp + Hann River gauges with factor of 4.7	53	1.8
	Jeannie River	106	3,638	Endeavour River at Flaggy + Annan at Beesbike	0	3.2
	Endeavour River	107	2,182	Endeavour River at Flaggy + Annan at Beesbike	27	3.5x + 21,000

NRM Region	Basin	AWRC No.*	Basin area (km <sup>2</sup> )	Relevant gauges	% of Basin covered by key gauges	Correction factor
<b>Wet Tropics</b>	Daintree River	108	2,107	Daintree River at Bairds + Bloomfield River at China Camp	56	1.6
	Mossman River	109	473	Mossman River at Mossman	22	2.3
	Barron River	110	2,188	Barron River at Myola	89	1.3
	Mulgrave-Russell River	111	1,983	Mulgrave River at Peets Bridge + Russell River at Bucklands	42	2.0x + 450,000
	Johnstone River	112	2,325	South Johnstone River at Upstream Central Mill + North Johnstone at Tung Oil	57	1.6x + 540,000
	Tully River	113	1,683	Tully River at Euramo	86	1.1
	Murray River	114	1,107	Murray River at Upper Murray	14	5.0x + 600,000
	Herbert River	116	9,844	Herbert River at Ingham	87	1.2
<b>Burdekin</b>	Black River	117	1,057	Black River at Bruce Highway + Bluewater Creek at Bluewater	32	3.1
	Ross River	118	1,707	Ross River at Aplins Weir + Alligator Creek at Allendale (from 2001–02). Previous upscale period uses Ross River Dam HW + Bohle at Hervey Range Rd + Alligator Creek with factor of 1.6x + 75,000	52	1.9
	Haughton River	119	4,051	Haughton River at Powerline + Barratta at Northcote	62	1.6
	Burdekin River	120	130,120	Burdekin River at Clare	100	1.0
	Don River	121	3,736	Don River at Reeves + Elliot River at Guthalungra + Euri Creek at Koonandah (from 1999/00). Previous upscale period uses Don + Elliot gauges with factor of 2.9x + 170,000	46	1.5x + 210,000
<b>Mackay-Whitsunday</b>	Proserpine River	122	2,494	O'Connell River at Staffords Crossing + Andromache River at Jochheims + St Helens Creek at Calen	0	3.6

NRM Region	Basin	AWRC No.*	Basin area (km <sup>2</sup> )	Relevant gauges	% of Basin covered by key gauges	Correction factor
	O'Connell River	124	2,387	O'Connell River at Staffords Crossing + Andromache River at Jochheims + St Helens Creek at Calen	29	3.5
	Pioneer River	125	1,572	Pioneer River at Dumbleton Weir TW	95	1.1
	Plane Creek	126	2,539	Sandy Creek at Homebush + Carmila Creek at Carmila	16	5.6x + 210,000
<b>Fitzroy</b>	Styx River	127	3,013	Waterpark Creek at Byfield*	0	5.7x + 260,000
	Shoalwater Creek	128	3,601	Waterpark Creek at Byfield*	0	6.6x + 300,000
	Water Park Creek	129	1,836	Waterpark Creek at Byfield	12	5.4x + 43,000
	Fitzroy River	130	142,552	Fitzroy River at The Gap	95	1.1
	Calliope River	132	2,241	Calliope River at Castlehope	57	1.9x + 95,000
	Boyne River	133	2,496	Calliope River at Castlehope*	0	2.1
<b>Burnett-Mary</b>	Baffle Creek	134	4,085	Baffle Creek at Mimdale	34	2.4x + 95,000
	Kolan River	135	2,901	Kolan River at Springfield + Gin Gin Creek at Brushy Creek	37	2.4x + 19,000
	Burnett River	136	33,207	Burnett River at Figtree Ck (from 1996/97). Previous upscale period uses Burnett River at Mount Lawless with factor of 1.2x + 84,000	92	1.1
	Burrum River	137	3,362	Gregory River at Leeson's + Elliott River at Dr Mays Crossing + Isis River at Bruce Highway	40	3.0x + 27,000
	Mary River	138	9,466	Mary River at Home Park	72	1.4
* Gauges used which are not in the basin area						

Current annual and pre-development TSS, DIN and PN load estimates were calculated for all basins using a systematic approach. The DIN loads for the basins of the Wet Tropics and Haughton Basin were calculated using the model originally developed in Lewis *et al.* (2014) which uses a combination

of the annual nitrogen fertiliser applied in each basin coupled with basin discharge (calculated as per previous description). DIN loads for the Burdekin, Pioneer and Fitzroy basins were taken from those measured in the Great Barrier Reef Catchment Loads Monitoring Program<sup>2</sup>. If the measured data for the most recent years in these basins were unavailable, a mean of the long-term annual mean concentration from the previous monitoring data were coupled with the annual discharge to calculate a load. DIN loads for the remaining basins were calculated using an annual mean concentration which was multiplied by the corresponding annual basin discharge calculations. The annual mean concentration for each basin was informed using a combination of available monitoring data and Source Catchments model outputs. The pre-development DIN loads were calculated using a combination of the estimates from the Source Catchments model as well as available monitoring data from 'pristine' locations.

The TSS and PN loads were similarly determined through a step-wise process. For the basins where the Great Barrier Reef Catchment Loads Monitoring Program captured >95% of the basin area (e.g. Burdekin, Pioneer and Fitzroy) the measured/reported TSS and PN loads were used. If the measured data for the most recent years were unavailable, a mean of the long-term annual mean concentration from the previous monitoring data was coupled with the annual discharge to calculate a load. For other basins with monitoring data, the range of annual mean concentrations were compiled and compared with the latest Source Catchment modelled values. From these data a 'best estimate' of an annual mean concentration was produced and applied with the annual discharge data to calculate loads. Finally, for the basins that have little to no monitoring data, the annual mean concentration from the Source Catchments modelled data was examined along with nearest neighbour monitoring data to determine a 'best estimate' concentration to produce the load. The pre-development TSS and PN loads were calculated using a combination of the annual mean concentrations from the Source Catchments model and available monitoring data from 'pristine' or 'reference' locations. The corresponding discharge was used (as calculated previously) to produce a simulation of the pre-development load for the water year.

---

<sup>2</sup> <https://www.reefplan.qld.gov.au/tracking-progress/paddock-to-reef/modelling-and-monitoring>

### 3 DRIVERS AND PRESSURES INFLUENCING WATER QUALITY IN 2023–24

#### 3.1 Coastal development including agriculture

The Wet Tropics, Burdekin, and Mackay-Whitsunday regions are characterised by a variety of land uses including agricultural (sugarcane, grazing, cropping, and other horticulture), mining, and urban development. Parts of the Cape York region are less developed than other Reef catchments. Land-based activities in this region are assumed to have a reduced impact on marine ecosystems (Waterhouse *et al.*, 2017) despite a history of widespread grazing and mining impacts. Specifically:

- Cape York
  - The Pascoe River catchment has an area of 2,088 km<sup>2</sup> with a high proportion (84%) of nature/conservation land use with ~15% closed grazing (QLUMP, 2015). However, there is no longer any active grazing within the Pascoe catchment (Polglase pers. comm., February 2023). Feral cattle and pigs, fire, and road erosion are the main pressures affecting water quality. These impacts are considered to be minimal in this focus region relative to other Cape York and Reef catchments (Cape York NRM and South Cape York Catchments, 2016).
  - The Stewart River catchment has an area of 2,770 km<sup>2</sup> and is mostly nature/conservation land use (94%) with ~2% current grazing land use (QLUMP, 2015). However, feral cattle continue to graze much of the catchment area. Current and legacy cattle grazing impacts and road erosion are current pressures affecting sediment loads within the catchment.
  - The Normanby Basin is 24,550 km<sup>2</sup> and has a high proportion of nature/conservation land use (46%) and grazing (52%) (QLUMP, 2015). Additional lands have shifted from grazing to conservation since 2015, resulting in ~53% conservation land use and ~47% grazing. Horticulture accounts for only 1% of land use but has been expanding in the Laura and West Normanby sub-catchments. Current and historic cattle grazing, post-European initiation and acceleration of gully erosion, agricultural land clearing, alluvial mining, wildfires, and road erosion are the primary pressures affecting water quality across the Normanby catchment (Brooks *et al.*, 2013; Shellberg and Brooks, 2013; Cape York NRM and South Cape York Catchments, 2016; Spencer *et al.*, 2016). Horticulture in the Laura sub-catchment has also increased nutrient concentrations in the Laura River (Howley, 2020).
  - The Annan-Endeavour River Basin is 2,186 km<sup>2</sup> and has a high proportion of nature/conservation land use (52% as of 2015) and closed grazing (40%) (QLUMP, 2015). Additional grazing land has been converted to conservation land use since 2015 and approximately 80% of the Annan catchment is now under conservation or Aboriginal freehold. Sources of pollution in the Endeavour catchment include urban runoff from the township of Cooktown, cattle grazing, horticulture, and road erosion. Historic mining disturbances, cattle grazing impacts (current and historic), wildfires, and road erosion are the primary sources of pollution to the Annan River (Shellberg *et al.*, 2016).
- Wet Tropics
  - The Barron-Daintree focus region is primarily influenced by discharge from the Daintree, Mossman, and Barron catchments and, to a lesser extent, by other Wet Tropics rivers south of the focus region (Brodie *et al.*, 2013; Waterhouse *et al.*, 2017). The Daintree catchment is 2,107 km<sup>2</sup> and has a high proportion of protected areas (56% natural/minimal use lands and 32% forestry). The remaining area consists of 7% grazing and, to a lesser extent, sugarcane and urban areas. The Mossman catchment is 479 km<sup>2</sup> and consists of 76% natural/minimal use lands, 10% sugarcane, and smaller areas of grazing and urban land uses. The Barron catchment has an area of 2,189 km<sup>2</sup> and consists of 29% natural/minimal use lands, 31% grazing, 18% forestry, 11% cropping (including bananas and sugarcane), and smaller areas of dairy and urban land uses (Terrain NRM, 2015). The Barron River is the

- most hydrologically modified river in the Wet Tropics region and is heavily regulated by water supply infrastructure.
- The Russell-Mulgrave Basins contain a high proportion of upland National Park and forest (72%), with 13% of the area used for sugarcane production on the coastal floodplain (Terrain NRM, 2015). The Johnstone Basin is 2,326 km<sup>2</sup> and has a relatively high proportion of natural/minimal use lands (55%). The remaining area has 16% grazing, 12% sugarcane, and smaller areas of dairy (in the upper catchment), bananas and other crops, and urban land uses (Terrain NRM, 2015).
  - The Tully River Basin is 1,685 km<sup>2</sup> and has a high proportion of natural/minimal use lands (75%). The remaining area is comprised of 12% sugarcane, 4% bananas, 5% grazing, and smaller areas of forestry, other crops and urban land uses (Terrain NRM, 2015). The Murray River Basin has an area of 1,115 km<sup>2</sup> and has a high proportion of natural/minimal use lands (64%) (Terrain NRM, 2015). The remaining area is comprised of 14% sugarcane, 10% forestry, 6% grazing and smaller areas of bananas, other crops and urban land uses. The Herbert River Basin is 9,842 km<sup>2</sup> and consists of 27% natural/minimal use lands, 56% grazing, 8% sugarcane, and smaller areas of forestry (Terrain NRM, 2015).
  - The Burdekin region is one of the two large dry tropical catchment regions adjacent to the Reef. The region is primarily influenced by discharge from the Burdekin, Haughton, Ross and Black Rivers, with cattle grazing as the primary land use on over 95% of the catchment area (NQ Dry Tropics, 2016). There is also intensive irrigated sugarcane on the floodplains of the Burdekin and Haughton Rivers. Fluctuations in climate and cattle numbers greatly affect the state and nature of vegetation cover and, therefore, the susceptibility of soils to erosion and off-site transport of suspended sediments and associated nutrients.
  - The Mackay-Whitsunday region has a wet or mixed wet and dry tropical climate. The region is influenced by the Pioneer, Gregory, Proserpine, O'Connell, and Don Rivers. Catchment land use is dominated by agriculture broadly divided into grazing in the upper catchments (43%), sugarcane cultivation on the coastal plains (19%), and dispersed areas of nature conservation (19%) (Folkers *et al.*, 2014). In addition, there are expanding urban areas along the coast.

### 3.2 Climate and cyclone activity

Climate is a major driver of the condition of water quality and ecosystems and can vary substantially between years. It is heavily driven by the El Niño Southern Oscillation (ENSO) cycle. Climate models predict continued warming, increasing intensity of extreme rainfall events, fewer but more intense tropical cyclones, and more frequent and extreme La Niña and El Niño events (Schaffelke *et al.*, 2017).

The 2023–24 wet season had active cyclone influence in the inshore Reef (Figure 3-1), following on from low levels of cyclone activity in previous years. The two cyclones of note in the 2023–24 season were Tropical Cyclone Jasper and Tropical Cyclone Kirrily. Tropical Cyclone Jasper crossed the Queensland coast as a category 2 system near Wujal Wujal on 13 December 2023; it had previously peaked as a category 5 in the Coral Sea. Following its crossing, the cyclone quickly weakened to a tropical depression but stalled over Cape York for several days bringing major flooding for the rivers of the southern Cape York and northern Wet Tropics. Tropical Cyclone Kirrily crossed the coast near Townsville as a category 1 system on 25 January 2024; it peaked in the Coral Sea as a category 3 system but weakened greatly on its approach to the coast.




Figure 3-1: Trajectories of tropical cyclones affecting the Reef in the 2023–24 water year (1 October to 30 September) and in previous years.

### 3.2.1 Rainfall for the Reef, NRM regions, and basins

Queensland rainfall is highly variable on seasonal, inter-annual, and decadal timescales. Wet season (December to April) rainfall in 2023–24 was above the 30-year long-term average (1961 to 1990) in the Cape York, Wet Tropics and Burnett-Mary regions, similar to the average for the Burdekin and Fitzroy regions and below average for the Mackay Whitsunday region (Figure 3-2 and Figure 3-3).




Figure 3-2: Average daily wet season rainfall ( $\text{mm d}^{-1}$ ) in the Reef catchment (left) long-term daily average (1961–1990, produced by BoM), (centre) 2023–24 wet season, and (right) the difference between the long-term average and 2023–24 rainfall. Source data: Bureau of Meteorology (2024).




Figure 3-3: Difference between daily average wet season rainfall (December 2023–April 2024) and the long-term wet season rainfall average (from 1961–1990). Red and blue bars denote basins with rainfall below and above the long-term average, respectively. Note that the basins are ordered from north to south (left to right). Source data: Bureau of Meteorology (2024).

### 3.2.2 Freshwater discharge for the Reef, NRM regions, and basins

The volume of freshwater discharge across the Reef lagoon is closely related to rainfall during the wet season and has a significant influence on coastal water quality (Waterhouse *et al.*, 2024). The total annual freshwater discharge for all of the Reef basins, relative to long-term medians (based on water year, calculated using the methods described in Section 2.7), is shown in Figure 3-4. Discharge at the regional level is shown in Figure 3-5.

In 2023–24, the overall Reef catchment area had discharge 1.7 times the long-term median, which was the largest discharge since the 2018–19 water year. On a regional basis, the Cape York and Wet Tropics NRM regions had discharge well above their long-term medians (2.6 and 2.1 times above, respectively). Discharge from the Cape York region for the 2023–24 water year was the highest on record (in records extending back to the 1990–91 water year) and the discharge from the Wet Tropics was the highest since the 2010–11 water year. The Burnett-Mary NRM region also had above average discharge (1.7 times higher) while discharge from the Burdekin was 1.2 times the long-term median. Discharge in the Fitzroy region was very close to the long-term median. Discharge from the Mackay-Whitsunday NRM region was below the long-term median (0.8 times).

Annual discharge for each of the 35 Reef basins in 2023–24 is shown in Table 3-1 and compared to long-term median annual flows. With the exception of the Jacky Jacky Creek Basin, the basins within the Cape York NRM region had discharge greater than 1.5 times their long-term medians with the Normanby, Jeannie, and Endeavour Basins having discharge exceeding 3 times the long-term median. The Wet Tropics basins also experienced wetter-than-average conditions with discharge exceeding 3 times the respective long-term medians for the Daintree and Barron Basins, between 2–3 times the long-term median discharge in the Mossman and Herbert Basins, and the remaining Wet Tropics basins (Mulgrave-Russell, Johnstone, Tully, and Murray) had discharge between 1.5–2.0 times their long-term medians. The Burrum River Basin (Burnett-Mary NRM) recorded discharge that exceeded the long-term median by 3 times. The Waterpark Creek Basin (Fitzroy NRM region), Burnett and Mary

Basins (Burnett-Mary region) had discharges that were 2–3 times their respective long-term medians, while the Styx and Shoalwater Creek Basins (Fitzroy NRM) recorded discharge between 1.5–2.0 times the long-term medians. The basins in the Mackay-Whitsunday region all had discharge below their long-term medians.




Figure 3-4: Long-term total annual discharge in ML (water year: 1 October to 30 September) for the 35 main Reef basins. Source: DRMW (2024), <https://water-monitoring.information.qld.gov.au/>.




Figure 3-5: Corrected annual water year (1 October to 30 September) discharge from each NRM region (using the correction factors in Table 2-3) for 2003–04 to 2023–24 in (ML per year). Data derived from DRMW (2024).

Table 3-1: Annual water year discharge (ML) of the 35 main Reef basins (1 October 2020 to 30 September 2024, inclusive) and 30-year long-term (LT) median discharge (1990–91 to 2019–20). Colours indicate levels above the long-term median: yellow for 1.5 to 2 times, orange for 2 to 3 times and red greater than 3 times.

Basin	LT median	2020 - 2021	2021 - 2022	2022 - 2023	2023 - 2024
Jacky Jacky Creek	2,471,267	3,607,722	2,365,731	4,611,721	3,487,440
Olive Pascoe River	3,180,267	5,540,683	4,879,388	6,053,581	6,050,915
Lockhart River	1,538,839	2,680,976	2,360,994	2,929,152	2,927,862
Stewart River	758,172	1,419,942	569,738	1,366,633	1,100,668
Normanby River	3,864,344	6,149,878	3,562,637	11,791,399	16,300,347
Jeannie River	1,428,920	1,342,490	1,566,621	2,093,623	4,440,165
Endeavour River	1,583,881	1,489,348	1,734,492	2,310,900	4,877,431
Daintree River	1,918,174	1,834,774	2,519,318	4,685,640	9,176,968
Mossman River	604,711	654,566	800,754	815,267	1,745,893
Barron River	622,447	667,265	692,908	1,217,590	3,603,793
Mulgrave-Russell River	4,222,711	4,771,460	4,091,750	4,291,804	6,786,526
Johnstone River	4,797,163	5,324,040	4,712,174	5,385,426	8,157,637
Tully River	3,393,025	4,123,338	3,175,489	3,660,701	5,563,920
Murray River	1,484,246	1,947,050	1,269,280	1,526,232	2,595,878
Herbert River	3,879,683	6,842,168	3,283,590	4,919,143	8,516,360
Black River	293,525	429,282	273,677	353,756	526,432
Ross River	279,376	232,975	202,811	209,681	285,424
Haughton River	558,735	595,709	735,754	1,219,825	583,152
Burdekin River	4,406,780	8,560,072	5,442,976	9,702,259	5,745,479
Don River	496,485	510,906	383,927	999,723	372,511
Proserpine River	859,348	537,613	446,839	1,869,821	618,392
O'Connell River	835,478	522,680	434,427	1,817,882	601,214
Pioneer River	616,216	235,359	277,610	761,905	589,249
Plane Creek	1,058,985	600,958	489,222	1,440,350	632,961
Styx River	629,037	927,219	1,080,829	849,506	1,030,316
Shoalwater Creek	727,306	1,072,570	1,250,433	982,586	1,191,945
Water Park Creek	392,614	675,102	820,627	601,479	772,773
Fitzroy River	2,875,792	436,730	4,505,289	3,078,896	2,100,507
Calliope River	257,050	123,050	250,551	135,396	172,394
Boyne River	179,108	31,002	171,925	44,649	85,541
Baffle Creek	347,271	112,323	1,000,587	170,693	424,436
Kolan River	115,841	19,211	818,716	83,734	139,893
Burnett River	264,307	118,241	3,894,616	358,852	598,898
Burrum River	130,835	44,691	1,612,683	270,059	476,512
Mary River	908,873	420,909	10,139,380	673,298	1,806,668
Sum of basins	60,746,947	64,602,302	71,817,742	83,283,163	104,086,501

## 4 SATELLITE REMOTE SENSING OF MARINE WATER QUALITY

This section presents results from satellite remote sensing of wet season water quality.

### 4.1 Satellite remote sensing of Reef water types

To illustrate wet season influence on coastal water quality and identify potential risk to ecosystems, satellite-derived map products were produced for the Reef. This includes frequency maps predicting the areas affected by the Reef WT1–2 combined (Figure 4-1) or the three Reef water types individually (Figure 4-2) from December 2023 to April 2024 (the 2023–24 wet season).

#### 4.1.1 Areas affected

The extent and frequency of the occurrence of combined Reef WT1 and WT2 was variable across regions, cross-shelf, and between years, reflecting the concentrations and intensity of the river discharge and resuspension events (Figure 4-1). A well-documented inshore to offshore gradient is documented (for example, Devlin *et al.*, 2013, 2015), with coastal areas experiencing the highest frequency of the Reef WT1 and mid-shelf and offshore areas less frequently exposed to the Reef WT1 (Figure 4-2).

**Frequency of occurrence:** The frequencies of occurrence of the combined Reef WT1 and WT2 were lower than the mean long-term frequencies in the southern Reef, including the Mackay-Whitsunday, Fitzroy, and Burnett-Mary regions (Figure 4-1a,e,f), indicating drier conditions. Frequencies were greater in Cape York and the Wet Tropics and closer to the long-term average in the Burdekin region, but with some local variability. These results agreed with the rainfall distribution in 2024 (Figure 3-2). The frequencies of occurrence measured across the Tully focus region were above the long-term average (Figure 4-1g).

**Reef area exposed:** In 2023–24 only 4% of the Reef was exposed to the Reef WT1, 19% of the Reef was exposed to the Reef WT2 and 61% of the Reef was exposed to the Reef WT3 (Figure 4-3b and Table C-10). The area exposed to the Reef WT1 was slightly above both the long-term and coral recovery period percentages (3% of the Reef) and only the inshore (enclosed coastal and open coastal) Reef waters were exposed (Figure 4-3c: 75% and 22% of the total enclosed coastal and open coastal waterbody areas, respectively). The area exposed to the Reef WT2 was slightly above both the long-term and coral recovery period percentages (17 and 16% of the Reef) and 93% of the total open coastal, 61% of the total enclosed coastal and 25% of the total mid-shelf waterbody areas were exposed.




Figure 4-1: Map showing the frequency of the Reef WT1–2 combined in the a) long-term (20 wet seasons since 2002–03) b) representative coral recovery period (2011–12 to 2016–17, 6 wet seasons or 132 weeks), c) typical wet-year composite, d) typical dry-year wet season composites and e) 2023–24 wet season (22 weeks). The 2023–24 frequency maps were produced using Sentinel-3 images and the FU colour scale. Previous wet seasons (prior to 2021) and reference period composites have been produced using both MODIS satellite imagery and the wet season colour scale (Waterhouse *et al.*, 2021) and Sentinel-3 satellite imagery and the Forel-Ule colour scale (from 2021). The highest frequency is shown in orange and the lowest frequency is shown in blue. f) Difference map showing areas with an increase (in blue) and decrease (in brown) in exposure to Reef WT1–2 in 2023–24 against long-term trends (calculated as (e) 2024 minus (a) long-term). g) Plots on the right show the frequency values recorded along three transects extending from the Tully, Burdekin, and Pioneer Rivers to the external boundaries of the Marine Park and illustrate the differences in the spatial distribution and frequency of occurrence between the different representative periods. OC: open coastal, M: mid-shelf and O: offshore marine water body boundaries.




Figure 4-2: Map showing the frequency of the Reef WT1 (FU10–21), WT2 (FU6–9), and WT3 (FU4–5) in the 2023–24 wet season (22 weeks). The highest frequency is shown in orange and the lowest frequency is shown in blue. These maps are used in the exposure assessment to represent the spatial likelihood of exposure of each of the wet season water types in 2023–24.




Figure 4-3: Areas (km<sup>2</sup>) and percentages (%) of the Reef lagoon (total 348,839 km<sup>2</sup>) affected by the different Reef water types including division by waterbodies (enclosed coastal, open coastal, mid-shelf, and offshore) and waters affected by the Reef WT1–2 combined, and the three Reef water types individually during the current wet season and for a range of reference periods (24: 2023–24 wet season, LT: long-term, CR: Coral Recovery, W: Wet years, and D: Dry years). The data are presented in detail in Table C-10.

As described in Section 2.6, the multi-year reference maps (long-term, wet, and dry years) were reviewed in 2023 to include 20 years of satellite imagery (Figure 4-1). In 2023–24, as in the three previous years, the Reef area exposed to Reef WT3 (61% of the Reef) was greater than the long-term average (53% of the Reef) and the ‘wet’ year’s area (58% of the Reef). This result is related to large areas exposed to Reef WT3 measured in the mid-shelf and offshore Reef (96% and 48% of the total mid-shelf and offshore waterbody areas, respectively). This result is not fully understood but is likely an indication of shelf upwelling in the central and southern Reef areas and is not related to direct riverine influence. Image classification by optical type does not directly elucidate the cause of variations in water colour, and Reef WT3 in particular (but also, to some extent, Reef WT1 and WT2 in some coastal areas) is sometimes due to oceanographic processes not related to catchment runoff. This should be further investigated in a future case study by comparing Reef WT3 areas with sea-surface temperature climatology (for example, Wijffels *et al.*, 2018). Furthermore, major reprocessing of the MODIS and Sentinel 3-OLCI radiance was undertaken in 2018 and 2021, respectively. This could have influenced the result of the classification of the colour classes, particularly for the clearest waters. Reef WT3 is associated with low land-based contaminant concentrations and has a low magnitude score in the Reef exposure assessment (Figure 4-4 and Figure 4-5). While Reef WT3 areas were larger than in the reference periods, this did not result in increasing the potential risk offshore as 100% of the offshore areas were classified as no/very low potential risk in the 2023–24 exposure assessment (Figure 4-6 and Figure 4-7).

#### 4.1.2 Composition of Reef water types

All mean concentrations of water quality parameters were reviewed in 2023 using water quality data (surface samples) collected from 2004 to April 2023, and the archive of weekly MODIS-Aqua and Sentinel-3 water type maps. Detailed summaries of water quality parameters for the long-term period (20 wet seasons) and reporting year are provided in [Appendix B](#). Boxplots of long-term water quality parameters are shown in Figure 4-4 and are fully described below.

Mean long-term concentrations of water quality parameters showed similar patterns between focus regions, with maximum concentrations measured in Reef WT1 and minimum concentrations in Reef WT3 (Figure 4-4). However, there were distinct differences in the concentrations of individual parameters across regions. Across years, the frequency of sampling in flood events as well as the location, timing, and number of samples historically collected in each region is a major influence on these results. Thus, the *magnitude scores* for the exposure maps are calculated using the mean long-term concentrations of water quality parameters across the whole of the Reef (Section 4.1.3 and Figure 4-5).

The long-term mean TSS, Chl-*a*, PP, and PN concentrations ( $\pm$  standard deviation) were above the Reef-wide wet season guideline values (GVs) in the Reef WT1 and WT2, with Chl-*a* only slightly above in Reef WT2 (Chl-*a* =  $0.65 \pm 0.75 \mu\text{g L}^{-1}$ ) and only the long-term mean PN concentration was above the wet season GV in the Reef WT3 (PN =  $28.30 \pm 21.31 \mu\text{g L}^{-1}$ ) (Figure 4-5). Using these data, magnitude scores in the exposure mapping were calculated as the proportional exceedance of the GV, and negative magnitude scores capped to zero (Figure 4-5). Magnitude scores have no defined ecological significance but are used in the risk framework as a relative measure to assign potential risk grading for each Reef water type (refer Section 4.1.3).




Figure 4-4: Long-term (2004–2023) concentrations of water quality parameters and Secchi disk depth boxplots for each Reef Water Type (WT1, WT2, and WT3). Water types were extracted from the MODIS-Aqua (2004–2020) and Sentinel-3 (2021–2023) weekly satellite databases. The mean is plotted as a cross (x) and its numerical value is indicated in text. The interquartile range is delimited by the box and the median by the line inside the box. Whiskers indicate variability outside the upper and lower quartiles. Data beyond the whiskers range are considered outliers and are not plotted.




Figure 4-5: Mean long-term concentrations of water quality parameters (top) and magnitude scores across the three Reef water types (bottom). Red lines show the Reef-wide wet season GV's (Appendix B Table B-4). Magnitude scores are calculated as the proportional exceedance of the guideline:  $magnitude_{water\ type} = ([Poll.]_{water\ type} - GV)/GV$  and Poll. = TSS, Chl-a, PP, or PN. Negative magnitude scores are scored as zero.

#### 4.1.3 Potential exposure risk to Reef ecosystems

This section presents the area (km<sup>2</sup>) and percentage (%) of coral reefs and seagrass meadows affected by different categories of exposure (or potential risk) based on satellite-derived Reef water types.

The exposure categories are not validated against ecological health data and represent relative potential risk categories for seagrass and coral reef ecosystems. The areas and percentages of ecological communities affected by the different categories of exposure were calculated as a relative measure between regions and the long-term average.

**Reef-wide:** The area exposed to a potential risk in 2023–24 was spatially limited relative to the scale of the Reef with 90% exposed to no or very low potential risk (Table 4-1 and Figure 4-6e). This result is similar to the long-term patterns (91% of the Reef). Approximately 9% of the Reef was exposed to combined potential risk categories II–IV, which is still a relatively large area at approximately 31,673 km<sup>2</sup>. However, only 1% of the Reef was in the highest exposure category (IV) and only 2% of the Reef was in category III (Table 4-1); the total area of these categories combined was 10,329 km<sup>2</sup>. These patterns were very similar to the long-term patterns (Table 4-1). Patterns were also similar across marine regions, with more than 84% of each region classified as no / very low risk and less than 4% classified as category III or category IV, respectively (Figure 4-7b). It is important to note that while these percentages are relatively small, the total areas are still significant, especially when considering specific habitat areas.

**Reef waterbodies:** Only the inshore Reef waters, including the enclosed (macro-tidal enclosed coastal and enclosed coastal waterbodies combined) and open coastal (macro-tidal open coastal and open coastal waterbodies combined) were exposed to the highest categories of potential risk (III and IV, Figure 4-7a). However, open coastal waters were largely exposed to the lowest category of potential risk only (no/very low risk = 44% and II: 45%) and only 11% and 1% of the open coastal waters were exposed to the potential risk category III and IV. The enclosed coastal waters had the largest proportion of waters classified as higher relative risk, with 50% of the combined inshore waters exposed to risk category IV. Approximately 77% (<3,600 km<sup>2</sup>) of Reef seagrasses occur in the inshore waters, but only 4% (<900 km<sup>2</sup>) of

coral reefs (Appendix C-6). The mid-shelf and offshore waterbodies were largely classified as no/very low potential risk (95% of the mid-shelf and 100% of the offshore waters) (Figure 4-7a).

Table 4-1: Areas (km<sup>2</sup>) and percentages (%) of the Reef, Reef-wide coral reefs, and Reef-wide surveyed seagrass affected by different risk categories of exposure during the 2023–24 wet season and the long-term (2003–2022). The last three rows show the differences between % affected in 2023–24 and the long-term average (■: increase, ■: decrease, and ■: no change, difference <5%). Areas south of the Marine Park (Hervey Bay) are not included.

Total surface area covered		Total		Potential Risk category				Total area exposed II–IV
				No / very low	Lowest		Highest	
					I	II	III	
Reef-wide	area	348,839	2024	317,166	21,344	6,200	4,129	31,673
			LT	317,183	23,596	4,247	3,813	31,657
	%	100%	2024	90%	6%	2%	1%	9%
			LT	91%	7%	1%	1%	9%
Reef-wide coral reefs	area	24,914	2024	23,285	1,253	226	150	1,629
			LT	24,072	564	179	99	842
	%	100%	2024	93%	5%	1%	1%	7%
			LT	97%	2%	1%	0%	3%
Reef-wide surveyed seagrass	area	4,660	2024	1,328	1,706	950	675	3,331
			LT	1,373	1,943	609	734	3,287
	%	100%	2024	29%	37%	20%	14%	71%
			LT	29%	42%	13%	16%	71%
Difference (2023–24 to long-term average)	Reef-wide			<-1%	-1%	1%	<1%	<-1%
	Reef-wide coral reefs			-3%	3%	<1%	<1%	3%
	Reef-wide surveyed seagrass			-1%	-5%	7%	-1%	1%

Similar cross-shore patterns were observed across Reef marine regions and all mid-shelf and offshore waterbodies were largely classified as no/very low potential risk (Figure 4-7c). Mid-shelf waterbodies in the Wet Tropics region had the greatest exposure to potential risk category II (24% of the Wet Tropics mid-shelf waterbody), followed by the Cape York region (18% of the Cape-York mid-shelf waterbody). The Wet Tropics and Cape York region open coastal waterbodies had the greatest exposure to risk categories III (25% and 19% of the Wet Tropics and Cape York open coastal waterbodies, respectively), followed by the Burdekin region (15% of the Burdekin region open coastal waterbody). In the other Reef regions, <10% of the open coastal waterbodies were exposed to risk categories III. Differences across regions are further described in the Regional Results section below (Section 4.2).




Figure 4-6: Map showing the reclassified surface exposure in the a) long-term (20 wet seasons since 2002–03), b) representative coral recovery period (2012–2017, 132 weeks), c) typical wet-year and d) typical dry-year wet season composites and e) 2023–24 wet seasons (22 weeks). Relative potential risk categories range from I: no/low risk to IV: highest relative risk. f) Difference map showing areas with an increase (in red, ⬇️) and decrease (in purple, ⬆️) in risk category in 2023–24 against long-term trends (calculated as (e) 2024 minus (a) long-term).




Figure 4-7: Percentage of the a) Reef waterbodies, b) Reef regions, c) regional Reef waterbodies, d) seagrass and e) coral habitats affected by different risk categories of exposure during the 2023–24 wet season. Water body classifications are shown along the x-axes: enclosed coastal (EC), open coastal (OC), mid-shelf (Mid), and offshore (Off).

**Reef habitats (coral reefs and seagrasses):** In 2023–24, it was estimated that:

- Approximately 7% of coral reefs (or 1,629 km<sup>2</sup>) were exposed to combined potential risk categories II–IV (Table 4-1). However, <2% were in the highest exposure categories IV and III combined and only the enclosed coastal and open coastal coral reef habitats were exposed, equating to 377 km<sup>2</sup>. The total enclosed coastal coral reef area affected by the highest exposure categories was 89% (57% to cat. IV and 32% to cat. III, Figure 4-7e). Only 5% of the open coastal reefs were exposed to cat. IV and 27% to category III. Mid-shelf and offshore coral reefs were only exposed to the lowest risk category II or to no potential risk. The coral areas exposed to potential risk categories III and IV in 2023–24 were similar to the long-term patterns (<2% of the coral reefs, Table 4-1).
- Approximately 71% of seagrasses (or 3331 km<sup>2</sup>) were exposed to combined potential risk categories II–IV. Approximately 14% (675 km<sup>2</sup>) were in the highest exposure category (IV) and 20% were in category III (950 km<sup>2</sup>) and only the enclosed coastal and open coastal seagrass habitats were exposed (Figure 4-7d). The total enclosed coastal seagrass area affected by the highest exposure categories was 85% (45% to cat. IV and 40% to cat. III, Figure 4-7d). 17% of the open coastal seagrasses were exposed to cat. III and 1% only were exposed to the highest category IV. Mid-shelf and offshore seagrasses were only exposed to the lowest risk category II or to no potential risk. The seagrass areas exposed to combined potential risk categories II to IV in 2023–24 were similar to the long-term areas. There was, however, an increase in area exposed to the potential risk category III: +7%.

## 4.2 Regional exposure of coastal waters and ecosystems to wet season discharge

The results of the remote sensing analysis for each region are presented below. This provides smaller-scale interpretation of the results, which can be highly variable between locations, thereby enhancing the relevance of the remote sensing products for regional managers.

### 4.2.1 Cape York region

As described for the Reef, a number of remote sensing products were generated to represent wet season water quality conditions in the Cape York region. These maps are presented in Figure 4-8, which presents the frequency of the combined Reef WT1–2, the frequency of Reef WT1, WT2, and WT3 individually; the exposure maps - each in the long-term and 2023–24 wet season; and a difference map showing areas exposed to an increased risk in 2023–24.

Sampling of the Cape York waters occurred during and between the main flood events. A full description of water quality patterns and flood plumes is available in [Section 5](#) of this report.




Figure 4-8: Long-term and current year remote sensing results for the Cape York region showing the a) frequency of combined Reef WT1–2; b) the frequency of Reef WT1, WT2 and WT3 individually regrouped into five likelihood categories [ $<0.2$  (Rare),  $0.2\text{--}0.4$ ,  $0.4\text{--}0.6$ ,  $0.6\text{--}0.8$  and  $0.8\text{--}1$  (very frequent)]; c) exposure to potential risk - each in the long-term (bottom) and 2023–24 wet season (top). d) Difference map showing any areas with an increase (in red,  $\oplus$ ) or decrease (in purple,  $\ominus$ ) in risk category in 2023–24 against long-term trends [calculated as (c, top) exposure in 2024 minus (c, bottom) long-term]. Note that optical water types – especially the Reef WT3– do not always correspond to direct catchment discharge influence and can also be due to oceanographic processes (see definitions in Table 2-2).

Table 4-2 presents the areas (km<sup>2</sup>) and percentages (%) of Cape York region, coral reef, and seagrass areas affected by different categories of exposure (or potential risk) based on satellite-derived Reef water types. The exposure categories are not validated against ecological health data and represent relative potential risk categories for seagrass and coral reef ecosystems. Category I (no/very low risk) represents waters with ambient or detectable, but low concentrations of the water quality parameters investigated and therefore low risk of any detrimental ecological effect. The areas and percentages of ecological communities affected by the different categories of exposure were calculated as a relative measure between regions and the long-term average.

In 2023–24, it was estimated that:

- **Cape York region:** Approximately 89% of Cape York was not exposed to a potential risk, which was below the long-term patterns (93%, Table 4-2). Approximately 11% (or about 10,300 km<sup>2</sup>) of the Cape York region was exposed to combined potential risk categories II–IV, with 1% (1032 km<sup>2</sup>) of the Cape York region in the highest exposure category (IV) and 2% (2,007 km<sup>2</sup>) in category III.
- **Cape York waterbodies:** The mid-shelf and offshore waterbodies were largely exposed to no/very low risk (82% and 99% of the Cape York mid-shelf and offshore waterbodies, Figure 4-7c). Only the enclosed coastal and open coastal Cape York waters were exposed to the highest categories of potential risk (III and IV). The area exposed corresponded to 40% (cat. III) and 44% (cat. IV) of the total Cape York enclosed coastal area (Figure 4-7c) and 19% (cat. III) and 2% (cat. IV) of the open coastal areas.
- **Cape York habitats:**
  - **Coral reefs:** Approximately 89% of coral reefs in the Cape York region were not exposed to a potential risk, which was below the long-term trends (93%). About 1% of corals were in the highest exposure category (IV) and 1% in category III (combined 179 km<sup>2</sup>) and they were all inshore and enclosed coastal reefs (Figure 4-9a). Approximately 1% and 2% (<300 km<sup>2</sup>) of the Cape York corals reefs occur in the enclosed and open coastal waters, respectively (Appendix C-6).  
The coral area exposed to higher potential risk corresponded to 36% (cat. III) and 55% (cat. IV) of the total enclosed coastal coral reef area in Cape York, and to 31% (cat. III) and 10% (cat. IV) of the total open coastal coral reef area (Figure 4-9a). Mid-shelf reefs were exposed to the lower risk category II or to no/very low risk (16% and 84% of the total mid-shelf coral reef area in Cape York). 94% of the Cape York offshore reefs were classified as no/very low risk.
  - **Seagrasses:** Approximately 66% (or 1,751 km<sup>2</sup>) of seagrasses in the Cape York region were exposed to combined potential risk categories II–IV (Table 4-2). 9% (252 km<sup>2</sup>) of seagrasses were in the highest exposure category (IV) and 18% were in category III (484 km<sup>2</sup>), and they were all inshore and enclosed coastal seagrasses (Figure 4-9b). A total of 27% and 40% (~1,800 km<sup>2</sup>) of the Cape York seagrass occur in the enclosed and inshore waters respectively (Appendix C-6).  
The seagrass area exposed to higher potential risk corresponded to 46% (cat. III) and 32% (cat. IV) of the total enclosed coastal seagrass area in Cape York and to 14% (cat. III) and 2% (cat IV) of the total open coastal seagrass area (Figure 4-9b). Mid-shelf and offshore seagrasses were largely classified as no/very low risk (88% and 77% of the Cape York mid-shelf and offshore seagrasses).
  - **Comparison to long-term trends:** The coral areas exposed to highest potential risk categories III and IV were similar to the long-term patterns (<3% of the coral reefs). There was however an increase in the total coral area exposed to the lowest potential risk category (II: +7%). There was also an increase in the total seagrass area exposed to the middle potential risk category (III: +8%).

Table 4-2: Areas (km<sup>2</sup>) and percentages (%) of the Cape York region, Cape York coral reefs, and Cape York surveyed seagrass affected by different categories of exposure during the 2023–24 wet season and the long-term (2003–2022).

The last three rows show the differences between % affected in 2023–24 and the long-term average (■: increase, ■: decrease, ■: no change, difference <5%).

decrease,  , no change, difference <5%).

Total surface area covered		Total		Potential Risk category				Total area exposed II–IV
				No / Very low	Lowest		Highest	
					I	II	III	
Cape York region	area	96,316	2024	86,009	7,268	2,007	1,032	10,307
			LT	89,387	5,351	1,026	553	6,929
	%	100%	2024	89%	8%	2%	1%	11%
			LT	93%	6%	1%	1%	7%
Cape York coral reefs	area	10,375	2024	9,283	912	98	81	1,092
			LT	10,030	209	92	44	345
	%	100%	2024	89%	9%	1%	1%	11%
			LT	97%	2%	1%	0%	3%
Cape York surveyed seagrass	area	2,655	2024	904	1,015	484	252	1,751
			LT	1,175	1,012	265	203	1,480
	%	100%	2024	34%	38%	18%	9%	66%
			LT	44%	38%	10%	8%	56%
Difference (2023–24 to long-term average)	Cape York Region			-4%	2%	1%	<1%	4%
	Cape York coral reefs			-7%	7%	<1%	<1%	7.2%
	Cape York surveyed seagrass			-10%	<1%	8%	2%	10%




Figure 4-9: Percentage of the Cape York region a) coral reef and b) surveyed seagrass habitats affected by different risk categories of exposure during the 2023–24 wet season. Water body classifications are shown along the x-axes: enclosed coastal (EC), open coastal (OC), mid-shelf (Mid), and offshore (Off).

#### 4.2.2 Wet Tropics region

As described for the Reef, a number of remote sensing products were generated to represent wet season water quality conditions in the Wet Tropics region. These maps are presented in

Figure 4-10, which presents the frequency of the combined Reef WT1–2, the frequency of Reef WT1, WT2, and WT3 individually; the exposure maps – each in the long-term and 2023–24 wet season; and a difference map showing areas exposed to an increased risk in 2023–24.




Figure 4-10: Long-term and 2023–24 remote sensing results for the Wet Tropics region showing the a) frequency of combined Reef WT1–2 II; b) the frequency of Reef WT1, WT2, and WT3 individually regrouped into five likelihood categories [ $<0.2$  (Rare),  $0.2–0.4$ ,  $0.4–0.6$ ,  $0.6–0.8$  and  $0.8–1$  (very frequent)]; c) exposure to potential risk - each in the long-term (bottom) and 2023–24 wet season (top). d) Difference map showing any areas with an increase (in red,  $\oplus$ ) or decrease (in purple,  $\ominus$ ) in risk category in 2023–24 against long-term trends [calculated as (c, top) exposure in 2024 minus (c, bottom) long-term]. Note that optical water types – especially Reef WT3– do not always correspond to direct catchment discharge and can also be due to oceanographic processes (see definitions in Table 2-2).

Table 4-3 presents the areas (km<sup>2</sup>) and percentage (%) of Wet Tropics region, coral reef, and seagrass areas affected by different categories of exposure (or potential risk) based on satellite-derived wet season water maps.

The exposure categories are not validated against ecological health data and represent relative potential risk categories for both seagrass and coral reef ecosystems. Category I (no/very low risk) represents waters with ambient or detectable but low concentrations of the water quality parameters investigated and therefore low risk of any detrimental ecological effect. The areas and percentages of ecological communities affected by the different categories of exposure were calculated as a relative measure between regions and the long-term average.

In 2023–24, it was estimated that:

- Wet-Tropics wide: 84% of the Wet Tropics region was not exposed to a potential risk, which was below the long-term patterns (88%, Table 4-3). 16% (or about 5,073 km<sup>2</sup>) of the Wet Tropics region was exposed to combined potential risk categories II–IV. However, only 1% (443 km<sup>2</sup>) of the region was in the highest exposure category (IV) and 3% was in category III (1,036 km<sup>2</sup>).
- Wet Tropics waterbodies: only the enclosed coastal and open coastal Wet Tropics waters were exposed to the highest categories of potential risk (III and IV). The open coastal area exposed corresponded to 25% (cat. III) and 2% (cat. IV) of the total Wet Tropics inshore area (Figure 4-7c). A total of 34% and 60% of the enclosed coastal areas were exposed to categories III and IV, respectively. The mid-shelf and offshore Wet Tropics waterbodies were largely exposed to no/very low risk (76% and 99% of the mid-shelf and offshore waterbodies) and 24% of the mid-shelf Wet tropics waterbody was also exposed to the lowest category of risk (II, 24%).
- Wet Tropics habitats:
  - **Coral reefs:** 3% of coral reefs in the Wet Tropics region were exposed to a potential risk (combined potential risk categories II–IV, Table 4-3). Less than 1% of coral were in the highest exposure category (IV), 1% were in the category III (combined 37 km<sup>2</sup>) and they were all enclosed coastal or open coastal reefs (Figure 4-11a). Only 3% (~80 km<sup>2</sup>) of the Wet Tropics corals occur in the inshore waters (Appendix C-6).

The open coastal coral area exposed to higher potential risk corresponded to 45% (cat. III) and 12% (cat. IV) of the total open coastal reef area in the Wet Tropics (Figure 4-11a), which were larger areas than the previous 2022–23 wet season (+16% and +3%, respectively). A total of 22% and 76% of the enclosed coastal areas were exposed to categories III and IV respectively (compared to 53% exposed to cat. IV last year). Mid-shelf and offshore reefs were largely exposed to no potential risk (>90% of the total mid-shelf and offshore reef areas in the Wet Tropics).

- **Seagrasses:** A total of 98% (or 229 km<sup>2</sup>) of seagrasses in the Wet Tropics region were exposed to a potential risk (combined potential risk categories II–IV, Table 4-3). A total of 29% (67 km<sup>2</sup>) of seagrasses were in the highest exposure category (IV) and 38% (89 km<sup>2</sup>) were in category III, and they were all inshore seagrasses (Figure 4-11b). 98% (~230 km<sup>2</sup>) of the Wet Tropics seagrass occur in the inshore waters (Appendix C-6).

The open coastal seagrass area exposed to potential risk corresponded to 36% (cat. III) and 2% (cat. IV) of the total inshore coastal seagrass in the Wet Tropics (Figure 4-11b). A total of 41% and 48% of the total enclosed coastal seagrass were exposed to categories III and IV, respectively. Mid-shelf seagrasses were classified as no/very low risk (66% of the Wet Tropics mid-shelf seagrasses) or as the lowest category of risk (II, 33% of the Wet Tropics mid-shelf seagrasses).

- **Comparison with long-term trends:** While the total area in the Wet Tropics region exposed to combined potential risk categories II–IV in 2023–24 was above the average long-term areas (+4%), this did not significantly impact the ecosystem exposure results. The total coral and seagrass areas exposed to the risk categories II–IV were similar to the long-term patterns (changes  $\leq 1\%$ ). There was furthermore a decrease in the seagrass area exposed to the highest potential risk categories (IV: -12% ) toward the lowest potential risk category (II: +9%).

Table 4-3: Areas (km<sup>2</sup>) and percentages (%) of the Wet Tropics region, Wet Tropics coral reefs, and Wet Tropics surveyed seagrass affected by different risk categories of exposure during the 2023–24 wet season and the long-term (2003–2022). The last three rows show the differences between % affected in 2023–24 and the long-term average (■: increase, ■: decrease, ■: no change, difference <5%).

decrease,  : No change, difference <0.7%.

Total surface area covered		Total		Potential Risk category				Total area exposed II–IV
				No / Very low	Lowest		Highest	
					I	II	III	
Wet Tropics region	area	31,976	2024	26,903	3,594	1,036	443	5,073
			LT	28,022	2,912	605	436	3,953
	%	100%	2024	84%	11%	3%	1%	16%
			LT	88%	9%	2%	1%	12%
Wet Tropics coral reefs	area	2,425	2024	2,348	41	24	13	78
			LT	2,349	41	29	7	77
	%	100%	2024	97%	2%	1%	1%	3%
			LT	97%	2%	1%	0%	3%
Wet Tropics surveyed seagrass	area	232	2024	4	72	89	67	229
			LT	5	51	82	95	227
	%	100%	2024	2%	31%	38%	29%	98%
			LT	2%	22%	35%	41%	98%
Difference (2023–24 to long-term average)	Wet Tropics region			-4%	2%	1%	<1%	4%
	Wet Tropics coral reefs			<-1%	<1%	<-1%	<1%	<1%
	Wet Tropics surveyed seagrass			<-1%	9%	3%	-12%	<1%




Figure 4-11: Percentage of the Wet Tropics region a) coral reef and b) surveyed seagrass habitats affected by different risk categories of exposure during the 2023–24 wet season. Water body classifications are shown along the x-axes: enclosed coastal (EC), open coastal (OC), mid-shelf (Mid), and offshore (Off).

### 4.2.3 Burdekin region

As described for the Reef, a number of remote sensing products were generated to represent wet season water quality conditions in the Burdekin region. These maps are presented in Figure 4-12, which presents the frequency of the combined Reef WT1–2; the frequency of Reef WT1, WT2, and WT3 individually; the exposure maps – each in the long-term and 2023–24 wet season; and a difference map showing areas exposed to an increased risk in 2024.




Figure 4-12: Long-term and current year remote sensing results for the Burdekin region showing the a) frequency of combined Reef WT1–2; b) the frequency of Reef WT1, WT2, and WT3 individually regrouped into five likelihood categories [ $<0.2$  (Rare),  $0.2\text{--}0.4$ ,  $0.4\text{--}0.6$ ,  $0.6\text{--}0.8$  and  $0.8\text{--}1$  (very frequent)]; c) exposure to potential risk - each in the long-term (bottom) and 2023–24 wet season (top). d) Difference map showing any areas with an increase (in red,  $\uparrow$ ) or decrease (in purple,  $\downarrow$ ) in risk category in 2023–24 against long-term trends [calculated as (c, top) exposure in 2024 minus (c, bottom) long-term]. e) Close ups to the Magnetic Island region. Note that optical water types – especially the Reef WT3 – do not always correspond to direct catchment discharge and can also be due to oceanographic processes (see definitions in Table 2-2).

Table 4-4 presents the areas (km<sup>2</sup>) and percentage (%) of Burdekin region, coral reef, and seagrass areas affected by different categories of exposure (or potential risk) based on satellite-derived Reef water types.

The exposure categories are not validated against ecological health data and represent relative potential risk categories for seagrass and coral reef ecosystems. Category I (no/very low risk) represents waters with ambient or detectable but low concentrations of the water quality parameters investigated and therefore low risk of any detrimental ecological effect. The areas and percentages of ecological communities affected by the different categories of exposure were calculated as a relative measure between regions and the long-term average.

In 2023–24, it was estimated that:

- Burdekin-wide: 91% of the region was not exposed to a potential risk, similar to long-term patterns (90%, Table 4-4). 9% (or about 4,200 km<sup>2</sup>) of the Burdekin region was exposed to combined potential risk categories II–IV. However, only 1% (613 km<sup>2</sup>) of the region was in the highest exposure category (IV) and 2% (924 km<sup>2</sup>) was in category III.
- Burdekin waterbodies: only the enclosed coastal and open coastal Burdekin waters were exposed to the highest categories of potential risk (III and IV). The open coastal area exposed was however spatially limited and corresponded to 15% (cat. III) and 1% (cat. IV) of the total Burdekin open coastal area (Figure 4-7c). 28% and 64% of the enclosed coastal areas were exposed to categories III and IV, respectively. The mid-shelf and offshore Burdekin waterbodies were largely exposed to no / very low risk (>99% of both waterbodies).
- Burdekin habitats:
  - Coral reefs: Approximately 1% of coral reefs in the Burdekin region were exposed to combined potential risk categories II–IV, with less than 1% in the highest exposure categories IV and III (combined 14 km<sup>2</sup>, Table 4-4). Only 1% (<40 km<sup>2</sup>) of the Burdekin corals occur in the inshore waters (Appendix C-6).
  - The open coastal coral area exposed to higher potential risk was limited and corresponded to 30% (cat. III) and 1% (cat. IV) of the total open coastal reefs area in the Burdekin region (
  - Figure 4-13a). A total of 40% and 54% of the enclosed coastal areas were exposed to categories III and IV respectively. Mid-shelf and offshore coral reefs were exposed to no risk (>99% in both waterbodies).
  - Seagrasses: 87% (or 616 km<sup>2</sup>) of seagrasses in the Burdekin region were exposed to combined potential risk categories II–IV. 20% (139 km<sup>2</sup>, Table 4-4) of seagrasses were in the highest exposure category (IV) and 32% (227 km<sup>2</sup>) were in category III, and they were all inshore seagrasses (Figure 4-13b). A total of 99% (~700 km<sup>2</sup>) of the Burdekin seagrasses occur in the inshore waters (Appendix C-6).
  - The open coastal seagrass area exposed to higher potential risk was limited and corresponded to 33% (cat. III) and 1% (cat. IV) of the total inshore seagrass area in the Burdekin region (
  - Figure 4-13b). A total of 30% and 67% of the enclosed coastal seagrass areas were exposed to categories III and IV respectively. Mid-shelf seagrasses were largely exposed to no / very low risk (84% of the Burdekin mid-shelf seagrasses), and to the lowest risk category II (16% of the Burdekin mid-shelf seagrasses).
  - **Comparison to long-term trends:** The coral and seagrass areas in the Burdekin region exposed to combined potential risk categories II–IV in 2023–24 were similar to the average long-term areas ( $\pm <1\%$  change). There was however an increase

in the seagrass area exposed to potential risk category III (+10%) from the potential risk category II (-8%). This was linked to an increased frequency of exposure to Reef WT1 south of Magnetic Island, where large seagrass meadows are located (Figure 4-12e).

Table 4-4: Areas (km<sup>2</sup>) and percentages (%) of the Burdekin region, Burdekin coral reefs, and Burdekin surveyed seagrass affected by different risk categories of exposure during the 2023–24 wet season and the long-term (2003–2022). The last three rows show the differences between % affected in 2023–24 and the long-term average (■: increase, ■: decrease, ■: no change, difference <5%).

Total surface area covered		Total		Potential Risk category				Total area exposed II–IV
				No / very low	Lowest		Highest	
					I	II	III	
Burdekin region	area	47,009	2024	42,833	2,639	924	613	4,176
			LT	42,281	3,363	747	617	4,728
	%	100%	2024	91%	6%	2%	1%	9%
			LT	90%	7%	2%	1%	10%
Burdekin coral reefs	area	2,966	2024	2,932	20	11	3	34
			LT	2,924	28	12	3	42
	%	100%	2024	99%	1%	0.4%	0.1%	1%
			LT	99%	1%	0.4%	0%	1%
Burdekin surveyed seagrass	area	708	2024	92	251	227	139	616
			LT	88	311	154	156	621
	%	100%	2024	13%	35%	32%	20%	87%
			LT	12%	44%	22%	22%	88%
Difference (2023–24 to long-term average)	Burdekin region			1%	-2%	<1%	<-1%	-1%
	Burdekin coral reefs			<1%	<-1%	<-1%	<1%	<-1%
	Burdekin surveyed seagrass			1%	-8%	10%	-2%	-1%




Figure 4-13: Percentage of the Burdekin region a) coral reef and b) surveyed seagrass habitats affected by different risk categories of exposure during the 2023–24 wet season. Water body classifications are shown along the x-axes: enclosed coastal (EC), open coastal (OC), mid-shelf (Mid), and offshore (Off).

#### 4.2.4 Mackay-Whitsunday region

As described for the Reef, a number of remote sensing products were generated to represent wet season water quality conditions in the Mackay-Whitsunday region. These maps are presented in Figure 4-14, which presents the frequency of the combined Reef WT1–2; the frequency of Reef WT1, WT2, and WT3 individually; the exposure maps in the long-term and 2023–24 wet season; and a difference map showing areas exposed to an increased risk in 2023–24.




Figure 4-14: Long-term and current year remote sensing results for the Mackay-Whitsunday region showing the a) frequency of Reef WT1–2; b) the frequency of Reef WT1, WT2, and WT3 individually regrouped into five likelihood categories [ $<0.2$  (Rare),  $0.2\text{--}0.4$ ,  $0.4\text{--}0.6$ ,  $0.6\text{--}0.8$  and  $0.8\text{--}1$  (very frequent)]; c) exposure to potential risk - each in the long-term (bottom) and 2023–24 wet season (top). d) Difference map showing areas with an increase (in red,  $\uparrow$ ) or decrease (in purple,  $\downarrow$ ) in risk category in 2023–24 against long-term trends [calculated as (c, top) exposure in 2024 minus (c, bottom) long-term]. Note that optical water types – especially the Reef WT3 – do not always correspond to direct catchment discharge and can also be due to oceanographic processes (see definitions in Table 2-2).

Table 4-5 presents the areas (km<sup>2</sup>) and percentage (%) of Mackay-Whitsunday region, coral reef, and seagrass areas affected by different categories of exposure (or potential risk) based on satellite-derived Reef water types.

The exposure categories are not validated against ecological health data and represent relative potential risk categories for seagrass and coral reef ecosystems. Category I (no/very low risk) represents waters with ambient or detectable but low concentrations of the water quality parameters investigated and therefore low risk of any detrimental ecological effect. The areas and percentages of ecological communities affected by the different categories of exposure were calculated as a relative measure between regions and the long-term average.

Table 4-5: Areas (km<sup>2</sup>) and percentages (%) of the Mackay-Whitsunday region, Mackay-Whitsunday coral reefs, and Mackay-Whitsunday surveyed seagrass affected by different risk categories of exposure during the 2023–24 wet season and the long-term (2003–2022). The last three rows show the differences between % affected in 2023–24 and the long-term average (■: increase, ■: decrease, ■: no change, difference ≤5%).

Mackay-Whitsunday		Total		Potential Risk category				Total area exposed II–IV
				No / very low	Lowest	Highest		
						I	II	
Mackay-Whitsunday region	area	48,957	2024	44,331	3,413	768	445	4,626
			LT	42,449	5,602	471	434	6,507
	%	100%	2024	91%	7%	2%	1%	9%
			LT	87%	11%	1%	1%	13%
Mackay-Whitsunday coral reefs	area	3,216	2024	3,095	76	35	11	122
			LT	3,019	166	25	7	197
	%	100%	2024	96%	2%	1%	0%	4%
			LT	94%	5%	1%	0%	6%
Mackay-Whitsunday surveyed seagrass	area	307	2024	54	141	55	57	253
			LT	10	186	36	76	297
	%	100%	2024	18%	46%	18%	19%	82%
			LT	3%	60%	12%	25%	97%
Difference (2023–24 to long-term average)	Mackay-Whitsunday region			4%	-4%	1%	0%	-4%
	Mackay-Whitsunday coral reefs			2%	-3%	0%	0%	-2%
	Mackay-Whitsunday surveyed seagrass			14%	-14%	6%	-6%	-14%




Figure 4-15: Percentage of the Mackay-Whitsunday region a) coral reef and b) surveyed seagrass habitats affected by different risk categories of exposure during the 2023–24 wet season. Water body classifications are shown along the x-axes: enclosed coastal (EC), open coastal (OC), mid-shelf (Mid), and offshore (Off).

In 2023–24, it was estimated that:

- Mackay-Whitsunday wide: 91% of the region was not exposed to a potential risk, above the long-term patterns (87%, Table 4-5). A total of 9% of the Mackay-Whitsunday region was exposed to combined potential risk categories II–IV (or about 4600 km<sup>2</sup>). However, only 1% (445 km<sup>2</sup>) of the region was in the highest exposure category (IV) and 2% (768 km<sup>2</sup>) in category III.
- Mackay-Whitsunday waterbodies: only the enclosed coastal and open coastal waters were exposed to the highest categories of potential risk (III and IV, (Figure 4-7c). The open coastal area exposed was however spatially limited and corresponded to 4% (cat. III) of the total Mackay-Whitsunday inshore area. A total of 37% and 51% of the enclosed coastal areas were exposed to categories III and IV, respectively. The mid-shelf and offshore Mackay-Whitsunday waterbodies were not exposed to potential risk.
- Mackay-Whitsunday habitats:
  - **Coral reefs:** Approximately 4% (or 122 km<sup>2</sup>) of coral reefs in the Mackay-Whitsunday region were exposed to combined potential risk categories II–IV (Table 4-5). However, less than 1% of coral were in the highest exposure category (IV) and 1% in category III (combined 46 km<sup>2</sup>), and they were all enclosed coastal or open coastal reefs (Figure 4-15a). A total of 9% (<300 km<sup>2</sup>) of the Mackay-Whitsunday corals occur in the inshore waters (Appendix C-6). The open coastal coral area exposed to higher potential risk was spatially limited and corresponded to 10% (cat. III) and 1% (cat. IV) of the total open coastal reef area in the Mackay-Whitsunday region. A total of 39% and 33% of the enclosed coastal areas were exposed to categories III and IV, respectively. Mid-shelf and offshore reefs were not exposed to a potential risk.
  - **Seagrasses:** All of the surveyed seagrass beds in the Mackay-Whitsunday region are located in the inshore area (Appendix C-6). Approximately 82% of seagrasses in the Mackay-Whitsunday region were exposed to combined potential risk categories II–IV (253 km<sup>2</sup>, Table 4-5). A total of 19% (57 km<sup>2</sup>) of seagrasses were in the highest exposure category (IV) and 18% (55 km<sup>2</sup>) were in category III. The open coastal seagrass area exposed to higher potential risk was spatially limited and corresponded to 4% (cat. III) of the total open coastal seagrass area in the Mackay-Whitsunday region Figure 4-15b. Approximately 41% and 50% of the enclosed coastal areas were exposed to categories III and IV, respectively.

- **Comparison with long-term trends:** The coral areas in the Mackay-Whitsunday region exposed to combined potential risk categories II–IV in 2023–24 were very similar to the long-term areas (-2% change). There was a decrease in the seagrass area exposed to the potential risk category IV (-6%) toward the potential risk category III (+6%), and from the lowest potential risk categories (II: -14%) toward the no/very Low risk category (I: +14%).

#### 4.2.5 Fitzroy and Burnett-Mary regions

In 2023–24, water quality monitoring in the Fitzroy region continued in accordance with the MMP monitoring design via a separately funded project, and the results of this are included as [Appendix D](#). There is still no formal water quality monitoring program in the Burnett-Mary region that is reported as part of the Paddock to Reef program. It should be noted that exposure maps have a higher degree of uncertainty in the Fitzroy and Burnett-Mary regions than in those described above due to limited validation from *in situ* monitoring.

Discharge in the Fitzroy region was very close to the long-term median and discharge in the Burnett Mary region was above the long-term median (1.7 times long-term median) (Figure 3-5).

As with all regions, the exposure categories are not validated against ecological health data and represent relative potential risk categories for seagrass and coral reef ecosystems. Category I (no/very low risk) represents waters with ambient or detectable but low concentrations of the water quality parameters investigated and therefore low risk of any detrimental ecological effect. The areas and percentages of ecological communities affected by the different categories of exposure were calculated as a relative measure between regions and the long-term average.

#### Fitzroy

Table 4-6 presents the areas (km<sup>2</sup>) and percentage (%) of Fitzroy region, coral reef, and seagrass areas affected by different categories of exposure (or potential risk) based on satellite-derived wet season water maps. In 2023–24, it was estimated that:

- Fitzroy-wide: 92% of the Fitzroy region was not exposed to a potential risk, similar to long-term patterns (91%, Table 4-6). 8% (or about 6,500 km<sup>2</sup>) of the Fitzroy region was exposed to combined potential risk categories II–IV. However, only 2% (1,455 km<sup>2</sup>) of the region was in the highest exposure category (IV) and 1% (1,272 km<sup>2</sup>) in category III.
- Fitzroy waterbodies: only the enclosed coastal and open coastal Fitzroy waters were exposed to the highest categories of potential risk (III and IV). The open coastal area exposed was however spatially limited and corresponded to 8% (cat. III) and 1% (cat. IV) of the total Fitzroy inshore area (Figure 4-7c). 26% and 69% of the enclosed coastal areas were exposed to categories III and IV, respectively. The offshore and mid-shelf Fitzroy waterbodies were not exposed to a potential risk.
- Fitzroy habitats:
  - **Coral reefs:** Approximately 3% of coral reefs in the Fitzroy region were exposed to combined potential risk categories II–IV (Table 4-6). 1% of coral were in the highest exposure category (IV) and 1% in category III (combined 68 km<sup>2</sup>), and they were all enclosed coastal or mid-shelf reefs (Figure 4-16a). Only 4% (<200 km<sup>2</sup>) of the Fitzroy corals occur in the inshore waters (Appendix C-6). The open coastal coral area exposed to higher potential risk was limited and corresponded to 25% (cat. III) and 1% (cat. IV) of the total open coastal coral reef area in the Fitzroy. Approximately 12% and 82% of the enclosed coastal areas were exposed to categories III and IV respectively. All of the mid-shelf and offshore reefs were classified as no / very low risk.

- **Seagrasses:** Approximately 67% (or about 320 km<sup>2</sup>) of seagrasses in the Fitzroy region were exposed to combined potential risk categories II–IV (Table 4-6), which was under the long-term trends). 23% (109 km<sup>2</sup>) of seagrasses were in the highest exposure category (IV) and 12% (56 km<sup>2</sup>) were in category III, and they were all inshore seagrasses (Figure 4-16b). Approximately 81% (<400 km<sup>2</sup>) of the Fitzroy seagrasses occur in the inshore waters (Appendix C-6).  
The open coastal seagrass area exposed to higher potential risk was limited and corresponded to 3% (cat. III) of the total open coastal seagrass area in the Fitzroy region (no open coastal seagrasses were exposed to the higher risk category IV). 33% and 73% of the enclosed coastal areas were exposed to categories III and IV respectively and 100% of the mid-shelf areas were exposed to the lowest risk category II.
- **Comparison with long-term trends:** The coral and seagrass areas exposed to highest potential risk categories II to IV were similar to the long-term patterns. There was a decrease in the seagrass area exposed to the risk cat. IIV and II (-8% and 19% toward the no/very low risk category (+23%).

Table 4-6: Areas (km<sup>2</sup>) and percentages (%) of the Fitzroy region, Fitzroy coral reefs, and Fitzroy surveyed seagrass affected by different risk categories of exposure during the 2023–24 wet season and the long-term (2003–2022). The last three rows show the differences between % affected in 2023–24 and the long-term average (red: increase, blue: decrease, green: no change, difference ≤5%). Areas south of the Marine Park (Hervey Bay) are not included.

Total surface area covered		Total		Potential Risk category				Total area exposed II–IV	
				No / Very low	Lowest		Highest		
					I	II	III		IV
Fitzroy region	area	86,869	2024	80,351	3,791	1,272	1,455	6,518	
			LT	78,805	5,265	1,189	1,610	8,064	
	%	100%	2024	92%	4%	1%	2%	8%	
			LT	91%	6%	1%	2%	9%	
Fitzroy coral reefs	area	4,881	2024	4,725	88	39	28	155	
			LT	4,719	113	14	35	161	
	%	100%	2024	97%	2%	1%	1%	3%	
			LT	97%	2%	0%	1%	3%	
Fitzroy surveyed seagrass	area	478	2024	159	155	56	109	319	
			LT	49	243	41	145	429	
	%	100%	2024	33%	32%	12%	23%	67%	
			LT	10%	51%	9%	30%	90%	
Difference (2023–24 to long-term average)	Fitzroy region			2%	-2%	<1%	<-1%	-2%	
	Fitzroy coral reefs			<1%	<-1%	1%	<-1%	<-1%	
	Fitzroy surveyed seagrass			23%	-19%	3%	-8%	-23%	

Fitzroy




Figure 4-16: Percentage of the Fitzroy region a) coral reef and b) surveyed seagrass habitats affected by different risk categories of exposure during the 2023–24 wet season. Water body classifications are shown along the x-axes: enclosed coastal (EC), open coastal (OC), mid-shelf (Mid), and offshore (Off).

## Burnett-Mary

Table 4-7 presents the areas (km<sup>2</sup>) and percentage (%) of Burnett-Mary region, coral reef, and seagrass areas affected by different categories of exposure (or potential risk) based on satellite-derived wet season water maps.

In 2023–24, it was estimated that:

- **Burnett-Mary wide:** Approximately 97% of the Burnett-Mary region was not exposed to a potential risk, which was similar long-term patterns (96%, Table 4-7). 3% of the Burnett-Mary region (or about 974 km<sup>2</sup>) was exposed to combined potential risk categories II–IV, with <1% in the highest exposure category (IV) and 1% in category III (combined 334 km<sup>2</sup>).
- **Burnett-Mary waterbodies:** only the enclosed coastal and open coastal Burnett-Mary waters were exposed to the highest categories of potential risk (III and IV). The open coastal area exposed corresponded to 8% (cat. III) and 2% (cat. IV) of the total Burnett-Mary inshore area (Figure 4-7c). A total of 43% and 42% of the enclosed coastal areas were exposed to categories III and IV respectively. >99% of the mid-shelf and offshore Burnett-Mary waterbodies were exposed to no / very low risk.
- **Burnett-Mary habitats:**
  - **Coral reefs:** Approximately 2% of coral reefs in the Burnett-Mary region were exposed to combined potential risk categories II–IV (Table 4-7). <1% of coral reefs were exposed to the highest risk categories IV (less than 1 km<sup>2</sup>) and 2% to cat. III (5 km<sup>2</sup>) these were all enclosed coastal or open coastal reefs (Figure 4-17a). Only 2% (<10 km<sup>2</sup>) of the Burnett-Mary corals occur in the inshore waters (Appendix C-6).  
The open coastal coral area exposed to potential risk category III and IV corresponded to 80% and 7% of the total enclosed coastal and open coastal coral reef area in the Burnett-Mary region. The enclosed coastal area exposed to potential risk category III and IV corresponded to 87% and 4% of the total enclosed coastal area in the Burnett-Mary region. All of the mid-shelf coral reefs were exposed to no / very low risk. There are no offshore reefs in the Burnett-Mary region.
  - **Seagrasses:** Approximately 57% (or 148 km<sup>2</sup>) of seagrasses in the Burnett-Mary region were exposed to combined potential risk categories II–IV (Table 4-7). 16% (41 km<sup>2</sup>) of seagrasses were in the highest exposure category (IV) and 14% (35 km<sup>2</sup>) were in category III and they were all enclosed coastal or open coastal seagrasses (Figure 4-17b). A total of 71% (<200 km<sup>2</sup>) of the Burnett-Mary corals occur in the inshore waters (Appendix C-6).  
The open coastal seagrass area exposed to higher potential risk corresponded to only 3% (cat. III) of the total inshore seagrass area in the Burnett-Mary region. A total of 36% and 45% of the enclosed coastal seagrass areas were exposed to categories III and IV respectively 100% of the mid-shelf seagrasses in the Burnett-Mary region were exposed to the lowest risk category II.
  - **Comparison to long-term trends:** The coral areas in the Burnett-Mary region exposed to combined potential risk categories II–IV in 2023–24 were similar to long-term areas. There was a decrease in the seagrass area exposed to the risk cat. II (-26% toward the no/very low risk category (+26%).

Table 4-7: Areas (km<sup>2</sup>) and percentages (%) of the Burnett-Mary region, Burnett-Mary coral reefs, and Burnett-Mary surveyed seagrass affected by different risk categories of exposure during the 2023–24 wet season and the long-term (2003–2022). The last three rows show the differences between % affected in 2023–24 and the long-term average (red: increase, blue: decrease, green: no change, difference ≤5%). Areas south of the Marine Park (Hervey Bay) are not included.

Total surface area covered		Total		Potential Risk category				Total area exposed II–IV
				No / Very low	Lowest		Highest	
					I	II	III	
Burnett-Mary region	area	37,713	2024	36,739	640	194	141	974
			LT	36,238	1,103	209	163	1,475
	%	100%	2024	97%	2%	1%	0%	3%
			LT	96%	3%	1%	0%	4%
Burnett-Mary coral reefs	area	285	2024	279	1	5	0	6
			LT	279	3	3	0	6
	%	100%	2024	98%	0%	2%	0%	2%
			LT	98%	1%	1%	0%	2%
Burnett-Mary surveyed seagrass	area	259	2024	111	72	35	41	148
			LT	43	140	30	46	217
	%	100%	2024	43%	28%	14%	16%	57%
			LT	17%	54%	12%	18%	83%
Difference (2023–24 to long-term average)	Burnett-Mary region			1%	-1%	<-1%	<-1%	-1%
	Burnett-Mary coral reefs			<-1%	-1%	1%	<1%	0%
	Burnett-Mary surveyed seagrass			26%	-26%	2%	-2%	-26%

Burnett-Mary




Figure 4-17: Percentage of the Burnett-Mary region a) coral reef and b) surveyed seagrass habitats affected by different risk categories of exposure during the 2023–24 wet season. Water body classifications are shown along the x-axes: enclosed coastal (EC), open coastal (OC), mid-shelf (Mid), and offshore (Off).

### 4.3 Satellite remote sensing summary and discussion

#### *Water type frequency maps (Sentinel-3 data)*

Sentinel-3 satellite images of the Reef and the Forel-Ule colour scale were used to produce Reef water types instead of the MODIS imagery and the wet season colour scale. FU equivalent water types were defined by grouping the FU colour classes 1–3 as “Reef WT4” (equivalent to marine waters in the WS scale), FU colour classes 4–5 as “Reef WT3” (equivalent to WS Tertiary water type), FU colour classes 6–9 as “Reef WT2” (equivalent to wet season secondary water type) and  $FU \geq 10$  as “Reef WT1” (equivalent to wet season primary water type), as defined in Petus *et al.* (2019) and Table 2-2.

Except for the ‘coral recovery period’, reference maps (long-term, wet, and dry frequency maps) were all updated in the 2022–23 technical report (Gruber *et al.*, 2024) using a combination of MODIS-Aqua (wet seasons 2003 to 2020) and Sentinel-3 (wet seasons 2021 to 2023) satellite water type maps. The previous update was in the 2018–19 reporting (Gruber *et al.*, 2020). The long-term reference maps now include 20 years of satellite imagery (2003 to 2022).

All mean long-term concentrations of water quality parameters were also reviewed for the 2022–23 technical report to improve the accuracy of the water type characterisation, building on the additional field data that is collected every wet season. TSS (80 samples) and Chl-*a* (110 samples) data collected as part of a pilot program in partnership with the Reef Trust Partnership Crown of Thorns Starfish Control Program (Waterhouse *et al.*, 2023) were also included in the analyses. The dataset, largely collected in the mid-shelf to offshore waterbodies, helped progress the characterisation of the Reef WT4 even though much more data are still needed (Table 2-2). The colour class category and water type corresponding to the location and week of acquisition of each water quality sample were extracted from the archive of MODIS-Aqua (wet seasons from 2004) and Sentinel-3 (wet seasons from 2021) weekly colour class maps.

This year’s results are in agreement with previous results and confirm that Sentinel-3 satellite data and the FU scale are useful for mapping Reef optical water types. Sentinel maps showed an inshore-to-offshore spatial pattern similar to the well-documented MODIS patterns (for example, Waterhouse *et al.*, 2021), with the highest frequency of the Reef WT1 (typically enriched in sediment and dissolved organic matter, brownish turbid waters) in the inshore waterbody, and more particularly in the enclosed coastal waters. Mid-shelf waterbodies were most frequently exposed to the Reef WT2 and WT3, and offshore waterbodies were most frequently exposed to the Reef WT3 (typically with low land-based contaminant concentrations and including the influence of oceanographic processes).

Only 4% of the Reef was exposed to Reef WT1 waters during the 2023–24 wet season, and only the inshore waters were exposed. 20% was exposed to Reef WT2, which is similar to (slightly above) the pattern for the long-term and representative coral recovery periods. As observed in previous years, the Reef area exposed to Reef WT3 in 2023–24 was larger than all reference periods (61% of the Reef), including the ‘wet’ years (58% of the Reef). This result is related to anomalously large Reef WT3 areas measured in the mid-shelf and offshore, which are almost certainly due to oceanographic processes such as upwelling rather than direct catchment discharge influence. This should be further investigated in a future case study by comparing the Reef WT3 maps with sea surface temperature climatology (for example, Wijffels *et al.*, 2018) or by using the eReefs model to investigate whether the Reef WT3 waters are due to processes not influenced by catchment discharges. Oceanographic processes that influence water colour might in turn be influenced by climate change, which would require further investigation.

*Exposure maps (Sentinel-3 and field water quality data)*

Reef WT3 waters are associated with low land-based contaminant concentrations with only the long-term mean PN concentration above the wet season GV in the Reef WT3, and a low magnitude score in the Reef exposure assessment (Figure 4-5). While Reef WT3 areas in 2023–24 were larger than expected, this did not result in increasing the Reef-wide potential risk. The total Reef area exposed to a potential risk in 2023–24 was spatially limited and similar to the long-term patterns. Ninety percent of the Reef was exposed to no/very low potential risk and only 3% (but about 10,000 km<sup>2</sup>) of the Reef was in the highest exposure categories III and IV.

The offshore and mid-shelf and waterbodies were largely classified as no/very low potential risk (100% and 95%, respectively), and this pattern was observed in all Reef regions. Open coastal waters were largely exposed to the lowest category of risk (II, 45% of the open coastal waterbody) or to no/very low risk (44% of the open coastal waterbody), and only 11% and 1% of the total Reef open coastal waters were exposed to the highest potential risk categories III and IV. The Reef enclosed coastal waters had the highest relative potential risk, with 26% and 50% of the enclosed coastal waters exposed to categories III and IV, respectively. This, however, represent a very small proportion of the total size of the Reef (less than 2% of the Reef). The Wet Tropics, Cape York and Burdekin region open coastal waterbodies had the greatest exposure to risk categories III (25 % and 19% and 15% of the Wet Tropics, Cape York and Burdekin open coastal waterbodies), with all other regions between 4% and 8% (Figure 4-7).

As a result, mid-shelf and offshore Reefs habitats (surveyed seagrass and coral reefs) were either exposed to the lowest risk category II or to no potential risk. Open coastal seagrasses and coral reefs were largely exposed to the lowest category of risk (II, 63% and 49% of the total Reef seagrass and coral areas, respectively). Enclosed coastal habitats were the most at risk, with 89% (less than 1% of the total coral reef area of the Reef) and 85% (~24% of the total seagrass area in the Reef) of the total enclosed coastal seagrasses and corals in the Reef classified as combined category III–IV. Enclosed coastal areas are shallow regions of the Reef and it is likely that wind-driven resuspension (some of which was originally derived from river discharge in previous events) may influence the TSS concentrations and resulting exposure results in this very inshore region.

Regional areas exposed to a potential risk (combined risk categories II–IV) were largely similar to the long-term patterns in the southern regions, but a 4% increase was observed in the total areas exposed in both the Cape York and Wet Tropics regions (Figure 4-18).

In the Cape York region, the total coral and seagrass area exposed to a potential risk (II–IV) were over the long-term trends (+7% and +10% respectively), which was consistent with the high discharge measured in 2023–24. There was also an increase in coral areas exposed to the lowest risk category (II+7%), and an increase in seagrass areas exposed to the risk category III (+8%).

While the total areas in the Wet Tropics region exposed to a potential risk (combined risk categories II–IV) were greater than the average long-term areas (+4% changes), this did not impact the ecosystem exposure results. The total coral and seagrass areas exposed to the risk categories II–IV were similar to the long-term patterns (changes ≤1%). There was furthermore a decrease in seagrass area exposed to the highest potential risk categories (IV: -12%) toward the lowest risk category II (+9%) which is likely related to localised WT1 movements north of Hinchinbrook Island and may include the influence of different patterns/timing of sediment resuspension or some bottom influence as this is a shallow area (Figure 4-10d).

There was also an increase in the seagrass areas exposed to potential risk category III in the Burdekin region (III: +10%), which was related to a higher frequency of primary waters

measured south of Magnetic Island (Figure 4-12e). However, the total seagrass area exposed to a potential risk (II-IV) was similar to the long-term trends.

In the Mackay-Whitsundays, Fitzroy and Burnett-Mary regions, the total seagrass area exposed to a potential risk (II-IV) were under the long-term trends (-14%, -23%, and -26%, respectively). There was a decrease in the seagrass area exposed to the lowest potential risk (cat. II) toward the no/very low risk category in the Mackay-Whitsunday, Fitzroy, and Burnett-Mary regions and smaller areas of seagrass meadows were exposed to potential risk category IV in both the Mackay-Whitsunday (-6%) and Fitzroy regions (-8%)". This is consistent with the lower discharge measured in 2023–24 in these regions.

It should be noted there are several caveats to the exposure maps:

- Reef-wide water quality GVs are applied rather than site or waterbody-specific GVs.
- This assessment does not take into account the current condition of Reef ecosystems and long-term impacts on these communities. For example, it is recognised that inshore communities may have adapted to the most turbid Reef water types and exposure history; therefore, the highest risk of an ecological response could be during large events when Reef WT1 and WT2 extend into otherwise low exposure (more offshore) areas.
- Reporting the areas of coral reefs and seagrass in the highest potential exposure categories cannot be assessed in terms of ecological relevance at this stage and is included as a comparative measure between regions and between years.
- One-week exposures are reported. The ecological consequence of exposure of this duration is not presently known.
- The degree of validation against *in situ* data varies between regions, with limited water quality data in the Fitzroy and Burnett-Mary regions.
- It is impossible to fully separate the direct influence of riverine plume from wind- and wave-driven sediment resuspension in optical satellite images, and this may particularly influence exposure results in the shallow enclosed coastal Reef waters. Similarly, it is impossible at this stage to separate catchment versus oceanographic processes in offshore Reef WT3 waters.

Satellite methods and tools developed through the MMP to map Reef water types have now proved to be efficient for the mapping of water quality trends. However, there is a need to keep integrating spatial and temporal information obtained from the water type maps and *in situ* water quality measurements with environmental data to better understand physical influences that can lead to light reduction and water colour changes in both wet and dry seasons, and from the inshore to offshore Reef areas. Multivariate statistical analyses would be useful to gain further understanding of these processes. Furthermore, it would be interesting to collect extra samples in the transition zone between Reef WT1 and WT2 in the future to better understand drivers of water colour variability there and further characterise concentrations and productivity in this region of flood plumes. Furthermore, there might also be a need to discard water quality samples collected in the enclosed coastal waters in the characterisation of the water type composition (Section 2.6.2) and the calculations of the exposure scores (Figure 4-5), as GVs for enclosed coastal waters are different from other areas of the Reef. This however would discard important water quality information collected in the flood plumes where the highest turbidity is typically measured. Separating the enclosed coastal waters information in the exposure assessment process and applying higher water quality guideline values to these samples should however be investigated in a future case study to improve the accuracy of the risk classification.

Finally, there is a major gap in the availability of *in situ* water quality data in the Burnett-Mary region and in mid-shelf and offshore waterbodies across the Reef. These data are essential for improving confidence in the remote sensing products across all regions and waterbodies. The pilot study to investigate options for water quality sampling as part of the Reef Trust Partnership Crown of Thorns Starfish Control Program (Waterhouse *et al.*, 2023) is a good example of the opportunities that exist to expand these water quality datasets. The pilot study also highlighted the potential of using a Smartphone app, the Eye on Water (<https://www.eyeonwater.org/>) to collect vessel-based Forel-Ule colour information. Using the Eye on Water app concomitantly to the water quality data allowed retrieval of water colour information at the exact site location, even when satellite images are obscured by clouds. It thus increases the number of data available to match up concentrations of water quality parameters and the four Reef water types and can help improve the characterisation of water quality concentrations across Reef water types. Using the Smartphone water colour data in combination to the satellite FU data could, in the future, improve the mapping of water quality patterns in the Reef and should be investigated for greater integration in the MMP.




Figure 4-18: Areas (km<sup>2</sup>) and percentages (%) of the Reef and Reef regions, coral reefs, and surveyed seagrass affected by different risk categories of exposure during the 2023–24 wet season and the long-term (2003–2022). The left figures show the differences between % affected in 2023–24 and the long-term average (red: increase, blue: decrease, green: no change, difference ≤5%). Note the different x-axis scale for the Fitzroy and Burnett-Mary difference plots.

## 5 FOCUS REGION WATER QUALITY AND WATER QUALITY INDEX

The following sections provide detailed analysis of key water quality variables in nine focus regions in the context of local environmental drivers, specifically focused on the annual water quality condition and long-term trends. Monitoring results from the duration of the MMP (since 2005) are used to provide context for interpreting recent monitoring. For each of the focus regions, the following information is included and discussed (except Cape York where data are presented differently, as some aspects of monitoring in this region differ from other regions):

- a map of monitoring locations,
- time-series of the combined discharge from local rivers that influence the focus region,
- regional trends in key water quality parameters since 2005,
- presentation of the long-term trend and annual condition of ambient water quality relative to guideline values (GVs) using the WQ Index, and
- results from flood event monitoring (if conducted during the year).

Site-specific data and additional information are presented in Appendix C and include:

- Appendix C-1 Figure C-1: Time-series of chlorophyll and turbidity measured by moored FLNTUSB instruments;
- Appendix C-2 Figure C-2: Time-series of temperature and salinity measured by moored Sea-Bird Electronics instruments;
- Appendix C-3 Table C-1: Cape York, Wet Tropics, Burdekin and Mackay-Whitsunday: Summary statistics for each water quality variable from each monitoring location for the current water year;
- Appendix C-3 Table C-2: Annual summaries of moored FLNTUSB turbidity measurements for each monitoring location, including percentage exceedances of GV; and
- Table C-3 to C-8: Summary of water quality data (collected as part of the JCU event-based sampling) across the Reef colour classes and water types.

### 5.1 Cape York region

The Cape York region is divided into four focus regions: the Pascoe, Stewart, Normanby, and Annan-Endeavour. The monitoring results are presented separately for each.

Water quality monitoring commenced in the Cape York region as part of the MMP in January 2017. Twenty-one sites in four focus regions (Figure 5-1) are sampled four to five times per year during ambient conditions. Additional event samples are collected depending on the location and accessibility of flood plumes at these and additional sites ([Appendix A](#)). Ambient sampling primarily occurs between 1 November to 30 April (wet season) due to strong trade winds ( $>25 \text{ km h}^{-1}$ ) preventing access during the winter months.




Figure 5-1: Water quality sampling sites in the Cape York region shown with water body boundaries. River datasets for map courtesy Grill *et al.* (2019).

The 2023–24 water year is the eighth year of sampling for the Cape York region. In consultation between CYWP, AIMS, and the Reef Authority, the laboratory analysis methods and the number of sites sampled in Cape York changed in 2020 (Moran *et al.*, 2023). Because of this change, long-term trends cannot yet be assessed and the annual condition index is only calculated from this time onwards. Water quality results within each focus region have been assessed relative to distance from river mouths and compared against the Eastern Cape York Water Quality Guidelines for the enclosed coastal, open coastal, mid-shelf and offshore water bodies (State of Queensland, 2020). For comparison with the GVs, water quality results have been categorised as ambient wet season, ambient dry season, or event based on an evaluation of the river hydrograph at the time of sampling, antecedent rainfall, salinity measurements, and field observations. The annual condition Water Quality Index has also been calculated for each focus region. This Index is based on the current year only and is not a comparison against previous data (see Section 2.5).

The 2023–24 water year was an above average discharge year for Cape York, particularly for the Normanby Basin and Annan catchment, where Tropical Cyclone Jasper in mid-December 2023 brought extreme rainfall and flood levels exceeded historic records by over 3 m in the upper Normanby and Annan rivers. The marine flood plumes following TC Jasper extended from the northern Wet Tropics to north of the Stewart transect (Figure 5-2), and beyond the outer reefs- inundating a large portion of the northern Great Barrier Reef for over one week, and potentially impacting water quality through the following wet season.




Figure 5-2: Combined Sentinel satellite images showing the extent of flood plumes and maximum turbidity (FU units) observed over the week of 18–24 December 2023, following TC Jasper.

### 5.1.1 Pascoe

The Pascoe focus region is influenced primarily by discharge from the Pascoe and Olive Rivers. Six sampling sites (Figure 5-3) are located along two transects to the northeast and southeast from the Pascoe River mouth out to Eel Reef and past Middle Reef (locally known as Blue Bells). Floodwaters have been observed flowing in both directions depending on wind and other local conditions. Enclosed coastal waters near the mouth of the Pascoe and site PRS01 are highly turbid due to tidal and wind-driven resuspension of shallow sediments.




Figure 5-3: Water quality sampling sites in the Pascoe River transect with water body boundaries.

The Pascoe River transect was sampled three times under ambient wet season conditions and twice under ambient dry season conditions from October 2023–May 2024 (Figure 5-4). NR05 along the northern transect near Eel Reef was not sampled due to logistical challenges related to tides and weather.

Total discharge for the year was approximately twice the annual median discharge (Figure 5-5), with the first flood event peaking on 23 January 2024 (Figure 5-4). Significant rainfall continued until the end of April.




Figure 5-4: Daily discharge and sampling dates for the Pascoe River (gauge 102102A) for the 2023–24 water year. Red dots represent sampling dates.




Figure 5-5: Long-term discharge for the Olive and Pascoe Rivers. Daily (blue) from Pascoe R. at Garraway Creek (gauge 102102A) and water year (1 October to 30 September, red symbols) for the combined Olive Pascoe Rivers discharge volumes are shown. Red dashed line represents long-term median of the combined annual discharge.

The total discharge and modelled loads estimated for the 2023–24 water year from the Pascoe catchment (upscaled from the Garraway gauge) are shown in Figure 5-6. The discharge and loads calculated for the 2023–24 water year from the Pascoe catchment (not including the Olive catchment) were 1.9-fold above the long-term median. Over the 18-year period from 2006–07:

- discharge ranged from 425 GL (2015–16) to 3,770 GL (2018–19)
- TSS loads ranged from 19 kt (2015–16) to 194 kt (2018–19)
- DIN loads ranged from 34 t (2015–16) to 275 t (2018–19)
- PN loads ranged from 68 t (2015–16) to 1,068 t (2018–19).




Figure 5-6: Modelled loads of (A) TSS, DIN, and PN and (B) discharge for the Pascoe catchment (note Pascoe catchment only, does not include the Olive catchment) from 2006 to 2024. The loads reported here are a combination of 'best estimates' based on 'up-scaled' discharge data from gauging stations and monitoring data for the 2014–15, 2016–17, 2017–18, 2018–19, 2019–20, 2020–21 and 2022–23 water years and an average of the annual mean concentrations for these seven water years applied to the remaining dataset. Dotted line represents the long-term median for basin discharge. Note the different scales on the two y-axes.

### Ambient water quality

According to modelled estimates, total TSS and DIN loads nearly doubled over the 2023–24 water year compared to the previous year, despite similar total discharge (Figure 5-6). As a result, the overall WQ Index for the Pascoe region declined from "good" in 2022–23 to "moderate", driven by declines in the productivity and particulate sub-indicators (Figure 5-8).

All sample results are plotted against distance from the river mouth in Figure 5-7.




Figure 5-7: Water quality parameters (surface and subsurface samples) and Secchi depth over distance (km) from the river mouth for the Pascoe River focus region, all 2023–24 ambient monitoring samples. Water body classifications are shown along the x-axes: enclosed coastal (EC), open coastal (OC) and mid-shelf (MS). Note the y-axes are logarithmic scales. Fitted lines are generalised additive models.

Comparison of the 2023–24 ambient results with previous years and the GVs (Table C-1) highlights that:

- The annual condition WQ Index declined from good to moderate, due to declines in the productivity and particulate sub-indicators;
- Mean and median TSS concentrations were below the wet season and annual GVs at most sites;
- Mean Secchi depths exceeded the GV at 3 out of 5 sites;
- Despite an increase in river discharge, the water clarity sub-indicator improved compared to 2022–23 wet season but remained below previous years (Figure 5-8);
- Chl-a median concentrations exceeded the GVs at open coastal sites, consistent with previous years, but did not exceed the guidelines at SR05 in the mid-shelf or enclosed coastal water bodies;

- Median  $\text{NO}_x$  concentrations were more than double the GV at all open coastal and mid-shelf water bodies, with concentrations 4 to 5 times the GV at some sites. Mean and median concentrations also were almost double those from the previous year, contributing to the decline in the productivity sub-indicator; and
- PN exceeded the GV at PRN02, PRS2.5 and PRBB in the open coastal and mid-shelf water bodies. PP exceeded the GV at PRN02 and PRS2.5.




Figure 5-8: The annual condition WQ Index for the Pascoe focus region for 2023–24. The WQ Index uses two formulations but only the annual condition index (based on post-2015 sampling design) is currently available for the Cape York region. WQ Index colour coding: ◆ – 'very good'; ◆ – 'good'; ◆ – 'moderate'; ◆ – 'poor'; ◆ – 'very poor'. Indicators or sub-indicators that are used to calculate the WQ Index are shown as coloured lines on each plot. Error bars (vertical black lines) on the WQ Index represent the 95% quantile intervals. Calculations for index formulations are described in [Appendix B](#).

### Event water quality

The first flood event of the year for the Pascoe focus region reached peak discharge at the Garraway gauge on 23 January 2024 (Figure 5-4), and event samples were collected on 25 January. TSS remained relatively low across the plume (maximum  $12 \text{ mg L}^{-1}$ ). However, DIN concentrations were highly elevated along the southern transect in the enclosed coastal and open coastal waterbody ( $4 \mu\text{g L}^{-1}$ ). PN, DOC and POC concentrations were also elevated along the southern transect. Chl-*a* concentrations remained relatively low, potentially due to low light availability at the time of sampling, as indicated by low Secchi depths ( $<4 \text{ m}$ ) in the enclosed and open coastal waterbody. High TDN concentrations were measured just outside the visible plume line, near PRBB (Figure 5-9).




Figure 5-9: Sentinel satellite images showing average magnitude Pascoe River flood event and sampling locations on 25 January 2024.

### 5.1.2 Stewart

The Stewart focus region is influenced primarily by discharge from the Stewart River. During flood conditions it can also be influenced by floodwater from the Normanby and Kennedy Rivers and potentially by runoff from coastal creeks and mudflats.

Four sampling sites for the Stewart River are located in a transect from the river mouth to mid-shelf reefs, representing a gradient in water quality (Figure 5-10). The transect was sampled five times (four times during ambient wet conditions and once during ambient dry conditions) between November 2023 and May 2024 (Figure 5-11; Table A-1). There were no major flood events in the Stewart River over the 2023–24 wet season, however there was significant influence from the Normanby Basin. Satellite images from late December (TC Jasper) through to early April show that floodwater from the Normanby and Kennedy Rivers frequently inundated the Stewart transect (Figure 5-24).




Figure 5-10: Water quality sampling sites in the Stewart River transect with water body boundaries.




Figure 5-11: Daily discharge and sampling dates for the Stewart River (gauge 104001A) for the 2023–24 wet season.




Figure 5-12: Long-term discharge for the Stewart River (gauge 104001A – Telegraph Road). Daily (blue) and water year (1 October to 30 September, red circles) discharge volumes shown. Red dashed line represents long-term median annual discharge.

The combined discharge and modelled loads estimated for the 2023–24 water year from the Stewart Basin are shown in Figure 5-13. The discharge and loads calculated for the 2023–24 water year from the Stewart Basin were above (1.5 times) the long-term median. Over the 18-year period from 2006–07:

- discharge ranged from 289 GL (2014–15) to 3,002 GL (2018–19)
- TSS loads ranged from 8.7 kt (2014–15) to 90 kt (2018–19)
- DIN loads ranged from 13 t (2014–15) to 135 t (2018–19)
- PN loads ranged from 40 t (2014–15) to 420 t (2018–19).




Figure 5-13: Loads of (A) TSS, DIN and PN, and (B) discharge for the Stewart Basin from 2006 to 2024. The loads reported here are based on the best estimates of annual mean concentration informed by nearest neighbour monitoring and by the Source Catchments modelling data and applied to each water year. Dotted line represents the long-term median for basin discharge. Note the different scales on the two y-axes.

### Ambient water quality

The Stewart River ambient condition sampling results are plotted against the distance from the river mouth in Figure 5-14 and are compared against the GVs for each water body (Table C-1). TSS concentrations generally decreased with distance from the river mouth, while Secchi depth increased. Wet season Chl-*a* concentrations increased with distance, while dry season concentrations decreased with distance. Nutrients relationships with distance varied and did not always show clear trends.




Figure 5-14: Water quality parameters (all surface and subsurface samples for the 2023–24 season) and Secchi depth over distance (km) from river mouth for the Stewart River focus region during ambient wet season and dry season conditions. Water body classifications are shown along the x-axes: enclosed coastal (EC), open coastal (OC), and mid-shelf (MS). Note the y-axes are logarithmic scales. Fitted lines are generalised additive models.

Comparison of the 2023–24 ambient results with previous years and the water quality GVs ([Appendix C-3](#), [Appendix C-5](#)) highlights that:

- The annual condition WQ Index for the Stewart region scored ‘moderate’ overall, similar to 2022–23; however, there was a significant decline in the productivity sub-indicator (Figure 5-15);
- Median TSS concentrations met the annual and wet season GVs at all sites;
- Secchi depths did not meet the GVs at all sites except for SR02 in the enclosed coastal water body;
- Mean Chl-a concentrations did not meet the wet season GVs at SR03 or SR04 in the open coastal and mid-shelf water bodies. Concentrations increased compared to the

2022–23 wet season, likely due to the increased influence from Normanby Basin flood water;

- Median  $\text{NO}_x$  concentrations were more than double the annual and wet season GVs, contributing to the decline in the productivity sub-indicator;
- Median  $\text{PO}_4$  concentrations exceeded the annual and wet season GVs in the open coastal and mid-shelf water bodies, and significantly increased compared to 2022–23 concentrations. Similar to previous years, concentrations were consistently higher at depth than within surface samples;
- Median PP concentrations were below the annual or wet season GVs at all sites except for SR03;
- Median PN results were mixed, exceeding annual or wet season GVs at SR03 and SR04, but meeting GVs at SR02 and SR05.




Figure 5-15: Annual condition version of the WQ Index for the Stewart focus region. The WQ Index uses two formulations but only the annual condition Index (based on post-2015 sampling design) is currently available for the Cape York region. WQ Index colour coding: ◆ – 'very good'; ◆ – 'good'; ◆ – 'moderate'; ◆ – 'poor'; ◆ – 'very poor'. Indicators or sub-indicators that are used to calculate the WQ Index are shown as coloured lines on each plot. Error bars (vertical black lines) on the WQ Index represent the 95% quantile intervals. Calculations for index formulations are described in [Appendix B](#).

### Event water quality

There were no major flood events in the Stewart River over the 2023–24 wet season and no targeted flood monitoring.

### 5.1.3 Normanby

The Normanby focus region is influenced by discharge from the Normanby, Laura, Kennedy, Hann, Mossman, Morehead, and Annie Rivers via three distributaries: the North Kennedy, Normanby, and Bizant (the Normanby Basin). Five sampling sites are located along a transect from the Normanby River mouth to Corbett Reef in the offshore water body (Figure 5-16). Site CI01 is located near Cliff Isles ('Marrpa' in Lama Lama language). Four additional event-only sites are: NR01 at the Normanby River mouth, two sample sites located near the Kennedy River and one near the Bizant River mouth in the enclosed coastal water body.




Figure 5-16: Water quality sampling sites in the Normanby Basin focus region with water body boundaries.

The Normanby transect was sampled six times (spread over seven days) from December 2023 to June 2024 (Figure 5-17). Only one of these sampling trips was scheduled as an event sampling trip (26 March 2024); however, due to consistent above-average discharge conditions through the wet season, typical event conditions influenced parts of the Princess

Charlotte Bay during most regular sampling trips (particularly along the Kennedy River side and site CI01). Additional samples were collected from event sample locations near the Bizant and Kennedy River mouths on 31 December 2023, approximately 10 days after peak flooding from TC Jasper. Floodwaters from the Normanby Basin also influenced samples within the Stewart transect (Figure 5-24). Long-term discharge is shown in Figure 5-18.




Figure 5-17: Daily discharge and sampling dates for the Normanby River (gauge 105107A) for the 2023–24 wet season. Note there is a 2 to 3-day travel time between the gauge and coastal waters, and thus event samples were collected earlier in the flood stage than shown on the hydrograph.




Figure 5-18: Long-term discharge for the Normanby River (gauge 105107A – Kalpower Crossing). Daily (blue line) and water year (1 October to 30 September, red symbols) discharge volumes shown. Method for estimation is described in Table 2-3.

The discharge for the 2023–24 water year from the Normanby Basin was well above the long-term median at 4.2-fold and was the highest on record (Figure 5-19). Our modelled load estimates suggest that TSS and DIN exports in the 2023–24 water year were the highest on record, although monitoring data (once available) will need to validate this finding. In any case, the discharge and loads would be amongst the highest over the history of the MMP and represent back-to-back large flow years from the basin. Over the 18-year period from 2006–07:

- discharge ranged from 2,314 GL (2011–12) to 16,300 GL (2023–24)
- TSS loads ranged from 55 kt (2014–15) to 645 kt (2023–24)
- DIN loads ranged from 42 t (2011–12) to 520 t (2023–24)
- PN loads ranged from 124 t (2009–10) to 2,470 t (2018–19).




Figure 5-19: Modelled loads of (A) total suspended solids, dissolved inorganic (DIN) and particulate nitrogen (PN) and (B) discharge for the Normanby Basin. The loads reported here are a combination of 'best estimates' based on 'up-scaled' discharge and monitoring data from the Normanby River at Kalpowar gauging station (covers ~50% of the basin area). The dotted line represents the long-term median for basin discharge. Note the different scales on the two y-axes.

### Ambient water quality

Due to the extensive flooding over the wet season, there was some freshwater influence over the January, February, and March ambient sampling trips (Figure 5-17). Ambient water quality results are plotted against distance from the Normanby river mouth (or Kennedy for CI01) in Figure 5-20. Ambient results are compared against the GVs for each water body in Table C-1.

As shown in Figure 5-20, TSS, PN, and PP concentrations generally declined with increasing distance from the river mouth, while Secchi depth increased (water clarity improved). Wet season Chl-a and NO<sub>x</sub> concentrations did not show clear trends with distance but during the dry season, Chl-a decreased with distance from the river mouth.




Figure 5-20: Water quality parameters (surface and subsurface) and Secchi depth over distance (km) from river mouth for the Normanby focus region, all ambient 2023–24 sampling dates. Water body classifications are shown along the x-axes: enclosed coastal (EC), open coastal (OC), and mid-shelf (MS). Note the y-axes are logarithmic scales. Fitted lines are generalised additive models.

Comparison of the 2023–24 ambient results with previous years and the GV (Table C-9) highlights that:

- Overall, the annual condition WQ Index score for the Normanby remained 'moderate' but declined to close to a 'poor' score due to declines in the scores for the productivity and particulates sub-indicators (Figure 5-21);
- The water clarity sub-indicator score increased from 'poor' to 'moderate'. Water clarity at Princess Charlotte Bay has consistently been 'poor' or 'very poor' over the MMP

monitoring period as the Bay is relatively shallow, with frequent resuspension of muddy sediments in addition to frequent flooding;

- TSS concentrations met the GV at all sites except for NR06 in the offshore water body. Previous years samples have generally had higher TSS in sub-surface samples than at the surface, likely due to benthic sediment resuspension from strong currents at depth. Over the 2023–24 wet season, higher concentrations were measured in surface waters at some locations, due to flood water influence;
- Secchi depth did not meet GV at any sites except for NR02 in the enclosed coastal zone;
- Median  $\text{NO}_x$  and  $\text{PO}_4$  concentrations exceeded the annual and wet season GV at all open coastal, mid-shelf and offshore sites;
- Median Chl-a concentrations exceeded the wet season GV at open coastal, mid-shelf and offshore sites, and increased compared to 2022–23 at most sites; and
- Median PN and PP concentrations exceeded the GV at most sites and concentrations increased compared to 2022–23.




Figure 5-21: Annual condition version of the WQ Index for the Normanby focus region. The WQ Index uses two formulations but only the annual condition index (based on post-2015 sampling design) is currently available for the Cape York region. WQ Index colour coding: ◆ – 'very good'; ◆ – 'good'; ◆ – 'moderate'; ◆ – 'poor'; ◆ – 'very poor'. Indicators or sub-indicators that are used to calculate the WQ Index are shown as coloured lines on each plot. Error bars (vertical black lines) on the WQ Index represent the 95% quantile intervals. Calculations for index formulations are described in [Appendix B](#).

### Event water quality

Normanby Basin rivers were in flood for most of the 2023–24 wet season (Figure 5-17), influencing water quality in parts of Princess Charlotte Bay (PCB) during ambient sampling trips. Additional event samples were collected from flood plumes near the Bizant and Kennedy river mouths during regular sampling on 31 December 2023, 10 days after the Normanby River reached historic peak river height and discharge levels ( $2333 \text{ m}^3 \text{ s}^{-1}$ ) as a result of TC Jasper. Flood plumes were evident at PCB for over 3 weeks and flowed far to the north, inundating an area over  $3500 \text{ km}^2$  and many coral reefs (Figure 5-23). Samples collected on 31 December had the highest event TSS concentrations on record for Normanby-Kennedy transects sites (since sampling began in 2018), with  $165 \text{ mg L}^{-1}$  at KR01 (2 km from Kennedy River mouth)

and  $20 \text{ mg L}^{-1}$  at KR02 (6 km from the river mouth) (Figure 5-22). The highest event nutrient concentrations on record for PCB were also recorded that day, including DIN ( $93.0 \text{ } \mu\text{g L}^{-1}$ ), PN ( $512.4 \text{ } \mu\text{g L}^{-1}$ ), POC ( $3948 \text{ } \mu\text{g L}^{-1}$ ) and PP ( $54.2 \text{ } \mu\text{g L}^{-1}$ ), as well as Chl-a ( $5.0 \text{ } \mu\text{g L}^{-1}$ ) (Figure 5-22), from an extra sampling location 6 km from the Normanby river mouth, between NR02 and KR01.

Targeted flood monitoring also occurred on the 26 March 2024, close to peak discharge from an above average magnitude Normanby Basin flood event (peak discharge  $1414 \text{ m}^3 \text{ s}^{-1}$ ; Figure 5-17). As with previous 2023–24 flood events, flood plumes from this event travelled to the north well beyond Cliff Isles and the Stewart River transect and inundating Corbett Reef and numerous reefs further north (Figure 5-24). TSS peaked at  $52 \text{ mg L}^{-1}$  at KR01 and declined with increasing salinity while Secchi depths increased along the salinity gradient (Figure 5-22). Maximum DIN concentrations ( $>30 \text{ } \mu\text{g L}^{-1}$ ) were recorded near the Kennedy and Normanby River mouths (Chl-a peaked at  $0.8 \text{ } \mu\text{g L}^{-1}$ ) and generally declined with increasing salinity despite increasing Secchi depth measurements.




Figure 5-22: Water quality data under the influence of flood plumes from the Normanby and Kennedy rivers on 31 December 2023 and 26 March 2024 including TSS, Secchi depth, chlorophyll-a (Chl-a) and dissolved inorganic nitrogen (DIN).




Figure 5-23: Satellite true colour image of Kennedy and Normanby River flood plume and sample locations on 31 December 2023 (left). The corresponding processed imagery on the right panel shows the Forel Ule classes.




Figure 5-24: Satellite true colour image of Kennedy and Normanby River flood plume and sample locations on 26 March 2024 (left). The corresponding processed imagery on the right panel shows the Forel Ule classes.

#### 5.1.4 Annan-Endeavour

The Annan-Endeavour focus region is influenced primarily by discharge from the Endeavour and Annan Rivers. Five sampling sites are located along transects from the two river mouths to mid-shelf reefs, representing a gradient in water quality (Figure 5-25). Additional sites ER01 and AR01 are sampled during events. In addition to manual sampling, dataloggers monitor continuous chlorophyll fluorescence, turbidity, and conductivity at Dawson Reef 6 km from the mouth of the Annan River and Forrester Reef 30 km north of the Endeavour River mouth (Figure 5-25).

The Annan and Endeavour transect was sampled for ambient wet season conditions five times between November 2023–March 2024. Event samples were collected following TC Jasper on 20 December 2023 and again on 24 March 2024 (Figure 5-27). Five samples of TSS, salinity and Chl-*a* were collected adjacent to Dawson Reef and Forrester Reef dataloggers to estimate TSS and Chl-*a* concentrations from logger measurements of turbidity and chlorophyll fluorescence, respectively.




Figure 5-25: Water quality sampling sites in the Annan-Endeavour region shown with water body boundaries.

The estimated total discharge from the Endeavour Basin for the 2023–24 water year was 3.1 times the long-term median, the highest on record (Table 3-1, Figure 5-28). The combined discharge and modelled loads estimated for the 2023–24 water year from the Endeavour Basin were also estimated to be the highest on record and are shown in Figure 5-26. In fact, preliminary estimates for the measured loads suggest that the TSS and PN loads reported here are likely underestimated. Over the 18-year period from 2006–07:

- discharge ranged from 753 GL (2019–20) to 4,877 GL (2023–24)
- TSS loads ranged from 38 kt (2019–20) to 244 kt (2023–24)
- DIN loads ranged from 34 t (2019–20) to 219 t (2023–24)
- PN loads ranged from 105 t (2019–20) to 683 t (2023–24).




Figure 5-26: Loads of (A) total suspended solids, dissolved inorganic (DIN) and particulate nitrogen (PN) and (B) discharge for the Endeavour Basin from 2006 to 2024. The loads reported here are the best estimates of annual mean concentration informed by nearest neighbour monitoring and by the Source Catchments modelling data applied to each water year. Dotted line represents the long-term median for basin discharge. Note the different scales on the two y-axes.




Figure 5-27: Daily discharge and sampling dates for the Endeavour Basin using combined values from the Annan River (gauge 107003A) and Endeavour River gauge (107001B) for the 2023–24 wet season.




Figure 5-28: Long-term discharge for the Endeavour Basin using combined values from the Annan River (gauge 107003A - Beesbike) and Endeavour River (gauge 107001B - Flaggy). Daily (blue) and water year (1 October to 30 September, red symbols) discharge volumes shown. Red dashed line represents long-term median of the combined annual discharge. Method for estimation is described in Table 2-3.

### Ambient water quality

Both ambient and event water quality results were plotted against distance from the mouths of the Annan or Endeavour River (Figure 5-29). Ambient mean and median values for each parameter are compared against the Eastern Cape York regional guidelines for the open coastal (ER02, ER03, AR02, AR03, and Dawson Reef), mid-shelf (AE04), and offshore (Forrester Reef) water bodies in Table C-1.

Trends associated with distance from the river mouth are difficult to discern due to the influence of two rivers with varying discharge on the sample sites. However, TSS and PN concentrations generally declined with distance from the rivers, while Secchi depth increased (Figure 5-29).




Figure 5-29: Water quality parameters (surface and subsurface samples) and Secchi depth over distance from river mouth (km) for the Endeavour Basin focus region during ambient conditions (2023–24 water year). Note that data includes samples collected at varying distances from two river mouths (Annan and Endeavour), with each site plotted at the distance from the closest river. Complex regression lines result from this combination of data and varying river influences. Water body classifications are shown along the x-axes: open coastal (OC) and mid-shelf (MS). Note the y-axes are logarithmic scales. Fitted lines are generalised additive models.

Comparison of the 2023–24 ambient results with previous years and the water quality GVs (Table C-1) highlights that:

- Overall, the annual condition WQ Index scored 'good' for the Annan-Endeavour focus region (Figure 5-30), similar to the 2022–23 water year. However, there was a decline in the productivity and particulates sub-indicator scores;
- Median TSS met the GVs at all sites;
- Mean Secchi depth did not meet the GV at any open coastal water body sites, but did meet the GV at AE04 in the mid-shelf water body;
- Chl-a met GVs at all sites;

- $\text{NO}_x$  exceeded the annual and wet season GVs at all sites, with some median values more than double the GVs and 2022–23 median.  $\text{PO}_4$  met the GVs at all sites except for AE04 in the mid-shelf water body;
- Median PN concentrations exceeded the relevant GVs at AR02, AR03, and AE04 but met the GVs at ER02 and ER03. PP met the relevant GVs at all sites;
- The median wet season turbidity and chlorophyll values calculated from the datalogger at Dawson Reef exceeded the respective GVs for the open coastal water body. Median values have declined since 2018, but increased over the 2023–24 wet season (Table 5-1); and
- The median wet season turbidity at Forrester Reef calculated from the datalogger was below the relevant GV. Wet season median chlorophyll concentration matched the GV for the mid-shelf water body and was almost double the previous wet season median (Table 5-2).




Figure 5-30: Annual condition version of the WQ Index for the Annan-Endeavour focus region. The WQ Index uses two formulations but only the annual condition index (based on post-2015 sampling design) is currently available for the Cape York region. WQ Index colour coding: ◆ – 'very good'; ◆ – 'good'; ◆ – 'moderate'; ◆ – 'poor'; ◆ – 'very poor'. Indicators or sub-indicators that are used to calculate the WQ Index are shown as coloured lines on each plot. Error bars (vertical black lines) on the WQ Index represent the 95% quantile intervals. Calculations for index formulations are described in [Appendix B](#).

Table 5-1: Median wet season turbidity and chlorophyll concentrations from a Wetlabs FLNTU datalogger at Dawson Reef from the 2018–19 water year until present and guideline values (GV) from the open coastal water body.

Dawson Reef	2018–19	2019–20	2020–21	2021–22	2022–23	2023–24	GV
Median turbidity wet season (NTU)	0.87	0.60	0.59	0.54	0.60	0.85	0.8
Median chlorophyll wet season ( $\mu\text{g L}^{-1}$ )	0.31	0.20	0.20	0.11	0.15	0.27	0.46

Table 5-2: Median wet season turbidity and chlorophyll concentrations from a Wetlabs FLNTU datalogger at Forrester Reef from the 2019–20 water year until present and guideline values (GV) from the mid shelf water body.

<b>Forrester Reef</b>	<b>2019 –20</b>	<b>2020 –21</b>	<b>2021 –22</b>	<b>2022 –23</b>	<b>2023– 24</b>	<b>GV</b>
Median turbidity wet season (NTU)	0.29	0.23	0.31	0.26	0.27	0.5
Median chlorophyll wet season ( $\mu\text{g L}^{-1}$ )	0.19	0.13	0.15	0.12	0.26	0.26

### *Event water quality*

River discharge for the Annan and Endeavour Rivers over the 2023–24 water year was three times above the mean annual discharge (Figure 5-26), primarily due to flooding associated with TC Jasper. Flooding in the Annan catchment peaked at over 14 m in the upper catchment on 18 December 2023, approximately 3 m higher than historic DNRM flood peak records (since 1969). The Annan River flood event was categorised as a 1-in-1000-year flood, and was accompanied by extensive landslides and debris flows, as well as severe bank erosion and loss of riparian vegetation, which all contributed to extreme sediment loads discharged to the coastal waters from the Annan catchment and other rivers to the south (Howley *et al.* 2024). The Endeavour River also experienced extreme flooding associated with the Jasper event, but river heights did not surpass historic flood levels. Targeted event sampling for the Annan-Endeavour sites occurred on 20 December 2023, 2 days after peak discharge in the upper Annan catchment (Figure 5-27; Figure 5-31). Flood plumes at that time from the combined Annan, Endeavour, Bloomfield, Daintree, and other Wet Tropics rivers reached from the river mouths beyond the outer reefs and north past Cape Melville, inundating coral reefs and seagrass meadows across the entire southeastern Cape York region (Figure 5-2)




Figure 5-31: Satellite true colour image of flood plumes associated with Annan and Endeavour rivers and Tropical Cyclone Jasper plus sample locations on 20 December 2023 (left). The corresponding processed imagery on the right panel shows the Forel Ule classes.

Samples collected from the TC Jasper flood plume had some of the highest TSS and DIN concentrations and lowest Secchi depths measured over the MMP sampling period (since January 2018) from sites AR02, AR03, and AE04. Conversely, these sites had some of the lowest Chl-*a* concentrations over the MMP period, likely due to low water clarity.

Continuous monitoring in the Annan estuary showed that turbidity peaked near 2000 NTU on 18 December (Howley *et al.*, 2024). On the same day, MMP dataloggers at Dawson Reef, 6 km from the Annan River mouth (Figure 5-25), measured peak turbidity of 200 NTU, compared to previous maximum event turbidity measurements between 10 to 40 NTU. Chl-*a* peaked at  $3.0 \mu\text{g L}^{-1}$  on 24 December 2023 and was above the GV ( $0.46 \mu\text{g L}^{-1}$ ) for 13 days. Salinity dropped to 5 at Dawson Reef on 18 December 2023 and remained below 20 until 24 December, causing coral reef mortality (Howley *et al.* 2024). Approximately 30 km to the northeast at Forrester Reef, turbidity peaked at 13 NTU while salinity dropped to 5 on 19 December 2023 and remained below 25 for 3 days. Chl-*a* at Forrester Reef peaked at  $3.0 \mu\text{g L}^{-1}$  on 24 December 2023.

Additional event sampling occurred on 24 March 2024 following an average magnitude flood event. Turbidity at Dawson Reef on this day reached 56 NTU while TSS was  $86 \text{ mg L}^{-1}$  near the Annan River mouth.

Sample results from both event sampling trips are plotted against salinity in Figure 5-32. Typical for flood plumes, TSS, PN, PP, and DIN concentrations decreased along the salinity gradient, while Secchi depths and Chl-*a* generally increased.




Figure 5-32: Flood event water quality data from the Annan-Endeavour transect on 20 December 2023 (blue circles) and 24 March 2024 (red circles) including total suspended solids (TSS), Secchi depth, chlorophyll a (Chl-a), dissolved inorganic nitrogen (DIN), particulate nitrogen (PN), and particulate phosphorus (PP) plotted over the salinity gradient. Surface samples are plotted as blue circles and squares and depth samples as red circles and squares.

## 5.2 Wet Tropics region

The Wet Tropics region is divided into three focus regions which are dominated by the Barron and Daintree Rivers (Barron-Daintree), the Russell and Mulgrave Rivers (Russell-Mulgrave) and the Tully River. The results on the pressures and monitoring findings are presented separately for each focus region.

### 5.2.1 Barron-Daintree

This focus region contains the six sites of the 'Cairns Transect', which are sampled three times a year (Figure 5-33). This sampling design and frequency did not change in 2015 (unlike all other focus regions), as these sites are part of a long-term AIMS time-series (the 'Cairns Transect').




Figure 5-33: Sampling sites in the Barron-Daintree focus region shown with water body boundaries.

The combined discharge and loads calculated for the 2023–24 water year from the Barron, Daintree, and Mossman Basins were around 4.6 times higher than the long-term median values, the highest on record (Figure 5-34; Table 3-1). Over the 19-year period from 2005–06:

- discharge ranged from 1,855 GL (2019–20) to 14,527 GL (2023–24)
- TSS loads ranged from 183 kt (2019–20) to 1,546 kt (2023–24)
- DIN loads ranged from 211 t (2019–20) to 1,728 t (2023–24)
- PN loads ranged from 493 t (2019–20) to 4,079 t (2023–24).

Of the three focus regions within the Wet Tropics NRM region, the Barron, Daintree and Mossman Basins commonly contribute the lowest discharge and consistent loads compared to the two focus regions to the south (i.e., Russell-Mulgrave and Johnstone Basins and the Tully-Murray and Herbert Basins). However, in the 2023–24 water year modelled loads for the Barron, Daintree, and Mossman Basins were the highest seen over the monitoring period (Figure 5-34) and contributed similar discharge and loads to those focus regions to the south.




Figure 5-34: Combined discharge from rivers in the Barron-Daintree region including the Daintree River (gauge 108002A – Bairds), Bloomfield River (108003A – China Camp), Mossman River (gauge 109001A – Mossman) and Barron River (gauge 110001D – Myola). Daily (blue) and water year (1 October to 30 September, red symbols) discharge volumes shown. Red dashed line represents long-term median of the combined annual discharge.




Figure 5-35: Loads of (A) TSS, DIN and PN and (B) discharge for the Barron, Daintree, and Mossman Basins from 2006–2024. The loads reported here are a combination of ‘best estimates’ for each basin based on ‘up-scaled discharge data from gauging stations, monitoring data (Barron River), the DIN model developed in Lewis *et al.* (2014) and annual mean concentrations and discharge from monitoring data or Source Catchments modelling data. The dotted line represents the long-term median for basin discharge. Note the different scales on the two y-axes.

The very high discharge in the Barron-Daintree region was due to large rainfall associated with TC Jasper in December 2023 (Figure 5-34). Floodwaters impacted the inshore region throughout the Wet Tropics over the weeks following the passage of TC Jasper and were dispersed cross-shelf, which may have contributed to the patterns in the WQ Index scores for the Barron-Daintree region and the Wet Tropics more broadly.

Coastal hydrodynamics for the Wet Tropics region were available through the eReefs GBR4 hydrodynamic model (Skerratt *et al.* 2019) and excerpts from the model are shown in Figure 5-36 to Figure 5-39. Prevailing coastal currents were in a general northerly direction in the days prior to TC Jasper landfall (Figure 5-36) and as the cyclone crossed the Queensland coast (Figure 5-37). Net southerly transport occurred following the cyclone’s passage (Figure 5-38). Acute localised flooding impacts were visible from the Daintree and Barron Rivers around Cairns and to a lesser extent from the Russell-Mulgrave, Johnston, and Tully rivers two days after the cyclone, with reduced salinities (~34 PSU) (Figure 5-38). Ten days after the

passage of the cyclone, south-easterly transport patterns persisted, moving fresh water across the shelf (Figure 5-39). Although some southerly transport around Cape Grafton occurred, model runs suggested that the majority of floodwaters were transported cross-shelf which likely reduced the influence of the Barron and Daintree Rivers on the Russell-Mulgrave and Tully regions further to the south (Figure 5-39). The Russell-Mulgrave and Tully River discharges were relatively minor compared to Barron-Daintree (Table 3-1). Acute impacts of TC Jasper were observed in water quality of the Cape York NRM Region (Section 5.1) but were not generally observed in the Wet Tropics NRM Region and the associated WQ Index scores (discussion throughout Section 5.2). These patterns are corroborated by results from satellite remote sensing products (Section 4.2) for the Wet Topics region.




Figure 5-36: eReefs hydrodynamic model showing coastal hydrodynamics and drivers for the Wet Tropics region on 9 December 2023, prior to landfall of TC Jasper. Data: eReefs CSIRO GBR4 Hydrodynamic Model v2.0




Figure 5-37: eReefs hydrodynamic model showing coastal hydrodynamics and drivers for the Wet Tropics region on 13 December 2023, as TC Jasper crossed the Queensland coast near Wujal Wujal. Data: eReefs CSIRO GBR4 Hydrodynamic Model v2.0




Figure 5-38: eReefs hydrodynamic model showing coastal hydrodynamics and drivers for the Wet Tropics region on 15 December 2023, two days after TC Jasper crossed the Queensland coast. Data: eReefs CSIRO GBR4 Hydrodynamic Model v2.0




Figure 5-39: eReefs hydrodynamic model showing coastal hydrodynamics and drivers for the Wet Tropics region on 23 December 2023, ten days after TC Jasper crossed the Queensland coast. Data: eReefs CSIRO GBR4 Hydrodynamic Model v2.0

*Ambient water quality and the in situ Water Quality Index*

Distinct long-term trends (since 2005) were observed in some water quality variables, while others showed little change (Figure 5-40). Site-specific statistics and comparison to GVs for all variables are available in [Appendix C-5](#).

Concentrations of Chl-a remained generally stable, fluctuating around local GVs from 2005 until 2015 (Figure 5-40a). Over the period 2015–2024, mean concentration of Chl-a decreased overall (conditions improved). Chl-a in 2023–24 was below (met) the local GVs.

Secchi depth gradually declined (i.e., water clarity worsened) from 2005 until 2016 (Figure 5-40b). Over the period 2015–2024, mean Secchi depth remained stable (not improved or declined overall) despite a small oscillation over this 9-year period. Secchi depth in 2023–24 was below (not meeting) the local GVs at all sites except Green Island (C11).

Concentrations of TSS have fluctuated above and below the GVs since monitoring began in 2005 (Figure 5-40c). Over the period 2015–2024, mean concentration of TSS decreased (conditions improved). TSS in 2023–24 was below (met) the local GVs at all sites. Trends in TSS are different to those observed for Secchi depth. Although these are both indicators for water clarity, differences can be expected due to differences in methodology and GVs.

Concentrations of NO<sub>x</sub> generally increased and remained above the local GVs from 2005 until 2015 (Figure 5-40d). Over the period 2015–2024, there was a potential trend for improvement in conditions (NO<sub>x</sub> declined), but it was not a significant change at this time. Oscillations in concentration were visible over the most recent 9-year period, which may develop into a significant change (improvement) in the coming years if trends continue. NO<sub>x</sub> in 2023–24 was above (did not meet) the local GVs at all sites except Fairlead Buoy (C8).

Concentrations of PO<sub>4</sub> were generally stable and close to the local GVs from 2005 until 2015 (Figure 5-40e). Over the period 2015–2024, mean concentration of PO<sub>4</sub> varied, decreasing in concentration until around 2022 and then increasing again to 2024. The concentration of PO<sub>4</sub> in 2023–24 was close to (or slightly higher) compared with 2015 but there is currently no significant change in concentration between these time points. PO<sub>4</sub> in 2023–24 was above (exceeded) the local GVs for all sites, which represented a deterioration in conditions compared to 2022–23 when PO<sub>4</sub> was just below (met) the local GVs.

Concentrations of PN remained stable and well below the local GVs since monitoring began in 2005 (Figure 5-40f). Over the period 2015–2024, mean concentration of PN remained very stable (not improved or declined overall). PN in 2023–24 was below (met) the local GVs.

Concentrations of PP remained relatively stable and close to the local GVs since monitoring began in 2005 (Figure 5-40g). Over the period 2015–2024, mean concentration of PP decreased (conditions improved) although the data show that there has been some oscillation over this 9-year period. PP in 2023–24 was below (met) the local GVs at some sites and above (exceeded) the local GVs at others.

Concentrations of POC remained relatively stable since monitoring began in 2005. There was a slight increase (deterioration in conditions) between 2005 and 2017 and a slight decrease (improvement in conditions) between 2017 and 2022 (Figure 5-40h). There was a small upward trend since 2022 (slight deterioration). Between 2015 and 2024, concentrations have increased slightly but this did not represent a significant change between these time points.

Concentrations of DOC increased substantially since 2005, although concentrations have stabilised in recent years (Figure 5-40i).

The WQ Index is calculated using two different formulations to communicate the a) long-term trend in water quality (based on the pre-2015 sampling design) and b) annual condition of water quality (based on the post-2015 sampling design, which increased the power to detect change). For the Barron-Daintree focus region, no additional sites were added in 2015, with

sampling still conducted three times per year. Section 2.5 and Appendix B contain details of the calculations for both Index formulations.

The long-term WQ Index showed a small trend (i.e., changing by a single grade) of decline in water quality from 2005–2018, driven by Chl-*a* and PP indicators (Figure 5-41a). Since 2018, this trend has reversed and water quality now shows an overall trend of improvement. This improving trend was driven by improvements in Chl-*a*, water clarity, and PP indicators.

The annual condition WQ Index scored water quality as ‘moderate’ during the 2015–18 water years and ‘good’ during the past six water years, including 2023–24 (Figure 5-41b). There was a slight increase in the annual condition score this year (change within a single grade), driven by improvements in the productivity and water clarity indicators. This version of the Index scores water quality parameters against GVs relevant to the season when samples are collected (wet versus dry GVs).

It is important to note that the two versions of the WQ Index are designed to answer separate questions and therefore differences in scores between the versions are expected.




Figure 5-40: Temporal trends in water quality variables for the Barron-Daintree focus region: a) chlorophyll a (Chl-a), b) Secchi depth, c) total suspended solids (TSS), d) nitrate/nitrite ( $\text{NO}_x$ ), e) phosphate ( $\text{PO}_4$ ), f) particulate nitrogen (PN), g) particulate phosphorus (PP), h) particulate organic carbon (POC) and i) dissolved organic carbon (DOC). Generalised additive mixed effect models (trends) are represented by blue lines with shaded areas defining 95% confidence intervals of those trends and black dots represent observed data (depth weighted averages). These trends and data are accounting for the effects of wind, waves, tides, and seasons after applying x-z detrending. Dashed horizontal reference lines indicate annual guideline values for open coastal waters.




Figure 5-41: The Water Quality Index (WQ Index) for the Barron-Daintree focus region. The WQ Index uses two formulations to communicate: a) long-term trend (based on pre-2015 sampling design) and b) the annual condition (based on post-2015 sampling design). WQ Index colour coding: ● / ◆ – 'very good'; ○ / ◇ – 'good'; ● / ◆ – 'moderate'; ○ / ◇ – 'poor'; ● / ◆ – 'very poor'. Indicators or sub-indicators that are used to calculate the WQ Index are shown as coloured lines on each plot. Error bars (vertical black lines) on the WQ Index represent the 95% quantile intervals. Calculations for both formulations are described in [Appendix B](#).

### Event water quality

The Barron River at Myola gauge (Figure 5-42) peaked at 14.15 m on 17 December 2023, well above the major flood level of 10 m (430,000 ML d<sup>-1</sup>), which coincided with the crossing of TC Jasper on 13 December 2023. Three other flow events exceeded the minor flood level (5.0 m) on 19 January 2024 (5.99 m), 20 March 2024 (5.74 m), and 27 March 2024 (5.87 m). The total discharge for the Barron Basin for 2023–2024 was 3,600,000 ML which is 5.8 times above the long-term median (Table 3-1). Monitoring of the Barron River flood plume occurred on 28 December 2023 across the southern section of the Cairns Transect 11 days after the major flood peak.




Figure 5-42: Barron River at Myola flow gauge for the 2023–24 wet season. The date of event sampling by JCU is marked as an orange dot.

Additional event monitoring sites were visited in the southern section of the Cairns Transect to sample the large Barron River flood in December 2024 and capture a broader area of the salinity gradient (Figure 5-43). Samples were collected from both the top (~30 cm below the surface) and bottom of the water column (~1-2 m above the seabed), with additional sites collected to capture a broader salinity gradient. The sampling was conducted 11 days after the flood peak and the satellite images show that the plume had dissipated considerably over that period (Figure 5-44).




Figure 5-43: Satellite true colour image on 28 December 2023 (left panel) showing the flood plume sampling sites from the Barron River. The corresponding processed imagery on the right panel shows the Forel Ule classes.




Figure 5-44: True colour satellite images of the Barron flood plume from 24 December 2023 (left panel) and from 28 December 2023 (right panel).

As the sampling was conducted 11 days after the flood peak (earlier sampling was prevented by reduced site access, weather, and sea state), the plume was well-dispersed throughout the waters off Cairns as shown by the satellite images (Figure 5-44). However, this timeline of sampling (i.e., ~1 to 2 weeks after the peak) allowed the primary productivity of the offshore waters to be likely near peak levels following deposition and/or dispersion of the particulate material, and was assessed to determine the degree to which flood waters influenced the area in the weeks following the peak flow. Due to the mixing of the waters and the ‘flashiness’ of the river flows (i.e., the elevated river flow quickly receded), the majority of samples were collected from the >25 salinity zone. The concentrations of TSS, DIP, PN, and PP were similar to those measured at comparable salinities from the Russell-Mulgrave and Tully flood plumes (Figure 5-45 and Figure 5-46). However,  $\text{NO}_x$  concentrations in the Barron plume were lower than at comparable salinities from the Russell-Mulgrave and Tully River flood plumes (Figure 5-45b), and Chl-a concentrations (Figure 5-45c) in four of the depth samples (ranging from 1 to  $4 \mu\text{g L}^{-1}$ ) were much higher compared to the samples from the Russell-Mulgrave and Tully flood plumes. This result suggests that the  $\text{NO}_x$  had largely been consumed by algae and the algae had begun to sink to the seafloor. Pesticide grab samples were also collected in the transect, and concentrations of diuron, metolachlor, and imidacloprid were measured at very low levels and at salinities close to seawater (Figure 5-47). While the results for diuron and metolachlor showed a distinct difference between the surface and depth samples, the results for imidacloprid showed variability across the whole depth profile. This most likely reflects the persistence of imidacloprid in marine waters, and is discussed further in Kaserzon *et al.*, (in prep).




Figure 5-45: Water quality data from the Cairns Transect region under the influence of flood plumes from the Barron River on 19 and 28 December 2023 including A) total suspended solids (TSS), B) oxidised nitrogen ( $\text{NO}_x$ ), and C) chlorophyll a plotted over the salinity gradient. Surface samples are plotted as blue circles and squares and depth samples as red circles and squares.




Figure 5-46: Water quality data from the Cairns Transect region under the influence of flood plumes from the Barron River on 19 and 28 December 2023 including A) dissolved inorganic phosphorus (DIP), B) particulate nitrogen (PN), and C) particulate phosphorus (PP) plotted over the salinity gradient. Surface samples are plotted in blue circles and squares and depth samples as red circles and squares.




Figure 5-47: Water quality data from the Cairns Transect region under the influence of flood plumes on 28 December 2023 including the herbicides diuron (A) and metolachlor (B) and the insecticide imidacloprid (C) plotted over the salinity gradient. Surface samples are plotted as blue circles and depth samples as red circles.

### 5.2.2 Russell-Mulgrave

The Russell-Mulgrave focus region is primarily influenced by discharge from the Russell-Mulgrave and Johnstone Basins and, to a lesser extent, by other rivers south of the focus region such as the Burdekin (Brodie *et al.*, 2013; Waterhouse *et al.*, 2017). Three sites were sampled three times per year in this focus region until the end of 2014. Following the implementation of the revised MMP water quality sampling design in 2015, 12 monitoring sites are sampled in this focus region up to 10 times per year, with five sites sampled during both the dry and wet seasons and seven additional sites sampled during major flood events (Table A-1). The monitoring sites form a transect from the river mouth to mid-shelf waters, representing a gradient in water quality. Five sites are in the open coastal water body, five sites are located in the mid-shelf water body, one site is in mid-estuarine waters, and one site is in enclosed coastal waters (Figure 5-48).




Figure 5-48: Sampling sites in the Russell-Mulgrave focus region, shown with the water body boundaries.

The combined discharge volume of the Russell-Mulgrave and Johnstone Rivers for the 2023–24 water year was 1.7-fold higher than the long-term median (Figure 5-49).

The combined discharge and loads calculated for the 2023–24 water year from the Russell-Mulgrave and Johnstone Basins were in the higher range to that recorded over the past decade (Figure 5-50). Over the 19-year period from 2005–06:

- discharge ranged from 6,318 GL (2014–15) to 15,813 GL (2010–11)
- TSS loads ranged from 350 kt (2014–15) to 896 kt (2010–11)
- DIN loads ranged from 835 t (2014–15) to 2,722 t (2010–11)
- PN loads ranged from 1,177 t (2014–15) to 3,005 t (2010–11).

Of the three focus regions within the Wet Tropics NRM region, the Russell-Mulgrave and Johnstone Basins collectively contribute similar discharge and loads to the Tully-Murray and Herbert Basins during low to average discharge years, although the latter basins contribute higher values (particularly DIN) during the high discharge years such as in 2008–09 and 2010–11 water years.




Figure 5-49: Combined discharge for rivers in the Russell-Mulgrave region including the North Johnstone (gauge 112004A – Tung Oil) and South Johnstone gauge 112101B – Central Mill), Russell (gauge 111101D – Bucklands) and Mulgrave (gauge 111007A – Peets Bridge) Rivers. Daily (blue) and water year (1 October to 30 September, red symbols) discharge is shown. Red dashed line represents the long-term median of the combined annual discharge.




Figure 5-50: Loads of (A) TSS, DIN and PN and (B) discharge for the Russell, Mulgrave and Johnstone Basins from 2006 to 2024. The loads reported here are a combination of 'best estimates' for each basin based on 'up-scaled discharge data from gauging stations, monitoring data (Johnstone River), the DIN model developed in Lewis *et al.* (2014) and annual mean concentrations and discharge from monitoring data or Source Catchments modelling data. Dotted line represents the long-term median for basin discharge. Note the different scales on the two y-axes.

### *Ambient water quality and the in situ Water Quality Index*

Water quality showed trends along the sampling transect (cross-shelf gradient in northerly direction). Sites located nearest to the river mouth (distance from river mouth = 0 km) had high concentrations of  $\text{NO}_x$  and particulate nutrients (PN and PP), which declined with distance away from the river mouth, reaching low levels in mid-shelf waters (Figure 5-51, Table C-2). Concentrations of Chl-a and TSS showed a similar pattern to nutrient concentrations and tended to decline with distance from the river mouth. Secchi depths were low at sites near the river mouth (water clarity was poor) and increased (water clarity improved) with distance from the river mouth. These spatial patterns are generally consistent with those that are typically observed in the region.

Seasonal differences in  $\text{NO}_x$ , TSS, and PN are also typical, where concentrations (especially near the river mouth) are much higher during the wet than the dry season. These seasonal

differences tended to become less pronounced further offshore (e.g., concentrations of PN during wet and dry seasons are similar in mid-shelf waters). In 2023–24 typical seasonal differences were observed for  $\text{NO}_x$  and PN; however, TSS concentrations were similar between wet and dry season near the river mouth, likely due to coastal processes and wind-driven resuspension. Concentrations of PP were similar between wet and dry seasons. Concentrations of Chl-a were higher in the wet season compared to the dry season, and Secchi depths were generally similar between the wet and dry seasons (Figure 5-51).




Figure 5-51: Water quality variables measured during ambient wet and dry season sampling in 2023–24 along the Russell-Mulgrave focus region transect. Chlorophyll a (Chl-a), nitrate/nitrite ( $\text{NO}_x$ ), total suspended solids (TSS), Secchi depth, particulate nitrogen (PN), and particulate phosphorus (PP) are shown with distance from the Russell-Mulgrave River mouth. Water body classifications are shown along the x-axes: open coastal (OC) and mid-shelf (MS). Note the y-axes are logarithmic scales. Fitted lines are generalised additive models.

Distinct long-term trends (since 2005) were observed in some water quality variables, while others showed little change (Figure 5-52). Site-specific statistics and comparison to GVs for all variables are available in [Appendix C-5](#).

Concentrations of Chl-*a* fluctuated above and below the GVs since monitoring began in 2005 (Figure 5-52a). Over the period 2015–2024, mean concentration of Chl-*a* decreased overall. Chl-*a* in 2023–24 was below (met) the local GVs at all sites. Chlorophyll fluorescence measured by FLNTU instruments (Figure 5-52a) likewise fluctuated around GVs since monitoring began in 2007. The differences between FLNTU chlorophyll fluorescence and Chl-*a* concentration reflect differences in sampling location and measurement method.

Secchi depth gradually declined (i.e., water clarity worsened) from 2005 until 2015. Over the period 2015–2024, mean Secchi depth increased (i.e., water clarity improved) in a steady trend (Figure 5-52b). Despite this trend for improvement, Secchi depth in 2023–24 was below (did not meet) the local GVs.

Concentrations of TSS fluctuated above and below the GVs since monitoring began in 2005. Over the period 2015–2024, mean concentration of TSS declined until around 2022 and increased in recent years. Although there appears to be an improvement in TSS since 2022, values have not significantly increased when compared to 2015 (Figure 5-52c). TSS concentrations in 2023–24 were below (met) the local water quality GVs.

Turbidity remained relatively stable and close to the GVs since monitoring began in 2005 (Figure 5-52d). Over the period 2015–2024, turbidity remained stable (not improved or declined overall) despite small oscillations over this period. Turbidity in 2023–24 was above (did not meet) the GVs at some sites (RM10 and RM7) but was below (met) GVs at others (RM1 and RM8).

Concentrations of NO<sub>x</sub> steadily increased and remained above the local GVs from 2005 until 2015 (Figure 5-52e). Over the period 2015–2024, mean concentrations of NO<sub>x</sub> steadily decreased (conditions improved). NO<sub>x</sub> in 2023–24 was above (did not meet) the local GVs.

Concentrations of PO<sub>4</sub> fluctuated above the local GVs from 2005 until 2015 (Figure 5-52f). Over the period 2015–2024, mean concentration of PO<sub>4</sub> increased (conditions declined) despite a large oscillation during this 8-year period. PO<sub>4</sub> in 2023–24 was above (did not meet) the local GVs.

Concentrations of PN were below (met) local GVs but increased over the period 2005–2015 (Figure 5-52g). Over the period 2015–2018 concentrations increased to levels above local GVs and were more stable in recent years. Although there appears to be a slight increase in concentration (deterioration in condition) between 2015 and 2024, this trend does not represent a significant change between these time points. PN in 2023–24 was above (did not meet) the local GVs at some sites but met the GVs at RM1 and RM7.

Concentrations of PP remained relatively stable and close to the local GVs since monitoring began in 2005 (Figure 5-52h). Over the period 2015–2024, mean concentration of PP remained stable (not improved or declined overall) despite showing some minor variability. PP in 2023–24 was above (did not meet) the local GVs at some sites but met the GVs at RM1, RM3, and RM7.

Concentrations of POC increased dramatically over the period 2005–2017 (Figure 5-52i). Over the period 2015–2024, mean concentration of POC decreased, as has been observed in most other focus regions (Figure 5-52j).

The WQ Index is calculated using two different formulations to communicate the a) long-term trend in water quality (based on the pre-2015 sampling design) and b) annual condition of water quality (based on the post-2015 sampling design, which increased the power to detect change). Section 2.5 and Appendix B contain details of the calculations for both Index formulations.

For the Russell-Mulgrave region, the long-term WQ Index has shown a small trend (i.e., changing within a grade) of decline in water quality from 2009–2019, which stabilised around 2020 (Figure 5-53a). This downward trend was generally driven by trends in PN, PP, and Chl-

a indicators. Over the last four years, this trend appears to have reversed and water quality now shows signs of overall improvement. This improving trend is driven by improvements in Chl-*a*, PN, and PP indicators.

The annual condition WQ Index has scored water quality as ‘moderate’ since its inception (2015–present) (Figure 5-53b). This version of the Index scores water quality parameters against GVs relevant to the season when samples are collected (wet versus dry GVs).

It is important to note that the two versions of the WQ Index are designed to answer separate questions and therefore differences in scores between the versions are expected.




Figure 5-52: Temporal trends in water quality variables for the Russell-Mulgrave focus region: a) chlorophyll *a* (Chl-*a*), b) Secchi depth, c) total suspended solids (TSS), d) turbidity, e) nitrate/nitrite ( $\text{NO}_x$ ), f) phosphate ( $\text{PO}_4$ ), g) particulate nitrogen (PN), h) particulate phosphorus (PP), i) particulate organic carbon (POC) and j) dissolved organic carbon (DOC). Generalised additive mixed effect models (trends) are represented by blue lines with shaded areas defining 95% confidence intervals of those trends and black dots represent observed data (depth weighted averages). These trends and data are accounting for the effects of wind, waves, tides, and seasons after applying x-z detrending. Trends of records from ECO FLNTUSB instruments are represented in red, and individual records can be found in Figure C-1. Dashed horizontal reference lines indicate annual guidelines for open coastal waters.




Figure 5-53: The Water Quality Index (WQ Index) for the Russell-Mulgrave focus region. The WQ Index uses two formulations to communicate the a) long-term trend (based on pre-2015 sampling design) and b) annual condition (based on post-2015 sampling design). WQ Index colour coding: ● / ◆ – 'very good'; ○ / ◇ – 'good'; ● / ◆ – 'moderate'; ○ / ◇ – 'poor'; ● / ◆ – 'very poor'. Indicators or sub-indicators that are used to calculate the WQ Index are shown as coloured lines on each plot. Error bars (vertical black lines) on the WQ Index represent the 95% quantile intervals. Calculations for both formulations are described in [Appendix B](#).

### Event water quality

The Mulgrave River at Peets Bridge gauge peaked at 8.98 m on 17 December 2023, well above the major flood level of 8.0 m, which coincided with the crossing of TC Jasper on 13 December 2023. Another four minor flood levels (5.0 m) were exceeded over the wet season including on 23 February 2024 (5.70 m), 8 March 2024 (5.26 m), 29 March 2024 (6.70 m), and 16 April 2024 (6.29 m). The Russell River at Bucklands gauge peaked at 7.92 m on 17 December 2023 just below the major flood level of 8.0 m. One other flow event peaked above the moderate flood level (7.0 m) on 8 March 2024 (7.41 m). Another three minor flood levels (6.0 m) were exceeded at this gauge site over the wet season including on 24 February 2024 (6.44 m), 19 March 2024 (6.65 m), and 29 March 2024 (6.59 m). The combined gauge peak flows on 17 December 2024 reached 200,000 ML d<sup>-1</sup> (Figure 5-54). The total discharge for the Russell Mulgrave Basin for 2023–2024 was 6,800,000 ML which is 1.6 times above the long-

term median (Table 3-1). Water quality monitoring of the Russell Mulgrave River flood plume occurred on 19 December 2023, two days after the major flood peak.




Figure 5-54: Russell-Mulgrave River (Mulgrave River at Peets Bridge + Russell River at Bucklands flow gauges) for the 2023–24 wet season. The date of offshore event sampling by JCU is marked as an orange dot.

The standard event monitoring sites off the Russell-Mulgrave region were all sampled at the top (~30 cm below the surface) of the water column with additional sites collected to capture more of the salinity gradient (Figure 5-55). Samples from the bottom of the water column (~1–2 m above the seabed) were collected from the standard ambient monitoring sites to examine the vertical mixing of the plume. Selected results from the water column profiling are shown in Figure 5-56. The profiles of salinity show the increasing influence of the river flood plume in the upper ~5 to 10 m of the water column at the sites offshore from the Russell-Mulgrave River (Figure 5-56). The more distant sites from the river (e.g. RM1 at Fitzroy Island and RM2) still showed a plume influence in the upper 10 m of the water column (Figure 5-55 and Figure 5-56). The light profiles show increasing PAR available at greater water column depths from the river mouth to the more distant sites offshore (Figure 5-56).




Figure 5-55: Satellite true colour image on 19 December 2023 (left panel) from the Wet Tropics NRM region showing the flood plume sampling sites from the Russell-Mulgrave (RM labels) and the Tully (TUL labels) Rivers. The corresponding processed imagery on the right panel shows the Forel Ule classes.




Figure 5-56: Map of the offshore Russell-Mulgrave standard event sampling sites (left); and plots of the water column profiles (right) of salinity (black lines) and light (orange lines with  $K_D$  values) at the selected sites shown on the map.

The sampling was conducted two days after the flood peak (earlier sampling was prevented by limited site access due to road closures) and observations and satellite imagery show that the plume was largely constrained within the inshore waterbody (Figure 5-55). There was still considerable discharge from the river and (with the additional samples collected) the sampling was able to capture excellent coverage across the salinity gradient from freshwater (0 salinity) to seawater (~35 salinity). The TSS concentrations were all below 20 mg L<sup>-1</sup> with a peak of 18.5 mg L<sup>-1</sup> at 6.8 salinity (Figure 5-57a). These TSS concentrations are within the range of the measurements from previous wet seasons at the upstream Mulgrave River at Deeral GBR Catchment Loads Monitoring Program site.

As the sampling captured the early stages of the plume, there was less opportunity for the development of primary productivity which is reflected in the both the NO<sub>x</sub> (Figure 5-57b) and Chl-*a* (Figure 5-57c) concentrations. The NO<sub>x</sub> concentrations closely follow a conservative mixing trend over the salinity gradient consistent with previous plume sampling that included samples near the plume peak, while the Chl-*a* values were highly variable but all below 0.5 µg L<sup>-1</sup>. Interestingly, the DIP (Figure 5-58a) and PN (Figure 5-58-b) concentrations did not display conservative mixing behaviour over the salinity gradient, which may reflect desorption/adsorption processes respectively occurring in the plume particularly in the 0 to 20 salinity values. Conversely, the behaviour of PP (Figure 5-58c) generally followed a pattern consistent with conservative mixing.

Both diuron and metolachlor displayed conservative mixing behaviour in the Russell Mulgrave plume while imidicloprid concentrations were more variable (Figure 5-59). The relatively lower concentrations (near detection limits) of imidacloprid may explain this variability. The concentrations of diuron, metolachlor, and imidachloprid were below available guideline values, although metolachlor exceeded a proposed 99% guideline value for freshwaters (King *et al.* 2017); it is anticipated this value will soon be revised and available for marine systems to better ascertain risk.




Figure 5-57: Water quality data from the Russell-Mulgrave region under the influence of flood plumes on 19 December 2023 including A) total suspended solids (TSS), B) Oxidised nitrogen (NO<sub>x</sub>), and C) chlorophyll a plotted over the salinity gradient. Surface samples are plotted as blue circles and depth samples as red circles.




Figure 5-58: Water quality data from the Russell-Mulgrave region under the influence of flood plumes on 19 December 2023 including A) dissolved inorganic phosphorus (DIP), B) particulate nitrogen (PN), and C) particulate phosphorus (PP) plotted over the salinity gradient. Surface samples are plotted as blue circles and depth samples as red circles.




Figure 5-59: Water quality data from the Russell-Mulgrave region under the influence of flood plumes on 19 December 2023 including the herbicides A) diuron and B) metolachlor and C) the insecticide imidacloprid plotted over the salinity gradient. Surface samples are plotted as blue circles and depth samples as red circles.

### 5.2.3 Tully

The Tully focus region is primarily influenced by discharge from the Tully-Murray and Herbert Rivers and, to a lesser extent, by the Burdekin River in large flow years (Brodie *et al.*, 2013).

One site was sampled in this focus region three times per year until the end of 2014. Following the implementation of the revised MMP water quality sampling design in 2015, 11 monitoring

sites are sampled in this focus region up to 10 times per year, with six sites sampled during both the dry and wet seasons and five additional sites sampled during major flood events (Table A-1). The monitoring sites form a transect from the river to mid-shelf waters, representing a gradient in water quality. Seven sites are in the open coastal water body, one is located in the mid-shelf water body, one site is in mid-estuarine waters, and two sites are in lower estuarine waters (Figure 5-60).




Figure 5-60: Sampling sites in the Tully focus region, shown with the water body boundaries.

The combined discharge and loads calculated for the 2023–24 water year from the Tully, Murray, and Herbert Basins were around 1.9-fold higher than the long-term median (Figure 5-62). Over the 19-year period from 2005–06:

- discharge ranged from 4,491 GL (2014–15) to 24,166 GL (2010–11)
- TSS loads ranged from 260 kt (2014–15) to 1,827 kt (2010–11)
- DIN loads ranged from 1,082 t (2014–15) to 5,875 t (2010–11)
- PN loads ranged from 796 t (2014–15) to 5,307 t (2010–11).

Of the three focus regions within the Wet Tropics NRM region, the Tully, Murray, and Herbert Basins collectively contribute similar discharge and TSS and PN loads to the Russell, Mulgrave, and Johnstone Basins during low-to-moderate discharge years. However, the Tully, Murray, and Herbert Basins contribute higher values during the high discharge years such as in 2008–09 and 2010–11 as well as generally higher DIN loads in the average to above-average discharge years.




Figure 5-61: Combined discharge from rivers in the Tully region including the Tully (gauge 113006A – Euramo), Murray (gauge 114001A – Upper Murray) and Herbert (gauge 116001F – Ingham) Rivers. Daily (blue) and water year (1 October to 30 September, red) discharge is shown. Red dashed line represents the long-term median of the combined annual discharge. Please note as this is the combined discharge, high flows in one river will not necessarily be visible in the graph.




Figure 5-62: Loads of (A) TSS, DIN and PN and (B) discharge for the Tully, Murray, and Herbert Basins from 2006 to 2024. The loads reported here are a combination of 'best estimates' for each basin based on 'up-scaled discharge data from gauging stations, monitoring data (Tully and Herbert Rivers), the DIN model developed in Lewis *et al.* (2014) and annual mean concentrations and discharge from monitoring data or Source Catchments modelling data. The dotted line represents the long-term median for basin discharge. Note the different scales on the two y-axes.

### *Ambient water quality and the in situ Water Quality Index*

Water quality showed trends along the sampling transect (cross-shelf gradient in northerly direction). Sites located nearest to the river mouth (distance from river mouth = 0 km) had high concentrations of  $\text{NO}_x$  and particulate nutrients (PN and PP), which declined with distance away from the river mouth, reaching low levels in mid-shelf waters (Figure 5-63, Table C-2). Concentrations of Chl-*a* and TSS showed a similar pattern to nutrient concentrations and tended to decline with distance from the river mouth. Secchi depths were low at sites near the river mouth (water clarity was poor) and increased (water clarity improved) with distance from the river mouth. These spatial patterns are generally consistent with those that are typically observed in the region.

This year, seasonal differences were observed, where concentrations (especially near the river mouth) were higher during the wet than the dry season. Concentrations of  $\text{NO}_x$ , Chl-*a* and particulate nutrients (PP and PN) showed generally similar trends between wet and dry

seasons with higher concentration inshore compared with offshore and higher concentrations in the wet season compared with the dry season. Concentrations of TSS were higher in the wet season at enclosed coastal sites but were similar between the wet and dry seasons at sites in mid-shelf waters. Secchi depths were similar (water clarity was similar) across the transect during both the wet and dry seasons (Figure 5-63).




Figure 5-63: Water quality variables measured during ambient and event sampling in 2023–24 along the Tully focus region transect. Chlorophyll *a* (Chl-*a*), nitrate/nitrite (NO<sub>x</sub>), total suspended solids (TSS), Secchi depth, particulate nitrogen (PN), and particulate phosphorus (PP) are shown with distance from the Tully River mouth. Water body classifications are shown along the x-axes: open coastal (OC) and mid-shelf (MS). Note the y-axes are logarithmic scales. Fitted lines are generalised additive models.

Distinct long-term trends (since 2005) were observed in some water quality variables, while others showed little change (Figure 5-64). Site-specific statistics and comparison to GVs for all variables are available in [Appendix C-5](#).

Concentrations of Chl-a showed large fluctuations above and below the GVs since monitoring began in 2005 (Figure 5-64a). Over the period 2015–2024, mean concentration of Chl-a remained stable (not improved or declined overall) despite a large oscillation over this period. Chl-a in 2023–24 was below (met) the local GVs. Chlorophyll fluorescence measured by FLNTU instruments (Figure 5-64a) did not reflect these trends and was generally stable and well above (not meeting) GVs since monitoring began in 2007. The differences between FLNTU chlorophyll fluorescence and Chl-a concentration reflect differences in sampling location and measurement method.

Secchi depth gradually declined (i.e., water clarity worsened) from 2005 until 2015 and remained below (did not meet) the GVs. Over the period 2015–2024, mean Secchi depth increased (improved in condition) (Figure 5-64b). Secchi depth in 2023–24 was below (did not meet) the local GVs.

Concentrations of TSS fluctuated above and below the GVs since monitoring began in 2005 (Figure 5-64c). Over the period 2015–2024, mean concentration of TSS showed a slight decreasing trend, but this change was not significant between these time points. There was an oscillation in TSS over this 9-year period with a general trend of improvement between 2015 and 2022 and a deterioration in recent years. TSS concentrations in 2023–24 were below (met) the local water quality GVs.

Turbidity remained relatively stable and well above the GVs since monitoring began in 2005 (Figure 5-64d). Over the period 2015–2024, turbidity increased (declined in condition). Turbidity in 2023–24 was above (did not meet) the GVs.

Concentrations of NO<sub>x</sub> steadily increased and remained above the local GVs from 2005 until 2015 (Figure 5-64e). Over the period 2015–2024, mean concentration of NO<sub>x</sub> showed a large decrease (improvement in condition). NO<sub>x</sub> in 2023–24 was above (did not meet) the local GVs.

Concentrations of PO<sub>4</sub> fluctuated generally above the local GVs from 2005 until 2015 (Figure 5-64f). Over the period 2015–2024, mean concentration of PO<sub>4</sub> increased (condition declined) despite a large oscillation during this 9-year period. PO<sub>4</sub> in 2023–24 was generally above (did not meet) the local GVs.

Concentrations of PN generally increased since monitoring began in 2005 and started exceeding local GVs around 2015 (Figure 5-64g). Between 2015–2024, mean concentration of PN remained stable; however, PN concentrations oscillated over this 9-year period. PN in 2023–24 was just below (met) the local GVs.

Concentrations of PP have remained relatively stable and close to the local GVs since monitoring began in 2005 (Figure 5-64h). Over the period 2015–2024, mean concentration of PP has remained stable (not improved or declined overall) despite showing some minor variability over this 8-year period. PP in 2023–24 was below (met) the local GVs at some sites but was above (not meeting) the local GVs at other sites within the region.

Concentrations of POC showed variability over the period 2005–2017 (Figure 5-64i). Over the period 2015–2024, mean concentration of POC decreased overall, similar to all other focus regions. Concentrations of DOC increased over the period 2007–2015 but have stabilised since 2015 (Figure 5-64j).

The WQ Index is calculated using two different formulations to communicate the a) long-term trend in water quality (based on the pre-2015 sampling design) and b) annual condition of water quality (based on the post-2015 sampling design, which increased the power to detect change). Section 2.5 and Appendix B contain details of the calculations for both Index formulations.

The long-term WQ Index for the Tully region has shown the most variability of any focus region since the inception of the MMP. The period 2007–2014 was characterised by a large (i.e., changing over two grades) decline in water quality, driven by declines in all indicators with the exception of NO<sub>x</sub> (Figure 5-65a). The period 2014–2020 showed some variability, but water quality did not improve or decline overall. Over the last four years, a trend of improvement in water quality was observed and the long-term WQ Index scored moderate for 2023–24 (Figure 5-65a). This improving trend is driven by improvements in Chl-*a*, PN, PP, and water clarity indicators.

The annual condition WQ Index scored water quality as ‘moderate’ or ‘poor’ since its inception (2015–present). For the 2023–24 year, the Tully region received a ‘moderate’ score (Figure 5-65b). This version of the Index scores water quality parameters against GVs relevant to the season when samples are collected (wet versus dry GVs). It is important to note that the two versions of the WQ Index are designed to answer separate questions and therefore differences in scores between the versions are expected.




Figure 5-64: Temporal trends in water quality variables for the Tully focus region: a) chlorophyll a (Chl-a), b) Secchi depth, c) total suspended solids (TSS), d) turbidity, e) nitrate/nitrite ( $\text{NO}_x$ ), f) phosphate ( $\text{PO}_4$ ), g) particulate nitrogen (PN), h) particulate phosphorus (PP), i) particulate organic carbon (POC) and j) dissolved organic carbon (DOC). Generalised additive mixed effect models (trends) are represented by blue lines with shaded areas defining 95% confidence intervals of those trends and black dots represent observed data (depth weighted averages). These trends and data are accounting for the effects of wind, waves, tides, and seasons after applying x-z detrending. Trends of records from ECO FLNTUSB instruments are represented in red, and individual records can be found in Figure C-1. Dashed horizontal reference lines indicate annual guidelines for open coastal waters.




Figure 5-65: The Water Quality Index (WQ Index) for the Tully focus region. The WQ Index uses two formulations to communicate: a) long-term trend (based on pre-2015 sampling design) and b) the annual condition (based on post-2015 sampling design). WQ Index colour coding: ● / ◆ – 'very good'; ● / ◆ – 'good'; ● / ◆ – 'moderate'; ● / ◆ – 'poor'; ● / ◆ – 'very poor'. Indicators or sub-indicators that are used to calculate the WQ Index are shown as coloured lines on each plot. Error bars (vertical black lines) on the WQ Index represent the 95% quantile intervals. Calculations for both formulations are described in [Appendix B](#).

### Event water quality

The Tully at Euramo gauge peaked at 8.88 m on 17 December 2023 (moderate flood) (Figure 5-66) just below the major flood level of 9.0 m (86,000 ML d<sup>-1</sup>), which coincided with the crossing of TC Jasper on 13 December 2023. Other flood peaks above the moderate flood level (8.0 m) during the 2023–24 season occurred on 25 February 2024 (8.66 m peak) and 10 March 2024 (8.30 m peak). The Tully River had three additional floods that exceeded the minor flood level (6.0 m) which peaked on 18 January 2024 (6.46 m), 30 March 2024 (7.12 m), and 16 April 2024 (6.04 m). The total discharge for the Tully Basin for 2023–2024 was 5,600,000 ML which is 1.6 times above the long-term median (Table 3-1). Monitoring of the Tully River flood plume occurred on 16 December 2023, one day prior to the major flood peak with another sampling trip on 21 March 2024 which fell between a moderate peak on 10 March 2024 and a minor peak on 30 March 2024.




Figure 5-66: Tully River at Euramo flow gauge for the 2023–24 wet season. The dates of event sampling by JCU are marked as orange dots.

The standard event monitoring sites for the Tully region were all sampled at the surface (~30 cm below the surface) of the water column, with additional sites collected to capture the salinity gradient (Figure 5-67). Samples from the bottom of the water column (~1–2 m above the seabed) were collected from the standard ambient monitoring sites to examine the vertical mixing of the plume. The satellite image from 19 December 2023 when the sampling was undertaken is shown in Figure 5-55. This shows highly turbid waters just offshore from the coast including around Dunk and Bedarra Islands. The plume on 16 December 2023 was largely confined to the coast, with some plume influence beginning to impinge on the mid-shelf area. The satellite image from the second sampling period on 21 March 2024 shows relatively lower turbidity in the waters near the coast, but with a more developed plume further offshore onto the mid-shelf (Figure 5-67).




Figure 5-67: Satellite true colour image on 13 March 2024 (left panel) showing the flood plume sampling sites from the Tully focus region. The corresponding processed imagery on the right panel shows the Forel Ule classes from 15 March 2024.

The December sampling was conducted one day before the flood peak, and observations from satellite imagery show that the plume was largely constrained within the inshore waterbody (Figure 5-55). With the additional samples collected, the sampling was able to capture solid coverage across the salinity gradient from freshwater (0 salinity) to seawater (~35 salinity) for both the December and March events.

The TSS concentrations were considerably higher in the December event ( $72 \text{ mg L}^{-1}$  at the end-of-river site compared to the March sampling of  $12 \text{ mg L}^{-1}$  at the same site) and this trend was generally reflected over the entire salinity gradient (Figure 5-68a) with the exception of elevated TSS concentration of  $67 \text{ mg L}^{-1}$  at 12 salinity in March 2024; this sample was from the Hull River mouth which is shallow (3 m) and so could be explained by either resuspension or river input (Figure 5-68a). TSS concentrations in the deep samples were generally higher than the corresponding surface samples which likely reflects settling through the water column or resuspension of the benthic sediments.

As the December sampling captured the early period of the plume there was less opportunity for the development of primary productivity due to light requirements for growth, which is reflected in the both the  $\text{NO}_x$  (Figure 5-68b) and Chl-a (Figure 5-68c) concentrations. In the samples from December, the  $\text{NO}_x$  concentrations closely followed a conservative mixing trend over the salinity gradient consistent with previous plume sampling where samples were collected near the plume peak, while the Chl-a values were generally below  $0.5 \text{ } \mu\text{g L}^{-1}$  between the 0 and 20 salinity zone, and mostly between  $0.5$  to  $1.0 \text{ } \mu\text{g L}^{-1}$  between the 20 and 35 salinity zone. The lower TSS concentrations recorded in the March sampling likely

provided more available light for primary production. Hence, while the  $\text{NO}_x$  concentrations in March were generally lower than the corresponding concentrations at similar salinities from the December sampling,  $\text{NO}_x$  concentrations did not follow conservative mixing in the March event and were greatly reduced by 25 salinity. With the exception of some outliers, the Chl-*a* concentrations were higher (mostly in the 0.5 to 1.0  $\mu\text{g L}^{-1}$  range) in the lower salinity ranges (5 to 25 salinity zone) compared to the December samples (Figure 5-68c).

The concentrations of DIP (Figure 5-69a), PN (Figure 5-69b), and PP (Figure 5-69c) did not display conservative mixing behaviour over the salinity gradient which may reflect desorption/adsorption processes respectively occurring in the plume. There is also evidence for the influence of potential sediment resuspension, with some elevated concentrations in the depth samples (which notably was not observed in the corresponding  $\text{NO}_x$  concentrations).

There was less pesticide sampling of the Tully plume compared to the Russell Mulgrave plume, so limited interpretation can be made on pesticide behaviour over the salinity gradient. Concentrations of diuron and imidacloprid in the December sampling were higher in the Tully plume freshwater sections (Figure 5-70) compared to the Russell Mulgrave, with imidacloprid around the proposed freshwater guideline from King *et al.* (2017). This result is expected due to the sampling times of the plumes, with the Tully sampled just before the peak flood and the Russell Mulgrave sampled just after the peak. As expected, all pesticide concentrations were much lower in the samples taken in March.




Figure 5-68: Water quality data from the Tully region under the influence of flood plumes on 16 December 2023 and 21 March 2024 including A) total suspended solids (TSS), B) Oxidised nitrogen (NO<sub>x</sub>), and C) chlorophyll a plotted over the salinity gradient. Surface samples are plotted as blue circles and squares and depth samples as red circles and squares.




Figure 5-69: Water quality data from the Tully region under the influence of flood plumes on 16 December 2023 and 21 March 2024 including A) dissolved inorganic phosphorus (DIP), B) particulate nitrogen (PN), and C) particulate phosphorus (PP) plotted over the salinity gradient. Surface samples are plotted as blue circles and squares and depth samples as red circles and squares.




Figure 5-70: Water quality data from the Tully region under the influence of flood plumes on 16 December 2023 and the 21 March 2024 including the herbicides A) diuron and B) metolachlor and C) the insecticide imidacloprid plotted over the salinity gradient. Surface samples are plotted as blue circles and depth samples as red circles.

### 5.3 Burdekin region

Three sites were sampled in this focus region three times per year until the end of 2014. Following the implementation of the revised MMP water quality sampling design in 2015, 15 sites are now sampled in this focus region up to nine times per year. Six sites are sampled during both the dry and wet seasons and nine additional sites are sampled during major flood events (Table A-1). The monitoring sites are located along a transect away from the river mouth in a north-westerly direction, representing a gradient in water quality. Eight sites are in open coastal waters, two sites are in the mid-shelf water body, and five sites are in enclosed coastal waters (Figure 5-71).




Figure 5-71: Sampling sites in the Burdekin focus region, shown with the water body boundaries.

The total discharge for the Burdekin region (Burdekin and Haughton Basins) in 2023–24 was 1.3 times above the long-term median (Figure 5-72; Table 3-1). The combined discharge and loads calculated for the 2023–24 water year from the Burdekin and Haughton Basins were in the average range over the past decade (Figure 5-73).

Over the 19-year period from 2005–06:

- discharge ranged from 1,036 GL (2014–15) to 37,470 GL (2010–11)
- TSS loads ranged from 290 kt (2013–14) to 15,024 kt (2007–08)
- DIN loads ranged from 275 t (2014–15) to 4,019 t (2010–11)
- PN loads ranged from 586 t (2013–14) to 22,083 t (2007–08).

During the very large discharge years (2007–08, 2008–09, 2010–11 and 2018–19), the Burdekin and Haughton Basins (dominated by the Burdekin Basin) produced by far the highest loads of TSS and PN compared to any of the other focus regions. In contrast, the DIN loads are either similar to or lower than the basins of the Wet Tropics and Mackay-Whitsunday regions during the high discharge years and much lower during the lower discharge years.




Figure 5-72: Total discharge from rivers in the Burdekin region including the Black (gauge 117002A – Bruce Highway), Haughton (gauge 119003A – Powerline), Burdekin (gauge 120006B – Clare), Elliot (gauge 121002A – Guthalungra) and Don (gauge 121003A – Reeves) Rivers and Bluewater (gauge 117003A – Bluewater), Barratta (gauge 119101A – Northcote) and Euri (gauge 121004A – Koonandah) Creeks. Daily (blue) and water year (1 October to 30 September, red) discharge is shown. Red dashed line represents the long-term median annual discharge.




Figure 5-73: Loads of (A) TSS, DIN and PN and (B) discharge for the Burdekin and Haughton Basins from 2006 to 2024. The loads reported here are a combination of 'best estimates' for each basin based on 'up-scaled' discharge data from gauging stations, monitoring data (Burdekin River), and annual mean concentrations and discharge from monitoring data or Source Catchments modelling data. Dotted line represents the long-term median for basin discharge. Note the different scales on the two y-axes.

### *Ambient water quality and the in situ Water Quality Index*

Water quality showed trends along the sampling transect (cross-shelf gradient in northerly direction). Sites located nearest to the river mouth (distance from river mouth = 0 km) had high concentrations of  $\text{NO}_x$  and particulate nutrients (PN and PP), which declined with distance away from the river mouth, reaching low levels at sites furthest from the river mouth (Figure 5-74, Table C-2). Concentrations of Chl-a and TSS showed a similar pattern to nutrient concentrations and tended to decline with distance from the river mouth. Secchi depths were low at sites near the river mouth (water clarity was poor) and increased (water clarity improved) with distance from the river mouth. These spatial patterns are generally consistent with those that are typically observed in the region.

This year, seasonal differences in Chl-a,  $\text{NO}_x$ , and PN, were observed, where concentrations (especially near the river mouth) were higher during the wet than the dry season. For  $\text{NO}_x$ , these seasonal differences tended to become less pronounced further offshore (e.g.,

concentrations during wet and dry seasons were similar at sites furthest from the river mouth) while the opposite was observed for PN. Secchi depth and concentrations of TSS were variable between seasons, with generally poorer conditions observed in the inshore region and generally better conditions observed further from the river mouth during the dry season compared with the wet season (Figure 5-74).




Figure 5-74: Water quality variables measured during ambient and event sampling in 2023–24 along the Burdekin focus region transect. Chlorophyll *a* (Chl-*a*), nitrate/nitrite ( $\text{NO}_x$ ), total suspended solids (TSS), Secchi depth, particulate nitrogen (PN), and particulate phosphorus (PP) are shown with distance from the Burdekin River mouth. Water body classifications are shown along the x-axes: Enclosed coastal (EC) and open coastal (OC). Note the y-axes are logarithmic scales. Fitted lines are generalised additive models.

Distinct long-term trends (since 2005) were observed in some water quality variables, while others showed little change (Figure 5-75). Site-specific statistics and comparison to GVs for all variables are available in [Appendix C-5](#).




Figure 5-75: Temporal trends in water quality variables for the Burdekin focus region: a) chlorophyll a (Chl-a), b) Secchi depth, c) total suspended solids (TSS), d) turbidity, e) nitrate/nitrite ( $\text{NO}_x$ ), f) phosphate ( $\text{PO}_4$ ), g) particulate nitrogen (PN), h) particulate phosphorus (PP), i) particulate organic carbon (POC) and j) dissolved organic carbon (DOC). Generalised additive mixed effect models (trends) are represented by blue lines with shaded areas defining 95% confidence intervals of those trends and black dots represent observed data (depth weighted averages). These trends and data are accounting for the effects of wind, waves, tides, and seasons after applying x-z detrending. Trends of records from ECO FLNTUSB instruments are represented in red, and individual records can be found in Figure C-1. Dashed horizontal reference lines indicate annual guidelines for open coastal waters.

Concentrations of Chl-a showed large fluctuations above and below the GVs since monitoring began in 2005 (Figure 5-75a). Over the period 2015–2024, mean concentration of Chl-a remained stable (not improved or declined overall) despite a large oscillation over this 9-year period. Chl-a in 2023–24 was just below (met) the local GVs. Chlorophyll fluorescence measured by FLNTU instruments (Figure 5-75a) does not reflect these trends and has been fluctuating above GVs since monitoring began in 2007. The differences between FLNTU chlorophyll fluorescence and Chl-a concentration reflect differences in sampling location and measurement method.

Secchi depth remained relatively stable and below (did not meet) the GVs since monitoring began in 2005 (Figure 5-75b). Over the period 2015–2024, mean Secchi depth remained stable (not improved or declined overall) despite small oscillations over this 9-year period. Secchi depth in 2023–24 was below (did not meet) the local GVs.

Concentrations of TSS fluctuated above and below the GVs since monitoring began in 2005. Over the period 2015–2024, mean concentration of TSS oscillated, but concentrations of TSS did not significantly increase or decrease over this 9-year period (Figure 5-75c). TSS concentrations in 2023–24 were below (met) the local water quality GVs.

Turbidity remained relatively stable and above the GVs since monitoring began in 2005 (Figure 5-75d). Over the period 2015–2023, turbidity remained stable (i.e., water clarity did not significantly change) despite small oscillations over this 9-year period. Turbidity in 2023–24 was above (did not meet) the local water quality GVs.

Concentrations of  $\text{NO}_x$  steadily increased and remained well above the local GVs from 2005 until 2015 (Figure 5-75e). Over the period 2015–2024, mean concentration of  $\text{NO}_x$  decreased.  $\text{NO}_x$  in 2023–24 was above (did not meet) the local GVs.

Concentrations of  $\text{PO}_4$  were generally stable and well above the local GVs from 2005 until 2015 (Figure 5-75f). Over the period 2015–2022, mean concentration of  $\text{PO}_4$  steadily decreased (improved) but then increased (condition deteriorated) again in recent years, resulting in no overall significant trend for the 9-year period from 2015–2024.  $\text{PO}_4$  in 2023–24 was above (did not meet) the local water quality GVs.

Concentrations of PN showed large fluctuations above and below the GVs since monitoring began in 2005 (Figure 5-75g). Over the period 2015–2024, mean concentration of PN increased (condition deteriorated). PN in 2023–24 was just below (met) the local GVs for the Burdekin region.

Concentrations of PP remained relatively stable and generally below the local GVs since monitoring began in 2005 (Figure 5-75h). Over the period 2015–2024, mean concentration of PP remained stable (not improved or deteriorated overall). PP in 2023–24 was below (met) the local GVs for the Burdekin region.

Concentrations of POC showed large oscillations since monitoring began in 2005 (Figure 5-75i). Over the period 2015–2024, mean concentration of POC decreased, similar to all other focus regions. Concentrations of DOC increased over the period 2005–2017 but have stabilised in recent years (Figure 5-75j).

The WQ Index is calculated using two different formulations to communicate the a) long-term trend in water quality (based on the pre-2015 sampling design) and b) annual condition of water quality (based on the post-2015 sampling design, which increased the power to detect change). Section 2.5 and Appendix B contain details of the calculations for both Index formulations.

The long-term WQ Index showed a small (i.e., changing by a single grade) overall decline in water quality from the period 2010–2018, driven by PN and PP indicators (Figure 5-76a). The Index then stabilised for three years from 2018–2020 and showed a trend of improvement over the period 2021–2023 (Figure 5-76a). In 2023–24 the long-term Index was stable.

The annual condition WQ Index generally scored water quality as ‘moderate’ since its inception (2015–present), with the 2023–24 year receiving a ‘good’ score, which is only the second time a ‘good’ score has been achieved in this region (Figure 5-76b). This version of the Index scores water quality parameters against GVIs relevant to the season when samples are collected (wet versus dry GVIs).

It is important to note that the two versions of the WQ Index are designed to answer separate questions and therefore differences in scores between the versions are expected.




Figure 5-76: The Water Quality Index (WQ Index) for the Burdekin focus region. The WQ Index uses two formulations to communicate: a) long-term trend (based on pre-2015 sampling design) and b) the annual condition (based on post-2015 sampling design). WQ Index colour coding: ● / ◆ – ‘very good’; ○ / ◇ – ‘good’; ● / ◆ – ‘moderate’; ○ / ◇ – ‘poor’; ● / ◆ – ‘very poor’. Indicators or sub-indicators that are used to calculate the WQ Index are shown as coloured lines on each plot. Error bars (vertical black lines) on the WQ Index represent the 95% quantile intervals. Calculations for both formulations are described in [Appendix B](#).

### Event water quality

No event sampling was conducted in the 2023–24 wet season in the Burdekin focus region.

## 5.4 Mackay-Whitsunday region

The Mackay-Whitsunday region comprises four major river basins: the Proserpine, O’Connell, Pioneer, and Plane Basins. The region may also be influenced by runoff from the Fitzroy River during extreme events or through longer-term transport and mixing.

Three sites were sampled in this focus region three times per year until the end of 2014. From 2015, 11 sites have been sampled in this focus region up to five times per year, with five sites sampled during both the dry and wet seasons and six additional sites sampled during major flood events (Table A-1). The sites are located along a transect from the O’Connell River mouth to open coastal waters, representing a gradient in water quality. Ten sites are within open coastal waters and one site is in enclosed coastal waters (Figure 5-77).




Figure 5-77: Sampling sites in the Mackay-Whitsunday focus region, shown with the water body boundaries.

Annual discharge for the Mackay-Whitsunday region in the 2023–24 water year was below the long-term median levels (Figure 5-78).

The combined discharge and loads calculated for the 2023–24 water year from the Proserpine, O’Connell, Pioneer and Plane Basins (Figure 5-79) were in the lower range of values recorded over the past decade. Over the 19-year period from 2005–06:

- discharge ranged from 919 GL (2014–15) to 17,425 GL (2010–11)
- TSS loads ranged from 120 kt (2014–15) to 3,163 kt (2010–11)
- DIN loads ranged from 242 t (2014–15) to 3,814 t (2010–11)
- PN loads ranged from 356 t (2014–15) to 8,564 t (2010–11).




Figure 5-78: Combined discharge from rivers in the Mackay-Whitsunday region including the O'Connell (gauge 124001B – Stafford's Crossing), Andromache (gauge 124003A – Jochheims) and Pioneer (gauge 125016A – Dumbleton Weir T/W) Rivers and St. Helens (gauge 124002A – Calen), Sandy (gauge 126001A – Homebush) and Carmila (gauge 126003A – Carmila) Creeks. Daily (blue) and water year (1 October to 30 September, red) discharge is shown. Red dashed line represents the long-term median of the combined annual discharges. See Table 2-3 for a list of flow gauge data used. Please note as this is the combined discharge, high flows in one river will not necessarily be visible in the graph.




Figure 5-79: Loads of (A) TSS, DIN and PN and (B) discharge for the Proserpine, O'Connell, Pioneer, and Plane Basins from 2006 to 2024. The loads reported here are a combination of 'best estimates' for each basin based on 'up-scaled' discharge data from gauging stations, monitoring data (O'Connell and Pioneer Rivers and Sandy Creek), and annual mean concentrations and discharge from monitoring data or Source Catchments modelling data. Dotted line represents the long-term median for basin discharge. Note the different scales on the two y-axes.

*Ambient water quality and the in situ Water Quality Index*

Water quality showed trends along the sampling transect (O'Connell River mouth to open coastal waters). The site located in the enclosed coastal water body (distance from river mouth = 0 km) had high concentrations of Chl-*a*, TSS, and particulate nutrients (PN and PP), which declined with distance away from the river mouth (Figure 5-80, Table C-2). Secchi depths were low at sites near the river mouth (water clarity was poor) and increased (water clarity improved) with distance from the river mouth. NO<sub>x</sub> was variable across the transect. These spatial patterns (except NO<sub>x</sub>) are generally consistent with those that are typically observed in the region.

This year, some seasonal differences in Chl-*a*, NO<sub>x</sub>, and TSS were observed. Concentrations of Chl-*a* were greater during the wet than the dry season while NO<sub>x</sub>, and TSS varied across the transect. For TSS, concentrations were greater during the dry season at inshore sites and the seasonal differences tended to become less pronounced further offshore (e.g., concentrations during wet and dry seasons were similar at sites furthest from the river mouth). NO<sub>x</sub> concentrations were similar at sites which were close to the river mouth and furthest from the river mouth but were higher during the wet season at sites at intermediate distances. Secchi depths and concentrations of PN and PP generally didn't show seasonal differences along the transect (Figure 5-80).




Figure 5-80: Water quality variables measured during ambient and event sampling in 2023–24 along the Mackay-Whitsunday focus region transect. Chlorophyll *a* (Chl-*a*), nitrate/nitrite ( $\text{NO}_x$ ), total suspended solids (TSS), Secchi depth, particulate nitrogen (PN), and particulate phosphorus (PP) are shown with distance from the O'Connell River mouth. Water body classifications are shown along the x-axes: Enclosed coastal (EC) and open coastal (OC). Note the y-axes are logarithmic scales. Fitted lines are generalised additive models.

Distinct long-term trends (since 2005) were observed in some water quality variables, while others showed little change (Figure 5-81). Site-specific statistics and comparison to GVs for all variables are available in [Appendix C-5](#).




Figure 5-81: Temporal trends in water quality variables for the Mackay-Whitsunday focus region: a) chlorophyll a (Chl-a), b) Secchi depth, c) total suspended solids (TSS), d) turbidity, e) nitrate/nitrite ( $\text{NO}_x$ ), f) phosphate ( $\text{PO}_4$ ), g) particulate nitrogen (PN), h) particulate phosphorus (PP), i) particulate organic carbon (POC) and j) dissolved organic carbon (DOC). Generalised additive mixed effect models (trends) are represented by blue lines with shaded areas defining 95% confidence intervals of those trends and black dots represent observed data (depth weighted averages). These trends and data are accounting for the effects of wind, waves, tides, and seasons after applying x-z detrending. Trends of records from ECO FLNTUSB instruments are represented in red, and individual records can be found in Figure C-1. Dashed horizontal reference lines indicate annual guidelines for open coastal waters.

Concentrations of Chl-a were generally stable and well above the GVs over the period 2005–2015 (Figure 5-81a). Over the period 2015–2024, mean concentration of Chl-a decreased (improved) overall and in 2023–24 was below (met) the local GVs. Chlorophyll fluorescence measured by FLNTU instruments (Figure 5-81a) was relatively stable and well above GVs since monitoring began in 2007. The differences between FLNTU chlorophyll fluorescence and Chl-a concentration reflect differences in sampling location and measurement method.

Secchi depth gradually decreased (i.e., water clarity worsened) over the period 2005–2015 and was below (did not meet) GVs since monitoring began in 2005 (Figure 5-81b). Over the period 2015–2024, mean Secchi depth gradually increased (i.e., water clarity improved). Secchi depth in 2023–24 was below (did not meet) the local GVs.

Concentrations of TSS fluctuated above and below the GVs since monitoring began in 2005, although tend to be more stable than other focus regions. Over the period 2015–2024, mean concentration of TSS declined (water clarity improved) in a steady trend (Figure 5-81c). TSS concentrations in 2023–24 were below (met) the local water quality GVs at most sites.

Turbidity oscillated but remained above the GVs since monitoring began in 2005 (Figure 5-81d). Over the period 2015–2024, turbidity decreased overall (i.e., water clarity improved) despite oscillations over this 9-year period. Turbidity in 2023–24 was above (did not meet) the GVs at all sites except the O’Connell River mouth.

Concentrations of NO<sub>x</sub> steadily increased and remained well above the local GVs from 2005 until 2015 (Figure 5-81e). Over the period 2015–2024, mean concentration of NO<sub>x</sub> decreased, driven predominantly by changes in recent years. NO<sub>x</sub> in 2023–24 was above (did not meet) the local GVs in the Mackay-Whitsunday region but was just below (met) the local GVs at WHI1 and WHI5.

Concentrations of PO<sub>4</sub> were generally stable and well above the local GVs from 2005 until 2015 (Figure 5-81f). Over the period 2015–2024, mean concentration of PO<sub>4</sub> did not show an overall significant trend although there has been an oscillation (improvement and then deterioration in condition) over this 9-year period. PO<sub>4</sub> in 2023–24 was just above (did not meet) the local GVs.

Concentrations of PN showed fluctuations above and below the GVs since monitoring began in 2005 (Figure 5-81g). Over the period 2015–2024, mean concentration of PN oscillated (deteriorated until around 2018 and then improved in condition in recent years). However, these changes do not represent an overall significant trend for the 9-year period. PN in 2023–24 was below (met) the local water quality GVs.

Concentrations of PP increased and were generally above local GVs from the period 2005–2015 (Figure 5-81h). Over the period 2015–2024, mean concentration of PP decreased (improved). PP in 2023–24 was just above (did not meet) the local water quality GVs.

Concentrations of POC showed a large oscillation since monitoring began in 2005 (Figure 5-81i). Over the period 2015–2024, mean concentration of POC decreased, similar to all other focus regions. Concentrations of DOC increased over the period 2005–2017 but have stabilised in recent years (Figure 5-81j).

The WQ Index is calculated using two different formulations to communicate the a) long-term trend in water quality (based on the pre-2015 sampling design) and b) annual condition of water quality (based on the post-2015 sampling design, which increased the power to detect change). Section 2.5 and Appendix B contain details of the calculations for both Index formulations.

The long-term WQ Index showed a small (i.e., changed by a single grade) overall decline in water quality from period 2007–2018, driven by water clarity, PN, and PP indicators (Figure 5-82a). This trend then stabilised and has reversed in recent years. Between 2020 and 2023

a slight trend of improvement in water quality was observed, which stabilised in 2023–24 (Figure 5-82a).

The annual condition WQ Index scored water quality as ‘moderate’ or ‘poor’ since its inception (2015–present), with the 2023–24 year receiving a ‘moderate’ score (Figure 5-82b). This version of the Index scores water quality parameters against GVs relevant to the season when samples are collected (wet versus dry GVs).

It is important to note that the two versions of the WQ Index are designed to answer separate questions and therefore differences in scores between the versions are expected.




Figure 5-82: The Water Quality Index (WQ Index) for the Mackay-Whitsunday focus region. The WQ Index uses two formulations to communicate: a) long-term trend (based on pre-2015 sampling design) and b) the annual condition (based on post-2015 sampling design). WQ Index colour coding: ● / ◆ – ‘very good’; ● / ◆ – ‘good’; ● / ◆ – ‘moderate’; ● / ◆ – ‘poor’; ● / ◆ – ‘very poor’. Indicators or sub-indicators that are used to calculate the WQ Index are shown as coloured lines on each plot. Error bars (vertical black lines) on the WQ Index represent the 95% quantile intervals. Calculations for both formulations are described in [Appendix B](#).

*Event water quality*

No event sampling was conducted during the 2023–24 wet season in the Mackay-Whitsunday focus region.

## 6 CONCLUSIONS

In 2023–24, the overall Reef catchment area had discharge of 1.7 times the long-term median, the largest discharge since the 2018–19 water year. On a regional basis, the Cape York and Wet Tropics NRM regions had discharge well above their respective long-term medians (2.6 and 2.1 times above, respectively) while central and southern catchments experienced near-median discharge.

There were two cyclones which affected the Reef during 2023–24 and several tropical low pressure systems also resulted in high rainfall and river discharge events in some locations. Tropical Cyclone Jasper impacted the Cape York region and particularly the Annan and Endeavour catchments from 13–19 December 2023 while Tropical Cyclone Kirrily impacted the Burdekin region around Townsville from 25–26 January 2024. As a result of TC Jasper, discharge from the Cape York region for the 2023–24 water year was the highest on record (in records extending back to the 1990–91 water year) and the discharge from the Wet Tropics was the highest since the 2010–11 water year. In contrast to the Cape York region, impacts to the Wet Tropics from TC Jasper appear to be limited in their extent, possibly due to south-easterly dispersal of freshwater from the Barron and Daintree rivers around Cape Grafton and across the shelf. This likely contributed to patterns seen in the long-term and annual condition WQ Index scores as well as the patterns observed from remote sensing products (Table 6-1) for the Wet Tropics region.

Central and southern Reef catchments experienced moderate rainfall and discharge, despite impacts from TC Kirrily, which brought only limited rainfall to the region. Discharge from the Burdekin was 1.2 times the long-term median and the Fitzroy was close to the long-term median. The discharge from the Mackay-Whitsunday NRM region was below (0.8 times) the long-term median while the Burnett-Mary NRM region received above average discharge (1.7 times the long-term median).

In 2023–24, the long-term WQ Index showed trends of stability in water quality in all regions where this score is able to be generated (this does not include the Cape York region). The annual condition WQ Index scored water quality as either ‘good’ or ‘moderate’ in all focus regions in 2023–24. There was a decline in the annual index score to ‘moderate’ for the Cape York region, as a result of impacts from TC Jasper.

Recent trend analysis based on the previous nine years of monitoring data (presented in the GAMMs) indicates that many water quality indicators are showing signs of stability or improvement in all focus regions. The only exceptions to this trend were PN in the Burdekin focus region and PO<sub>4</sub> in the Russell-Mulgrave and Tully focus regions, which showed a trend of deterioration over the last nine years. In the Mackay-Whitsunday region, all water quality indicators showed a trend of improvement since 2015. These findings are likely a product of near-median river discharge over the last ~5 years in most focus regions, with few major flood events or cyclones impacting most of the Reef catchments in recent years prior to 2023–24, noting that Cape York long-term trends are not assessed yet as there are not enough data for a robust long-term assessment.






In 2023–24, TSS was the only indicator that met guideline values (GVs) in all focus regions. Chl-*a* met GV in all regions except for the Normanby and Stewart focus regions in Cape York. PN and PP met GV in most focus regions. PO<sub>4</sub>, NO<sub>x</sub>, and Secchi depth did not meet GV in most focus regions, despite recent trends of improvement in some of these indicators (especially NO<sub>x</sub>). This year, trend analysis showed that NO<sub>x</sub> improved since 2015 in all focus regions (with the exception of the Barron-Daintree), although it remains well above GV and continued improvement is needed before GV may be met.























This report shows some acute impacts to water quality conditions resulting from TC Jasper, especially in the Cape York region, but presents some positive results and trends of









improvement for many parameters in the Reef lagoon for the 2023–24 water year. It is important to interpret these trends cautiously, and further work is needed to determine to what extent oceanographic and climatic drivers contribute to observed trends. Associated trends in catchment load reductions from the Great Barrier Reef Catchment Loads Monitoring Program (Water Quality & Investigations, 2024) are an important line of evidence that need to be established before trends in inshore water quality can be related to catchment land use practices.







The main findings for each NRM region are highlighted below and the results are separated into ambient (routine sampling during wet and dry seasons) and event-based monitoring (sampling during flood events). Table 6-1 provides a high-level summary by NRM region.















In conclusion, the inshore water quality component of the Marine Monitoring Program continues to provide a high value, long-term dataset to inform management of the Reef. The multiple lines of evidence utilised in the Program include *in situ* water chemistry sample collection, *in situ* water quality time-series' from dataloggers, remote sensing analysis, and marine modelling, which all assist in providing a high degree of confidence in the interpretation of the results. The Program's outputs have a wide application across scientific, management, government, and public audiences. The Program's findings were used extensively in the recent update of the 2022 Scientific Consensus Statement (Waterhouse *et al.*, 2024), the Great Barrier Reef Outlook Report 2024 (Great Barrier Reef Marine Park Authority, 2024), and the current review of the Reef water quality targets. Continuation of the Program and strengthening of linkages with other Paddock to Reef components will further enhance the value of the Marine Monitoring Program.

Table 6-1: Summary of results for some of the primary indicators measured in the MMP Inshore Water Quality program, 2023–24. \*Arrows indicate difference relative to long-term patterns:  area exposed in 2023–24 similar (difference ≤ 5%) to long-term patterns,  decrease in area exposed (difference > 5%),  increase in area exposed (difference > 5%),  coral reef,  seagrass.

NRM region	Drivers and Pressures		Remote sensing mapping and modelling		Water Quality Index	
	Cyclone activity (timing)	River discharge (relative to long-term median; <1.5 is blue, 1.5-2 is orange and >2 is red)	Area (in %) exposed to a potential risk* (categories II-IV)  [and difference relative to long term]	Area (in %) exposed to the <u>highest</u> potential risk (categories III and IV)*  [and difference relative to long term]	Annual 2023–24	Long-term
Reef-wide	NA	1.7	<p>Reef: 9% exposed [ &lt; +1%]</p> <p> 7% exposed [ +3%]</p> <p> 71% exposed [ +1%]</p> <p>• Note: [ +7%] in seagrass area exposed to cat. of risk III. Likely related to [] in Cape York region.</p>	<p>Reef: 3% exposed [ +1%]</p> <p> 2% exposed [ &lt; +1%]</p> <p> 35% exposed [ +6%]</p> <p>• Only inshore Reef waters and habitats, with the largest proportion in the enclosed coastal waters.</p>	NA	NA
Cape York	TC Jasper, 13–17 December 2023	2.6	<p>Cape York: 11% exposed [ +4%]</p> <p> 11% exposed [ +7%]</p> <p> 66% exposed [ +10%]</p>	<p>Cape York: 3% [ +1%]</p> <p> 2% exposed [ &lt; +1%]</p> <p> 28% exposed [ +6%]</p>	Moderate	NA

	Drivers and Pressures		Remote sensing mapping and modelling		Water Quality Index	
NRM region	Cyclone activity (timing)	River discharge (relative to long-term median; <1.5 is blue, 1.5-2 is orange and >2 is red)	Area (in %) exposed to a potential risk* (categories II-IV)  [and difference relative to long term]	Area (in %) exposed to the <u>highest</u> potential risk (categories III and IV)*  [and difference relative to long term]	Annual 2023–24	Long-term
			<ul style="list-style-type: none"> <li><b>Note:</b> there was an increase in coral areas exposed to the lowest risk category (II+7%), and an increase in seagrass areas exposed to the risk category III (+8%).</li> </ul>	<ul style="list-style-type: none"> <li><b>Note:</b> Only <i>inshore</i> Cape York waters and habitats are exposed, with the largest proportion in the enclosed coastal waters.</li> </ul>		
<b>Wet Tropics</b>	TC Jasper, 13–17 December 2023	2.1	<p>Wet Tropics: 16% exposed [→ +4%]</p> <p> 3% exposed [→ &lt;1%]</p> <p> 98% exposed [→ &lt;1%]</p> <ul style="list-style-type: none"> <li><b>Note:</b> There was a decrease in seagrass area exposed to the highest potential risk categories (IV: -12%) toward the lowest risk category II (+9%) likely related to localised WT1 movements north of Hinchinbrook Island</li> </ul>	<p>Wet Tropics: 5% [→ +1%]</p> <p> 2% exposed [→ +1%]</p> <p> 67% exposed [↘ -9%]</p> <ul style="list-style-type: none"> <li><b>Note:</b> Only <i>inshore</i> Wet Tropics waters and habitats are exposed, with the largest proportion in the enclosed coastal waters.</li> </ul>	Good	<b>Declined</b> between 2008–2018, <b>improved</b> past 5 years and showing signs of <b>stability</b> in all focus regions in 2023–24
<b>Burdekin</b>	TC Kirrily, 25–26 January 2024	1.2	<p>Burdekin: 9% exposed [→ -1%]</p> <p> 1% exposed [→ &lt;-1%]</p> <p> 87% exposed [→ -1%]</p>	<p>Burdekin: 3% exposed [→ -1%]</p> <p> &lt;1% exposed [→ &lt;-1%]</p> <p> 52% exposed [↗ +8%]</p>	Good	<b>Declined</b> gradually from 2010–2018, <b>stable</b> over the last 3 years

NRM region	Drivers and Pressures		Remote sensing mapping and modelling		Water Quality Index	
	Cyclone activity (timing)	River discharge (relative to long-term median; <1.5 is blue, 1.5-2 is orange and >2 is red)	Area (in %) exposed to a potential risk* (categories II-IV)  [and difference relative to long term]	Area (in %) exposed to the <u>highest</u> potential risk (categories III and IV)*  [and difference relative to long term]	Annual 2023–24	Long-term
			<p>• <b>Note:</b> There was an increase in the seagrass areas exposed to potential risk category III in the Burdekin region, which was related to a higher frequency of primary waters measured south of Magnetic Island. However, the total seagrass area exposed to a potential risk (II-IV) was similar to the long-term trends.</p>	<p>• <b>Note:</b> Only inshore Burdekin waters and habitats are exposed, with the largest proportion in the enclosed coastal waters.</p>		
<b>Mackay-Whitsunday</b>	NA	0.8	<p>Mackay-Whitsundays: 9% exposed [➡ -4%]</p> <p> 4% exposed [➡ -2%]</p> <p> 82% exposed [➡ -14%]</p> <p><b>Note:</b> There was a decrease in seagrass area exposed to the cat. of risk II and IV (-14% and -6%)</p>	<p>Mackay-Whitsundays: 2% exposed [➡ -1%]</p> <p> 1% exposed [➡ &lt;-1%]</p> <p> 36% exposed [➡ &lt;-1%]</p> <p><b>Note:</b> Only <u>inshore</u> Mackay-Whitsunday waters and habitats are exposed, with the largest proportion in the enclosed coastal waters.</p>	Moderate	<b>Declined</b> 2008–2017, <b>improved</b> over the period 2018 to 2024
<b>Fitzroy</b>	NA	1.0	<p>Fitzroy: 8% exposed [➡ -2%]</p> <p> 3% exposed [➡ &lt;-1%]</p>	<p>Fitzroy: 3% exposed [➡ &lt;-1%]</p> <p> 1% exposed [➡ &lt;1%]</p>	Good (see <a href="#">Appendix D</a> )	<b>Improved</b> from 2008–2015, <b>stable</b> over last 4

NRM region	Drivers and Pressures		Remote sensing mapping and modelling		Water Quality Index	
	Cyclone activity (timing)	River discharge (relative to long-term median; <1.5 is blue, 1.5-2 is orange and >2 is red)	Area (in %) exposed to a potential risk* (categories II-IV)  [and difference relative to long term]	Area (in %) exposed to the <u>highest</u> potential risk (categories III and IV)*  [and difference relative to long term]	Annual 2023–24	Long-term
			 67% exposed [  -23%]  <b>Note</b> There was a decrease in seagrass area exposed to the cat. of risk II and IV (-19% and -8%)	 34% exposed [  <-4%]  <b>Note:</b> Only <u>inshore</u> Fitzroy waters and habitats are exposed, with the largest proportion in the enclosed coastal waters.		years (see <a href="#">Appendix D</a> )
Burnett-Mary	NA	1.7	Burnett-Mary: 3% exposed [  <-1%]   2% exposed [  <1%]   57% exposed [  -26%]  <b>Note</b> There was a decrease in seagrass area exposed to the cat. of risk II(-126%)	Burnett-Mary: 1% exposed [  <-1%]   2% exposed [  <1%]   29% exposed [  <1%]  <b>Note:</b> Only <u>inshore</u> Burnett-Mary waters and habitats are exposed, with the largest proportion in the enclosed coastal waters.	NA	NA

## 6.1 Cape York

The annual condition WQ Index for the Cape York region was 'moderate' for the 2023–24 water year. Despite the extreme impacts of TC Jasper, the Annan-Endeavour focus region had a 'good' score for the annual WQ Index, while the Normanby, Stewart, and Pascoe focus regions were 'moderate'. No long-term trends have been evaluated yet in the Cape York region but the annual score declined from the previous year, driven in all focus regions by declines in productivity and particulate sub-indicator scores.

Discharge from rivers in the Cape York region was between 2 to 4 times above the long-term median discharge for all focus regions, except for the Stewart which was close to the median. The Normanby saw flooding across Princess Charlotte Bay throughout most of the wet season, with floodwaters consistently flowing over 120 km to the north past the Stewart River transect and some freshwater influence was measured during most ambient Normanby sampling trips. Discharge in the Annan-Endeavour focus region was dominated by the extreme flooding associated with TC Jasper which resulted in extreme sediment loads discharged to the coast, and flood plumes extending beyond the outer reef and north to Cape Melville.

*Ambient water quality - Enclosed coastal, open coastal, and mid-shelf waters:*

- NO<sub>x</sub> exceeded the GVs at most sites for all Cape York focus regions and median concentrations increased compared median values from the 2022–23 wet season, contributing to declines in the productivity sub-indicator scores for all focus regions. In many cases median NO<sub>x</sub> values were doubled compared to 2022–23 medians.
- Chl-a met the GVs at almost all sites except for some Pascoe region sites. However, median Chl-a concentrations increased across the Stewart, Normanby, and Annan-Endeavour transects.
- Median Secchi depths were less than (did not meet) the GVs at all focus regions except for some Annan-Endeavour transect sites (those sites closer to the Endeavour River).
- TSS met the GVs at most sites, except for some within the Normanby region.
- PN, PP, and PO<sub>4</sub> comparisons against GVs were mixed, with some sites and focus regions meeting the GVs.
- Increases in median NO<sub>x</sub>, PO<sub>4</sub>, and Chl-a concentrations across the Stewart focus region were likely due to the influence from Normanby Basin flood plumes throughout the wet season.
- Median wet season turbidity and chlorophyll concentrations measured continuously (every 10 minutes) at Dawson Reef (6 km from the Annan River mouth) increased compared to the previous four wet seasons, and turbidity at Dawson Reef exceeded the GV for the open coastal water body.
- Median wet season turbidity concentrations measured continuously at Forrester Reef (30 km to the northeast from Dawson Reef), stayed stable compared to the previous four wet seasons and met the GVs for the mid-shelf water body, while chlorophyll concentrations doubled and matched the GV.

Table 6-2: Water quality indicator summary for Cape York. Performance relative to guideline values is shown as: generally exceeding (✗) or meeting (✓) guideline values across all sites. Mixed results across sites are indicated by both signs.

Water quality indicator	Pascoe	Stewart	Normanby	Annan-Endeavour
NO <sub>x</sub>	✗	✗	✗	✗
PO <sub>4</sub>	✗ ✓	✓	✗ ✓	✗ ✓
PN	✗	✗ ✓	✗ ✓	✗ ✓
PP	✓	✗ ✓	✗ ✓	✓
TSS	✓	✓	✗ ✓	✓
Secchi depth	✗	✗	✗	✗ ✓
Chl-a	✗ ✓	✓	✓	✓

### Event water quality

- TC Jasper, a 1-in-1000-year event in mid-December caused extreme flooding across the Annan and Normanby catchments, as well as rivers further south. Primarily due to this event, total discharge for the Annan River was 3 times the median annual discharge. Flood plumes from the combined Annan, Endeavour, Bloomfield, Daintree, and other Wet Tropics rivers reached from south of Cairns north past Cape Melville, and from the river mouths out beyond the outer reefs. Flooding in the Normanby also reached historic levels, and flood plumes in Princess Charlotte Bay were evident three weeks past peak Jasper discharge, flowing far to the north and inundating an area of over 3500 km<sup>2</sup>.
- Targeted flood sampling along the Annan-Endeavour focus region on 20 December, two days after peak discharge in the upper Annan catchment, showed some of the highest TSS and DIN concentrations and lowest Secchi depths measured over the MMP sampling period (since January 2018) from sites AR02, AR03, and AE04 along a transect from the Annan River mouth to the mid-shelf reef. Conversely, these sites had some of the lowest Chl-a concentrations over the historic MMP sampling period for this focus region, likely due to low water clarity.
- Turbidity measured continuously at Dawson Reef, 6 km from the Annan River mouth measured peak turbidity of 200 NTU, compared to previous maximum event turbidity measurements between 10 to 40 NTU. Median turbidity for the wet season at this site exceeded the GV, likely due to the influence of TC Jasper. Chlorophyll at Dawson Reef was above the GV for 13 days after the event, while salinity dropped to 5 and remained below 20 until 24 December, causing coral reef mortality as observed by the MMP water quality sampling team.
- TC Jasper event samples were collected from flood plumes near the Bizant and Kennedy river mouths during regular sampling on 31 December 2023, 10 days after

the Normanby River peaked at Kalpowar Crossing. Samples collected on 31 December had the highest event TSS concentrations on record for Normanby-Kennedy transects sites (since sampling began in 2018), with 165 mg L<sup>-1</sup> at KR01 (2 km from Kennedy River mouth) and 20 mg L<sup>-1</sup> at KR02 (6 km from the river mouth). The highest DIN, PN, POC, PP, and Chl-a concentrations on record for PCB were also recorded that day, approximately 6 km from the Normanby river mouth.

- Flooding in the Normanby and Kennedy Rivers in January, February, March, and April influenced water quality in Princess Charlotte Bay as well as the Stewart transect, where there was no major local flooding but ambient samples showed increases as much as double or more in Chl-a, NO<sub>x</sub>, and PO<sub>4</sub> concentrations.
- Land use has not changed significantly across the Pascoe catchment over recent years, and there were no extreme flood events. Other factors that may have contributed to increased TSS and DIN loads include the timing of rainfall and fires across the catchment. Wildfires burnt most of the upper Pascoe catchment late in the 2023 dry season ([www.firenorth.org.au/nafi3](http://www.firenorth.org.au/nafi3)). Late dry season fires can increase sediment and nutrient run-off to adjacent rivers (Townsend and Douglas, 2000; Howley *et al.*, 2022). This may have contributed to increased sediment and nutrient loads and declines in the Pascoe region water quality.
- In the Cape York region, 89% of the area was not exposed to a potential risk category, in keeping with long-term patterns (93%), and only 1% (1,032 km<sup>2</sup>) of the region was exposed to the highest risk category IV. Approximately 11% (1,092 km<sup>2</sup>) of the region's coral reefs and 66% (1,751 km<sup>2</sup>) of the region's seagrasses were exposed to a potential risk (combined risk categories II–IV). The total coral and seagrass area exposed to a potential risk (II–IV) were over the long-term trends (+7% and +10% respectively), which was consistent with the high discharge measured in 2023–24. There was an increase in coral areas exposed to the lowest risk category (II+7%), and an increase in seagrass areas exposed to the risk category III (+8%). Only the inshore Cape York waters, seagrass, and coral habitats were exposed to the highest categories of potential risk (III and IV), with the largest proportion located in the region's enclosed coastal waters. Mid-shelf and offshore Cape York reefs and seagrasses were exposed to the lower potential risk category II or to no/very low risk.

## 6.2 Wet Tropics

The combined discharge from the Daintree, Mossman, and Barron basins was around 4.6 times greater than the long-term median. Discharge from the Russell-Mulgrave and Johnstone basins was 1.7 times higher than the long-term median, after near-median discharge in the previous three years. Discharge from the Tully-Murray-Herbert basins was 1.9 times the long-term median after discharges close to or below the long-term median discharge in the previous three years.

*Ambient water quality - Enclosed coastal, open coastal, and mid-shelf waters:*

- NO<sub>x</sub>, PO<sub>4</sub>, and Secchi depth generally did not meet water quality GVs for any focus region in the Wet Tropics (Table 6-3).
- TSS, Chl-*a*, PN, and PP met GVs for all focus regions in the Wet Tropics.
- Over the period from 2015–2024, NO<sub>x</sub>, Secchi depth, and Chl-*a* showed a trend of improvement in two of the three focus regions in the Wet Tropics.
- Many indicators showed a trend of stability (no net improvement or deterioration) across all focus regions of the Wet Tropics.
- PO<sub>4</sub> was the only indicator which showed a trend of deterioration, which occurred in the Russell-Mulgrave and Tully focus regions but not in the Barron-Daintree.
- Long term Water Quality Index scores have remained stable this year after showing a trend of improvement over the past two to five years (depending on the region). For the 2023–24 water year, the annual condition Water Quality Index score was 'good' despite high discharge from Wet Tropics rivers due to TC Jasper.

Table 6-3: Water quality indicator summary for the three focus regions of the Wet Tropics. Performance relative to guideline values is shown as: generally exceeding (✗) or meeting (✓) guideline values across all sites. The trend of the indicator (2015–present) is shown as: deteriorating (↘), improving (↗) or stable (→).

Water quality indicator	Barron-Daintree	Russell-Mulgrave	Tully
NO <sub>x</sub>	✗ →	✗ ↗	✗ ↗
PO <sub>4</sub>	✗ →	✗ ↘	✗ ↘
PN	✓ →	✓ →	✓ →
PP	✓ ↗	✓ →	✓ →
TSS	✓ ↗	✓ →	✓ →
Secchi depth	✗ →	✗ ↗	✗ ↗
Chl- <i>a</i>	✓ ↗	✓ ↗	✓ →

*Wet season and event water quality*

In the Wet Tropics, 84% of the region was not exposed to a potential risk, in keeping with long-term patterns, and only 1% (or 443 km<sup>2</sup>) of the region was exposed to the highest risk category IV. Approximately 3% (78 km<sup>2</sup>) of the region's coral reefs and 98% (227 km<sup>2</sup>) of the region's seagrasses were exposed to a potential risk (combined risk categories II–IV). While the total areas in the Wet Tropics region exposed to a potential risk (combined risk categories II–IV) were greater than the average long-term areas (+4% changes), this did not impact the ecosystem exposure results. The total coral and seagrass areas exposed to the risk categories II–IV were similar to the long-term patterns (changes ≤1%). There was furthermore a decrease in seagrass area exposed to the highest potential risk categories (IV: -12%) toward the lowest risk category II (+9%) which is likely related to localised WT1 movements north of Hinchinbrook Island and may include the influence of different patterns/timing of sediment resuspension or some bottom influence as this is a shallow area. Only the inshore Wet Tropics waters, seagrass, and coral habitats were exposed to the highest categories of potential risk (III and IV), with the largest proportion located in the region's enclosed coastal waters. Mid-shelf and offshore Wet Tropics reefs and mid-shelf Wet Tropics seagrasses were largely exposed to no/very low risk (>76%).

### 6.3 Burdekin

The combined discharge from the Burdekin and Haughton basins was around 1.3 times the long-term median, following three years of above-median discharge (2.2 times higher in 2022–23, 1.2 times higher in 2021–22 and 1.8 times higher in 2020–21).

*Ambient water quality - Enclosed coastal, open coastal, and mid-shelf waters:*

- NO<sub>x</sub>, Secchi depth, and PO<sub>4</sub> did not meet water quality GVs in the Burdekin region (Table 6-4).
- Chl-*a*, PN and PP were meeting GVs at most sites in the Burdekin region.
- TSS was meeting GVs at all sites in the Burdekin region
- Over the period from 2015–2024, NO<sub>x</sub> showed a trend of improvement in the Burdekin region. Most other indicators showed a trend of stability (no net improvement or deterioration). PN was the only indicator which showed a trend of deterioration.
- Water Quality Index scores showed a long-term trend of deterioration from 2010–2018 followed by a period of stability. There was an improvement between 2021–2023 driven by improvements in PN, PP, and Chl-*a* but this trend has stabilised in 2023–24. For the 2023–24 water year, the annual condition Water Quality Index score was 'good'.

Table 6-4: Water quality indicator summary for the Burdekin region. Performance relative to guideline values is shown as: generally exceeding (✖) or meeting (✓) guideline values across all sites. The trend of the indicator (2015–present) is shown as: deteriorating (↘), improving (↗) or stable (→).

Water quality indicator	Burdekin
NO <sub>x</sub>	✖ ↗
PO <sub>4</sub>	✖ →
PN	✓ ↘
PP	✓ →
TSS	✓ →
Secchi depth	✖ →
Chl- <i>a</i>	✓ →

#### *Wet season and event water quality*

- In the Burdekin region, approximately 91% of the area was not exposed to a potential risk in keeping with long-term patterns (90%), and only 1% (or 694 km<sup>2</sup>) of the region was exposed to the highest risk category IV. Less than 1% (34 km<sup>2</sup>) of the region's

coral reefs and 87% (616 km<sup>2</sup>) of the region's seagrasses were exposed to a potential risk (combined risk categories II–IV). There was an increase in the seagrass areas exposed to potential risk category III in the Burdekin region, which was related to a higher frequency of Reef WT1 waters measured south of Magnetic Island. However, the total seagrass area exposed to a potential risk (II–IV) was similar to the long-term trends. Only the inshore Burdekin waters, seagrass, and coral habitats were exposed to the highest categories of potential risk (III and IV), with the largest proportion located in the region's enclosed coastal waters. Mid-shelf Burdekin seagrasses and mid-shelf and offshore Burdekin reefs were largely exposed to no/very low risk (>99%).

## 6.4 Mackay-Whitsunday

The combined discharge from the Proserpine, O'Connell, Pioneer, and Plane Basins was close to the long-term median values following a relatively wet period in 2022–23 (around 2 times the long-term median), but with three years of well below-median discharge prior to that.

*Ambient water quality - Enclosed coastal and open coastal waters:*

- PN, TSS, and Chl-*a* met GVs at most sites within the Mackay-Whitsunday region.
- NO<sub>x</sub>, PO<sub>4</sub>, PP, and Secchi depth did not meet water quality GVs in the Mackay-Whitsunday region (Table 6-5).
- Over the period from 2015–2024, most indicators showed a trend of improvement in the Mackay-Whitsunday region except for PO<sub>4</sub> and PN, which showed a trend for stability. No indicators showed a trend of deterioration.
- Water Quality Index scores showed a long-term trend of deterioration from 2007–2017. The trend then stabilised and started to improve in 2021, showing slight improvement each year from 2021–2023, driven by improvements in PN and Chl-*a*. For the 2023–24 water year, the long-term Water Quality Index has stabilised and was scored as 'moderate' and the annual condition Water Quality Index score was 'moderate'.

Table 6-5: Water quality indicator summary for Mackay-Whitsunday. Performance relative to guideline values is shown as: generally exceeding (✖) or meeting (✔) guideline values across all sites. The trend of the indicator (2015–present) is shown as: deteriorating (↘), improving (↗) or stable (→).

Water quality indicator	Mackay-Whitsunday
NO <sub>x</sub>	✖ ↗
PO <sub>4</sub>	✖ →
PN	✔ →
PP	✖ ↗
TSS	✔ ↗
Secchi depth	✖ ↗
Chl- <i>a</i>	✔ ↗

*Wet season and event water quality*

- In the Mackay-Whitsunday region, 91% of the area was not exposed to a potential risk in keeping with long-term patterns (87%) and only 1% (or 445 km<sup>2</sup>) of the region was exposed to the highest risk category IV. Approximately 4% (1223 km<sup>2</sup>) of the region's coral reefs and 82% (253 km<sup>2</sup>) of the region's seagrasses were exposed to a potential risk. Marine habitat areas exposed to respective risk categories II, III, or IV were overall similar to the long-term areas but there was a shift in the seagrass area exposed from lowest potential risk (II: -16%) to no/very low potential risk (+13%) and from the greatest risk category IV (-6%) to the potential risk category III (+6). Only inshore waters, seagrasses, and coral habitats were exposed to the highest categories of potential risk (III and IV), with the largest proportion located in the region's enclosed coastal waters. Mid-shelf and offshore Mackay-Whitsunday reefs were all exposed to no/very low risk (100%).

## 7 REFERENCES

- Álvarez-Romero JG, Devlin MJ, Teixeira da Silva E, Petus C, Ban N, Pressey RJ, Kool J, Roberts S, Cerdeira WA, Brodie J (2013). A novel approach to model exposure of coastal-marine ecosystems to riverine flood plumes based on remote sensing techniques. *Journal of Environmental Management* 119:194-207.
- Australian and Queensland governments, (2003). Reef Water Quality Protection Plan for catchments adjacent to the Great Barrier Reef World Heritage Area. The State of Queensland and Commonwealth of Australia. Queensland Department of Premier and Cabinet, Brisbane. 43 pp.
- Australian and Queensland governments, (2018a). Reef 2050 Water Quality Improvement Plan 2017-2022. Queensland government, Brisbane, Australia.
- Australian and Queensland governments, (2018b). Paddock to Reef Integrated Monitoring, Modelling and Reporting Program: Program Design 2018-2022. Queensland government, Brisbane, Australia.
- Bainbridge ZT, Wolanski E, Álvarez-Romero JG, Lewis SE, Brodie JE (2012). Fine sediment and nutrient dynamics related to particle size and floc formation in a Burdekin River flood plume, Australia. *Marine Pollution Bulletin, The Catchment to Reef Continuum: Case studies from the Great Barrier Reef* 65, 236–248. doi:10.1016/j.marpolbul.2012.01.043
- Brinkman R, Herzfeld M, Andrewartha J, Rizwi F, Steinberg C, Spagnol S (2011). Hydrodynamics at the whole of GBR scale. AIMS Final Project Report MTSRF Project 2.5i.1, June 2011. Report to Reef and Rainforest Research Centre. Australian Institute of Marine Science, Townsville. 42 pp.
- Brodie J, Waterhouse J, Schaffelke B, Furnas M, Maynard J, Collier C, Lewis S, Warne M, Fabricius K, Devlin M, McKenzie L, Yorkston H, Randall L, Bennett J, Brando V (2013). Scientific Consensus Statement. Chapter 3: Relative risks to the Great Barrier Reef from degraded water quality The State of Queensland. Published by the Reef Water Quality Protection Plan Secretariat, July 2013. <http://www.reefplan.qld.gov.au/about/scientific-consensus-statement/water-quality-risks.aspx>
- Brooks A, Spencer J, Olley J, Pietsch T, Borombovits D, Curwen G, Shellberg J, Howley C, Gleeson A, Simon A, Bankhead N, Klimetz D, Eslami-Endargoli L, Bourgeault A (2013) An empirically-based sediment budget for the Normanby Basin: sediment sources, sinks, and drivers on the Cape York Savannah, Australian Rivers Institute, Griffith University, Final report for the Australian Government Caring for Our Country - Reef Rescue Program.
- Bureau of Meteorology [BOM] (2011). Climate averages, average monthly wind velocity over Australia. Australian Government, Bureau of Meteorology, viewed 7 December 2017, [http://www.bom.gov.au/jsp/ncc/climate\\_averages/wind-velocity/index.jsp?period=jan#maps](http://www.bom.gov.au/jsp/ncc/climate_averages/wind-velocity/index.jsp?period=jan#maps)
- Bureau of Meteorology [BOM] (2022). Rainfall data, accessed online October 2022, <http://www.bom.gov.au/climate/data/>
- Bureau of Meteorology [BOM] (2017). eReefs Catchments: simulations, nowcasts, and forecasts of water quantity and quality flowing to the Great Barrier Reef. Final Report, eReefs Project Phase 3, Great Barrier Reef Foundation, Brisbane, QLD Australia.
- Cape York NRM and South Cape York Catchments (2016). Eastern Cape York Water Quality Improvement Plan. Cape York Natural Resource Management and South Cape York Catchments, Cooktown, Queensland, Australia.

- Carstensen J, Klais R, Cloern, JE (2015). Phytoplankton blooms in estuarine and coastal waters: Seasonal patterns and key species. *Estuarine, Coastal and Shelf Science* 162: 98–109.
- Commonwealth of Australia (2018). Reef 2050 Long-Term Sustainability Plan. <http://www.environment.gov.au/marine/gbr/publications/reef-2050-long-term-sustainability-plan>.
- De'ath G, Fabricius KE (2008). Water quality of the Great Barrier Reef: distributions, effects on reef biota and trigger values for the protection of ecosystem health. Final Report to the Great Barrier Reef Marine Park Authority. Australian Institute of Marine Science, Townsville. 104 pp.
- De'ath G, Fabricius KE (2010). Water quality as a regional driver of coral biodiversity and macroalgae on the Great Barrier Reef. *Ecological Applications* 20: 840–850.
- Department of Environment and Resource Management (2009). Queensland Water Quality Guidelines, Version 3. 167 p. Available at [www.derm.qld.gov.au](http://www.derm.qld.gov.au). ISBN 978-0-9806986-0-2.
- Department of Regional Development, Manufacturing and Water (DRMW) (2024). River discharge data. <https://water-monitoring.information.qld.gov.au/> Accessed October 2024.
- Devlin M, McKinna LW, Álvarez-Romero J, Petus C, Abott B, Harkness P, Brodie J (2012). Mapping the pollutants in surface plume waters in the Great Barrier Reef, Australia. *Marine Pollution Bulletin* 65: 224–235. doi:10.1016/j.marpolbul.2012.03.001
- Devlin MJ, Teixeira da Silva E, Petus C, Wenger A, Zeh D, Tracey D, Álvarez-Romero J, Brodie J (2013). Combining water quality and remote sensed data across spatial and temporal scales to measure wet season chlorophyll-a variability: Great Barrier Reef lagoon (Queensland, Australia). *Ecological Processes* 2.
- Devlin M, Petus C, Teixeira da Silva E, Tracey D, Wolff N, Waterhouse J, Brodie J (2015). Water quality and river plume monitoring in the Great Barrier Reef: An Overview of Methods Based on Ocean Colour Satellite Data. *Remote Sensing* 7: 12909–12941.
- Folkers A, Rohde K, Delaney K, Flet, I (2014). Mackay-Whitsunday Water Quality Improvement Plan 2014–2021. Reef Catchments and Australian Government. [http://reefcatchments.com.au/files/2015/06/WATER-QUALITY-IMPROVEMENT-PLAN-MACKAY-WHITSUNDAY-ISAAC-2014-2021\\_DRAFT.pdf](http://reefcatchments.com.au/files/2015/06/WATER-QUALITY-IMPROVEMENT-PLAN-MACKAY-WHITSUNDAY-ISAAC-2014-2021_DRAFT.pdf)
- Great Barrier Reef Marine Park Authority (2010). Water Quality Guidelines for the Great Barrier Reef Marine Park. Revised Edition 2010. Great Barrier Reef Marine Park Authority, Townsville. 100pp.
- Great Barrier Reef Marine Park Authority (2014). Great Barrier Reef Region Strategic Assessment. Great Barrier Reef Marine Park Authority, Townsville.
- Great Barrier Reef Marine Park Authority (2019). [Great Barrier Reef Outlook Report 2019](#), Great Barrier Reef Marine Park Authority, Townsville.
- Great Barrier Reef Marine Park Authority (2024). Great Barrier Reef Outlook Report 2024, Reef Authority, Townsville.
- Great Barrier Reef Marine Park Authority (2024), Great Barrier Reef Marine Monitoring Program Quality Assurance and Quality Control Manual 2022–23, Great Barrier Reef Marine Park Authority, Townsville, 189pp.
- Grill G, Lehner B, Thieme M. *et al.* (2019). Mapping the world's free-flowing rivers. *Nature* 569, 215–221. <https://doi.org/10.1038/s41586-019-1111-9>

- Gruber R, Waterhouse J, Logan M, Petus C, Tracey D, Lewis S, Howley C, Tonin H, Skuza M, Doyle J, Costello P, Davidson J, Gunn K, Wright M, Zagorskis I, Kroon F, Neilen A (2018). Marine Monitoring Program: Annual Report for inshore water quality monitoring 2016-2017. Report for the Great Barrier Reef Marine Park Authority, Great Barrier Reef Marine Park Authority, Townsville.
- Gruber R, Waterhouse J, Logan M, Petus C, Howley C, Lewis S, Tracey D, Langlois L, Tonin H, Skuza M, Costello P, Davidson J, Gunn K, Lefevre C, Moran D, Robson B, Shanahan M, Zagorskis I, Shellberg J, Neilen A (2020). Marine Monitoring Program: Annual Report for Inshore Water Quality Monitoring 2018–19. Report for the Great Barrier Reef Marine Park Authority, Great Barrier Reef Marine Park Authority, Townsville, 262pp.
- Gruber, R., Waterhouse, J., Petus, C., Howley, C., Lewis, S., Moran, D., James, C., Logan, M., Bove, U., Brady, B., Choukroun, S., Connellan, K., Davidson, J., Mellors, J., O'Callaghan, M., O'Dea, C., Shellberg, J., Tracey, D., Zagorskis, I., (2024). Great Barrier Reef Marine Monitoring Program Inshore Water Quality Monitoring: Annual Report 2022–23. Great Barrier Reef Marine Park Authority, Townsville. 298 pp.
- Howley C, (2020) Natural and Anthropogenic Drivers of Water Quality in the Normanby Basin and Princess Charlotte Bay, Northern Australia. PhD Dissertation. Griffith University, Australian Rivers Institute, School of Environment, Brisbane, Qld (199pp).
- Howley C, Blackman K, Bock E, Shellberg J, Jones B, (2022). Monitoring Fire Impacts on Ground Cover and Sediment Run-Off in a Northern GBR Catchment, Report to the Queensland Water Modelling Network, Department of Environment and Science. Howley C, Scobell L, Albert-Mitchell O, Shellberg J, Rosendale B (2024). Cyclone Jasper Environmental Impact Technical Investigation and Community Engagement Report for the Cape York Region: Annan & Bloomfield Catchments Focus. Report produced by Cape York Water Partnership for the Queensland Government Department of Environment, Science and Innovation.
- Jenks GF, Caspall FC (1971). Error on Choroplethic Maps: Definition, Measurement Reduction. *Annals of the Association of American Geographers* 61: 217-44.
- Kaserzon S, Shiels R, Elisei G, Paxman C, Li Y, Carswell C, Xia S, Prasad P, Gallen M, Reeks T, Thompson K, Taucare G, Marano K, Gorji SG, Mueller J (2024). Marine Monitoring Program Annual Report for Inshore Pesticide Monitoring: 2022–23. Report for the Great Barrier Reef Marine Park Authority, Great Barrier Reef Marine Park Authority, Townsville.
- Kaserzon S, Shiels R, Elisei G, Paxman C, Li Y, Carswell C, Xia S, Prasad P, Gallen M, Reeks T, Thompson K, Taucare G, Marano K, Gorji SG, Mueller J (in prep). Marine Monitoring Program Annual Report for Inshore Pesticide Monitoring: 2023–24. Report for the Great Barrier Reef Marine Park Authority, Great Barrier Reef Marine Park Authority, Townsville.
- King OC, Smith RA, Mann RM, Warne MSt.J (2017) Proposed aquatic ecosystem protection guideline values for pesticides commonly used in the Great Barrier Reef catchment area: Part 1 (amended) - 2,4-D, Ametryn, Diuron, Glyphosate, Hexazinone, Imazapic, Imidacloprid, Isoxaflutole, Metolachlor, Metribuzin, Metsulfuron-methyl, Simazine, Tebuthiuron. Department of Environment and Science. Brisbane, Queensland, Australia. 296 pp.
- Kuhnert PM, Liu Y, Henderson B, Dambacher J, Lawrence E, Kroon FJ (2015). Review of the Marine Monitoring Program (MMP). Final Report for the Great Barrier Reef Marine Park Authority, CSIRO, Australia: 278
- Lewis S, Brodie J, Endo G, Lough J, Furnas M, Bainbridge Z (2014). Synthesizing historical land use change, fertiliser and pesticide usage and pollutant load data in the regulated catchments to quantify baseline and changing Loads exported to the Great Barrier Reef.

- Centre for Tropical Water & Aquatic Ecosystem Research (TropWATER) Technical Report 14/20, James Cook University, Townsville, 105 pp.
- Lloyd-Jones LR, Kuhnert PM, Lawrence E, Lewis SE, Waterhouse J, Gruber RK, Kroon FJ (2022). Sampling re-design increases power to detect change in the Great Barrier Reef's inshore water quality. *PLOS ONE* 17(7): e0271930. <https://doi.org/10.1371/journal.pone.0271930>
- Margvelashvili N, Andrewartha J, Baird M, Herzfeld M, Jones E, Mongin M, Rizwi F, Robson BJ, Skerratt J, Wild-Allen K, Steven A (2018). Simulated fate of catchment-derived sediment on the Great Barrier Reef shelf. *Marine Pollution Bulletin* 135: 954-962.
- McCloskey GL, Baheerathan R, Dougall C, Ellis R, Bennett FR, Waters D, Darr S, Fentie B, Hateley LR, Askildsen M (2021). Modelled estimates of fine sediment and particulate nutrients delivered from the Great Barrier Reef catchments. *Marine Pollution Bulletin* 165: 112163 <https://doi.org/10.1016/j.marpolbul.2021.112163>.
- McKenzie LJ, Collier CJ, Langlois LA, Brien H and Yoshida RL (2024). Marine Monitoring Program: Annual Report for Inshore Seagrass Monitoring 2022–23. Report for the Great Barrier Reef Marine Park Authority, Great Barrier Reef Marine Park Authority, Townsville. 178pp.
- Moran D, Robson B, Gruber R, Waterhouse J, Logan M, Petus C, Howley C, Lewis S, James C, Tracey D, Mellors J, O'Callaghan M, Bove U, Davidson J, Glasson K, Jaworski S, Lefevre C, Nordborg M, Vasile R, Zagorskis I, Shellberg J. (2023). Marine Monitoring Program: Annual Report for Inshore Water Quality Monitoring 2021–22. Report for the Great Barrier Reef Marine Park Authority, Great Barrier Reef Marine Park Authority, Townsville.
- NQ Dry Tropics (2016). Burdekin Region Water Quality Improvement Plan 2016, NQ Dry Tropics, Townsville. <https://www.nqdrytropics.com.au/wqip2016/>
- Novoa S, Wernand MR, Van der Woerd HJ, (2013). The Forel-Ule scale revisited spectrally: preparation protocol, transmission measurements and chromaticity. *Journal of the European Optical Society-Rapid Publications* 8, 13057.
- Petus C, Devlin M, Thompson A, McKenzie L, Teixeira da Silva E, Collier C, Tracey D, Martin K (2016). Estimating the exposure of coral reefs and seagrass meadows to land-sourced contaminants in river flood plumes of the Great Barrier Reef: validating a simple Satellite Risk Framework with Environmental Data. *Remote Sensing* 8: 210.
- Petus C, Devlin M, da Silva E, Lewis S, Waterhouse J, Wenger A, Bainbridge Z, Tracey D (2018) Defining wet season water quality target concentrations for ecosystem conservation using empirical light attenuation models: a case study in the Great Barrier Reef (Australia). *Journal of Environmental Management* 213: 1-16.
- Petus C, Waterhouse J, Lewis S, Vacher M, Tracey D, Devlin M. (2019). A flood of information: Using Sentinel-3 water colour products to assure continuity in the monitoring of water quality trends in the Great Barrier Reef (Australia). *Journal of Environmental Management* 248: 109255.
- Pinheiro JC, Bates DM (2000). *Mixed-effects models in S and S-PLUS*, Statistics and Computing Series, Springer-Verlag, New York, NY.
- Puignou Lopez O, Lewis S, James C, Davis A, Mackay S. (in review). Hydrology of the Great Barrier Reef catchment area along a latitudinal gradient: Upscaling discharge to reflect catchment inputs. *Journal of Hydrology*.
- Queensland Land Use Mapping Program [QLUMP] (2015). Land use mapping for the Cape York NRM region, prepared by DNRM.

- R Core Team (2022). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL: <https://www.R-project.org/>
- Schaffelke B, Carleton J, Skuza M, Zagorskis I, Furnas MJ (2012). Water quality in the inshore Great Barrier Reef lagoon: Implications for long-term monitoring and management. *Marine Pollution Bulletin* 65:249-260. DOI: 10.1016/j.marpolbul.2011.10.031
- Shellberg J, Brooks A (2013). Alluvial Gully Prevention and Rehabilitation Options for Reducing Sediment Loads in the Normanby Catchment and Northern Australia. Australian Rivers Institute, Griffith University, Final report for the Australian Government Caring for Our Country - Reef Rescue Program.
- Shellberg JG, Spencer J, Brooks AP, Pietsch TJ (2016). Degradation of the Mitchell River fluvial megafan by alluvial gully erosion increased by post-European land use change, Queensland, Australia. *Geomorphology*, 266, 105-120.
- Skerratt JH, Mongin M, Baird ME, Wild-Allen KA, Robson BJ, Schaffelke B, Davies CH, Richardson AJ, Margvelashvili N, Soja-Wozniak M, Steven ADL (2019). Simulated nutrient and plankton dynamics in the Great Barrier Reef (2011–2016). *Journal of Marine Systems*, 192, 51-74. <http://dx.doi.org/10.1016/j.jmarsys.2018.12.006>
- Spencer J, Brooks A, Curwen G, Tews K (2016). A Disturbance Index Approach for Assessing Water Quality Threats in Eastern Cape York. A report to South Cape York Catchments and Cape York NRM for the Cape York Water Quality Improvement Plan, by the Australian Rivers Institute, Griffith University, 42 pp
- State of Queensland (2020). Environmental Protection (Water and Wetland Biodiversity) Policy 2019: Environmental Values and Water Quality Objectives, Eastern Cape York Basins <https://environment.des.qld.gov.au/management/water/policy/cape-york-eastern-basins>
- Steven AD, Baird ME, Brinkman R, Ca, NJ, Cox SJ, Herzfeld M, Hodge J, Jones E, King E, Margvelashvili N, Robillot C, (2019). eReefs: An operational information system for managing the Great Barrier Reef. *Journal of Operational Oceanography*, pp.1-17.
- Terrain NRM (2015). Wet Tropics Water Quality Improvement Plan 2015-2020. Terrain NRM, Innisfail. <https://www.wettropicsplan.org.au/regional-themes/water/wqip/>
- Thai P, Paxman C, Prasad P, Elisei G, Reeks T, Eaglesham G, Yeh R, Tracey D, Grant S, Mueller J (2020). Marine Monitoring Program: Annual report for inshore pesticide monitoring 2018-2019. Report for the Great Barrier Reef Marine Park Authority, Great Barrier Reef Marine Park Authority, Townsville, 69pp.
- Thompson A, Davidson J, Logan M, Thompson C (2024). Marine Monitoring Program Annual Report for Inshore Coral Reef Monitoring: 2022–23. Report for the Great Barrier Reef Marine Park Authority, Great Barrier Reef Marine Park Authority, Townsville.149 pp.
- Townsend, SA, Douglas, MM (2000). The effect of three fire regimes on stream water quality, water yield and export coefficients in a tropical savanna (northern Australia). *Journal of Hydrology*, 229(3-4): 118-137.
- Van der Woerd JH, Wernand RM (2018). Hue-angle product for low to medium spatial resolution optical satellite sensors. *Remote Sensing* 10, 180.
- Van der Woerd JH, Wernand RM, Peters M, Brockmann C (2016). True colour analysis of natural waters with SeaWiFS, MODIS, MERIS and OLCI by SNAP, Ocean Optics conference. At Victoria BC Canada XXIII.
- Water Quality & Investigations (2024). Great Barrier Reef Catchment Loads Monitoring Program: Loads and yields for sediment and nutrients, and Pesticide Risk Metric results

- (2021–2022) for rivers that discharge to the Great Barrier Reef, Department of Environment, Science and Innovation, Brisbane, Australia.
- Waterhouse J, Brodie J, Tracey D, Smith R, Vandergragt M, Collier C, Petus C, Baird M, Kroon F, Mann R, Sutcliffe T, Waters D, Adame F (2017). Scientific Consensus Statement 2017: A synthesis of the science of land-based water quality impacts on the Great Barrier Reef, Chapter 3: The risk from anthropogenic pollutants to Great Barrier Reef coastal and marine ecosystems. State of Queensland, 2017.
- Waterhouse J, Burton J, Garzon-Garcia A, Lewis S, Brodie J, Bainbridge Z, Robson R, Burford MA, Gruber RK, Dougall C (2018). Synthesis of knowledge and concepts - Bioavailable Nutrients: Sources, delivery and impacts in the Great Barrier Reef, July 2018. Supporting Concept Paper for the Bioavailable Nutrients Workshop, 15 March 2018. Reef and Rainforest Research Centre, 84pp.
- Waterhouse J, Gruber R, Logan M, Petus C, Howley C, Lewis S, Tracey D, James C, Mellors J, Tonin H, Skuza M, Costello P, Davidson J, Gunn K, Lefevre C, Moran D, Robson B, Shanahan M, Zagorskis I, Shellberg J (2021). Marine Monitoring Program: Annual Report for Inshore Water Quality Monitoring 2019–20. Report for the Great Barrier Reef Marine Park Authority, Great Barrier Reef Marine Park Authority, Townsville.
- Waterhouse J, Petus C, Lewis S, James C, O'Callaghan M, Tracey D, Doyle J, Patel F, Uthicke S (2023). Pilot study to investigate options for water quality sampling as part of the Reef Trust Partnership Crown of Thorns Starfish Control Program, Final Report July 2023. Report for the Reef Trust Partnership. TropWATER Report No. 23/50. 92 pp. James Cook University, Townsville.
- Waterhouse J, Pineda M-C, Sambrook K, Newlands M, McKenzie L, Davis A, Pearson R, Fabricius K, Lewis S, Uthicke S, Bainbridge Z, Collier C, Adame F, Prosser I, Wilkinson S, Bartley R, Brooks A, Robson B, Diaz-Pulido G, Reyes C, Caballes C, Burford M, Thorburn P, Weber T, Waltham N, Star M, Negri A, Warne M St J, Templeman S, Silburn M, Chariton A, Coggan A, Murray-Prior R, Schultz T, Espinoza T, Burns C, Gordon I, Devlin M (2024). 2022 Scientific Consensus Statement: Conclusions. In Waterhouse J, Pineda M-C, Sambrook K (Eds) 2022 Scientific Consensus Statement on land-based impacts on Great Barrier Reef water quality and ecosystem condition. Published by C2O Consulting on behalf of the Australian Government's Department of Climate Change, Energy, the Environment and Water (DCCEEW) and the Queensland Government's Office of the Great Barrier Reef and World Heritage (OGBRWH).
- Wells SC, Cole SJ, Moore RJ, Black KB, Khan U, Hapuarachchi P, Gamage N, Hasan M, MacDonald A, Bari M, Tuteja NK (2018). Forecasting the water flows draining to the Great Barrier Reef using the G2G Distributed Hydrological Model. Technical Report (Contract No. 112-2015-16), Centre for Ecology & Hydrology Wallingford, OX10 8BB, UK.
- Wernand MR, Hommersom A, Van der Woerd HJ (2012). MERIS-based ocean colour classification with the discrete Forel–Ule scale. *Ocean Science Discussions* 9, 2817–2849.
- Wernand MR, Van der Woerd HJ, Gieskes WWC (2013). Trends in ocean colour and chlorophyll concentration from 1889 to present. *PLoS One*, 0063766.
- Wijffels S, Beggs H, Griffin C, Middleton J, Cahill M, King E, Jones E, Feng M, Benthuyssen J, Steinberg C, Sutton P (2018). A fine spatial-scale sea surface temperature atlas of the Australian regional seas (SSTAARS): Seasonal variability and trends around Australasia and New Zealand revisited. *Journal of Marine Systems*. 187. 10.1016/j.jmarsys.2018.07.005.
- Wood SN (2006). Generalized additive models: An introduction with R. Chapman & Hall/CRC Publisher, City.

Wood SN (2011). Fast stable restricted maximum likelihood and marginal likelihood estimation of semiparametric generalized linear models. *Journal of the Royal Statistical Society (B)* 73: 3-36.

## APPENDIX A: WATER QUALITY SITE LOCATIONS AND FREQUENCY OF MONITORING

Table A-1: Description of the water quality sites sampled by AIMS, JCU and CYWP during 2023–24. The proposed number of visits is shown in black text, while the actual number of visits is shown in parentheses in red text. Actual visits can differ from proposed due to poor weather limiting site access.

Site location	Logger Deployment		Ambient sampling at fixed sites: proposed (actual)		Event-based sampling
NRM region	Turbidity and chlorophyll	Salinity	Number of times site is visited/year by AIMS	Number of times site is visited/year by JCU/ CYWP	Additional surface-sampling/year by JCU/ CYWP
<b>Cape York</b>					
<b><i>Normanby-Kennedy transect</i></b>					
Kennedy mouth (KR01)					2
Kennedy inshore (KR02)					2
Cliff Islands (CI01)				4 (Sampling 2 depths) (3)	1
Bizant River mouth (BR01)					
Normanby River mouth (NR01)					1
Normanby inshore (NR02)				4 (Sampling 2 depths) (4)	1
NR03 (NR03)				4 (Sampling 2 depths) (4)	1
NR04 (NR04)				4 (Sampling 2 depths) (4)	1
NR05 (NR05)				4 (Sampling 2 depths) (4)	
Corbett Reef (NR06)				4 (Sampling 2 depths) (4)	1
<b><i>Pascoe transect</i></b>					
Pascoe mouth north (PRN01)					
Pascoe mouth south (PRS01)				5 (Sampling 2 depths) (5)	1
PRN02 (PRN02)				5 (Sampling 2 depths) (5)	1
PRN03 (PRN03)					1
PRN04 (PRN04)				5 (Sampling 2 depths) (0)	1
Eel Reef (PRN05)					
Eel Reef North (PRN06)					
PRS2.5 (PRS2.5)				5 (Sampling 2 depths) (5)	1
Middle Reef (PRBB)				5 (Sampling 2 depths) (5)	1
Eel Reef South (PR-S05)				5 (Sampling 2 depths) (5)	1

Site location	Logger Deployment		Ambient sampling at fixed sites: proposed (actual)		Event-based sampling
NRM region	Turbidity and chlorophyll	Salinity	Number of times site is visited/year by AIMS	Number of times site is visited/year by JCU/ CYWP	Additional surface-sampling/year by JCU/ CYWP
<b>Annan and Endeavour transect</b>					
Annan mouth (AR01)					2
Walker Bay (AR02b)				5 (Sampling 2 depths) (5)	2
Dawson Reef (AR03b)	✓			5 (Sampling 2 depths) (5)	2
Endeavour mouth (ER01)					1
Endeavour north shore (ER02b)				5 (Sampling 2 depths) (5)	1
Endeavour offshore (ER03)				5 (Sampling 2 depths) (5)	1
Egret and Boulder Reef (AE04)				5 (Sampling 2 depths) (5)	1
Forrester Reef (ER06)	✓				
<b>Stewart transect</b>					
Stewart mouth (SR01)					
SR02 (SR02)				5 (Sampling 2 depths) (5)	
SR03 (SR03)				5 (Sampling 2 depths) (5)	
Burkitt Island (SR04)				5 (Sampling 2 depths) (5)	
Hannah Island (SR05)				5 (Sampling 2 depths) (5)	
<b>Wet Tropics</b>					
<b>Cairns Long-term transect</b>					
Cape Tribulation (C1)*			3 (Sampling 2 depths) (3)		1
Port Douglas (C4)*			3 (Sampling 2 depths) (3)		1
Double Island (C5)*			3 (Sampling 2 depths) (3)		1
Yorkey's Knob (C6)*			3 (Sampling 2 depths) (3)		1
Fairlead Buoy (C8)*			3 (Sampling 2 depths) (3)		1
Green Island (C11)*			3 (Sampling 2 depths) (3)		1
<b>Russell-Mulgrave Focus Region</b>					
Fitzroy Island West (RM1)	✓		5 (Sampling 2 depths) (5)		2

Site location	Logger Deployment		Ambient sampling at fixed sites: proposed (actual)		Event-based sampling
NRM region	Turbidity and chlorophyll	Salinity	Number of times site is visited/year by AIMS	Number of times site is visited/year by JCU/ CYWP	Additional surface-sampling/year by JCU/ CYWP
RM2 (RM2)					1
RM3 (RM3)			5 (Sampling 2 depths) (5)	5 (Sampling 2 depths) (5)	1
RM4 (RM4)					1
High Island East (RM5)					1
Normanby Island (RM6)					1
Frankland Group West (Russell Island) (RM7)*	✓		5 (Sampling 2 depths) (5)	5 (Sampling 2 depths) (5)	1
High Island West (RM8)*	✓	✓	5 (Sampling 2 depths) (5)	5 (Sampling 2 depths) (5)	1
Palmer Point (RM9)					1
Russell-Mulgrave River mouth mooring (RM10)	✓	✓	5 (Sampling 2 depths) (5)	5 (Sampling 2 depths) (5)	1
Russell-Mulgrave River mouth (RM11)					1
Russell-Mulgrave junction [River] (RM12)					1
<b>Tully Focus Region</b>					
King Reef (TUL1)					
East Clump Point (TUL2)			5 (Sampling 2 depths) (5)	5 (Sampling 2 depths) (5)	2
Dunk Island North (TUL3)*	✓	✓	5 (Sampling 2 depths) (5)	5 (Sampling 2 depths) (5)	2
South Mission Beach (TUL4)					2
Dunk Island South East (TUL5)			5 (Sampling 2 depths) (5)	5 (Sampling 2 depths) (5)	2
Between Tam O'Shanter and Timana (TUL6)			5 (Sampling 2 depths) (5)	5 (Sampling 2 depths) (5)	2
Hull River mouth (TUL7)					2
Bedarra Island (TUL8)			5 (Sampling 2 depths) (5)	5 (Sampling 2 depths) (5)	2
Triplets (TUL9)					2
Tully River mouth mooring (TUL10)	✓	✓	5 (Sampling 2 depths) (5)	5 (Sampling 2 depths) (5)	2
Tully River (TUL11)					1
<b>Burdekin</b>					

Site location	Logger Deployment		Ambient sampling at fixed sites: proposed (actual)		Event-based sampling
NRM region	Turbidity and chlorophyll	Salinity	Number of times site is visited/year by AIMS	Number of times site is visited/year by JCU/ CYWP	Additional surface-sampling/year by JCU/ CYWP
<b>Burdekin Focus Region</b>					
Pelorus and Orpheus Island West (BUR1)*	✓		4 (Sampling 2 depths) (4)	5 (Sampling 2 depths) (5)	
Pandora Reef (BUR2)*	✓		4 (Sampling 2 depths) (4)	5 (Sampling 2 depths) (5)	
Cordelia Rocks (BUR3)					
Magnetic Island (Geoffrey Bay) (BUR4)*	✓		4 (Sampling 2 depths) (4)	5 (Sampling 2 depths) (5)	
Inner Cleveland Bay (BUR5)					
Cape Cleveland (BUR6)					
Haughton 2 (BUR7)			4 (Sampling 2 depths) (4)	5 (Sampling 2 depths) (4)	
Haughton River mouth (BUR8)				0 (Sampling 2 depths) (1)	
Barratta Creek (BUR9)					
Yongala IMOS NRS (BUR10)	✓	✓	4 (Sampling 2 depths) (4)		
Cape Bowling Green (BUR11)					
Plantation Creek (BUR12)					
Burdekin River mouth mooring (BUR13)	✓	✓	4 (Sampling 2 depths) (4)	5 (Sampling 2 depths) (5)	
Burdekin Mouth 2 (BUR14)					
Burdekin Mouth 3 (BUR14)					
<b>Mackay-Whitsunday</b>					
<b>Whitsunday focus Region</b>					
Double Cone Island (WHI1)*	✓		5 (Sampling 2 depths) (5)		
Hook Island W (WHI2)					
North Molle Island (WHI3)					

Site location	Logger Deployment		Ambient sampling at fixed sites: proposed (actual)		Event-based sampling
NRM region	Turbidity and chlorophyll	Salinity	Number of times site is visited/year by AIMS	Number of times site is visited/year by JCU/ CYWP	Additional surface-sampling/year by JCU/ CYWP
Pine Island (WHI4)*	✓	✓	5 (Sampling 2 depths) (5)		
Seaforth Island (WHI5)	✓		5 (Sampling 2 depths) (5)		
OConnell River mouth (WHI6)	✓	✓	5 (Sampling 2 depths) (5)		
Repulse Islands dive mooring (WHI7)			5 (Sampling 2 depths) (5)		
Rabbit Island NE (WHI8)					
Brampton Island (WHI9)					
Sand Bay (WHI10)					
Pioneer River mouth (WHI11)					

\*Sites which were part of the MMP sampling design from 2005–2015.

## APPENDIX B: WATER QUALITY MONITORING METHODS

### B-1 Comparison with Reef Water Quality guideline values

The Water Quality Guidelines provide a useful framework to interpret the water quality measurements obtained through the MMP. Table B-1 gives a summary of the guideline values (GVs) for water quality variables in four cross-shelf water bodies (Great Barrier Reef Marine Park Authority, 2010).

At present, the Water Quality Guidelines do not define GV's for dissolved inorganic nutrients (nitrate and phosphate) in the Reef lagoon as these nutrients are rapidly cycled through uptake and release by biota and are variable on small spatial and temporal scales (Furnas *et al.*, 2005, 2011). Due to this high variability, their concentrations did not show as clear spatial patterns or correlations with coral reef attributes as the other water quality parameters that were included in The Guidelines and are considered to be more representative of nutrient availability integrated over time (De'ath and Fabricius, 2010). However, the Queensland Water Quality Guidelines (Department of Environment and Resource Management [DERM], 2009) identify GV's for dissolved inorganic nutrients in marine water bodies. Guideline values for dissolved inorganic nutrients and turbidity (in enclosed coastal waters) were drawn from Queensland Water Quality Guidelines or provided by the Reef Authority. Site-specific GV's for all water quality variables are shown in Appendix C Table C-9.

Table B-1: Guidelines values for four cross-shelf water bodies, provided by the Reef Authority. Values come from the Water Quality Guidelines for the Great Barrier Reef Marine Park (Great Barrier Reef Marine Park Authority, 2010). Guidelines for some values come from other sources, as indicated below.

Parameter	Unit	Enclosed coastal		Open coastal		Mid-shelf		Offshore	
		Wet Tropics	Central Coast	Wet Tropics	Central Coast	Wet Tropics	Central Coast	Wet Tropics	Central Coast
Chlorophyll <i>a</i> *	µg L <sup>-1</sup>	2.0	2.0	0.45	0.45	0.45	0.45	0.40	0.40
Particulate nitrogen*	µg L <sup>-1</sup>	n/a	n/a	20.0	20.0	20.0	20.0	17.0	17.0
Particulate phosphorus*	µg L <sup>-1</sup>	n/a	n/a	2.8	2.8	2.8	2.8	1.9	1.9
Suspended solids*	mg L <sup>-1</sup>	5.0	15.0	2.0	2.0	2.0	2.0	0.7	0.7
Turbidity	NTU	10.0 <sup>QLD</sup>	6.0 <sup>QLD</sup>	1.5**	1.5**	1.5**	1.5**	<1 <sup>QLD</sup>	<1 <sup>QLD</sup>
Secchi depth	m	1.0	1.5	10.0	10.0	10.0	10.0	17.0	17.0
NO <sub>x</sub>	µg L <sup>-1</sup>	10.0 <sup>QLD</sup>	3.0 <sup>QLD</sup>	0.35***	0.35***	0.35***	0.35***	2.0 <sup>QLD</sup>	2.0 <sup>QLD</sup>
PO <sub>4</sub>	µg L <sup>-1</sup>	5.0 <sup>QLD</sup>	6.0 <sup>QLD</sup>	4.0 <sup>QLD</sup>	6.0 <sup>QLD</sup>	4.0 <sup>QLD</sup>	6.0 <sup>QLD</sup>	4.0 <sup>QLD</sup>	5.0 <sup>QLD</sup>

<sup>QLD</sup> Values are Queensland Water Quality Guidelines (Department of Environment and Resource Management [DERM], 2009). Please note these are 80<sup>th</sup> percentile guidelines.

\* Seasonal adjustments to these parameters are used to produce seasonal (wet and dry) guidelines for producing satellite exposure maps (Table B-4).

\*\* Turbidity trigger value (1.5 NTU) was derived for the MMP reporting by transforming the suspended solids GV's (2 mg L<sup>-1</sup>) using an equation based on a comparison between direct water samples and instrumental turbidity readings (see QA/QC Reports and Schaffelke *et al.*, 2009).

\*\*\* NO<sub>x</sub> GV's for open coastal and mid-shelf sites are provided by the Reef Authority.

### B-2 Calculation of the Water Quality Index

In the Great Barrier Reef Report Cards published prior to 2016, water quality assessments were based on the MMP broad-scale monitoring using ocean colour remote sensing imagery that covers a larger area than the fixed sampling locations reported here (Brando *et al.*, 2011). However, the current design of the MMP focuses on interpreting trends in site-specific water quality within key focus regions.

The Water Quality Index (WQ Index) is an interpretation tool developed by AIMS to visualise trends in the suite of water quality variables measured and to compare monitored water quality to existing Water Quality Guidelines (Department of Environment and Resource Management, 2009; Great Barrier Reef Marine Park Authority, 2010). The WQ Index uses a set of five key indicators:

- Water clarity,
- Chl-*a* concentrations,
- PN concentrations,
- PP concentrations, and
- NO<sub>x</sub> concentrations.

These five indicators are a subset of the comprehensive suite of water quality variables measured in the MMP inshore water quality program. They have been selected because GVs are available for these measures, and they can be considered as relatively robust indicators that integrate a number of bio-physical processes in the coastal ocean.

TSS concentration, turbidity, and Secchi depth are indicators of the clarity of the water, which is influenced by a number of factors, including wind, waves, tides, and river inputs of particulate material. Chl-*a* concentration is widely used as a proxy for phytoplankton biomass as a measure of the productivity of a system or its eutrophication status and is used to indicate nutrient availability (Brodie *et al.*, 2007). Particulate nutrients (PN, PP) are an indicator of nutrient stocks in the water column (predominantly bound in phytoplankton and other organic particles as well as adsorbed to fine sediment particles) but are less affected by small-scale variability in space and time than dissolved nutrients (Furnas *et al.*, 2005, 2011). Nitrate is included as an indicator of dissolved nutrient concentrations in the coastal zone, which tend to be rapidly used by phytoplankton. Guideline values for NO<sub>x</sub> were provided by the Reef Authority as available NO<sub>x</sub> GVs from the Queensland Water Quality Guidelines (Department of Environment and Resource Management [DERM], 2009) are the 80<sup>th</sup> percentiles, which are considered to be high and not representative of values normally found in the Reef lagoon.

The WQ Index is calculated using two different methods due to changes in the MMP design that occurred in 2015, as well as concerns that the Index was not responsive to changes in environmental pressures of each year. The changes in design included increased number of sites, increased sampling frequency and a higher sampling frequency during December to April to better represent wet season variability. Thus, statistical comparisons between MMP data from 2005–15 and data from 2015–onwards must account for these changes. The two versions of the WQ Index have different purposes.

**Long-term trend:** This version of the WQ Index is based on the pre-2015 MMP sampling design and uses only the original sites (located in the open coastal water body) and three sampling dates per year. This sampling design had low temporal and spatial resolution and was aimed at detecting Long-term trends in inshore water quality. To compensate for less frequent sampling, four-year running means are used to reduce the effect of sampling date on the Index. Monitoring data are compared against broad water body GVs that do not include wet and dry season GVs (Table B-1). Steps in the calculation of this version of the WQ Index are:

1. Calculate four-year mean values for each of the seven indicators (i.e., all values from 2005–08, 2006–09, 2007–10, etc).

2. Calculate the proportional deviations (ratios) of these running mean values (V) from the associated guideline value (GV) (Table B-1) as the difference of binary logarithms of values and guidelines:

$$\text{Ratio} = \log_2(GV) - \log_2(V)$$

for values where exceeding the GV constitutes a “fail”. For values where exceeding the GV constitutes a “pass” (i.e., Secchi depth), the right side of this equation would be reversed.

3. Binary logarithm transformations are useful for exploring data on powers of 2 scales, and thus are ideal for generating ratios of two numbers in a manner that will be symmetrical around 0. Ratios of 1 and -1 signify a doubling and a halving, respectively, compared to the guideline. Hence, a ratio of 0 indicates a running mean that is the same as its GV, ratios <0 signify running means that exceeded the GV and ratios >0 signify running means that complied with the GV.
4. Ratios exceeding 1 or -1 (more than twice or half the GV) are capped at 1 to bind the WQ Index scales to the region -1 to 1.
5. A combined water clarity ratio is generated by averaging the ratios of TSS and turbidity from loggers (where available).
6. The WQ Index for each site per four-year period is calculated by averaging the ratios of PP, PN, NO<sub>x</sub>, Chl-a, and water clarity.
7. In accordance with other Great Barrier Reef Report Card indicators, the WQ Index scores (ranging from -1 to 1) are converted to a ‘traffic light’ colour scheme for reporting whereby:
  - < -2/3 to -1 equates to ‘very poor’ and is coloured red
  - < -1/3 to -2/3 equates to ‘poor’ and is coloured orange
  - < 0 to -1/3 equates to ‘moderate’ and is coloured yellow
  - 0 to 0.5 equates to ‘good’ and is coloured light green
  - 0.5 to 1 equates to ‘very good’ and is coloured dark green.
8. For the focus region summaries, the Index scores of all sampling locations within a focus region (for example, all sites in the Tully focus region) are averaged (median) and converted into the colour scheme as above. For regional summaries, the Index scores of all sampling locations within a region (for example, all sites in the Wet Tropics region) are averaged (median) and converted as above.

**Annual condition:** This version of the WQ Index is based on the post-2015 MMP sampling design and uses all samples from open coastal and mid-shelf water bodies each year. (Note that the WQ Index in reports prior to the 2018–19 report included enclosed coastal sites, see below). Due to high spatial and temporal sampling, a running mean is not used. Monitoring data are compared against site-specific GVs that include wet and dry season GVs (Table C-9). Steps in the calculation of this version of the WQ Index are:

1. For each of the seven indicators, the annual, wet, and dry season (aggregations) means and medians (statistic) are calculated per year.
2. Guidelines from the Reef Authority are consulted to select the appropriate aggregation (annual, wet, or dry season) and statistic (mean or median) for each site and indicator (Table C-9).
3. Calculate the proportional deviations (ratios) of these aggregation statistics from the associated GVs as the difference of base 2 logarithms of values and GVs:

$$\text{Ratio} = \log_2(GV) - \log_2(V)$$

for values where exceeding the GV constitutes a “fail”. For values where exceeding the GV constitutes a “pass” (i.e., Secchi depth), the right side of this equation would be reversed.

4. Ratios exceeding 1 or -1 (more than twice or half the GV) are capped at 1 to bind the WQ Index scales to the region -1 to 1.
5. Ratios of several indicators are combined to create a hierarchical structure. Three groups were created by averaging ratios as follows:
  - water clarity (average of Secchi depth, TSS, and turbidity from loggers ratios),
  - productivity (average of Chl-a and NO<sub>x</sub> ratios), and
  - particulate nutrients (average of PN and PP ratios).
6. The WQ Index for each site is calculated by averaging the ratios of water clarity, productivity, and particulate nutrients.
7. In accordance with other Reef Report Card indicators, the WQ Index scores (ranging from -1 to 1) are converted to a ‘traffic light’ colour scheme for reporting whereby:
  - < -2/3 to -1 equates to ‘very poor’ and is coloured red
  - < -1/3 to -2/3 equates to ‘poor’ and is coloured orange
  - < 0 to -1/3 equates to ‘moderate’ and is coloured yellow
  - 0 to 0.5 equates to ‘good’ and is coloured light green
  - 0.5 to 1 equates to ‘very good’ and is coloured dark green.
8. For the focus region summaries, the Index scores of all sampling locations within a focus region (for example, all sites in the Tully focus region) are averaged and converted into the colour scheme as above. For regional summaries, the Index scores of all sampling locations within a region (for example, all sites in the Wet Tropics region) are averaged and converted as above.
9. As of the 2018–19 report, this version of the Index now includes error bars, which propagate error in the Index via bootstrapping. Aggregation uncertainty is propagated through the spatial (site > focus region > region) and measure (measure > sub-indicator > indicator) hierarchies by repeatedly re-sampling (100 times with replacement) and aggregating bootstrapping. Each aggregation yields 100 estimates of each mean, and thus error bars represent the 95% quantile confidence intervals.

The annual condition version of the WQ Index has only been calculated since 2016 and is subject to future revision and refinement.

### B-3 Monitoring of Reef water quality trends using remote sensing data

Remote sensing imagery is a useful assessment tool in the monitoring of turbid water masses and river flood plumes (hereafter river plumes) in the Reef lagoon. Ocean colour imagery provides synoptic-scale information regarding the movement, frequency of occurrence and composition of turbid waters in the Reef lagoon. Combined with *in situ* water quality sampling and modelling, the use of remote sensing is a valid and practical way to estimate wet season marine conditions as well as the extent and frequency of water type exposure on Reef ecosystems, including river plumes and resuspension events.

Until 2020, marine areas exposed to wet season water types were mapped using MODIS true colour images and a wet season (WS) water colour classification method, composed of 6 colours. This method is extensively presented in Álvarez-Romero *et al.* (2013) and used in, for example, Devlin *et al.* (2013) and Petus *et al.* (2014b, 2016, 2018 and 2019). Since 2020–21, the use of Sentinel-3 Ocean Land Colour Instrument (OLCI) satellite imagery and another colour scale (the Forel-Ule (FU) colour scale) was adopted, as the quality of the MODIS images was declining (Petus *et al.*, 2019).

The FU colour scale is an historical colour scale standard to determine the colour and classifies worldwide bodies of water (Novoa *et al.*, 2013). It is composed of 22 colours; going from indigo blue to cola brown, and is applicable for all natural waters (inland, estuarine, inshore and offshore) and all environmental conditions, including wet and dry season conditions (Wernand *et al.*, 2012, 2013; Van der Woerd *et al.*, 2016; Van der Woerd and Wernand, 2018). MODIS-Aqua WS and Sentinel-3 FU colour class maps showed very similar patterns over the 2017–18 wet season in a case study focusing on Wet Tropics and Burdekin regions of the Reef (Petus *et al.*, 2019 and Figure 2-2b and Table 2-2). This suggested that Sentinel-3 FU water colour products can be used to assure continuity in the monitoring of Reef water quality trends.

#### Production of the Reef water type maps

Previous methods used Daily MODIS Level-0 data acquired from the NASA Ocean Colour website, spectrally enhanced (from red-green-blue to hue-saturation-intensity colour system) and classified to six colour categories through a supervised classification using spectral signatures from typical wet season water masses types (including river plumes) in the Reef lagoon (Álvarez-Romero *et al.* 2013).

For this report, Reef water type maps were produced using daily Sentinel-3 OLCI Level 2 (hereafter, Sentinel-3 or S3) imagery reclassified to 21 distinct colour classes defined by their colour properties and using the Forel-Ule colour classification scale.

- Sentinel-3 imagery of the study area was downloaded on the EUMETSAT Data centre (URL: <https://www.eumetsat.int/eumetsat-data-centre>). Sentinel-3 are atmospherically corrected
- The imagery was processed with the FU Satellite Toolbox implemented in the Sentinel Application Platform (<https://step.esa.int/main/toolboxes/snap/>) and using automated tools (python scripts and ArcGIS toolboxes) developed through MMP funding

The FU satellite algorithm converts satellite normalised multi-band reflectance information into a discrete set of FU numbers using uniform colourimetric functions (Van der Woerd *et al.*, 2016, Van der Woerd and Wernand, 2018). The derivation of the colour of natural waters is based on the calculation of Tristimulus values of the three primaries (X, Y, Z) that specify the colour stimulus of the human eye. The algorithm is validated by a set of hyperspectral measurements from inland, coastal and marine waters (Van der Woerd *et al.*, 2016, Van der Woerd and Wernand, 2018). Technical details about the FU scale algorithm are synthesised through the European citclops (URL: <http://www.citclops.eu/>) and Eye on Water project webpages (<https://www.eyeonwater.org/>).

## Production of weekly Reef water type maps

Weekly Reef water type composites are then created to minimise the image area contaminated by dense cloud cover and intense sun glint (Álvarez-Romero *et al.*, 2013).

- The maximum FU value of each pixel/week is used to keep the colour class with the highest turbidity level for each wet season week.
- The weekly composite maps are then cleaned to remove single or small clusters of cells sometimes misclassified by the FU satellite algorithm in the offshore regions of the Great Barrier Reef (including, for example, around coral reefs due to bottom interference and residual glint contamination). The aim of cleaning is to minimise the image area contaminated by dense cloud cover and intense sun glint, and to remove shallow water interference around reefs. In all cases the effect of these phenomena can be that offshore waters are misclassified as, for example, Reef WT1 waters (FU  $\geq 10$ , previously primary waters). To minimise these effects an automated process is applied to the rasters that has the effect of sequentially infilling contiguous water-type areas one colour class at a time from FU1 through to FU21 using Python 2.7.3 (Python Software Foundation, 2012) and ArcGIS 10.7 (ESRI, 2019). Infilling was achieved using the following steps: 1) Raster to Polygon conversion (not simplified), 2) Union (no gaps) then 3) removal, using Erase, of an external polygon, and 4) Polygon to Raster conversion. This process generates a separate raster mask (values 1 or 0) for each colour class, and the final cleaned raster is created by adding the component raster masks. Whilst this process is effective at removing noise offshore it can occasionally have the effect of removing areas of turbid coastal and plume water if they are not directly connected to the coast. To counter this, a final step is included in the cleaning process whereby waters classified as FU classes  $\geq 10$  i.e., in the cleaned raster are replaced with pixels of FU classes  $\geq 10$  in the original raster, using Con (Spatial Analyst). Thus, pixels adjacent to the coast that are classified as highly turbid water are kept and pixels within otherwise contiguous water types offshore are removed.

## Production of annual, multi-annual, and typical wet and dry Reef frequency maps

- Four distinct Reef water type (WT) are defined by grouping the FU colour classes 1–3 (Reef WT4, equivalent to marine waters in the WS scale), FU colour classes 4–5 (Reef WT3, equivalent to WS Tertiary water type), FU colour classes 6–9 (Reef WT2, equivalent to WS Secondary water type) and FU  $\geq 10$  (Reef WT1, equivalent to wet season primary water type), as defined in Petus *et al.* (2019). The Reef water types are fully described in Table 2-2.
- Weekly Reef water type composites are thus overlaid in ArcGIS (i.e., presence/absence of one Reef water type), normalised (0–1) to compute each year a seasonal normalised frequency maps of occurrence of each Reef water type individually and of the Reef WT1–2 combined. Pixel (or cell) values of these maps range from 0 to 1; with a value of 1 meaning that one pixel has been exposed 22 weeks out of 22 weeks of the wet season.
- Annual frequency maps are then overlaid in ArcGIS to create multi-annual normalised frequency composites of occurrence of Reef water types. Multi-annual composites are calculated over different time frames, using the archive of MODIS-Aqua (2002–03 to 2019–20) and Sentinel-3 (2020–21 to 2022–23) water type maps. In order to combine the MODIS-Aqua and Sentinel-3 frequency composites, the MODIS frequency rasters were resampled to the same spatial resolution as the Sentinel imagery (0.00329 decimal degrees) using the Nearest Interpolation methods in ArcMAP 10.6 (Resample tool, Data Management)

Multi-annual frequency composites include: (i) a long-term period (2002–03 to 2021–22: 20 wet seasons) and (ii) a typical recovery period for Reef corals (2012–2017).

Composite frequency maps are also produced to represent typical wet year and dry year conditions. To account for broad-scale spatial variability in wet season river flows, wet- and dry-year maps are first produced separately by averaging frequency maps from the four wettest and driest years in each NRM region (Table B-2). The wet-year maps for each NRM region are combined into a single composite Reef-wide map using the maximum value of the input rasters. This method captures wet-year plume conditions across the entire Reef even if the most significant plume events originate outside the NRM (for example, if Fitzroy plumes are dominant in the Mackay-Whitsunday region the top-quartile discharges from the Fitzroy are already included in the composite raster). Conversely, the dry-year maps are combined into a Reef-wide composite map using the minimum value of the input rasters, which thus represents the least extensive plume from an average of the driest years in each NRM region.

Except for the coral recovery period, reference maps (long-term, wet and dry frequency maps) were all updated in 2023 to ensure they remain valid as a representative period and to improve their accuracy as more satellite data are available. The previous update was in 2019.

The daily, weekly, and wet season frequency maps are used to illustrate the wet season conditions for every year, to assess the extent of river flood plumes and resuspension events in the Reef and to compare seasonal with long-term trends, as well as trend in water composition during typical dry and wet years. Results are presented in the main report and in [Appendix C-6](#).

Table B-2: Wettest and driest years used to compute the typical wet and typical dry composite frequency maps in each NRM region. All years are in the top/bottom quartiles, except 2005 and 2007 for Cape York which are under the long-term median.

Region	Wet years				Dry years			
<b>Cape York</b>	2021	2006	2011	2019	2003	2016	2005	2007
<b>Wet Tropics</b>	2018	2009	2019	2011	2003	2020	2015	2005
<b>Burdekin</b>	2019	2008	2009	2011	2015	2016	2004	2003
<b>Mackay-Whitsunday</b>	2012	2008	2010	2011	2004	2015	2003	2006
<b>Fitzroy</b>	2008	2010	2013	2011	2006	2005	2007	2004
<b>Burnett-Mary</b>	2012	2022	2013	2011	2021	2006	2007	2005

## Susceptibility assessment

Frequency maps are compared with ecological health information collected through the coral reef and seagrass components of the MMP (e.g., McKenzie *et al.*, 2019, Thompson *et al.*, 2019) to better understand the susceptibility of the seagrass meadow and coral reef ecosystems to water quality conditions.

## Composition of Reef water types

The classification of four Reef water types allows mapping of large Reef waterbodies with different colour characteristics and concentrations of optically active components (TSS, CDOM, and Chl-*a*), water quality indicators (e.g. nutrients levels; Devlin *et al.*, 2015; Petus *et al.*, 2019), and light attenuation levels (Petus *et al.*, 2018) typically found in the Reef during the wet season (Table 2-2). Match-up of *in situ* concentrations of water quality parameters and the four Reef water types are performed to validate this concept and quantify the range and average of water quality concentrations found in each Reef water type.

Match-ups between sampled date and corresponding weekly Reef water type maps are performed at site location basis using the *extract tool* of the raster package (Hijmans *et al.*, 2015) with bilinear interpolation method in R 3.2.4 or the Extract Multi Values to Point (Spatial Analyst) in ArcMap (from 2020). The R tool interpolates from the values of the four nearest raster cells (R Core Team, 2019) while the ArcMap tool extract cell values at the exact location (used from 2020). Several land-based pollutants are investigated through match-ups between *in situ* data and the six colour class maps, including NO<sub>x</sub>, DIP, PP, PN, TSS, Chl-*a*, CDOM, and *K<sub>D</sub>* or Secchi depth. Boxplots of water quality parameters across water types (Figure 4-3) and MODIS WS colour classes (Table B-2) as well as the mean long-term concentrations of water quality parameters across the three wet season water types in all focus regions (Figure B-2) are presented. Work is currently underway to break down the turbid WT1 into a greater number of water types and identify Sentinel-3 Forel-Ule colour classes equivalent to the former MODIS colour classes 1 to 4.




Figure B-1: Long-term water quality (WQ) concentrations and Secchi depth boxplots for each wet season colour class. The mean is plotted as a cross (x) and its numerical value is indicated. The interquartile range is delimited by the box and the median by the line inside the box. Whiskers indicate variability outside the upper and lower quartiles. Data beyond the whiskers range are considered outliers and are not plotted. Long-term water quality values are reviewed and updated every 4 years (and/or in the case of extremely wet year or specific event patterns) to ensure the water type characterization remains valid as a representative period, and to improve its accuracy as more field data are collected every wet seasons. The last update was in 2019, using all field data available (from 2004–2019). Work is underway to update this figure.




Figure B-2: Mean long-term (2004–2023) concentrations of water quality parameters across the three wet season water types in all focus regions. Red lines show the Reef-wide wet season GV (Table B-4). The Burdekin region has the greatest average TSS, PP, and PN concentrations in the Reef water type 1, which exceeded the long-term Reef-scale average. The greatest mean NO<sub>x</sub> and CDOM concentrations are measured in the Reef water type 1 of the Wet Tropics and Cape York regions, respectively. The greatest mean Chl-a concentrations are measured in the Reef water type 1 of the Burdekin region, but concentrations are more uniform across region. Mean long-term concentrations of water quality parameters include samples collected from the enclosed coastal water body (Table B-1), where high TSS, PN, and PP concentrations are likely to contribute to exceedances of the Reef-wide GV.

Detailed summaries of water quality parameters (mean, standard deviation, minimum, maximum, and number of values for each parameter across colour classes and water types) for the long-term are provided in [Appendix C-4](#). Long-term water quality values are calculated using all surface data (<0.2 m) collected between December and April by JCU (since 2004), AIMS, and the CYWP (since 2016–17) in the whole of the Reef.

### Exposure maps and exposure assessment

Information on the long-term water chemistry concentrations measured in the Reef water type are compared to published water quality guideline values and, combined with frequency maps of occurrence of wet season colour classes, are used in a “*magnitude x likelihood*” risk

management framework to develop surface exposure maps (also referred to as potential risk maps in some Reef studies). Different frameworks have been used to estimate the exposure and potential risk from exposure and are described in Petus *et al.* (2014a, 2016), Waterhouse *et al.* (2017), Gruber *et al.* (2019), and used in the MMP reports before 2015–16. In a collaborative effort between the MMP monitoring providers (JCU water quality and seagrass teams and the AIMS coral monitoring team), an updated exposure assessment framework was developed in 2015–16 (modified from Petus *et al.*, 2016), where the ‘potential risk’ corresponds to an exposure to above guideline concentrations of land-based pollutant during wet season conditions and focuses on the TSS, Chl-a, PP, and PN concentrations.

- The ‘*magnitude of the exposure*’ corresponds to the long-term concentration of pollutants (proportional exceedance of the guideline) mapped through the Reef WT1, WT2 and WT3 (primary, secondary and tertiary water types).
- The ‘*likelihood of the exposure*’ is estimated by calculating the frequency of occurrence of each Reef water type. The exposure for each of the water quality parameters defined is as the proportional exceedance of the guideline multiplied by the likelihood of exposure in each of the Reef water type and calculated as below. For each cell (500 m x 500 m):

For each pollutant (Poll.) the exposure in the Reef WT1, WT2, WT3 (primary or secondary or tertiary):  $Poll\_expo_{water\ type}$  is calculated:

$$Poll\_expo_{water\ type} = magnitude_{water\ type} \times likelihood_{water\ type}$$

$$magnitude_{water\ type} = ([Poll.]_{water\ type} - guideline) / guideline$$

$$likelihood_{water\ type} = frequency_{water\ type}$$

where *water type* is the Reef WT1, WT2 or WT3 (primary, secondary and tertiary water types),  $[Poll.]_{water\ type}$  is long-term mean TSS, Chl-a, PN, or PP concentration measured in each respective wet season water types and *guideline* is the Reef-wide wet season GV from De’ath and Fabricius (2008) for TSS, Chl-a, PP, and PN (Table B-4).

Table B-3: Reef-wide wet season guideline values used to calculate the exposure score for satellite exposure maps. These guidelines are based on seasonal adjustments to Reef-wide annual guidelines (Great Barrier Reef Marine Park Authority, 2010), where wet season guidelines are +20% for TSS, PN, and PP, and +40% for Chl-*a* of annual guidelines (De'ath and Fabricius 2008).

Parameter	Unit	Reef-wide
Chlorophyll <i>a</i>	µg L <sup>-1</sup>	0.63
Particulate nitrogen	µg L <sup>-1</sup>	25
Particulate phosphorus	µg L <sup>-1</sup>	3.3
Suspended solids	mg L <sup>-1</sup>	2.4

These GVs are compared against the mean long-term concentrations to calculate the exposure score in the satellite exposure maps (proportional exceedance of the guideline). Mean long-term concentrations of water quality parameters are calculated using all available surface water quality data in all Reef marine regions and water bodies (Table B-4). The variability in the number of samples between regions and water types is primarily driven by the sampling design which was reviewed in 2014. The small number of samples in the Burnett-Mary region reflects the geographic extent of the MMP; with a majority of the samples collected by JCU in the 2011 and 2013 flood events when the design of the event monitoring was more opportunistic across the whole Reef. The relatively small number of samples in offshore waters reflects the geographic focus of the MMP design which is largely constrained to the open coastal and mid-shelf waters. The last update in the mean long-term concentrations was in the 2022–23 reporting year (Gruber *et al.*, 2024), using field data collected from 2004 to 2019. Note also that the long-term and Reef-wide concentrations of water quality parameters are used rather than the seasonal and/or regional mean concentrations in water type to avoid bias due to differential regional and seasonal sampling distribution.

Table B-4: Number of collected *in situ* samples used in exposure scoring by region and water type. Samples include all wet season (Dec–April) surface samples since 2004 (from JCU) and since the 2016–17 water year (AIMS and the CYWP) and up to April 2022.

Region	Reef Water type	Secchi	TSS	CHL	CDO M	NO <sub>x</sub>	DIP	PP	PN
Cape York	WT1	157	208	218	160	214	218	102	80
	WT2	225	295	301	180	301	301	188	170
	WT3	126	176	181	109	178	178	120	111
	Marine	8	13	13	4	13	13	5	4
Wet Tropics	WT1	185	406	399	388	356	356	57	58
	WT2	400	623	637	574	611	615	228	229
	WT3	203	289	296	239	273	274	143	143
	Marine	25	33	35	29	33	33	19	19
Burdekin	WT1	102	157	156	113	151	155	63	73
	WT2	202	258	260	194	258	260	99	106
	WT3	61	97	96	71	81	82	40	40
	Marine	21	33	39	23	28	29	20	19
Mackay-Whitsunday	WT1	28	45	42	43	45	45	20	20
	WT2	73	134	129	98	127	132	74	75
	WT3	20	39	39	27	33	34	27	27
	Marine	7	13	13	8	9	10	6	6
Fitzroy	WT1	22	103	104	78	105	105	17	17
	WT2	27	64	78	65	82	84	22	22
	WT3	8	20	25	11	16	17	8	8
	Marine	0	6	6	1	6	6	0	0
Burnett-Mary	WT1	7	16	16	7	7	16	0	0
	WT2	5	9	9	5	5	9	0	0
	WT3	0	2	2	0	0	0	0	0
	Marine	0	8	8	1	3	3	0	0
Reef-wide	WT1	501	935	935	789	878	895	259	248
	WT2	932	1383	1414	1116	1384	1401	611	602
	WT3	418	623	639	457	581	585	338	329
	Marine	61	106	114	66	92	94	50	48

For each pollutant, the total exposure ( $Poll\_expo$ ) is calculated as the exposure for each of the Reef water types:

$$Poll\_expo = Poll\_expo_{Primary} + Poll\_expo_{Secondary} + Poll\_expo_{Tertiary}$$

The overall exposure score ( $Score\_expo$ ) is calculated as the sum of the total exposure for each of the water quality parameters:

$$Score\_expo = TSS.exp + Chla.exp + PP.exp + PN.exp$$

For example, using the long-term mean Chl-a values measured during high flow conditions in the Reef WT1, WT2, and WT3:

$$Chla\_exp_{WT1} = \frac{1.45-0.63}{0.63} \times frequency_{water\ type\ (0-1, cell-specific)}$$

$$Chla\_exp_{WT2} = \frac{0.65-0.63}{0.63} \times frequency_{water\ type\ (0-1, cell-specific)}$$

$$Chla\_exp_{WT3} = 0 \text{ as Chl-a levels are below the guideline for Chl-a;}$$

The total exposure for Chl-a:

$$Chla\_expo = Chla\_expo_{WT1} + Chla\_expo_{WT2} + Chla\_expo_{WT3}$$

The overall exposure scores are then grouped into four potential classes (I–IV) based on a “Natural Break (or Jenks)” classification. Jenks is a statistical procedure, embedded in ArcGIS that analyses the distribution of values in the data and finds the most evident breaks in it (i.e., the steep or marked breaks; Jenks and Caspall, 1971). The Jenks classification determine the best arrangement of values into different classes by reducing the variance within classes and maximising the variance between classes.

The exposure classes are defined by applying the Jenks classification to the mean long-term exposure map (2003–2022), because this map presented the highest number of observations. Using the 2003–2018 mean exposure map, categories were defined as  $[>0-0.9] = \text{cat. I}$ ,  $[0.9-3.5] = \text{cat. II}$ ,  $[3.2-7.9] = \text{cat. III}$  and  $>7.9 = \text{cat. IV}$ . Category I and areas mapped as “exposure = 0 (no exposure)”, are re-grouped into a unique category I (no or very low exposure). These categories are to all exposure composites created (seasonal, coral recovery period, typical wet and dry periods).

The methods presented above are slightly different than methods used in previous wet season’s reports (2016–17 and 2017–18 wet seasons) where (i) seasonal mean concentrations of water quality parameters across water types were used to produce the seasonal exposure map and (ii) exposure maps were reclassified using four equally-distributed colour classes. Changes in 2019 (using only long-term mean concentrations of water quality parameters and a Jenk’s classification of the exposure maps) were made in response to: (i) concerns that water quality concentrations collected in a specific wet season would likely get biased toward the sample size and the location and timing of sampling in this particular wet season conditions and (ii) that the equally-distributed categories were not responsive enough to changes in environmental pressures of each year.

Exposure maps are produced for the whole of the Reef, for all focus regions and over different time frames: for the current reporting wet season (using the Sentinel-3 FU imagery), and over several multi-years period (using the archive of MODIS WS imagery): the long-term (2002–03 to 2021–22: 20 wet seasons), over a documented recovery period for coral reefs (2012–2017 period) and representation of typical wet-year and dry-year conditions. Except for the coral recovery period, reference maps (long-term, wet and dry frequency maps) were all updated in 2023 to ensure they remain valid as a representative period and to improve their accuracy as more satellite data are available. The last update was in 2019.

- Finally, assessments of ecosystem exposure are made through the calculation of the areas (km<sup>2</sup>) and percentages (%) of each region, coral reefs and seagrass meadows affected by different categories of exposure. The area and percentage are calculated as a relative measure between regions and waterbodies. The difference in percentages between the current year and in the long-term is also calculated. Figure B-3 presents the marine boundaries used for the Marine Park, each NRM region, the Reef waterbodies and the seagrass and coral reefs ecosystems. The area (km<sup>2</sup>) and percentages of seagrass and coral reefs in the Reef and regional waterbodies is indicated in Figure B-3. We assumed in this study that the seagrass shapefile can be used as a representation of the actual seagrass distribution. It is known, however, that absence on the composite map does not definitively equate to absence of seagrass and may also indicate un-surveyed areas.




Figure B-3: Boundaries used for the Marine Park, each NRM region, and the coral reefs and seagrass ecosystems. Coral reef and NRM layers derived from the Reef Authority, supplied 2013. Seagrass layer is a composite of surveys conducted by Department of Agriculture and Fisheries, QLD.


## B-4 References


- Álvarez-Romero JG, Devlin MJ, Teixeira da Silva E, Petus C, Ban N, Pressey RJ, Kool J, Roberts S, Cerdeira WA, Brodie J (2013). A novel approach to model exposure of coastal-marine ecosystems to riverine flood plumes based on remote sensing techniques. *Journal of Environmental Management* 119: 194-207.
- Brando VE, Blondeau-Patissier D, Schroeder T, Dekker AG, Clementson L (2011). Reef Rescue Marine Monitoring Program: Assessment of terrestrial run-off entering the Reef and inshore marine water quality monitoring using earth observation data. Final Report for 2010/11 Activities. CSIRO, Canberra. 201 pp.
- Brodie, J., De'ath, G., Devlin, M., Furnas, M., Wright, M., (2007). Spatial and temporal patterns of near-surface chlorophyll a in the Great Barrier Reef lagoon. *Marine and Freshwater Research* 58, 342–353.
- De'ath G, Fabricius KE (2008) Water quality of the Great Barrier Reef: distributions, effects on reef biota and trigger values for the protection of ecosystem health. Final Report to the Great Barrier Reef Marine Park Authority. Australian Institute of Marine Science, Townsville. 104 pp.
- De'ath G, Fabricius KE (2010). Water quality as a regional driver of coral biodiversity and macroalgae on the Great Barrier Reef. *Ecological Applications* 20: 840–850.
- Department of Environment and Resource Management (DERM) (2009). Queensland Water Quality Guidelines, Version 3. 167 p. Available at [www.derm.qld.gov.au](http://www.derm.qld.gov.au). ISBN 978-0-9806986-0-2.
- Devlin MJ, Wenger A, Petus C, da Silva ET, DeBose J, Álvarez-Romero J (2013). Reef Rescue Marine Monitoring Program. Final report of JCU activities 2011/12: flood plumes and extreme weather monitoring for the Great Barrier Reef Marine Park Authority (Report). James Cook University.
- Devlin M, Petus C, Teixeira da Silva E, Tracey D, Wolff N, Waterhouse J, Brodie J (2015). Water quality and river plume monitoring in the Great Barrier Reef: An Overview of Methods Based on Ocean Colour Satellite Data. *Remote Sensing* 7: 12909-12941.
- Furnas MJ, Mitchell AW, Skuza M, Brodie J (2005). In the other 90%: Phytoplankton responses to enhanced nutrient availability in the Great Barrier Reef lagoon. *Marine Pollution Bulletin* 51: 253-256.
- Furnas M, Alongi D, McKinnon D, Trott L, Skuza M (2011). Regional-scale nitrogen and phosphorus budgets for the northern (14°S) and central (17°S) Great Barrier Reef shelf ecosystem. *Continental Shelf Research* 31: 1967-1990. doi:10.1016/j.csr.2011.09.007
- Great Barrier Reef Marine Park Authority (GBRMPA) (2010). Water Quality Guidelines for the Great Barrier Reef Marine Park. Revised Edition 2010. Great Barrier Reef Marine Park Authority, Townsville. 100 pp.
- Gruber R, Waterhouse J, Logan M, Petus C, Howley C, Lewis S, Tracey D, Langlois L, Tonin H, Skuza M, Costello P, Davidson J, Gunn K, Lefevre C, Shanahan M, Wright M, Zagorskis I, Kroon F, Neilen A (2019). Marine Monitoring Program: Annual Report for Inshore Water Quality Monitoring 2017-18. Report for the Great Barrier Reef Marine Park Authority, Great Barrier Reef Marine Park Authority, Townsville, 304pp.
- Gruber R, Waterhouse J, Logan M, Petus C, Howley C, Lewis S, Tracey D, Langlois L, Tonin H, Skuza M, Costello P, Davidson J, Gunn K, Lefevre C, Moran D, Robson B, Shanahan M, Zagorskis I, Shellberg J, Neilen A (2020). Marine Monitoring Program: Annual Report for Inshore Water Quality Monitoring 2018–19. Report for the Great Barrier Reef Marine Park Authority, Great Barrier Reef Marine Park Authority, Townsville, 262pp.




- Hijmans RJ, Etten J van, Mattiuzzi M, Sumner M, Greenberg JA, Lamigueiro OP, Bevan A, Racine, EB, Shortridge A (2015). raster: Geographic Data Analysis and Modelling.
- Jenks GF and Caspall FC (1971). Error on Choroplethic Maps: Definition, Measurement Reduction. *Annals of the Association of American Geographers* 61: 217-44.
- McKenzie LJ, Collier CJ, Langlois LA, Yoshida RL, Uusitalo J, Smith N, Waycott M (2019). Marine Monitoring Program: Annual Report for inshore seagrass monitoring 2017-2018. Report for the Great Barrier Reef Marine Park Authority, Great Barrier Reef Marine Park Authority, Townsville, 188 pp.
- Petus C, Teixeira da Silva E, Devlin M, Álvarez-Romero J, Wenger A (2014a). Using MODIS data for mapping of water types within flood plumes in the Great Barrier Reef, Australia: towards the production of river plume risk maps for reef and seagrass ecosystems. *Journal of Environmental Management* 137: 163-177.
- Petus C, Collier C, Devlin M, Rasheed M, McKenna S (2014b). Using MODIS data for understanding changes in seagrass meadow health: A case study in the Great Barrier Reef (Australia). *Marine Environmental Research* 98: 68-85.
- Petus C, Devlin M, Thompson A, McKenzie L, Teixeira da Silva E, Collier C, Tracey D, Martin K (2016). Estimating the exposure of coral reefs and seagrass meadows to land-sourced contaminants in river flood plumes of the Great Barrier Reef: validating a simple Satellite Risk Framework with Environmental Data. *Remote Sensing*, 8, 210.
- Petus C, Devlin M, da Silva E, Lewis S, Waterhouse J, Wenger A, Bainbridge Z and Tracey D (2018) Defining wet season water quality target concentrations for ecosystem conservation using empirical light attenuation models: a case study in the Great Barrier Reef (Australia). *Journal of Environmental Management* 213: 1-16.
- Petus C, Waterhouse J, Lewis S, Vacher M, Tracey D, Devlin M. (2019). A flood of information: Using Sentinel-3 water colour products to assure continuity in the monitoring of water quality trends in the Great Barrier Reef (Australia). *Journal of Environmental Management* 248: 109255.
- R Core Team (2019). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL: <https://www.R-project.org/>
- Schaffelke B, Thompson A, Carleton J, Davidson J, Doyle J, Furnas M, Gunn K, Skuza M, Wright M, Zagorskis I (2009). Reef Rescue Marine Monitoring Program. Final Report of AIMS Activities 2008/09. Report for Reef and Rainforest Research Centre. Australian Institute of Marine Science, Townsville. 146 pp.
- Thompson A, Costello P, Davidson J, Logan M, Coleman G (2019). Marine Monitoring Program: Annual report for inshore coral reef monitoring 2017-18. Great Barrier Reef Marine Park Authority, Townsville. 127pp.
- Waterhouse J, Lønborg C, Logan M, Petus C, Tracey D, Lewis S, Tonin H, Skuza M, da Silva E, Carreira C, Costello P, Davidson J, Gunn K, Wright M, Zagorskis I, Brinkman R, Schaffelke B (2017) Marine Monitoring Program: Annual Report for inshore water quality monitoring 2015-2016. Report for the Great Barrier Reef Marine Park Authority, Great Barrier Reef Marine Park Authority, Townsville, 227pp.




## APPENDIX C: ADDITIONAL INFORMATION


### C-1 Time-series of turbidity and chlorophyll

















Figure C-1: Time-series of daily means of chlorophyll fluorescence and turbidity measured by moored ECO FLNTUSB instruments coloured dashed lines represent the water quality GV. Daily river discharge from the nearest river, daily wind speeds from the nearest weather stations, and daily tidal range from the nearest tidal gauge are also shown. Locations of loggers are shown in Figure 2-1, Section 5, and Figure D-1 and panels continue on additional pages below: a) Fitzroy West; b) High West; c) Russell-Mulgrave Mouth Mooring; d) Franklands West; e) Dunk North; f) Tully Mouth Mooring; g) Palms West; h) Pandora; i) Magnetic; j) Burdekin Mouth Mooring; k) Double Cone; m) Pine; n) Seaforth; o) O'Connell River, p) Repulse Island (discontinued in 2021), q) Barren, r) Keppels South, and t) Fitzroy River Mouth.

## C-2 Time-series of temperature and salinity




Figure C-2: Time-series of daily means of temperature and salinity derived from moored loggers (Sea-Bird Electronics SBE37s). Sub-figures represent instrument locations at: a) High West, b) Russel Mulgrave Mouth Mooring, c) Dunk North, d) Tully River Mouth Mooring, e) Burdekin Mouth Mooring, f) Pine, g) Repulse, h) O'Connell River mouth, i) Keppels South, and j) Fitzroy River mouth.

### C-3 Summary statistics for all sites

Table C-1: Summary statistics for water quality parameters at individual monitoring sites from 1 October 2023 to 30 September 2024. N = number of sampling occasions. See Section 2 for descriptions of each analyte and its abbreviation. Mean and median values that exceed available Water Quality Guidelines (DERM, 2009; Great Barrier Reef Marine Park Authority, 2010; State of Queensland, 2020) are shaded in red. Averages that exceed dry season guidelines are shaded in yellow. DOF is direction of failure ('H' = high values fail, while 'L' = low values fail).

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
Cape York	Pascoe	PRN02 (PRN02)	DIN ( $\mu\text{g L}^{-1}$ )	5	4.37	4.48	1.62	2.30	6.22	7.22					
			DOC ( $\mu\text{g L}^{-1}$ )	5	1108	1089	865	908	1205	1474					
			DON ( $\mu\text{g L}^{-1}$ )	5	82	80	64	70	88	109					
			DOP ( $\mu\text{g L}^{-1}$ )	5	3.84	3.41	3.41	3.41	4.21	4.77					
			Chl-a ( $\mu\text{g L}^{-1}$ )	5	0.27	0.32	0.14	0.14	0.37	0.37	H	Median	0.36	0.25	0.46
			NH <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	5	3.61	3.64	1.29	1.79	4.96	6.38					
			NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	5	0.76	0.70	0.34	0.50	0.95	1.29	H	Median	0.35	0.32	0.45
			PN ( $\mu\text{g L}^{-1}$ )	5	38.00	36.68	23.21	28.59	46.51	55.00	H	Mean		16	
			PN ( $\mu\text{g L}^{-1}$ )	5	38.00	36.68	23.21	28.59	46.51	55.00	H	Median	18		20
			PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	5	0.93	0.62	0.37	0.56	1.36	1.73	H	Median	1.4	1.86	0.93
			POC ( $\text{mg L}^{-1}$ )	5	177	176	119	159	199	231					
			PP ( $\mu\text{g L}^{-1}$ )	5	2.62	2.91	1.56	1.60	3.51	3.52	H	Mean		2.3	
			PP ( $\mu\text{g L}^{-1}$ )	5	2.62	2.91	1.56	1.60	3.51	3.52	H	Median	2.6		3
			Secchi (m)	5	2.90	2.90	2.36	2.54	3.26	3.44	L	Mean	10		
			Secchi (m)	5	2.90	2.90	2.36	2.54	3.26	3.44	L	Median			
			SiO <sub>4</sub> ( $\text{mg L}^{-1}$ )	5	155	139	80	90	225	240					
			TSS ( $\text{mg L}^{-1}$ )	5	1.24	0.81	0.41	0.57	1.42	3.01	H	Mean		1.6	

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			TSS (mg L <sup>-1</sup> )	5	1.24	0.81	0.41	0.57	1.42	3.01	H	Median	1.9		1.7
		Pascoe River mouth south (PRS01)	DIN (µg L <sup>-1</sup> )	5	22.07	8.54	7.17	8.09	29.65	56.87					
			DOC (µg L <sup>-1</sup> )	5	2026	1243	1102	1122	2256	4406					
			DON (µg L <sup>-1</sup> )	5	97	95	70	82	105	131					
			DOP (µg L <sup>-1</sup> )	5	3.16	2.79	1.67	2.04	4.09	5.20					
			Chl-a (µg L <sup>-1</sup> )	5	0.58	0.57	0.15	0.16	0.81	1.23	H	Median			0.7
			NH <sub>4</sub> (µg L <sup>-1</sup> )	5	7.56	6.30	4.34	4.76	10.70	11.70					
			NO <sub>x</sub> (µg L <sup>-1</sup> )	5	14.51	4.34	1.99	2.07	17.95	46.18	H	Median			1.5
			PN (µg L <sup>-1</sup> )	5	66.73	38.78	27.33	33.71	74.74	159.08	H	Mean			
			PN (µg L <sup>-1</sup> )	5	66.73	38.78	27.33	33.71	74.74	159.08	H	Median			
			PO <sub>4</sub> (µg L <sup>-1</sup> )	5	2.29	2.17	1.11	1.67	3.16	3.34	H	Median			3
			POC (mg L <sup>-1</sup> )	5	557	251	139	209	655	1533					
			PP (µg L <sup>-1</sup> )	5	6.16	4.55	1.91	2.27	8.28	13.80	H	Mean			
			PP (µg L <sup>-1</sup> )	5	6.16	4.55	1.91	2.27	8.28	13.80	H	Median			
			Secchi (m)	5	1.60	1.10	0.47	0.68	2.42	3.08	L	Mean			
			Secchi (m)	5	1.60	1.10	0.47	0.68	2.42	3.08	L	Median			3
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	5	2021	1474	323	1029	2621	4660					
			TSS (mg L <sup>-1</sup> )	5	14.09	6.62	2.19	2.76	22.43	36.46	H	Median			4
			DIN (µg L <sup>-1</sup> )	5	6.31	6.27	2.38	4.27	8.56	10.09					

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
		PRS05 (PRS05)	DOC ( $\mu\text{g L}^{-1}$ )	5	902	899	809	848	964	992					
			DON ( $\mu\text{g L}^{-1}$ )	5	78	77	71	74	81	89					
			DOP ( $\mu\text{g L}^{-1}$ )	5	3.92	4.03	3.13	3.68	4.26	4.49					
			Chl-a ( $\mu\text{g L}^{-1}$ )	5	0.24	0.26	0.13	0.20	0.30	0.32	H	Median	0.27		
			NH <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	5	3.59	4.17	1.75	2.59	4.43	5.02					
			NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	5	2.72	2.03	0.43	0.90	4.34	5.92	H	Median	0.35		
			PN ( $\mu\text{g L}^{-1}$ )	5	14.68	14.56	10.14	12.01	16.87	19.81	H	Mean			
			PN ( $\mu\text{g L}^{-1}$ )	5	14.68	14.56	10.14	12.01	16.87	19.81	H	Median	18		
			PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	5	2.63	2.09	1.08	1.54	4.10	4.34	H	Median	0.62		
			POC ( $\text{mg L}^{-1}$ )	5	70	71	47	61	77	92					
			PP ( $\mu\text{g L}^{-1}$ )	5	1.46	1.39	0.67	1.14	2.01	2.07	H	Mean			
			PP ( $\mu\text{g L}^{-1}$ )	5	1.46	1.39	0.67	1.14	2.01	2.07	H	Median	2		
			Secchi (m)	5	10.09	9.90	7.33	8.31	11.76	13.14	L	Mean	10		
			Secchi (m)	5	10.09	9.90	7.33	8.31	11.76	13.14	L	Median			
			SiO <sub>4</sub> ( $\text{mg L}^{-1}$ )	5	67	72	48	50	78	87					
			TSS ( $\text{mg L}^{-1}$ )	5	0.55	0.60	0.18	0.44	0.73	0.78	H	Mean			
			TSS ( $\text{mg L}^{-1}$ )	5	0.55	0.60	0.18	0.44	0.73	0.78	H	Median	1.5		
		PRS2.5 (PRS2.5)	DIN ( $\mu\text{g L}^{-1}$ )	5	6.97	6.44	2.72	3.94	10.49	11.28					
			DOC ( $\mu\text{g L}^{-1}$ )	5	1043	990	831	894	1097	1402					

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			DON ( $\mu\text{g L}^{-1}$ )	5	81	83	70	74	88	92					
			DOP ( $\mu\text{g L}^{-1}$ )	5	3.65	3.41	2.88	3.16	4.18	4.65					
			Chl-a ( $\mu\text{g L}^{-1}$ )	5	0.41	0.42	0.22	0.23	0.56	0.63	H	Median	0.36	0.25	0.46
			NH <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	5	4.17	4.76	2.19	2.65	5.49	5.78					
			NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	5	2.80	1.54	0.43	0.90	4.83	6.30	H	Median	0.35	0.32	0.45
			PN ( $\mu\text{g L}^{-1}$ )	5	28.62	27.37	21.97	24.86	32.93	35.95	H	Mean		16	
			PN ( $\mu\text{g L}^{-1}$ )	5	28.62	27.37	21.97	24.86	32.93	35.95	H	Median	18		20
			PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	5	2.17	2.32	1.18	1.92	2.57	2.85	H	Median	1.4	1.86	0.93
			POC (mg L <sup>-1</sup> )	5	140	135	100	125	149	191					
			PP ( $\mu\text{g L}^{-1}$ )	5	3.13	2.80	2.11	2.58	3.69	4.49	H	Mean		2.3	
			PP ( $\mu\text{g L}^{-1}$ )	5	3.13	2.80	2.11	2.58	3.69	4.49	H	Median	2.6		3
			Secchi (m)	5	3.94	4.30	3.04	3.16	4.48	4.72	L	Mean	10		
			Secchi (m)	5	3.94	4.30	3.04	3.16	4.48	4.72	L	Median			
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	5	230	128	66	81	343	530					
			TSS (mg L <sup>-1</sup> )	5	1.77	1.49	0.94	1.30	2.17	2.94	H	Mean		1.6	
			TSS (mg L <sup>-1</sup> )	5	1.77	1.49	0.94	1.30	2.17	2.94	H	Median	1.9		1.7
		Middle Reef (PRBB)	DIN ( $\mu\text{g L}^{-1}$ )	5	3.69	3.71	1.79	3.18	4.69	5.11					
			DOC ( $\mu\text{g L}^{-1}$ )	5	935	925	860	877	969	1042					
			DON ( $\mu\text{g L}^{-1}$ )	5	81	80	64	69	93	101					

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			DOP ( $\mu\text{g L}^{-1}$ )	5	4.37	4.34	3.59	3.69	5.02	5.20					
			Chl-a ( $\mu\text{g L}^{-1}$ )	5	0.30	0.31	0.09	0.15	0.43	0.52	H	Median	0.27		
			NH <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	5	2.63	2.94	1.23	1.78	3.46	3.75					
			NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	5	1.06	0.77	0.36	0.62	1.72	1.85	H	Median	0.35		
			PN ( $\mu\text{g L}^{-1}$ )	5	17.77	18.55	10.84	14.79	21.01	23.68	H	Mean			
			PN ( $\mu\text{g L}^{-1}$ )	5	17.77	18.55	10.84	14.79	21.01	23.68	H	Median	18		
			PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	5	1.89	1.86	1.08	1.55	2.20	2.76	H	Median	0.62		
			POC (mg L <sup>-1</sup> )	5	84	84	53	73	100	112					
			PP ( $\mu\text{g L}^{-1}$ )	5	1.53	1.58	0.66	0.95	2.12	2.34	H	Mean			
			PP ( $\mu\text{g L}^{-1}$ )	5	1.53	1.58	0.66	0.95	2.12	2.34	H	Median	2		
			Secchi (m)	5	7.56	8.20	5.10	5.70	8.80	10.00	L	Mean	10		
			Secchi (m)	5	7.56	8.20	5.10	5.70	8.80	10.00	L	Median			
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	5	78	81	33	53	103	120					
			TSS (mg L <sup>-1</sup> )	5	0.90	0.68	0.39	0.53	1.24	1.64	H	Mean			
			TSS (mg L <sup>-1</sup> )	5	0.90	0.68	0.39	0.53	1.24	1.64	H	Median	1.5		
	Stewart	Hannah Island (SR05)	DIN ( $\mu\text{g L}^{-1}$ )	5	5.56	5.25	3.13	3.88	7.37	8.17					
			DOC ( $\mu\text{g L}^{-1}$ )	5	1438	1124	1021	1042	1678	2326					
			DON ( $\mu\text{g L}^{-1}$ )	5	98	92	88	88	104	118					
			DOP ( $\mu\text{g L}^{-1}$ )	5	3.94	3.48	2.68	3.05	5.14	5.35					

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			Chl-a ( $\mu\text{g L}^{-1}$ )	5	0.42	0.27	0.13	0.18	0.47	1.06	H	Median	0.27		
			NH4 ( $\mu\text{g L}^{-1}$ )	5	3.99	3.57	2.72	3.28	4.75	5.65					
			NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	5	1.57	0.81	0.41	0.60	2.70	3.33	H	Median	0.35		
			PN ( $\mu\text{g L}^{-1}$ )	5	28.34	17.85	15.53	15.70	31.35	61.29	H	Mean			
			PN ( $\mu\text{g L}^{-1}$ )	5	28.34	17.85	15.53	15.70	31.35	61.29	H	Median	18		
			PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	5	2.80	2.77	2.20	2.29	3.07	3.67	H	Median	0.62		
			POC ( $\text{mg L}^{-1}$ )	5	139	82	70	71	156	314					
			PP ( $\mu\text{g L}^{-1}$ )	5	2.27	1.87	1.69	1.72	2.50	3.56	H	Mean			
			PP ( $\mu\text{g L}^{-1}$ )	5	2.27	1.87	1.69	1.72	2.50	3.56	H	Median	2		
			Secchi (m)	5	9.14	7.20	5.64	6.36	12.92	13.58	L	Mean	10		
			Secchi (m)	5	9.14	7.20	5.64	6.36	12.92	13.58	L	Median			
			SiO <sub>4</sub> ( $\text{mg L}^{-1}$ )	5	292	165	81	124	461	629					
			TSS ( $\text{mg L}^{-1}$ )	5	0.68	0.66	0.33	0.43	0.97	1.00	H	Mean			
			TSS ( $\text{mg L}^{-1}$ )	5	0.68	0.66	0.33	0.43	0.97	1.00	H	Median	1.5		
		Burkitt Island (SR04)	DIN ( $\mu\text{g L}^{-1}$ )	5	5.11	3.43	2.49	2.67	6.83	10.11					
			DOC ( $\mu\text{g L}^{-1}$ )	5	1316	989	912	947	1638	2093					
			DON ( $\mu\text{g L}^{-1}$ )	5	95	103	69	91	106	106					
			DOP ( $\mu\text{g L}^{-1}$ )	5	4.67	4.76	3.96	4.24	4.96	5.42					
			Chl-a ( $\mu\text{g L}^{-1}$ )	5	0.43	0.28	0.15	0.20	0.59	0.93	H	Median	0.36	0.25	0.46

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			NH <sub>4</sub> (µg L <sup>-1</sup> )	5	3.15	2.59	2.02	2.03	4.47	4.63					
			NO <sub>x</sub> (µg L <sup>-1</sup> )	5	1.96	0.84	0.47	0.64	2.37	5.47	H	Median	0.35	0.32	0.45
			PN (µg L <sup>-1</sup> )	5	27.74	20.86	15.88	16.27	31.25	54.44	H	Mean		16	
			PN (µg L <sup>-1</sup> )	5	27.74	20.86	15.88	16.27	31.25	54.44	H	Median	18		20
			PO <sub>4</sub> (µg L <sup>-1</sup> )	5	2.38	2.01	1.64	1.92	2.94	3.38	H	Median	1.4	1.86	0.93
			POC (mg L <sup>-1</sup> )	5	138	101	79	89	151	269					
			PP (µg L <sup>-1</sup> )	5	2.27	2.28	1.45	1.64	2.74	3.25	H	Mean		2.3	
			PP (µg L <sup>-1</sup> )	5	2.27	2.28	1.45	1.64	2.74	3.25	H	Median	2.6		3
			Secchi (m)	5	9.56	8.70	7.28	8.12	11.42	12.30	L	Mean	10		
			Secchi (m)	5	9.56	8.70	7.28	8.12	11.42	12.30	L	Median			
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	5	244	124	75	92	402	529					
			TSS (mg L <sup>-1</sup> )	5	0.93	0.64	0.55	0.61	1.10	1.75	H	Mean		1.6	
			TSS (mg L <sup>-1</sup> )	5	0.93	0.64	0.55	0.61	1.10	1.75	H	Median	1.9		1.7
		SR03 (SR03)	DIN (µg L <sup>-1</sup> )	5	4.74	3.15	2.29	2.44	6.44	9.38					
			DOC (µg L <sup>-1</sup> )	5	1296	1042	974	976	1605	1885					
			DON (µg L <sup>-1</sup> )	5	96	95	70	87	104	122					
			DOP (µg L <sup>-1</sup> )	5	4.00	4.96	1.33	3.00	5.11	5.57					
			Chl-a (µg L <sup>-1</sup> )	5	0.32	0.36	0.20	0.24	0.38	0.41	H	Median	0.36	0.25	0.46
			NH <sub>4</sub> (µg L <sup>-1</sup> )	5	2.95	2.45	1.72	2.04	3.81	4.73					

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			NO <sub>x</sub> (µg L <sup>-1</sup> )	5	1.79	0.70	0.40	0.57	2.63	4.65	H	Median	0.35	0.32	0.45
			PN (µg L <sup>-1</sup> )	5	24.90	21.50	19.99	20.75	25.51	36.72	H	Mean		16	
			PN (µg L <sup>-1</sup> )	5	24.90	21.50	19.99	20.75	25.51	36.72	H	Median	18		20
			PO <sub>4</sub> (µg L <sup>-1</sup> )	5	3.16	1.86	1.55	1.55	3.47	7.37	H	Median	1.4	1.86	0.93
			POC (mg L <sup>-1</sup> )	5	119	118	85	88	132	170					
			PP (µg L <sup>-1</sup> )	5	3.07	2.96	1.51	1.98	3.59	5.32	H	Mean		2.3	
			PP (µg L <sup>-1</sup> )	5	3.07	2.96	1.51	1.98	3.59	5.32	H	Median	2.6		3
			Secchi (m)	5	6.88	7.52	4.48	5.62	8.04	8.76	L	Mean	10		
			Secchi (m)	5	6.88	7.52	4.48	5.62	8.04	8.76	L	Median			
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	5	278	159	79	95	451	607					
			TSS (mg L <sup>-1</sup> )	5	1.26	1.25	0.45	0.69	1.86	2.08	H	Mean		1.6	
			TSS (mg L <sup>-1</sup> )	5	1.26	1.25	0.45	0.69	1.86	2.08	H	Median	1.9		1.7
		SR02 (SR02)	DIN (µg L <sup>-1</sup> )	5	5.78	2.93	2.16	2.22	10.09	11.50					
			DOC (µg L <sup>-1</sup> )	5	1508	1285	1033	1073	2035	2114					
			DON (µg L <sup>-1</sup> )	5	103	104	83	98	113	118					
			DOP (µg L <sup>-1</sup> )	5	4.16	4.08	2.57	3.31	5.19	5.65					
			Chl-a (µg L <sup>-1</sup> )	5	0.46	0.39	0.19	0.28	0.60	0.82	H	Median			0.4
			NH <sub>4</sub> (µg L <sup>-1</sup> )	5	3.24	2.12	1.61	1.61	4.36	6.52					
			NO <sub>x</sub> (µg L <sup>-1</sup> )	5	2.54	0.84	0.55	0.61	4.98	5.73	H	Median			1.5

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			PN ( $\mu\text{g L}^{-1}$ )	5	37.98	40.18	23.97	35.29	42.78	47.66	H	Mean			
			PN ( $\mu\text{g L}^{-1}$ )	5	37.98	40.18	23.97	35.29	42.78	47.66	H	Median			
			PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	5	2.08	2.57	0.71	1.46	2.80	2.85	H	Median			2
			POC ( $\text{mg L}^{-1}$ )	5	189	201	117	172	216	240					
			PP ( $\mu\text{g L}^{-1}$ )	5	5.84	4.03	2.33	3.20	6.36	13.27	H	Mean			
			PP ( $\mu\text{g L}^{-1}$ )	5	5.84	4.03	2.33	3.20	6.36	13.27	H	Median			
			Secchi (m)	5	3.78	5.10	1.36	1.56	5.36	5.54	L	Mean			
			Secchi (m)	5	3.78	5.10	1.36	1.56	5.36	5.54	L	Median			3.1
			SiO <sub>4</sub> ( $\text{mg L}^{-1}$ )	5	504	285	136	226	650	1222					
			TSS ( $\text{mg L}^{-1}$ )	5	6.14	1.00	0.90	0.97	8.29	19.57	H	Mean			
			TSS ( $\text{mg L}^{-1}$ )	5	6.14	1.00	0.90	0.97	8.29	19.57	H	Median			5
	Normanby	Corbett Reef (NR06)	DIN ( $\mu\text{g L}^{-1}$ )	4	4.32	4.85	1.78	3.12	5.74	6.13					
			DOC ( $\mu\text{g L}^{-1}$ )	4	1325	1153	1086	1089	1491	1803					
			DON ( $\mu\text{g L}^{-1}$ )	4	81	79	68	71	90	95					
			DOP ( $\mu\text{g L}^{-1}$ )	4	4.22	3.87	3.79	3.79	4.51	5.14					
			Chl-a ( $\mu\text{g L}^{-1}$ )	4	0.40	0.34	0.17	0.25	0.52	0.72	H	Median	0.26		
			NH <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	4	3.26	3.54	1.39	2.51	4.13	4.76					
			NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	4	1.06	0.61	0.33	0.38	1.56	2.41	H	Median	0.42		
			PN ( $\mu\text{g L}^{-1}$ )	4	22.60	24.97	13.80	18.83	27.32	28.09	H	Median	16		

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			PO <sub>4</sub> (µg L <sup>-1</sup> )	4	2.17	2.09	0.63	1.11	3.20	3.83	H	Median	0.39		
			POC (mg L <sup>-1</sup> )	4	114	123	74	93	139	143					
			PP (µg L <sup>-1</sup> )	4	1.86	1.95	1.18	1.57	2.20	2.44	H	Mean			
			PP (µg L <sup>-1</sup> )	4	1.86	1.95	1.18	1.57	2.20	2.44	H	Median	1.9		
			Secchi (m)	4	9.11	8.48	8.32	8.39	9.58	10.80	L	Mean	17		
			Secchi (m)	4	9.11	8.48	8.32	8.39	9.58	10.80	L	Median			
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	4	273	166	60	94	408	635					
			TSS (mg L <sup>-1</sup> )	4	0.64	0.61	0.41	0.51	0.76	0.92	H	Mean			
			TSS (mg L <sup>-1</sup> )	4	0.64	0.61	0.41	0.51	0.76	0.92	H	Median	0.5		
		NR05 (NR05)	DIN (µg L <sup>-1</sup> )	4	5.20	4.21	3.33	3.65	6.36	8.47					
			DOC (µg L <sup>-1</sup> )	4	1355	1161	1039	1086	1546	1943					
			DON (µg L <sup>-1</sup> )	4	84	80	68	71	96	107					
			DOP (µg L <sup>-1</sup> )	4	4.39	4.18	3.74	3.81	4.89	5.34					
			Chl-a (µg L <sup>-1</sup> )	4	0.59	0.46	0.22	0.27	0.85	1.12	H	Median	0.27		
			NH <sub>4</sub> (µg L <sup>-1</sup> )	4	3.19	2.98	1.91	2.20	4.10	4.75					
			NO <sub>x</sub> (µg L <sup>-1</sup> )	4	2.01	1.72	0.47	0.63	3.28	3.97	H	Median	0.35		
			PN (µg L <sup>-1</sup> )	4	25.53	22.26	15.66	18.97	30.79	39.99	H	Mean			
			PN (µg L <sup>-1</sup> )	4	25.53	22.26	15.66	18.97	30.79	39.99	H	Median	18		
			PO <sub>4</sub> (µg L <sup>-1</sup> )	4	2.31	2.71	1.03	1.80	2.99	3.03	H	Median	0.62		

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			POC (mg L <sup>-1</sup> )	4	145	148	79	93	198	207					
			PP (µg L <sup>-1</sup> )	4	2.44	2.06	1.44	1.52	3.21	3.97	H	Mean			
			PP (µg L <sup>-1</sup> )	4	2.44	2.06	1.44	1.52	3.21	3.97	H	Median	2		
			Secchi (m)	4	8.48	8.00	6.25	6.38	10.38	11.37	L	Mean	10		
			Secchi (m)	4	8.48	8.00	6.25	6.38	10.38	11.37	L	Median			
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	4	317	155	58	87	482	801					
			TSS (mg L <sup>-1</sup> )	4	1.04	0.92	0.51	0.72	1.32	1.74	H	Mean			
			TSS (mg L <sup>-1</sup> )	4	1.04	0.92	0.51	0.72	1.32	1.74	H	Median	1.5		
		NR04 (NR04)	DIN (µg L <sup>-1</sup> )	4	5.60	3.57	2.88	3.14	7.25	11.16					
			DOC (µg L <sup>-1</sup> )	4	1380	1403	1086	1237	1533	1643					
			DON (µg L <sup>-1</sup> )	4	82	84	76	80	86	86					
			DOP (µg L <sup>-1</sup> )	4	4.27	4.26	3.89	3.96	4.58	4.67					
			Chl-a (µg L <sup>-1</sup> )	4	0.70	0.69	0.31	0.33	1.07	1.12	H	Median	0.36	0.25	0.46
			NH <sub>4</sub> (µg L <sup>-1</sup> )	4	2.73	2.49	2.04	2.07	3.30	3.76					
			NO <sub>x</sub> (µg L <sup>-1</sup> )	4	2.87	1.23	0.51	0.61	4.47	7.52	H	Median	0.35	0.32	0.45
			PN (µg L <sup>-1</sup> )	4	29.49	30.03	18.87	23.82	35.38	39.35	H	Mean		16	
			PN (µg L <sup>-1</sup> )	4	29.49	30.03	18.87	23.82	35.38	39.35	H	Median	18		20
			PO <sub>4</sub> (µg L <sup>-1</sup> )	4	1.92	1.94	1.26	1.33	2.52	2.56	H	Median	1.4	1.86	0.93
			POC (mg L <sup>-1</sup> )	4	142	144	92	110	175	190					

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			PP ( $\mu\text{g L}^{-1}$ )	4	3.43	2.62	2.10	2.24	4.29	5.89	H	Mean		2.3	
			PP ( $\mu\text{g L}^{-1}$ )	4	3.43	2.62	2.10	2.24	4.29	5.89	H	Median	2.6		3
			Secchi (m)	4	8.75	9.25	5.41	6.62	11.08	11.40	L	Mean	10		
			Secchi (m)	4	8.75	9.25	5.41	6.62	11.08	11.40	L	Median			
			SiO <sub>4</sub> ( $\text{mg L}^{-1}$ )	4	236	149	80	98	340	514					
			TSS ( $\text{mg L}^{-1}$ )	4	1.54	1.43	0.93	0.94	2.09	2.30	H	Mean		1.6	
			TSS ( $\text{mg L}^{-1}$ )	4	1.54	1.43	0.93	0.94	2.09	2.30	H	Median	1.9		1.7
		Cliff Isles (CI01)	DIN ( $\mu\text{g L}^{-1}$ )	3	3.62	3.64	3.26	3.39	3.85	3.96					
			DOC ( $\mu\text{g L}^{-1}$ )	3	1805	1779	1690	1720	1885	1938					
			DON ( $\mu\text{g L}^{-1}$ )	3	108	106	100	102	113	116					
			DOP ( $\mu\text{g L}^{-1}$ )	3	4.49	3.56	3.42	3.47	5.33	6.21					
			Chl-a ( $\mu\text{g L}^{-1}$ )	3	0.48	0.41	0.36	0.38	0.57	0.64	H	Median	0.36	0.25	0.46
			NH <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	3	2.45	2.38	2.32	2.34	2.55	2.63					
			NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	3	1.17	0.98	0.85	0.90	1.40	1.61	H	Median	0.35	0.32	0.45
			PN ( $\mu\text{g L}^{-1}$ )	3	35.35	34.93	29.01	30.98	39.64	41.99	H	Mean		16	
			PN ( $\mu\text{g L}^{-1}$ )	3	35.35	34.93	29.01	30.98	39.64	41.99	H	Median	18		20
			PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	3	1.81	1.55	1.13	1.27	2.29	2.66	H	Median	1.4	1.86	0.93
			POC ( $\text{mg L}^{-1}$ )	3	222	189	158	168	268	308					
			PP ( $\mu\text{g L}^{-1}$ )	3	3.18	3.25	2.54	2.78	3.60	3.77	H	Mean		2.3	

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			PP ( $\mu\text{g L}^{-1}$ )	3	3.18	3.25	2.54	2.78	3.60	3.77	H	Median	2.6		3
			Secchi (m)	3	5.23	4.70	4.25	4.40	5.96	6.59	L	Mean	10		
			Secchi (m)	3	5.23	4.70	4.25	4.40	5.96	6.59	L	Median			
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	3	364	240	134	170	533	679					
			TSS (mg L <sup>-1</sup> )	3	1.13	1.01	0.81	0.88	1.36	1.53	H	Mean		1.6	
			TSS (mg L <sup>-1</sup> )	3	1.13	1.01	0.81	0.88	1.36	1.53	H	Median	1.9		1.7
		NR03 (NR03)	DIN ( $\mu\text{g L}^{-1}$ )	4	3.20	3.17	2.75	2.79	3.60	3.71					
			DOC ( $\mu\text{g L}^{-1}$ )	4	1640	1500	1060	1208	2016	2416					
			DON ( $\mu\text{g L}^{-1}$ )	4	91	90	80	82	100	102					
			DOP ( $\mu\text{g L}^{-1}$ )	4	4.45	4.57	3.52	3.87	5.08	5.22					
			Chl-a ( $\mu\text{g L}^{-1}$ )	4	0.40	0.32	0.21	0.26	0.51	0.72	H	Median	0.36	0.25	0.46
			NH <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	4	2.26	2.12	1.95	2.03	2.44	2.78					
			NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	4	0.94	0.67	0.61	0.62	1.15	1.65	H	Median	0.35	0.32	0.45
			PN ( $\mu\text{g L}^{-1}$ )	4	25.53	22.82	16.94	19.65	30.33	37.90	H	Mean		16	
			PN ( $\mu\text{g L}^{-1}$ )	4	25.53	22.82	16.94	19.65	30.33	37.90	H	Median	18		20
			PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	4	1.30	1.41	0.58	0.95	1.70	1.88	H	Median	1.4	1.86	0.93
			POC (mg L <sup>-1</sup> )	4	136	126	93	102	166	194					
			PP ( $\mu\text{g L}^{-1}$ )	4	2.62	2.32	1.84	1.87	3.25	3.82	H	Mean		2.3	
			PP ( $\mu\text{g L}^{-1}$ )	4	2.62	2.32	1.84	1.87	3.25	3.82	H	Median	2.6		3

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			Secchi (m)	4	6.36	6.33	4.85	5.30	7.41	7.93	L	Mean	10		
			Secchi (m)	4	6.36	6.33	4.85	5.30	7.41	7.93	L	Median			
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	4	290	258	94	139	428	532					
			TSS (mg L <sup>-1</sup> )	4	1.47	1.72	0.54	1.04	2.00	2.06	H	Mean		1.6	
			TSS (mg L <sup>-1</sup> )	4	1.47	1.72	0.54	1.04	2.00	2.06	H	Median	1.9		1.7
		Normanby inshore (NR02)	DIN (µg L <sup>-1</sup> )	4	4.04	2.31	2.31	2.31	5.09	8.20					
			DOC (µg L <sup>-1</sup> )	4	2143	1918	1079	1123	3073	3521					
			DON (µg L <sup>-1</sup> )	4	106	103	79	86	125	139					
			DOP (µg L <sup>-1</sup> )	4	4.74	4.88	3.83	4.18	5.35	5.45					
			Chl-a (µg L <sup>-1</sup> )	4	0.79	0.69	0.37	0.41	1.13	1.35	H	Median			0.7
			NH <sub>4</sub> (µg L <sup>-1</sup> )	4	2.58	1.94	1.83	1.86	3.04	4.23					
			NO <sub>x</sub> (µg L <sup>-1</sup> )	4	1.49	0.46	0.42	0.42	2.14	4.00	H	Median			1
			PN (µg L <sup>-1</sup> )	4	43.32	43.03	31.13	36.51	50.01	55.92	H	Mean			
			PN (µg L <sup>-1</sup> )	4	43.32	43.03	31.13	36.51	50.01	55.92	H	Median			
			PO <sub>4</sub> (µg L <sup>-1</sup> )	4	1.44	1.08	0.54	0.75	1.98	2.82	H	Median			2
			POC (mg L <sup>-1</sup> )	4	231	227	169	196	264	297					
			PP (µg L <sup>-1</sup> )	4	4.57	4.07	2.73	2.99	5.95	7.11	H	Mean			
			PP (µg L <sup>-1</sup> )	4	4.57	4.07	2.73	2.99	5.95	7.11	H	Median			
			Secchi (m)	4	2.64	2.60	1.86	2.04	3.22	3.47	L	Mean			

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			Secchi (m)	4	2.64	2.60	1.86	2.04	3.22	3.47	L	Median			1.5
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	4	368	218	67	144	533	879					
			TSS (mg L <sup>-1</sup> )	4	2.80	3.32	1.31	2.34	3.48	3.57	H	Mean			
			TSS (mg L <sup>-1</sup> )	4	2.80	3.32	1.31	2.34	3.48	3.57	H	Median			6
	Annan-Endeavour	Endeavour offshore (ER03)	DIN (µg L <sup>-1</sup> )	5	3.70	3.22	2.35	2.90	4.23	5.78					
			DOC (µg L <sup>-1</sup> )	5	981	976	945	949	1018	1020					
			DON (µg L <sup>-1</sup> )	5	77	70	66	68	89	93					
			DOP (µg L <sup>-1</sup> )	5	3.59	3.41	2.48	2.94	4.34	4.80					
			Chl-a (µg L <sup>-1</sup> )	5	0.21	0.17	0.16	0.17	0.24	0.30	H	Median	0.36	0.25	0.46
			NH <sub>4</sub> (µg L <sup>-1</sup> )	5	3.02	2.59	1.83	1.88	3.61	5.21					
			NO <sub>x</sub> (µg L <sup>-1</sup> )	5	0.67	0.56	0.32	0.45	0.78	1.25	H	Median	0.35	0.32	0.45
			PN (µg L <sup>-1</sup> )	5	19.36	17.43	16.86	17.24	20.95	24.35	H	Mean		16	
			PN (µg L <sup>-1</sup> )	5	19.36	17.43	16.86	17.24	20.95	24.35	H	Median	18		20
			PO <sub>4</sub> (µg L <sup>-1</sup> )	5	1.64	0.93	0.65	0.74	2.94	2.94	H	Median	1.4	1.86	0.93
			POC (mg L <sup>-1</sup> )	5	94	93	76	79	109	112					
			PP (µg L <sup>-1</sup> )	5	1.88	1.83	1.67	1.67	1.99	2.23	H	Mean		2.3	
			PP (µg L <sup>-1</sup> )	5	1.88	1.83	1.67	1.67	1.99	2.23	H	Median	2.6		3
			Secchi (m)	5	9.00	9.80	4.88	6.62	10.98	12.72	L	Mean	10		
			Secchi (m)	5	9.00	9.80	4.88	6.62	10.98	12.72	L	Median			

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	5	179	183	83	132	245	253					
			TSS (mg L <sup>-1</sup> )	5	0.60	0.57	0.50	0.52	0.65	0.78	H	Mean		1.6	
			TSS (mg L <sup>-1</sup> )	5	0.60	0.57	0.50	0.52	0.65	0.78	H	Median	1.9		1.7
		Endeavour north shore (ER02b)	DIN (µg L <sup>-1</sup> )	5	3.51	3.08	2.32	2.58	4.44	5.15					
			DOC (µg L <sup>-1</sup> )	5	1037	1049	950	955	1078	1154					
			DON (µg L <sup>-1</sup> )	5	77	68	67	67	90	94					
			DOP (µg L <sup>-1</sup> )	5	3.93	4.34	2.79	3.25	4.55	4.74					
			Chl-a (µg L <sup>-1</sup> )	5	0.24	0.22	0.17	0.18	0.31	0.35	H	Median	0.36	0.25	0.46
			NH <sub>4</sub> (µg L <sup>-1</sup> )	5	2.74	2.03	1.85	1.93	3.64	4.27					
			NO <sub>x</sub> (µg L <sup>-1</sup> )	5	0.77	0.77	0.46	0.59	0.95	1.08	H	Median	0.35	0.32	0.45
			PN (µg L <sup>-1</sup> )	5	21.39	19.25	16.89	17.35	25.06	28.42	H	Mean		16	
			PN (µg L <sup>-1</sup> )	5	21.39	19.25	16.89	17.35	25.06	28.42	H	Median	18		20
			PO <sub>4</sub> (µg L <sup>-1</sup> )	5	1.18	0.93	0.40	0.68	1.61	2.26	H	Median	1.4	1.86	0.93
			POC (mg L <sup>-1</sup> )	5	112	103	87	89	135	148					
			PP (µg L <sup>-1</sup> )	5	1.90	1.86	1.60	1.65	2.13	2.25	H	Mean		2.3	
			PP (µg L <sup>-1</sup> )	5	1.90	1.86	1.60	1.65	2.13	2.25	H	Median	2.6		3
			Secchi (m)	5	6.19	6.25	3.72	4.68	7.82	8.48	L	Mean	10		
			Secchi (m)	5	6.19	6.25	3.72	4.68	7.82	8.48	L	Median			
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	5	209	268	82	89	292	312					

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			TSS (mg L <sup>-1</sup> )	5	1.27	0.82	0.64	0.73	1.50	2.69	H	Mean		1.6	
			TSS (mg L <sup>-1</sup> )	5	1.27	0.82	0.64	0.73	1.50	2.69	H	Median	1.9		1.7
		Egret and Boulder Reef (AE04)	DIN (µg L <sup>-1</sup> )	5	4.16	2.28	2.16	2.25	5.57	8.53					
			DOC (µg L <sup>-1</sup> )	5	965	969	934	935	980	1007					
			DON (µg L <sup>-1</sup> )	5	78	73	67	71	84	92					
			DOP (µg L <sup>-1</sup> )	5	4.10	4.34	2.83	3.44	4.91	5.00					
			Chl-a (µg L <sup>-1</sup> )	5	0.26	0.25	0.16	0.18	0.30	0.41	H	Median	0.27		
			NH <sub>4</sub> (µg L <sup>-1</sup> )	5	2.68	1.86	1.46	1.65	3.91	4.52					
			NO <sub>x</sub> (µg L <sup>-1</sup> )	5	1.48	0.88	0.34	0.50	1.66	4.01	H	Median	0.35		
			PN (µg L <sup>-1</sup> )	5	18.64	18.03	14.63	17.05	20.18	23.29	H	Mean			
			PN (µg L <sup>-1</sup> )	5	18.64	18.03	14.63	17.05	20.18	23.29	H	Median	18		
			PO <sub>4</sub> (µg L <sup>-1</sup> )	5	1.83	1.86	0.96	1.05	2.21	3.05	H	Median	0.62		
			POC (mg L <sup>-1</sup> )	5	81	81	71	76	86	90					
			PP (µg L <sup>-1</sup> )	5	1.58	1.59	1.35	1.48	1.66	1.83	H	Mean			
			PP (µg L <sup>-1</sup> )	5	1.58	1.59	1.35	1.48	1.66	1.83	H	Median	2		
			Secchi (m)	5	11.49	10.40	6.44	8.06	14.64	17.91	L	Mean	10		
			Secchi (m)	5	11.49	10.40	6.44	8.06	14.64	17.91	L	Median			
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	5	106	103	57	60	133	178					
			TSS (mg L <sup>-1</sup> )	5	0.44	0.40	0.26	0.35	0.54	0.66	H	Mean			

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			TSS (mg L <sup>-1</sup> )	5	0.44	0.40	0.26	0.35	0.54	0.66	H	Median	1.5		
		Dawson Reef (AR03b)	DIN (µg L <sup>-1</sup> )	5	2.97	2.31	1.79	2.13	3.86	4.75					
			DOC (µg L <sup>-1</sup> )	5	1060	1030	974	1007	1111	1180					
			DON (µg L <sup>-1</sup> )	5	79	72	68	69	90	97					
			DOP (µg L <sup>-1</sup> )	5	4.52	3.72	2.66	3.22	4.92	8.08					
			Chl-a (µg L <sup>-1</sup> )	5	0.27	0.28	0.15	0.20	0.37	0.37	H	Median	0.36	0.25	0.46
			NH <sub>4</sub> (µg L <sup>-1</sup> )	5	1.90	1.61	1.39	1.55	2.46	2.51					
			NO <sub>x</sub> (µg L <sup>-1</sup> )	5	1.06	0.70	0.41	0.57	1.40	2.24	H	Median	0.35	0.32	0.45
			PN (µg L <sup>-1</sup> )	5	22.05	23.66	15.29	16.84	27.06	27.40	H	Mean		16	
			PN (µg L <sup>-1</sup> )	5	22.05	23.66	15.29	16.84	27.06	27.40	H	Median	18		20
			PO <sub>4</sub> (µg L <sup>-1</sup> )	5	1.49	1.08	0.81	0.90	1.77	2.88	H	Median	1.4	1.86	0.93
			POC (mg L <sup>-1</sup> )	5	112	100	72	75	144	169					
			PP (µg L <sup>-1</sup> )	5	1.94	1.83	1.29	1.56	2.10	2.92	H	Mean		2.3	
			PP (µg L <sup>-1</sup> )	5	1.94	1.83	1.29	1.56	2.10	2.92	H	Median	2.6		3
			Secchi (m)	5	8.36	8.00	6.58	6.82	9.54	10.86	L	Mean	10		
			Secchi (m)	5	8.36	8.00	6.58	6.82	9.54	10.86	L	Median			
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	5	254	261	83	185	351	388					
			TSS (mg L <sup>-1</sup> )	5	0.89	0.73	0.53	0.65	1.07	1.48	H	Mean		1.6	
			TSS (mg L <sup>-1</sup> )	5	0.89	0.73	0.53	0.65	1.07	1.48	H	Median	1.9		1.7

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
		Walker Bay (AR02b)	DIN ( $\mu\text{g L}^{-1}$ )	5	3.54	4.03	1.50	2.95	4.39	4.85					
			DOC ( $\mu\text{g L}^{-1}$ )	5	1051	1067	976	1006	1096	1110					
			DON ( $\mu\text{g L}^{-1}$ )	5	79	75	66	71	90	91					
			DOP ( $\mu\text{g L}^{-1}$ )	5	3.75	4.10	2.63	3.10	4.38	4.52					
			Chl-a ( $\mu\text{g L}^{-1}$ )	5	0.27	0.25	0.17	0.21	0.36	0.39	H	Median	0.36	0.25	0.46
			NH <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	5	2.65	2.87	1.07	2.39	3.25	3.65					
			NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	5	0.90	1.12	0.42	0.53	1.20	1.22	H	Median	0.35	0.32	0.45
			PN ( $\mu\text{g L}^{-1}$ )	5	23.63	20.58	17.89	19.70	26.87	33.13	H	Mean		16	
			PN ( $\mu\text{g L}^{-1}$ )	5	23.63	20.58	17.89	19.70	26.87	33.13	H	Median	18		20
			PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	5	1.15	0.77	0.37	0.56	1.46	2.57	H	Median	1.4	1.86	0.93
			POC (mg L <sup>-1</sup> )	5	125	108	87	102	143	183					
			PP ( $\mu\text{g L}^{-1}$ )	5	2.52	2.32	2.27	2.29	2.57	3.17	H	Mean		2.3	
			PP ( $\mu\text{g L}^{-1}$ )	5	2.52	2.32	2.27	2.29	2.57	3.17	H	Median	2.6		3
			Secchi (m)	5	6.03	6.50	5.03	5.12	6.60	6.90	L	Mean	10		
			Secchi (m)	5	6.03	6.50	5.03	5.12	6.60	6.90	L	Median			
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	5	179	151	122	133	239	252					
			TSS (mg L <sup>-1</sup> )	5	0.86	1.10	0.29	0.50	1.19	1.20	H	Mean		1.6	
			TSS (mg L <sup>-1</sup> )	5	0.86	1.10	0.29	0.50	1.19	1.20	H	Median	1.9		1.7

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
Wet Tropics	Barron-Daintree	Cape Tribulation (C1)	DIN ( $\mu\text{g L}^{-1}$ )	3	3.33	2.80	2.74	2.76	3.79	4.28					
			DOC ( $\mu\text{g L}^{-1}$ )	3	938	932	907	915	960	974					
			DON ( $\mu\text{g L}^{-1}$ )	3	97	93	84	87	107	114					
			DOP ( $\mu\text{g L}^{-1}$ )	3	4.54	4.41	4.34	4.37	4.69	4.83					
			Chl-a ( $\mu\text{g L}^{-1}$ )	3	0.20	0.26	0.11	0.16	0.26	0.26	H	Mean	0.45		
			Chl-a ( $\mu\text{g L}^{-1}$ )	3	0.20	0.26	0.11	0.16	0.26	0.26	H	Median		0.32	0.63
			NH <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	3	2.77	2.52	1.86	2.08	3.40	3.84					
			NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	3	0.56	0.46	0.30	0.35	0.75	0.90	H	Median	0.35		
			PN ( $\mu\text{g L}^{-1}$ )	3	13.39	13.39	11.34	12.02	14.75	15.43	H	Mean	20		
			PN ( $\mu\text{g L}^{-1}$ )	3	13.39	13.39	11.34	12.02	14.75	15.43	H	Median		16	25
			PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	3	2.53	3.02	1.35	1.90	3.25	3.37	H	Median	2		
			POC (mg L <sup>-1</sup> )	3	97	100	79	86	108	112					
			PP ( $\mu\text{g L}^{-1}$ )	3	2.47	3.09	1.15	1.79	3.28	3.37	H	Mean	2.8		
			PP ( $\mu\text{g L}^{-1}$ )	3	2.47	3.09	1.15	1.79	3.28	3.37	H	Median		2.3	3.3
			Secchi (m)	3	6.67	8.00	3.50	5.00	8.60	8.90	L	Mean	10		
			Secchi (m)	3	6.67	8.00	3.50	5.00	8.60	8.90	L	Median			
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	3	188	230	86	134	250	260					
			TSS (mg L <sup>-1</sup> )	3	0.84	0.84	0.76	0.78	0.90	0.92	H	Mean	2		
			TSS (mg L <sup>-1</sup> )	3	0.84	0.84	0.76	0.78	0.90	0.92	H	Median		1.6	2.4

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
		Port Douglas (C4)	DIN ( $\mu\text{g L}^{-1}$ )	3	3.23	3.50	2.24	2.66	3.86	4.04					
			DOC ( $\mu\text{g L}^{-1}$ )	3	902	893	840	858	944	970					
			DON ( $\mu\text{g L}^{-1}$ )	3	88	80	72	75	100	110					
			DOP ( $\mu\text{g L}^{-1}$ )	3	4.41	4.34	4.20	4.24	4.57	4.68					
			Chl-a ( $\mu\text{g L}^{-1}$ )	3	0.21	0.21	0.11	0.14	0.28	0.32	H	Median	0.3	0.32	0.63
			NH <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	3	2.72	3.22	1.55	2.11	3.43	3.54					
			NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	3	0.51	0.53	0.30	0.38	0.65	0.71	H	Median	0.31		
			PN ( $\mu\text{g L}^{-1}$ )	3	11.50	12.15	9.93	10.67	12.47	12.62	H	Median	14	16	25
			PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	3	2.48	2.48	1.57	1.87	3.08	3.38	H	Median	2		
			POC ( $\text{mg L}^{-1}$ )	3	78	76	57	63	92	100					
			PP ( $\mu\text{g L}^{-1}$ )	3	1.99	1.98	0.87	1.24	2.73	3.11	H	Median	2	2.3	3.3
			Secchi (m)	3	6.83	8.00	3.95	5.30	8.60	8.90	L	Median	13		
			SiO <sub>4</sub> ( $\text{mg L}^{-1}$ )	3	152	174	61	99	210	228					
			TSS ( $\text{mg L}^{-1}$ )	3	0.70	0.70	0.58	0.62	0.79	0.83	H	Median	1.2	1.6	2.4
		Double Island (C5)	DIN ( $\mu\text{g L}^{-1}$ )	3	3.38	3.64	2.79	3.07	3.75	3.80					
			DOC ( $\mu\text{g L}^{-1}$ )	3	921	878	859	866	967	1011					
			DON ( $\mu\text{g L}^{-1}$ )	3	92	89	86	87	97	101					
			DOP ( $\mu\text{g L}^{-1}$ )	3	4.67	4.72	4.10	4.30	5.05	5.21					
			Chl-a ( $\mu\text{g L}^{-1}$ )	3	0.22	0.18	0.17	0.18	0.26	0.31	H	Median	0.3	0.32	0.63

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			NH <sub>4</sub> (µg L <sup>-1</sup> )	3	2.96	3.22	2.50	2.74	3.24	3.25					
			NO <sub>x</sub> (µg L <sup>-1</sup> )	3	0.42	0.42	0.29	0.34	0.50	0.55	H	Median	0.31		
			PN (µg L <sup>-1</sup> )	3	12.72	13.26	11.08	11.81	13.74	13.98	H	Median	14	16	25
			PO <sub>4</sub> (µg L <sup>-1</sup> )	3	2.35	2.56	1.51	1.86	2.88	3.04	H	Median	2		
			POC (mg L <sup>-1</sup> )	3	86	84	72	76	96	102					
			PP (µg L <sup>-1</sup> )	3	2.19	2.18	2.05	2.10	2.28	2.33	H	Median	2	2.3	3.3
			Secchi (m)	3	9.50	7.50	5.25	6.00	12.60	15.15	L	Median	13		
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	3	178	143	124	130	218	256					
			TSS (mg L <sup>-1</sup> )	3	0.65	0.65	0.20	0.35	0.94	1.09	H	Median	1.2	1.6	2.4
		Green Island (C11)	DIN (µg L <sup>-1</sup> )	3	6.70	5.39	3.00	3.79	9.34	11.31					
			DOC (µg L <sup>-1</sup> )	3	879	887	838	855	905	913					
			DON (µg L <sup>-1</sup> )	3	93	87	86	86	97	103					
			DOP (µg L <sup>-1</sup> )	3	4.72	4.88	4.25	4.46	5.02	5.09					
			Chl-a (µg L <sup>-1</sup> )	3	0.15	0.13	0.08	0.10	0.20	0.24	H	Median	0.3	0.32	0.63
			NH <sub>4</sub> (µg L <sup>-1</sup> )	3	3.76	4.20	2.40	3.00	4.60	4.80					
			NO <sub>x</sub> (µg L <sup>-1</sup> )	3	2.94	1.19	0.59	0.79	4.74	6.51	H	Median	0.31		
			PN (µg L <sup>-1</sup> )	3	10.44	9.36	8.99	9.12	11.55	12.65	H	Median	14	16	25
			PO <sub>4</sub> (µg L <sup>-1</sup> )	3	3.23	3.41	2.85	3.04	3.45	3.48	H	Median	2		
			POC (mg L <sup>-1</sup> )	3	55	49	45	46	63	70					

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			PP ( $\mu\text{g L}^{-1}$ )	3	1.33	1.51	0.64	0.93	1.77	1.91	H	Median	2	2.3	3.3
			Secchi (m)	3	14.50	17.00	8.00	11.00	18.50	19.25	L	Median	13		
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	3	127	88	68	75	172	214					
			TSS (mg L <sup>-1</sup> )	3	0.32	0.32	0.30	0.31	0.33	0.34	H	Median	1.2	1.6	2.4
		Yorkey's Knob (C6)	DIN ( $\mu\text{g L}^{-1}$ )	3	3.93	3.43	2.83	3.03	4.73	5.38					
			DOC ( $\mu\text{g L}^{-1}$ )	3	1038	1038	1006	1017	1060	1071					
			DON ( $\mu\text{g L}^{-1}$ )	3	103	92	90	90	114	125					
			DOP ( $\mu\text{g L}^{-1}$ )	3	4.85	5.11	4.34	4.60	5.16	5.18					
			Chl-a ( $\mu\text{g L}^{-1}$ )	3	0.32	0.29	0.26	0.27	0.37	0.40	H	Mean	0.45		
			Chl-a ( $\mu\text{g L}^{-1}$ )	3	0.32	0.29	0.26	0.27	0.37	0.40	H	Median		0.32	0.63
			NH <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	3	3.21	2.87	2.52	2.64	3.71	4.13					
			NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	3	0.72	0.56	0.31	0.39	1.02	1.25	H	Median	0.35		
			PN ( $\mu\text{g L}^{-1}$ )	3	16.12	16.95	14.69	15.44	16.96	16.96	H	Mean	20		
			PN ( $\mu\text{g L}^{-1}$ )	3	16.12	16.95	14.69	15.44	16.96	16.96	H	Median		16	25
			PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	3	2.27	2.71	1.25	1.73	2.90	2.99	H	Median	2		
			POC (mg L <sup>-1</sup> )	3	130	149	98	115	149	149					
			PP ( $\mu\text{g L}^{-1}$ )	3	3.68	3.90	2.51	2.97	4.43	4.69	H	Mean	2.8		
			PP ( $\mu\text{g L}^{-1}$ )	3	3.68	3.90	2.51	2.97	4.43	4.69	H	Median		2.3	3.3
			Secchi (m)	3	3.17	2.50	2.50	2.50	3.70	4.30	L	Mean	10		

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			Secchi (m)	3	3.17	2.50	2.50	2.50	3.70	4.30	L	Median			
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	3	271	312	171	218	332	342					
			TSS (mg L <sup>-1</sup> )	3	1.55	1.55	0.82	1.07	2.04	2.28	H	Mean	2		
			TSS (mg L <sup>-1</sup> )	3	1.55	1.55	0.82	1.07	2.04	2.28	H	Median		1.6	2.4
		Fairlead Buoy (C8)	DIN (µg L <sup>-1</sup> )	3	2.80	2.52	1.98	2.16	3.38	3.81					
			DOC (µg L <sup>-1</sup> )	3	989	1004	908	940	1042	1060					
			DON (µg L <sup>-1</sup> )	3	101	95	92	93	108	114					
			DOP (µg L <sup>-1</sup> )	3	4.70	4.65	4.16	4.32	5.06	5.27					
			Chl-a (µg L <sup>-1</sup> )	3	0.33	0.38	0.22	0.27	0.41	0.42	H	Mean	0.45		
			Chl-a (µg L <sup>-1</sup> )	3	0.33	0.38	0.22	0.27	0.41	0.42	H	Median		0.32	0.63
			NH <sub>4</sub> (µg L <sup>-1</sup> )	3	1.93	1.89	1.67	1.74	2.10	2.21					
			NO <sub>x</sub> (µg L <sup>-1</sup> )	3	0.88	0.28	0.28	0.28	1.35	1.89	H	Median	0.35		
			PN (µg L <sup>-1</sup> )	3	17.56	16.65	15.32	15.76	19.17	20.43	H	Mean	20		
			PN (µg L <sup>-1</sup> )	3	17.56	16.65	15.32	15.76	19.17	20.43	H	Median		16	25
			PO <sub>4</sub> (µg L <sup>-1</sup> )	3	2.30	2.32	1.42	1.72	2.88	3.16	H	Median	2		
			POC (mg L <sup>-1</sup> )	3	148	143	121	129	166	178					
			PP (µg L <sup>-1</sup> )	3	3.72	3.52	2.53	2.86	4.54	5.04	H	Mean	2.8		
			PP (µg L <sup>-1</sup> )	3	3.72	3.52	2.53	2.86	4.54	5.04	H	Median		2.3	3.3
			Secchi (m)	3	3.83	3.00	2.55	2.70	4.80	5.70	L	Mean	10		

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			Secchi (m)	3	3.83	3.00	2.55	2.70	4.80	5.70	L	Median			
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	3	189	173	150	158	218	240					
			TSS (mg L <sup>-1</sup> )	3	2.28	2.28	0.84	1.32	3.23	3.71	H	Mean	2		
			TSS (mg L <sup>-1</sup> )	3	2.28	2.28	0.84	1.32	3.23	3.71	H	Median		1.6	2.4
	Russell-Mulgra ve	Fitzroy West (RM1)	DIN (µg L <sup>-1</sup> )	5	4.34	3.43	2.32	2.37	5.32	8.26					
			DOC (µg L <sup>-1</sup> )	5	930	913	861	865	968	1042					
			DON (µg L <sup>-1</sup> )	5	90	89	85	87	93	98					
			DOP (µg L <sup>-1</sup> )	5	4.20	4.26	3.44	4.00	4.51	4.78					
			Chl-a (µg L <sup>-1</sup> )	5	0.24	0.25	0.14	0.15	0.28	0.36	H	Mean	0.45		
			Chl-a (µg L <sup>-1</sup> )	5	0.24	0.25	0.14	0.15	0.28	0.36	H	Median		0.32	0.63
			NH <sub>4</sub> (µg L <sup>-1</sup> )	5	3.21	2.45	1.41	1.88	4.35	5.94					
			NO <sub>x</sub> (µg L <sup>-1</sup> )	5	1.13	0.98	0.33	0.48	1.45	2.44	H	Median	0.35		
			PN (µg L <sup>-1</sup> )	5	13.02	12.37	10.67	11.78	14.83	15.47	H	Mean	20		
			PN (µg L <sup>-1</sup> )	5	13.02	12.37	10.67	11.78	14.83	15.47	H	Median		16	25
			PO <sub>4</sub> (µg L <sup>-1</sup> )	5	2.79	2.86	1.90	2.04	3.56	3.56	H	Median	2		
			POC (mg L <sup>-1</sup> )	5	90	87	58	66	118	120					
			PP (µg L <sup>-1</sup> )	5	2.46	2.62	1.72	2.22	2.79	2.96	H	Mean	2.8		
			PP (µg L <sup>-1</sup> )	5	2.46	2.62	1.72	2.22	2.79	2.96	H	Median		2.3	3.3
			Secchi (m)	5	7.90	8.00	6.60	6.90	9.00	9.00	L	Mean	10		

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			Secchi (m)	5	7.90	8.00	6.60	6.90	9.00	9.00	L	Median			
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	5	155	102	61	65	236	311					
			TSS (mg L <sup>-1</sup> )	5	0.76	0.84	0.36	0.52	1.02	1.10	H	Mean	2		
			TSS (mg L <sup>-1</sup> )	5	0.76	0.84	0.36	0.52	1.02	1.10	H	Median		1.6	2.4
		RM3 (RM3)	DIN (µg L <sup>-1</sup> )	10	5.24	4.78	2.48	2.99	6.51	9.66					
			DOC (µg L <sup>-1</sup> )	10	1036	1049	829	873	1116	1293					
			DON (µg L <sup>-1</sup> )	10	89	88	75	81	96	109					
			DOP (µg L <sup>-1</sup> )	10	4.43	4.41	3.78	4.07	4.89	5.08					
			Chl-a (µg L <sup>-1</sup> )	10	0.26	0.24	0.18	0.19	0.34	0.37	H	Median	0.3	0.32	0.63
			NH <sub>4</sub> (µg L <sup>-1</sup> )	10	3.69	3.22	2.05	2.70	4.89	5.93					
			NO <sub>x</sub> (µg L <sup>-1</sup> )	10	1.55	0.53	0.30	0.34	2.47	4.58	H	Median	0.31		
			PN (µg L <sup>-1</sup> )	10	23.57	20.73	12.33	15.07	28.74	42.49	H	Median	14	16	25
			PO <sub>4</sub> (µg L <sup>-1</sup> )	10	2.69	2.52	1.12	1.86	3.72	4.20	H	Median	2		
			POC (mg L <sup>-1</sup> )	10	144	143	86	116	181	192					
			PP (µg L <sup>-1</sup> )	10	2.51	2.30	1.30	1.83	3.23	3.73	H	Median	2	2.3	3.3
			Secchi (m)	10	10.05	9.00	7.45	8.00	11.60	15.65	L	Median	13		
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	10	253	196	55	97	363	582					
			TSS (mg L <sup>-1</sup> )	10	1.39	1.12	0.39	0.69	1.48	3.32	H	Median	1.2	1.6	2.4
			DIN (µg L <sup>-1</sup> )	10	8.88	5.01	2.78	3.21	13.69	24.09					

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
		High West (RM8)	DOC ( $\mu\text{g L}^{-1}$ )	10	997	1016	896	903	1071	1111					
			DON ( $\mu\text{g L}^{-1}$ )	10	91	90	79	82	99	109					
			DOP ( $\mu\text{g L}^{-1}$ )	10	4.58	4.53	4.17	4.34	4.89	5.04					
			Chl-a ( $\mu\text{g L}^{-1}$ )	10	0.33	0.31	0.15	0.22	0.48	0.52	H	Mean	0.45		
			Chl-a ( $\mu\text{g L}^{-1}$ )	10	0.33	0.31	0.15	0.22	0.48	0.52	H	Median		0.32	0.63
			NH <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	10	4.75	3.47	1.96	2.64	6.26	10.71					
			NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	10	4.13	2.10	0.31	0.35	5.08	14.76	H	Median	0.35		
			PN ( $\mu\text{g L}^{-1}$ )	10	24.84	26.56	13.16	14.48	33.94	37.79	H	Mean	20		
			PN ( $\mu\text{g L}^{-1}$ )	10	24.84	26.56	13.16	14.48	33.94	37.79	H	Median		16	25
			PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	10	2.99	2.94	1.61	2.43	3.90	4.28	H	Median	2		
			POC (mg L <sup>-1</sup> )	10	146	143	83	115	186	202					
			PP ( $\mu\text{g L}^{-1}$ )	10	2.85	2.69	1.77	2.08	3.53	4.12	H	Mean	2.8		
			PP ( $\mu\text{g L}^{-1}$ )	10	2.85	2.69	1.77	2.08	3.53	4.12	H	Median		2.3	3.3
			Secchi (m)	10	8.10	7.00	5.23	5.90	12.00	12.55	L	Mean	10		
			Secchi (m)	10	8.10	7.00	5.23	5.90	12.00	12.55	L	Median			
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	10	350	313	83	133	415	854					
			TSS (mg L <sup>-1</sup> )	10	0.97	0.83	0.34	0.57	1.38	1.88	H	Mean	2		
			TSS (mg L <sup>-1</sup> )	10	0.97	0.83	0.34	0.57	1.38	1.88	H	Median		1.6	2.4
			DIN ( $\mu\text{g L}^{-1}$ )	10	11.46	8.52	2.43	3.44	21.72	23.33					

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
		Russell Mulgrave Mouth Mooring (RM10)	DOC ( $\mu\text{g L}^{-1}$ )	10	1096	1114	919	957	1207	1290					
			DON ( $\mu\text{g L}^{-1}$ )	10	95	92	85	87	106	110					
			DOP ( $\mu\text{g L}^{-1}$ )	10	4.43	4.18	3.83	3.87	4.61	5.78					
			Chl-a ( $\mu\text{g L}^{-1}$ )	10	0.37	0.37	0.23	0.26	0.47	0.53	H	Mean	0.45		
			Chl-a ( $\mu\text{g L}^{-1}$ )	10	0.37	0.37	0.23	0.26	0.47	0.53	H	Median		0.32	0.63
			NH <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	10	4.77	3.13	1.99	2.31	6.37	10.48					
			NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	10	6.69	6.11	0.28	0.45	12.03	15.58	H	Median	0.35		
			PN ( $\mu\text{g L}^{-1}$ )	10	33.47	30.52	14.58	18.90	47.30	56.90	H	Mean	20		
			PN ( $\mu\text{g L}^{-1}$ )	10	33.47	30.52	14.58	18.90	47.30	56.90	H	Median		16	25
			PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	10	3.04	3.14	1.63	2.14	4.03	4.28	H	Median	2		
			POC (mg L <sup>-1</sup> )	10	218	211	99	144	293	364					
			PP ( $\mu\text{g L}^{-1}$ )	10	4.48	4.54	2.71	3.55	5.09	6.19	H	Mean	2.8		
			PP ( $\mu\text{g L}^{-1}$ )	10	4.48	4.54	2.71	3.55	5.09	6.19	H	Median		2.3	3.3
			Secchi (m)	10	4.90	5.00	3.23	3.90	5.60	7.10	L	Mean	10		
			Secchi (m)	10	4.90	5.00	3.23	3.90	5.60	7.10	L	Median			
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	10	586	556	103	232	851	1116					
			TSS (mg L <sup>-1</sup> )	10	1.82	1.48	0.83	1.01	2.56	3.36	H	Mean	2		
			TSS (mg L <sup>-1</sup> )	10	1.82	1.48	0.83	1.01	2.56	3.36	H	Median		1.6	2.4
			DIN ( $\mu\text{g L}^{-1}$ )	10	3.63	3.57	1.96	2.46	4.31	6.01					

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
	Franklands West (RM7)		DOC ( $\mu\text{g L}^{-1}$ )	10	966	950	861	894	1059	1096					
			DON ( $\mu\text{g L}^{-1}$ )	10	88	89	67	74	96	111					
			DOP ( $\mu\text{g L}^{-1}$ )	10	4.74	4.65	4.10	4.30	5.02	5.69					
			Chl-a ( $\mu\text{g L}^{-1}$ )	10	0.31	0.30	0.14	0.18	0.40	0.53	H	Median	0.3	0.32	0.63
			NH <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	10	2.76	2.98	1.30	2.13	3.28	4.07					
			NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	10	0.86	0.72	0.28	0.34	1.02	2.15	H	Median	0.31		
			PN ( $\mu\text{g L}^{-1}$ )	10	22.48	21.45	11.23	13.92	30.61	35.94	H	Median	14	16	25
			PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	10	2.62	2.28	1.50	1.81	3.59	4.06	H	Median	2		
			POC (mg L <sup>-1</sup> )	10	138	160	66	92	167	179					
			PP ( $\mu\text{g L}^{-1}$ )	10	2.52	2.63	1.72	2.35	2.81	3.05	H	Median	2	2.3	3.3
			Secchi (m)	10	10.20	10.50	6.35	8.00	13.00	14.10	L	Median	13		
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	10	187	176	84	109	232	341					
			TSS (mg L <sup>-1</sup> )	10	0.98	0.74	0.47	0.63	1.31	1.98	H	Median	1.2	1.6	2.4
	Tully	Clump Point East (TUL2)	DIN ( $\mu\text{g L}^{-1}$ )	10	4.04	3.43	2.01	2.38	4.73	7.86					
			DOC ( $\mu\text{g L}^{-1}$ )	10	944	914	842	870	999	1136					
			DON ( $\mu\text{g L}^{-1}$ )	10	87	83	73	75	96	107					
			DOP ( $\mu\text{g L}^{-1}$ )	10	4.53	4.41	4.03	4.09	4.99	5.24					
			Chl-a ( $\mu\text{g L}^{-1}$ )	10	0.20	0.18	0.10	0.13	0.25	0.35	H	Median	0.3	0.32	0.63
			NH <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	10	3.37	3.05	1.73	2.04	4.00	6.19					

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			NO <sub>x</sub> (µg L <sup>-1</sup> )	10	0.67	0.39	0.28	0.28	0.79	1.80	H	Median	0.31		
			PN (µg L <sup>-1</sup> )	10	21.51	22.11	10.12	11.88	30.73	32.72	H	Median	14	16	25
			PO <sub>4</sub> (µg L <sup>-1</sup> )	10	2.43	2.40	1.00	1.52	3.25	3.90	H	Median	2		
			POC (mg L <sup>-1</sup> )	10	121	127	52	75	169	176					
			PP (µg L <sup>-1</sup> )	10	2.14	2.09	1.65	1.79	2.39	2.79	H	Median	2	2.3	3.3
			Secchi (m)	10	11.40	12.00	5.90	8.60	14.40	16.00	L	Median	13		
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	10	139	89	43	55	174	382					
			TSS (mg L <sup>-1</sup> )	10	0.65	0.54	0.17	0.34	0.92	1.29	H	Median	1.2	1.6	2.4
		Dunk North (TUL3)	DIN (µg L <sup>-1</sup> )	10	5.15	4.88	1.99	3.46	6.13	9.59					
			DOC (µg L <sup>-1</sup> )	10	1040	1072	915	959	1090	1157					
			DON (µg L <sup>-1</sup> )	10	87	84	73	83	93	104					
			DOP (µg L <sup>-1</sup> )	10	4.72	4.72	3.88	4.43	5.11	5.46					
			Chl-a (µg L <sup>-1</sup> )	10	0.30	0.21	0.16	0.18	0.40	0.62	H	Mean	0.45		
			Chl-a (µg L <sup>-1</sup> )	10	0.30	0.21	0.16	0.18	0.40	0.62	H	Median		0.32	0.63
			NH <sub>4</sub> (µg L <sup>-1</sup> )	10	3.99	3.68	1.71	2.62	5.10	7.13					
			NO <sub>x</sub> (µg L <sup>-1</sup> )	10	1.16	0.93	0.28	0.56	1.86	2.48	H	Median	0.35		
			PN (µg L <sup>-1</sup> )	10	22.23	22.53	11.54	15.21	27.96	32.88	H	Mean	20		
			PN (µg L <sup>-1</sup> )	10	22.23	22.53	11.54	15.21	27.96	32.88	H	Median		16	25
			PO <sub>4</sub> (µg L <sup>-1</sup> )	10	2.65	2.71	1.76	2.01	3.02	3.57	H	Median	2		

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			POC (mg L <sup>-1</sup> )	10	143	150	78	131	165	185					
			PP (µg L <sup>-1</sup> )	10	2.75	2.53	1.48	2.16	3.48	4.24	H	Mean	2.8		
			PP (µg L <sup>-1</sup> )	10	2.75	2.53	1.48	2.16	3.48	4.24	H	Median		2.3	3.3
			Secchi (m)	10	7.30	7.50	4.23	5.70	9.60	10.00	L	Mean	10		
			Secchi (m)	10	7.30	7.50	4.23	5.70	9.60	10.00	L	Median			
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	10	295	281	127	188	422	455					
			TSS (mg L <sup>-1</sup> )	10	1.39	1.17	0.81	0.92	1.75	2.35	H	Mean	2		
			TSS (mg L <sup>-1</sup> )	10	1.39	1.17	0.81	0.92	1.75	2.35	H	Median		1.6	2.4
		Dunk Island South East (TUL5)	DIN (µg L <sup>-1</sup> )	10	5.12	3.33	2.42	2.47	8.26	9.66					
			DOC (µg L <sup>-1</sup> )	10	985	981	869	902	1062	1127					
			DON (µg L <sup>-1</sup> )	10	93	87	72	80	101	131					
			DOP (µg L <sup>-1</sup> )	10	4.57	4.57	3.90	4.15	4.85	5.33					
			Chl-a (µg L <sup>-1</sup> )	10	0.31	0.30	0.14	0.18	0.38	0.52	H	Mean	0.45		
			Chl-a (µg L <sup>-1</sup> )	10	0.31	0.30	0.14	0.18	0.38	0.52	H	Median		0.32	0.63
			NH <sub>4</sub> (µg L <sup>-1</sup> )	10	3.96	3.01	1.96	2.08	6.50	6.76					
			NO <sub>x</sub> (µg L <sup>-1</sup> )	10	1.16	0.67	0.28	0.31	1.76	2.90	H	Median	0.35		
			PN (µg L <sup>-1</sup> )	10	19.86	20.39	11.23	13.08	25.82	29.86	H	Mean	20		
			PN (µg L <sup>-1</sup> )	10	19.86	20.39	11.23	13.08	25.82	29.86	H	Median		16	25
			PO <sub>4</sub> (µg L <sup>-1</sup> )	10	2.73	2.86	1.60	2.23	3.28	3.58	H	Median	2		

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			POC (mg L <sup>-1</sup> )	10	119	108	63	96	159	174					
			PP (µg L <sup>-1</sup> )	10	2.63	2.35	2.00	2.15	3.19	3.48	H	Mean	2.8		
			PP (µg L <sup>-1</sup> )	10	2.63	2.35	2.00	2.15	3.19	3.48	H	Median		2.3	3.3
			Secchi (m)	10	9.50	9.00	6.18	7.80	12.00	13.10	L	Mean	10		
			Secchi (m)	10	9.50	9.00	6.18	7.80	12.00	13.10	L	Median			
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	10	284	234	82	129	352	621					
			TSS (mg L <sup>-1</sup> )	10	1.47	1.22	0.68	0.90	1.66	3.05	H	Mean	2		
			TSS (mg L <sup>-1</sup> )	10	1.47	1.22	0.68	0.90	1.66	3.05	H	Median		1.6	2.4
		Between Tam O'Shanter and Timana (TUL6)	DIN (µg L <sup>-1</sup> )	10	5.77	5.25	2.02	4.18	7.60	10.44					
			DOC (µg L <sup>-1</sup> )	10	1157	1144	943	971	1309	1398					
			DON (µg L <sup>-1</sup> )	10	95	92	76	86	108	117					
			DOP (µg L <sup>-1</sup> )	10	4.65	4.57	3.63	3.96	5.26	5.88					
			Chl-a (µg L <sup>-1</sup> )	10	0.45	0.40	0.15	0.24	0.64	0.84	H	Mean	0.45		
			Chl-a (µg L <sup>-1</sup> )	10	0.45	0.40	0.15	0.24	0.64	0.84	H	Median		0.32	0.63
			NH <sub>4</sub> (µg L <sup>-1</sup> )	10	4.19	4.52	1.57	3.46	4.89	6.09					
			NO <sub>x</sub> (µg L <sup>-1</sup> )	10	1.58	0.86	0.28	0.39	2.74	4.35	H	Median	0.35		
			PN (µg L <sup>-1</sup> )	10	27.29	28.97	13.82	15.85	35.53	39.34	H	Mean	20		
			PN (µg L <sup>-1</sup> )	10	27.29	28.97	13.82	15.85	35.53	39.34	H	Median		16	25
			PO <sub>4</sub> (µg L <sup>-1</sup> )	10	2.65	2.56	1.93	2.14	3.05	3.47	H	Median	2		

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			POC (mg L <sup>-1</sup> )	10	184	193	104	136	230	245					
			PP (µg L <sup>-1</sup> )	10	3.43	2.99	2.16	2.62	4.42	5.57	H	Mean	2.8		
			PP (µg L <sup>-1</sup> )	10	3.43	2.99	2.16	2.62	4.42	5.57	H	Median		2.3	3.3
			Secchi (m)	10	5.50	6.00	2.23	3.70	7.10	8.33	L	Mean	10		
			Secchi (m)	10	5.50	6.00	2.23	3.70	7.10	8.33	L	Median			
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	10	522	488	126	297	736	1024					
			TSS (mg L <sup>-1</sup> )	10	1.91	1.22	0.68	0.93	2.41	4.88	H	Mean	2		
			TSS (mg L <sup>-1</sup> )	10	1.91	1.22	0.68	0.93	2.41	4.88	H	Median		1.6	2.4
		Bedarra (TUL8)	DIN (µg L <sup>-1</sup> )	10	5.41	5.18	2.18	3.47	6.76	10.02					
			DOC (µg L <sup>-1</sup> )	10	1132	1095	924	951	1302	1413					
			DON (µg L <sup>-1</sup> )	10	90	90	70	81	94	110					
			DOP (µg L <sup>-1</sup> )	10	4.31	4.14	3.44	3.55	4.80	5.73					
			Chl-a (µg L <sup>-1</sup> )	10	0.35	0.35	0.13	0.14	0.44	0.61	H	Mean	0.45		
			Chl-a (µg L <sup>-1</sup> )	10	0.35	0.35	0.13	0.14	0.44	0.61	H	Median		0.32	0.63
			NH <sub>4</sub> (µg L <sup>-1</sup> )	10	3.07	2.73	1.54	1.99	4.28	5.36					
			NO <sub>x</sub> (µg L <sup>-1</sup> )	10	2.35	2.08	0.39	0.55	3.72	5.57	H	Median	0.35		
			PN (µg L <sup>-1</sup> )	10	24.22	20.95	12.95	16.18	33.77	41.84	H	Mean	20		
			PN (µg L <sup>-1</sup> )	10	24.22	20.95	12.95	16.18	33.77	41.84	H	Median		16	25
			PO <sub>4</sub> (µg L <sup>-1</sup> )	10	2.56	2.63	1.52	1.98	3.11	3.39	H	Median	2		

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			POC (mg L <sup>-1</sup> )	10	147	142	81	107	184	219					
			PP (µg L <sup>-1</sup> )	10	3.28	2.93	2.16	2.51	3.95	5.06	H	Mean	2.8		
			PP (µg L <sup>-1</sup> )	10	3.28	2.93	2.16	2.51	3.95	5.06	H	Median		2.3	3.3
			Secchi (m)	10	6.78	7.00	4.23	4.90	8.20	9.00	L	Mean	10		
			Secchi (m)	10	6.78	7.00	4.23	4.90	8.20	9.00	L	Median			
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	10	606	529	123	374	861	1125					
			TSS (mg L <sup>-1</sup> )	10	2.14	0.93	0.46	0.64	4.37	5.33	H	Mean	2		
			TSS (mg L <sup>-1</sup> )	10	2.14	0.93	0.46	0.64	4.37	5.33	H	Median		1.6	2.4
		Tully River mouth mooring (TUL10)	DIN (µg L <sup>-1</sup> )	10	10.41	10.90	2.55	4.01	15.41	17.36					
			DOC (µg L <sup>-1</sup> )	10	1198	1233	957	998	1383	1476					
			DON (µg L <sup>-1</sup> )	10	98	98	82	89	106	116					
			DOP (µg L <sup>-1</sup> )	10	4.48	4.45	3.32	3.84	5.36	5.42					
			Chl-a (µg L <sup>-1</sup> )	10	0.64	0.64	0.20	0.28	1.08	1.12	H	Median	1.1		
			NH <sub>4</sub> (µg L <sup>-1</sup> )	10	4.08	3.66	1.65	2.84	4.96	7.70					
			NO <sub>x</sub> (µg L <sup>-1</sup> )	10	6.33	6.35	0.54	1.24	10.67	11.29	H	Median	3		
			PN (µg L <sup>-1</sup> )	10	38.62	38.23	18.71	24.13	50.19	62.10	H	Median			
			PO <sub>4</sub> (µg L <sup>-1</sup> )	10	2.70	2.48	2.08	2.17	3.02	4.01	H	Median	3		
			POC (mg L <sup>-1</sup> )	10	265	250	145	197	336	419					
			PP (µg L <sup>-1</sup> )	10	5.23	5.42	2.97	4.09	6.42	7.31	H	Median			

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			Secchi (m)	10	4.63	5.00	1.95	2.50	6.60	7.55	L	Median	1.6		
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	10	759	841	215	378	1004	1241					
			TSS (mg L <sup>-1</sup> )	10	3.70	2.94	1.12	1.60	4.49	9.18	H	Median	5		

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
Burdekin		Palms West (BUR1)	DIN (µg L <sup>-1</sup> )	9	4.97	4.62	3.09	3.79	6.12	7.57					
			DOC (µg L <sup>-1</sup> )	9	940	938	880	891	976	1009					
			DON (µg L <sup>-1</sup> )	9	92	93	78	87	98	110					
			DOP (µg L <sup>-1</sup> )	9	5.22	4.96	4.38	4.66	5.19	7.09					
			Chl-a (µg L <sup>-1</sup> )	9	0.29	0.27	0.20	0.24	0.33	0.41	H	Median	0.35	0.32	0.63
			NH <sub>4</sub> (µg L <sup>-1</sup> )	9	3.61	3.43	2.49	2.95	4.53	4.97					
			NO <sub>x</sub> (µg L <sup>-1</sup> )	9	1.37	1.30	0.52	0.62	1.86	2.70	H	Median	0.28		
			PN (µg L <sup>-1</sup> )	9	23.37	22.68	11.48	13.21	29.87	39.26	H	Median	12	16	25
			PO <sub>4</sub> (µg L <sup>-1</sup> )	9	2.40	2.25	1.64	2.06	2.94	3.27	H	Median	1		
			POC (mg L <sup>-1</sup> )	9	123	133	69	83	152	167					
			PP (µg L <sup>-1</sup> )	9	2.04	1.98	1.68	1.81	2.32	2.53	H	Median	2.2	2.3	3.3
			Secchi (m)	9	7.49	5.50	5.04	5.46	7.32	15.30	L	Mean	10		
			Secchi (m)	9	7.49	5.50	5.04	5.46	7.32	15.30	L	Median			

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	9	104	104	41	60	147	169					
			TSS (mg L <sup>-1</sup> )	9	0.87	0.94	0.37	0.57	1.15	1.29	H	Median	1.2	1.6	2.4
		Pandora (BUR2)	DIN (µg L <sup>-1</sup> )	9	5.46	5.08	3.56	4.42	6.82	7.66					
			DOC (µg L <sup>-1</sup> )	9	1014	1049	882	957	1063	1096					
			DON (µg L <sup>-1</sup> )	9	102	100	88	95	108	122					
			DOP (µg L <sup>-1</sup> )	9	4.92	4.96	4.04	4.46	5.48	5.67					
			Chl-a (µg L <sup>-1</sup> )	9	0.26	0.30	0.14	0.15	0.35	0.40	H	Median	0.35	0.32	0.63
			NH <sub>4</sub> (µg L <sup>-1</sup> )	9	4.34	4.10	2.24	3.14	5.80	6.59					
			NO <sub>x</sub> (µg L <sup>-1</sup> )	9	1.11	0.84	0.34	0.50	1.22	2.73	H	Median	0.28		
			PN (µg L <sup>-1</sup> )	9	24.98	20.41	13.06	13.32	35.83	37.59	H	Median	12	16	25
			PO <sub>4</sub> (µg L <sup>-1</sup> )	9	2.52	2.48	1.77	1.95	3.13	3.41	H	Median	1		
			POC (mg L <sup>-1</sup> )	9	142	147	81	94	186	203					
			PP (µg L <sup>-1</sup> )	9	2.98	2.73	1.52	2.12	3.65	4.97	H	Median	2.2	2.3	3.3
			Secchi (m)	9	5.12	4.70	2.48	3.68	6.70	9.10	L	Mean	10		
			Secchi (m)	9	5.12	4.70	2.48	3.68	6.70	9.10	L	Median			
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	9	175	125	78	101	280	343					
			TSS (mg L <sup>-1</sup> )	9	1.72	1.21	0.60	0.72	2.89	3.91	H	Median	1.2	1.6	2.4
		Magnetic (BUR4)	DIN (µg L <sup>-1</sup> )	9	4.94	3.50	2.53	2.87	5.21	10.96					
			DOC (µg L <sup>-1</sup> )	9	1071	1074	943	1002	1132	1222					

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values					
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet	
			DON (µg L <sup>-1</sup> )	9	99	97	84	89	102	123						
			DOP (µg L <sup>-1</sup> )	9	4.79	4.88	3.84	4.54	5.14	5.42						
			Chl-a (µg L <sup>-1</sup> )	9	0.53	0.50	0.24	0.40	0.59	0.97	H	Median	0.59	0.32	0.63	
			NH4 (µg L <sup>-1</sup> )	9	3.31	2.59	1.98	2.13	3.85	6.33						
			NO <sub>x</sub> (µg L <sup>-1</sup> )	9	1.63	1.05	0.43	0.60	1.83	4.65	H	Median	0.28			
			PN (µg L <sup>-1</sup> )	9	29.77	26.04	15.39	17.34	40.15	47.38	H	Median	17	16	25	
			PO <sub>4</sub> (µg L <sup>-1</sup> )	9	2.19	2.01	0.93	1.30	3.00	3.42	H	Median	1			
			POC (mg L <sup>-1</sup> )	9	183	165	112	123	251	272						
			PP (µg L <sup>-1</sup> )	9	3.48	3.50	2.47	2.92	3.75	4.88	H	Mean	2.8			
			PP (µg L <sup>-1</sup> )	9	3.48	3.50	2.47	2.92	3.75	4.88	H	Median		2.3	3.3	
			Secchi (m)	9	3.71	4.00	2.50	3.22	4.08	4.68	L	Median	4			
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	9	220	143	92	117	354	462						
			TSS (mg L <sup>-1</sup> )	9	1.90	1.84	1.02	1.12	2.23	3.45	H	Median	1.9	1.6	2.4	
		Haughton 2 (BUR7)		DIN (µg L <sup>-1</sup> )	8	4.58	3.75	2.04	2.27	7.17	8.48					
				DOC (µg L <sup>-1</sup> )	8	1057	1026	895	920	1178	1304					
			DON (µg L <sup>-1</sup> )	8	98	95	83	88	105	118						
			DOP (µg L <sup>-1</sup> )	8	4.69	4.72	3.58	4.24	5.39	5.62						
			Chl-a (µg L <sup>-1</sup> )	8	0.41	0.43	0.17	0.21	0.59	0.62	H	Mean	0.45			
			Chl-a (µg L <sup>-1</sup> )	8	0.41	0.43	0.17	0.21	0.59	0.62	H	Median		0.32	0.63	

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			NH <sub>4</sub> (µg L <sup>-1</sup> )	8	3.34	2.87	1.72	1.95	4.76	6.01					
			NO <sub>x</sub> (µg L <sup>-1</sup> )	8	1.24	0.88	0.32	0.32	2.37	2.50	H	Median	1		
			PN (µg L <sup>-1</sup> )	8	24.40	24.76	11.77	14.06	32.61	39.73	H	Median	13	16	25
			PO <sub>4</sub> (µg L <sup>-1</sup> )	8	2.49	2.32	1.30	1.77	3.39	3.94	H	Median	2		
			POC (mg L <sup>-1</sup> )	8	146	126	84	103	185	236					
			PP (µg L <sup>-1</sup> )	8	3.51	3.06	2.17	2.33	4.73	5.38	H	Median	2.1	2.3	3.3
			Secchi (m)	8	4.16	4.25	2.56	3.44	5.10	5.70	L	Mean	10		
			Secchi (m)	8	4.16	4.25	2.56	3.44	5.10	5.70	L	Median			
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	8	194	108	77	84	289	501					
			TSS (mg L <sup>-1</sup> )	8	1.67	1.40	0.52	0.97	1.50	3.83	H	Median	1.2	1.6	2.4
	Yongala (BUR10)		DIN (µg L <sup>-1</sup> )	4	2.88	2.77	2.09	2.15	3.56	3.83					
			DOC (µg L <sup>-1</sup> )	4	949	932	878	899	992	1042					
			DON (µg L <sup>-1</sup> )	4	86	85	79	81	90	93					
			DOP (µg L <sup>-1</sup> )	4	4.80	4.84	4.00	4.13	5.48	5.55					
			Chl-a (µg L <sup>-1</sup> )	4	0.25	0.24	0.09	0.10	0.40	0.43	H	Median	0.33	0.32	0.63
			NH <sub>4</sub> (µg L <sup>-1</sup> )	4	2.50	2.31	1.76	1.77	3.16	3.52					
			NO <sub>x</sub> (µg L <sup>-1</sup> )	4	0.38	0.37	0.29	0.30	0.45	0.48	H	Median	0.28		
			PN (µg L <sup>-1</sup> )	4	13.08	12.13	9.80	10.27	15.52	17.71	H	Median	14	16	25
			PO <sub>4</sub> (µg L <sup>-1</sup> )	4	1.70	1.66	0.95	1.02	2.37	2.51	H	Median	1		

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			POC (mg L <sup>-1</sup> )	4	85	75	56	57	108	127					
			PP (µg L <sup>-1</sup> )	4	1.58	1.49	1.02	1.18	1.95	2.26	H	Median	2	2.3	3.3
			Secchi (m)	4	15.50	15.75	12.88	14.00	17.10	17.78	L	Mean	10		
			Secchi (m)	4	15.50	15.75	12.88	14.00	17.10	17.78	L	Median			
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	4	50	42	24	28	69	88					
			TSS (mg L <sup>-1</sup> )	4	0.18	0.16	0.07	0.11	0.24	0.31	H	Median	0.8	1.6	2.4
		Haughton River mouth (BUR8)	DIN (µg L <sup>-1</sup> )	1	11.20	11.20	11.20	11.20	11.20	11.20					
			DOC (µg L <sup>-1</sup> )	1	1057	1057	1057	1057	1057	1057					
			DON (µg L <sup>-1</sup> )	1	101	101	101	101	101	101					
			DOP (µg L <sup>-1</sup> )	1	4.96	4.96	4.96	4.96	4.96	4.96					
			Chl-a (µg L <sup>-1</sup> )	1	0.70	0.70	0.70	0.70	0.70	0.70	H	Median	1		
			NH <sub>4</sub> (µg L <sup>-1</sup> )	1	6.72	6.72	6.72	6.72	6.72	6.72					
			NO <sub>x</sub> (µg L <sup>-1</sup> )	1	4.48	4.48	4.48	4.48	4.48	4.48	H	Median	4		
			PN (µg L <sup>-1</sup> )	1	55.72	55.72	55.72	55.72	55.72	55.72	H	Median			
			PO <sub>4</sub> (µg L <sup>-1</sup> )	1	2.48	2.48	2.48	2.48	2.48	2.48	H	Median	1		
			POC (mg L <sup>-1</sup> )	1	300	300	300	300	300	300					
			PP (µg L <sup>-1</sup> )	1	9.20	9.20	9.20	9.20	9.20	9.20	H	Median			
			Secchi (m)	1	1.00	1.00	1.00	1.00	1.00	1.00	L	Median	1.5		
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	1	243	243	243	243	243	243					

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			TSS (mg L <sup>-1</sup> )	1	18.02	18.02	18.02	18.02	18.02	18.02	H	Median	2		
		Burdekin River mouth mooring (BUR13)	DIN (µg L <sup>-1</sup> )	9	11.12	4.87	3.15	4.01	19.32	28.33					
			DOC (µg L <sup>-1</sup> )	9	1195	1200	959	986	1359	1503					
			DON (µg L <sup>-1</sup> )	9	101	100	93	97	104	112					
			DOP (µg L <sup>-1</sup> )	9	5.05	4.88	4.48	4.71	5.48	5.81					
			Chl-a (µg L <sup>-1</sup> )	9	0.70	0.69	0.47	0.60	0.81	0.94	H	Median	1		
			NH <sub>4</sub> (µg L <sup>-1</sup> )	9	6.53	3.99	2.49	3.10	10.34	13.93					
			NO <sub>x</sub> (µg L <sup>-1</sup> )	9	4.59	1.12	0.46	0.71	8.21	14.79	H	Median	4		
			PN (µg L <sup>-1</sup> )	9	45.59	49.53	19.44	24.05	62.96	75.31	H	Mean			
			PN (µg L <sup>-1</sup> )	9	45.59	49.53	19.44	24.05	62.96	75.31	H	Median			
			PO <sub>4</sub> (µg L <sup>-1</sup> )	9	3.31	3.41	0.98	1.39	4.86	6.77	H	Median	1		
			POC (mg L <sup>-1</sup> )	9	286	278	138	174	407	469					
			PP (µg L <sup>-1</sup> )	9	6.79	5.95	2.84	4.38	9.39	11.54	H	Median			
			Secchi (m)	9	2.36	2.10	0.78	1.32	3.40	4.30	L	Median	1.5		
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	9	428	265	114	128	883	940					
			TSS (mg L <sup>-1</sup> )	9	6.80	3.53	1.12	1.93	11.44	15.95	H	Median	2		

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
Mackay-Whitsunday		Double Cone (WHI1)	DIN ( $\mu\text{g L}^{-1}$ )	5	3.60	3.61	2.77	3.00	4.20	4.41					
			DOC ( $\mu\text{g L}^{-1}$ )	5	897	898	856	874	918	940					
			DON ( $\mu\text{g L}^{-1}$ )	5	85	86	80	83	87	87					
			DOP ( $\mu\text{g L}^{-1}$ )	5	4.55	4.57	3.64	4.10	4.83	5.62					
			Chl-a ( $\mu\text{g L}^{-1}$ )	5	0.33	0.32	0.21	0.21	0.38	0.52	H	Median	0.36	0.32	0.63
			NH <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	5	2.75	2.84	2.32	2.37	2.99	3.24					
			NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	5	0.85	0.77	0.29	0.31	1.31	1.56	H	Median	1		
			PN ( $\mu\text{g L}^{-1}$ )	5	15.10	14.25	11.88	12.65	17.29	19.40	H	Mean	14		
			PN ( $\mu\text{g L}^{-1}$ )	5	15.10	14.25	11.88	12.65	17.29	19.40	H	Median		16	25
			PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	5	2.34	2.09	1.38	1.56	2.73	3.93	H	Median	1		
			POC (mg L <sup>-1</sup> )	5	127	125	91	93	161	164					
			PP ( $\mu\text{g L}^{-1}$ )	5	2.79	2.60	1.91	2.08	3.27	4.07	H	Median	2.3	2.3	3.3
			Secchi (m)	5	5.50	6.00	4.00	4.00	6.30	7.20	L	Mean	10		
			Secchi (m)	5	5.50	6.00	4.00	4.00	6.30	7.20	L	Median			
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	5	63	61	42	46	75	92					
			TSS (mg L <sup>-1</sup> )	5	1.57	1.14	0.93	1.03	2.26	2.46	H	Median	1.4	1.6	2.4
		Pine (WHI4)	DIN ( $\mu\text{g L}^{-1}$ )	5	8.33	7.67	4.52	4.65	9.84	14.98					
			DOC ( $\mu\text{g L}^{-1}$ )	5	899	887	854	869	924	959					
			DON ( $\mu\text{g L}^{-1}$ )	5	90	95	78	84	97	97					

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			DOP ( $\mu\text{g L}^{-1}$ )	5	4.43	4.49	3.84	3.98	4.85	4.99					
			Chl-a ( $\mu\text{g L}^{-1}$ )	5	0.43	0.46	0.26	0.32	0.53	0.59	H	Median	0.36	0.32	0.63
			NH <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	5	3.94	4.10	2.59	3.12	4.67	5.24					
			NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	5	4.39	3.19	1.53	1.93	5.47	9.82	H	Median	1		
			PN ( $\mu\text{g L}^{-1}$ )	5	14.25	13.51	10.57	11.76	15.78	19.66	H	Mean	14		
			PN ( $\mu\text{g L}^{-1}$ )	5	14.25	13.51	10.57	11.76	15.78	19.66	H	Median		16	25
			PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	5	2.77	2.94	1.30	1.72	3.67	4.23	H	Median	1		
			POC (mg L <sup>-1</sup> )	5	117	115	83	96	128	161					
			PP ( $\mu\text{g L}^{-1}$ )	5	3.40	2.65	1.85	2.42	3.80	6.27	H	Median	2.3	2.3	3.3
			Secchi (m)	5	5.70	4.50	2.40	3.60	8.40	9.60	L	Mean	10		
			Secchi (m)	5	5.70	4.50	2.40	3.60	8.40	9.60	L	Median			
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	5	79	83	51	65	90	103					
			TSS (mg L <sup>-1</sup> )	5	2.69	1.75	0.50	0.79	3.44	7.00	H	Median	1.4	1.6	2.4
		Seaforth (WHI5)	DIN ( $\mu\text{g L}^{-1}$ )	5	4.82	4.62	2.40	2.78	6.78	7.50					
			DOC ( $\mu\text{g L}^{-1}$ )	5	911	914	842	867	941	992					
			DON ( $\mu\text{g L}^{-1}$ )	5	87	86	78	82	95	96					
			DOP ( $\mu\text{g L}^{-1}$ )	5	4.21	4.49	2.96	3.47	5.05	5.09					
			Chl-a ( $\mu\text{g L}^{-1}$ )	5	0.44	0.36	0.25	0.33	0.49	0.75	H	Median	0.36	0.32	0.63
			NH <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	5	2.74	2.63	1.84	2.11	3.34	3.80					

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			NO <sub>x</sub> (µg L <sup>-1</sup> )	5	2.07	0.70	0.55	0.64	4.05	4.42	H	Median	1		
			PN (µg L <sup>-1</sup> )	5	14.92	13.48	12.70	12.77	17.39	18.28	H	Mean	14		
			PN (µg L <sup>-1</sup> )	5	14.92	13.48	12.70	12.77	17.39	18.28	H	Median		16	25
			PO <sub>4</sub> (µg L <sup>-1</sup> )	5	2.43	1.94	1.63	1.86	2.83	3.90	H	Median	1		
			POC (mg L <sup>-1</sup> )	5	112	96	79	89	141	156					
			PP (µg L <sup>-1</sup> )	5	3.25	2.47	2.29	2.37	4.28	4.84	H	Median	2.3	2.3	3.3
			Secchi (m)	5	6.10	7.00	3.30	4.20	7.40	8.60	L	Mean	10		
			Secchi (m)	5	6.10	7.00	3.30	4.20	7.40	8.60	L	Median			
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	5	76	66	52	59	94	109					
			TSS (mg L <sup>-1</sup> )	5	1.55	1.07	0.74	0.78	2.36	2.80	H	Median	1.4	1.6	2.4
		OConnell River mouth (WHI6)	DIN (µg L <sup>-1</sup> )	5	4.17	2.98	2.28	2.41	6.45	6.71					
			DOC (µg L <sup>-1</sup> )	5	1593	1180	1006	1027	1791	2964					
			DON (µg L <sup>-1</sup> )	5	125	109	97	104	130	184					
			DOP (µg L <sup>-1</sup> )	5	4.85	5.03	3.92	4.29	5.31	5.68					
			Chl-a (µg L <sup>-1</sup> )	5	0.79	0.88	0.49	0.67	0.91	0.99	H	Median	1.3		
			NH <sub>4</sub> (µg L <sup>-1</sup> )	5	2.87	2.07	1.75	1.86	4.26	4.42					
			NO <sub>x</sub> (µg L <sup>-1</sup> )	5	1.30	0.91	0.53	0.55	2.03	2.45	H	Median	4		
			PN (µg L <sup>-1</sup> )	5	41.15	42.53	18.86	21.08	54.61	68.68	H	Mean			
			PN (µg L <sup>-1</sup> )	5	41.15	42.53	18.86	21.08	54.61	68.68	H	Median			

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			PO <sub>4</sub> (µg L <sup>-1</sup> )	5	4.46	3.79	2.57	3.08	5.17	7.68	H	Median	3		
			POC (mg L <sup>-1</sup> )	5	399	389	180	295	461	669					
			PP (µg L <sup>-1</sup> )	5	9.68	8.29	4.52	5.62	13.50	16.48	H	Median			
			Secchi (m)	5	1.70	2.00	0.60	0.90	2.20	2.80	L	Median	1.6		
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	5	528	217	54	99	666	1604					
			TSS (mg L <sup>-1</sup> )	5	12.34	4.90	2.56	3.18	17.73	33.31	H	Median	5		
		Repulse Islands dive mooring (WHI7)	DIN (µg L <sup>-1</sup> )	5	6.57	3.92	3.45	3.72	7.34	14.44					
			DOC (µg L <sup>-1</sup> )	5	1044	951	875	924	1121	1351					
			DON (µg L <sup>-1</sup> )	5	109	97	94	95	114	144					
			DOP (µg L <sup>-1</sup> )	5	5.02	4.80	3.73	4.48	5.67	6.41					
			Chl-a (µg L <sup>-1</sup> )	5	0.51	0.37	0.31	0.35	0.62	0.88	H	Mean	0.45		
			Chl-a (µg L <sup>-1</sup> )	5	0.51	0.37	0.31	0.35	0.62	0.88	H	Median		0.32	0.63
			NH <sub>4</sub> (µg L <sup>-1</sup> )	5	4.81	2.49	2.36	2.40	6.11	10.69					
			NO <sub>x</sub> (µg L <sup>-1</sup> )	5	1.76	1.44	0.50	0.83	2.09	3.96	H	Median	0.25		
			PN (µg L <sup>-1</sup> )	5	20.52	19.87	14.21	17.53	25.23	25.76	H	Median	18	16	25
			PO <sub>4</sub> (µg L <sup>-1</sup> )	5	2.68	1.86	1.52	1.66	4.13	4.23	H	Median	2		
			POC (mg L <sup>-1</sup> )	5	174	177	100	143	218	231					
			PP (µg L <sup>-1</sup> )	5	4.94	3.60	2.82	3.13	7.14	8.02	H	Median	2.1	2.3	3.3
			Secchi (m)	5	3.90	4.00	2.00	2.00	5.30	6.20	L	Mean	10		

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			Secchi (m)	5	3.90	4.00	2.00	2.00	5.30	6.20	L	Median			
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	5	163	80	71	71	230	363					
			TSS (mg L <sup>-1</sup> )	5	4.50	2.38	1.67	1.91	7.30	9.23	H	Median	1.6	1.6	2.4

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
Fitzroy		North Keppel Island (FTZ4)	DIN (µg L <sup>-1</sup> )	9	5.57	5.36	2.70	3.51	7.69	8.55					
			DOC (µg L <sup>-1</sup> )	9	1045	1084	937	1004	1091	1118					
			DON (µg L <sup>-1</sup> )	9	100	101	85	91	106	119					
			DOP (µg L <sup>-1</sup> )	9	4.75	4.72	3.84	4.30	5.16	5.67					
			Chl-a (µg L <sup>-1</sup> )	9	0.34	0.25	0.13	0.18	0.48	0.69	H	Mean	0.45		
			Chl-a (µg L <sup>-1</sup> )	9	0.34	0.25	0.13	0.18	0.48	0.69	H	Median		0.32	0.63
			NH <sub>4</sub> (µg L <sup>-1</sup> )	9	4.60	5.08	2.09	2.99	6.07	6.68					
			NO <sub>x</sub> (µg L <sup>-1</sup> )	9	0.97	0.74	0.32	0.47	1.12	2.55	H	Median	0.5		
			PN (µg L <sup>-1</sup> )	9	16.10	13.14	11.50	12.27	20.25	24.47	H	Median	15	16	25
			PO <sub>4</sub> (µg L <sup>-1</sup> )	9	1.05	0.93	0.62	0.67	1.22	1.97	H	Median	2		
			POC (mg L <sup>-1</sup> )	9	126	109	81	87	145	222					
			PP (µg L <sup>-1</sup> )	9	2.44	2.12	1.43	1.88	2.86	4.02	H	Median	2.5	2.3	3.3
			Secchi (m)	9	8.72	8.50	5.50	7.00	11.20	12.40	L	Median	10		

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	9	40	42	14	29	54	63					
			TSS (mg L <sup>-1</sup> )	9	0.69	0.48	0.21	0.38	0.93	1.68	H	Median	1	1.6	2.4
		Barren (FTZ1)	DIN (µg L <sup>-1</sup> )	9	6.14	6.27	2.84	3.62	8.26	10.28					
			DOC (µg L <sup>-1</sup> )	9	1115	1098	929	1029	1151	1365					
			DON (µg L <sup>-1</sup> )	9	105	98	86	89	110	140					
			DOP (µg L <sup>-1</sup> )	9	5.08	5.26	3.84	4.68	5.54	5.92					
			Chl-a (µg L <sup>-1</sup> )	9	0.27	0.19	0.11	0.13	0.38	0.59	H	Median	0.27	0.32	0.63
			NH <sub>4</sub> (µg L <sup>-1</sup> )	9	5.36	5.99	2.31	3.15	6.71	8.79					
			NO <sub>x</sub> (µg L <sup>-1</sup> )	9	0.78	0.49	0.28	0.34	0.63	2.29	H	Median	0.5		
			PN (µg L <sup>-1</sup> )	9	13.83	12.73	10.06	10.72	15.06	20.92	H	Median	12	16	25
			PO <sub>4</sub> (µg L <sup>-1</sup> )	9	1.15	1.01	0.34	0.53	1.83	2.15	H	Median	2		
			POC (mg L <sup>-1</sup> )	9	103	95	58	68	111	187					
			PP (µg L <sup>-1</sup> )	9	2.22	1.92	1.29	1.48	2.37	4.21	H	Median	1.9	2.4	3.4
			Secchi (m)	9	10.67	11.00	6.20	8.00	12.00	15.60	L	Median	12		
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	9	44	34	24	27	56	87					
			TSS (mg L <sup>-1</sup> )	9	0.65	0.24	0.11	0.17	1.06	2.14	H	Median	0.4	1.7	2.5
		Keppels South (FTZ2)	DIN (µg L <sup>-1</sup> )	9	6.14	5.11	3.34	4.33	8.31	10.09					
			DOC (µg L <sup>-1</sup> )	9	1118	1151	912	1005	1205	1272					
			DON (µg L <sup>-1</sup> )	9	105	100	81	90	116	141					

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			DOP ( $\mu\text{g L}^{-1}$ )	9	5.06	5.11	4.35	4.82	5.36	5.64					
			Chl-a ( $\mu\text{g L}^{-1}$ )	9	0.29	0.29	0.16	0.23	0.36	0.42	H	Mean	0.45		
			Chl-a ( $\mu\text{g L}^{-1}$ )	9	0.29	0.29	0.16	0.23	0.36	0.42	H	Median		0.32	0.63
			NH <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	9	5.07	4.76	2.65	3.89	6.52	7.46					
			NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	9	1.07	0.63	0.31	0.35	1.79	2.63	H	Median	0.5		
			PN ( $\mu\text{g L}^{-1}$ )	9	15.80	14.00	11.93	13.15	20.03	20.65	H	Mean	20		
			PN ( $\mu\text{g L}^{-1}$ )	9	15.80	14.00	11.93	13.15	20.03	20.65	H	Median		16	25
			PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	9	0.86	0.93	0.53	0.67	1.02	1.21	H	Mean	2		
			PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	9	0.86	0.93	0.53	0.67	1.02	1.21	H	Median			
			POC ( $\text{mg L}^{-1}$ )	9	115	113	84	93	136	153					
			PP ( $\mu\text{g L}^{-1}$ )	9	2.70	2.49	1.66	2.03	3.30	4.27	H	Mean	2.8		
			PP ( $\mu\text{g L}^{-1}$ )	9	2.70	2.49	1.66	2.03	3.30	4.27	H	Median		2.4	3.4
			Secchi (m)	9	8.78	8.00	5.60	7.10	10.80	12.90	L	Mean	10		
			Secchi (m)	9	8.78	8.00	5.60	7.10	10.80	12.90	L	Median			
			SiO <sub>4</sub> ( $\text{mg L}^{-1}$ )	9	48	50	21	27	60	84					
			TSS ( $\text{mg L}^{-1}$ )	9	1.30	0.46	0.29	0.40	2.15	3.97	H	Mean	2		
			TSS ( $\text{mg L}^{-1}$ )	9	1.30	0.46	0.29	0.40	2.15	3.97	H	Median		1.7	2.5
	Pelican (FTZ3)	DIN ( $\mu\text{g L}^{-1}$ )	9	6.10	5.36	4.14	4.67	7.68	9.09						
		DOC ( $\mu\text{g L}^{-1}$ )	9	1169	1185	1003	1107	1234	1312						

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
			DON ( $\mu\text{g L}^{-1}$ )	9	105	106	84	100	116	125					
			DOP ( $\mu\text{g L}^{-1}$ )	9	4.88	4.88	4.20	4.57	5.16	5.53					
			Chl-a ( $\mu\text{g L}^{-1}$ )	9	0.42	0.38	0.16	0.25	0.53	0.84	H	Mean	0.45		
			Chl-a ( $\mu\text{g L}^{-1}$ )	9	0.42	0.38	0.16	0.25	0.53	0.84	H	Median		0.32	0.63
			NH <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	9	5.29	4.87	3.32	3.82	6.49	7.98					
			NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	9	0.81	0.67	0.35	0.48	1.19	1.38	H	Median	0.5		
			PN ( $\mu\text{g L}^{-1}$ )	9	21.05	19.67	13.73	18.00	21.79	32.67	H	Mean	20		
			PN ( $\mu\text{g L}^{-1}$ )	9	21.05	19.67	13.73	18.00	21.79	32.67	H	Median		16	25
			PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	9	1.69	1.63	0.70	1.07	2.35	2.77	H	Mean	2		
			PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	9	1.69	1.63	0.70	1.07	2.35	2.77	H	Median			
			POC ( $\text{mg L}^{-1}$ )	9	169	159	94	129	171	296					
			PP ( $\mu\text{g L}^{-1}$ )	9	4.72	3.96	2.04	2.89	4.87	10.24	H	Mean	2.8		
			PP ( $\mu\text{g L}^{-1}$ )	9	4.72	3.96	2.04	2.89	4.87	10.24	H	Median		2.4	3.4
			Secchi (m)	9	4.25	3.75	1.53	2.90	5.30	7.78	L	Mean	10		
			Secchi (m)	9	4.25	3.75	1.53	2.90	5.30	7.78	L	Median			
			SiO <sub>4</sub> ( $\text{mg L}^{-1}$ )	9	77	77	48	64	88	111					
			TSS ( $\text{mg L}^{-1}$ )	9	3.57	1.92	0.32	0.53	5.22	10.16	H	Mean	2		
			TSS ( $\text{mg L}^{-1}$ )	9	3.57	1.92	0.32	0.53	5.22	10.16	H	Median		1.7	2.5
			DIN ( $\mu\text{g L}^{-1}$ )	9	6.21	5.92	3.95	4.85	7.44	9.25					

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
		Peak West (FTZ5)	DOC ( $\mu\text{g L}^{-1}$ )	9	1130	1150	979	1076	1187	1237					
			DON ( $\mu\text{g L}^{-1}$ )	9	101	104	84	92	110	117					
			DOP ( $\mu\text{g L}^{-1}$ )	9	4.71	4.80	4.10	4.20	5.11	5.34					
			Chl-a ( $\mu\text{g L}^{-1}$ )	9	0.48	0.46	0.26	0.33	0.55	0.85	H	Mean	0.45		
			Chl-a ( $\mu\text{g L}^{-1}$ )	9	0.48	0.46	0.26	0.33	0.55	0.85	H	Median		0.32	0.63
			NH <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	9	5.08	5.22	3.36	4.39	5.72	6.45					
			NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	9	1.13	0.56	0.29	0.34	1.37	3.58	H	Median	0.5		
			PN ( $\mu\text{g L}^{-1}$ )	9	24.12	20.73	15.02	18.97	30.69	37.14	H	Mean	20		
			PN ( $\mu\text{g L}^{-1}$ )	9	24.12	20.73	15.02	18.97	30.69	37.14	H	Median		16	25
			PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	9	1.99	1.70	0.59	0.91	3.11	3.53	H	Median	2		
			POC (mg L <sup>-1</sup> )	9	209	192	115	144	271	341					
			PP ( $\mu\text{g L}^{-1}$ )	9	5.53	4.82	2.20	3.36	7.33	9.75	H	Mean	2.8		
			PP ( $\mu\text{g L}^{-1}$ )	9	5.53	4.82	2.20	3.36	7.33	9.75	H	Median		2.4	3.4
			Secchi (m)	9	3.39	2.50	1.20	1.80	5.50	7.00	L	Mean	10		
			Secchi (m)	9	3.39	2.50	1.20	1.80	5.50	7.00	L	Median			
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	9	83	80	56	62	88	133					
			TSS (mg L <sup>-1</sup> )	9	6.38	5.33	1.59	3.12	10.19	13.72	H	Mean	2		
			TSS (mg L <sup>-1</sup> )	9	6.38	5.33	1.59	3.12	10.19	13.72	H	Median		1.7	2.5
			DIN ( $\mu\text{g L}^{-1}$ )	9	16.84	14.25	6.90	8.49	25.94	28.73					

Region	Subregion	Site	Measure	N	Mean	Median	Quantiles				Guideline Values				
							Q5	Q20	Q80	Q95	DOF	Location	Annual	Dry	Wet
		Fitzroy River Mouth (FTZ6)	DOC ( $\mu\text{g L}^{-1}$ )	9	1253	1249	1093	1153	1338	1409					
			DON ( $\mu\text{g L}^{-1}$ )	9	107	110	83	103	117	119					
			DOP ( $\mu\text{g L}^{-1}$ )	9	4.47	4.57	3.69	4.29	4.83	4.92					
			Chl-a ( $\mu\text{g L}^{-1}$ )	9	0.76	0.75	0.55	0.67	0.84	0.98	H	Median	1		
			NH <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	9	6.65	5.74	3.63	5.00	8.45	10.91					
			NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	9	10.19	9.35	0.68	3.98	15.81	22.45	H	Median	3		
			PN ( $\mu\text{g L}^{-1}$ )	9	34.63	31.02	21.10	27.19	41.25	52.34	H	Median			
			PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	9	5.55	5.96	2.11	3.50	7.05	8.58	H	Median	3		
			POC (mg L <sup>-1</sup> )	9	346	306	185	247	439	573					
			PP ( $\mu\text{g L}^{-1}$ )	9	10.05	7.73	4.57	6.55	14.83	18.41	H	Median			
			Secchi (m)	9	1.22	1.00	0.54	0.78	1.40	2.60	L	Median			
			SiO <sub>4</sub> (mg L <sup>-1</sup> )	9	154	124	87	94	210	278					
			TSS (mg L <sup>-1</sup> )	9	17.43	13.02	5.91	7.44	27.34	35.18	H	Median			

Table C-2: Summary of turbidity measurements from moored loggers (site locations in [Section 5](#) and Figure D-1 for the past three water years. N = number of daily means in the time-series; SE = standard error; '% d> Trigger' refers to the percentage of days each year with mean or median values above the site-specific water quality guideline values (Table C-9). Red shading indicates the annual means or medians that exceeded annual guideline values (Table C-9). Yellow shading indicates the means or medians that exceeded wet season guideline values for Cape York loggers which collect wet season data only (Table C-9). '% d> 5 NTU' refers to the percentage of days above 5 NTU, a threshold suggested by Cooper *et al.* (2007, 2008) above which hard corals are likely to experience photo-physiological stress.

Subregion	Site	Oct 2021 - Sept 2022						Oct 2022 - Sept 2023						Oct 2023 - Sept 2024					
		N	Annual Mean	SE	Annual Median	%d > Trigger	%d > 5 NTU	N	Annual Mean	SE	Annual Median	%d > Trigger	%d > 5 NTU	N	Annual Mean	SE	Annual Median	%d > Trigger	%d > 5 NTU
Cape York	Dawson	253	1.10	0.08	0.62	37.15	1.98	136	1.36	0.13	0.66	43.4	4.4	150	1.95	0.33	0.85	54.0	4.0
	Forrester	224	0.35	0.01	0.33	20.1	0.00	100	0.3	0.01	0.26	1	0	162	0.42	0.03	0.28	25.3	0.0
Johnstone Russell Mulgrave	Fitzroy West	235	0.94	0.05	0.79	27.66	0.85	277	1.58	0.06	1.19	64.98	2.17	226	1.14	0.13	0.74	33.19	1.77
	High West	365	0.91	0.03	0.84	33.70	0.00	93	0.91	0.05	0.79	29.03	0.00	144	1.06	0.04	0.89	43.75	0.00
	Russell Mulgrave Mouth Mooring	250	4.91	0.25	3.73	98.00	39.20	291	6.14	0.30	4.36	100.00	43.64	252	3.79	0.28	2.38	91.27	21.83
	Franklands West	364	0.77	0.03	0.65	60.99	0.00	360	0.95	0.04	0.76	69.72	0.56	251	0.87	0.06	0.64	54.98	0.80
Tully Herbert	Dunk North	365	2.40	0.15	1.37	78.90	10.68	347	3.59	0.22	1.72	89.91	21.61	167	2.84	0.28	1.55	79.64	11.98
	Tully River mouth mooring	365	4.88	0.18	4.00	50.14	33.70	365	5.76	0.24	4.26	52.33	42.19	202	4.68	0.25	3.96	49.01	23.76
Burdekin	Palms West	326	0.69	0.02	0.63	19.63	0.31	337	0.72	0.03	0.60	24.04	0.30	217	0.97	0.05	0.88	56.68	0.92
	Pandora	365	1.34	0.06	1.05	71.51	2.19	308	1.41	0.08	1.05	73.38	2.92	223	1.55	0.09	1.17	78.92	1.79
	Magnetic	365	1.89	0.12	1.12	38.63	6.30	301	2.42	0.13	1.43	55.48	13.29	249	2.60	0.18	1.75	71.89	8.43
	Burdekin River mouth mooring	352	8.80	0.37	6.68	71.88	60.51	365	10.33	0.39	8.96	78.08	67.67	145	6.67	0.64	4.13	50.34	42.76
Mackay Whitsunday	Double Cone	364	1.31	0.03	1.19	57.14	0.27	365	1.26	0.03	1.10	49.86	0.00	182	1.63	0.07	1.33	70.33	1.10
	Pine	365	2.25	0.09	1.70	72.88	8.49	251	2.52	0.11	2.02	80.08	10.76	246	2.05	0.11	1.60	63.82	4.88
	Seaforth	276	1.63	0.04	1.47	75.00	0.00	340	1.68	0.04	1.46	78.24	0.29	247	1.69	0.05	1.49	81.78	0.00
	OConnell River mouth	285	5.54	0.33	3.38	40.70	29.82	314	6.24	0.30	4.06	50.64	41.40	245	4.41	0.32	2.79	32.65	27.35
Fitzroy	Barren	365	0.57	0.03	0.40	70.14	0.00	365	0.30	0.01	0.25	30.96	0.00	172	0.56	0.04	0.48	79.65	0.00

Subregion	Site	Oct 2021 - Sept 2022						Oct 2022 - Sept 2023						Oct 2023 - Sept 2024					
		N	Annual Mean	SE	Annual Median	%d > Trigger	%d > 5 NTU	N	Annual Mean	SE	Annual Median	%d > Trigger	%d > 5 NTU	N	Annual Mean	SE	Annual Median	%d > Trigger	%d > 5 NTU
	Keppels South	307	0.90	0.05	0.57	15.31	0.33	365	0.83	0.04	0.57	13.42	0.27	233	1.13	0.08	0.61	19.74	2.15
	Fitzroy River Mouth	332	21.92	0.92	16.01		93.07	280	16.23	0.75	12.40		88.57	203	25.91	1.64	18.12		93.10

**C-4 Data used to generate remote sensing maps**

Table C-3: Summary of water quality data collected across the Sentinel-3 Reef water types (WT) as part of the wet season event sampling of the MMP. Multi-year samples were collected between December–April by AIMS and CYWP since 2016–17 and by JCU since 2003–04 and up to 2022–23.

Multi-year		TSS (mg L <sup>-1</sup> )	Chl- <i>a</i> (µg L <sup>-1</sup> )	CDOM (m <sup>-1</sup> )	Secchi depth (m)	DIN (µg L <sup>-1</sup> )	DIP (µg L <sup>-1</sup> )	PP (µg L <sup>-1</sup> )	PN (µg L <sup>-1</sup> )	
Reef region	WT1	mean	17.40	1.45	0.86	2.11	31.60	8.74	11.60	76.90
		SD	44.10	2.21	0.99	2.12	44.40	11.90	20.60	82.60
		min	0.00	0.07	0.00	0.00	0.24	0.00	0.00	0.00
		max	590.00	30.90	6.17	16.00	338.00	98.00	186.28	719.00
		count	935	935	789	501	878	895	259	248
	WT2	mean	4.63	0.65	0.25	4.61	12.10	4.40	3.75	37.50
		SD	7.00	0.75	0.41	2.77	23.50	5.22	4.01	23.40
		min	0.00	0.02	0.00	0.20	0.12	0.00	0.00	0.00
		max	130.00	12.50	3.50	17.00	363.00	63.00	58.00	210.50
		count	1383	1414	1116	932	1384	1401	611	602
	WT3	mean	2.35	0.37	0.09	8.22	6.51	2.76	2.46	28.30
		SD	3.94	0.40	0.18	4.14	10.50	3.24	3.80	21.30
		min	0.00	0.02	0.00	0.50	0.12	0.31	0.00	0.00
		max	31.00	5.34	2.00	22.00	77.00	21.00	49.00	268.00
		count	623	639	457	418	581	585	338	329
	WT4	mean	1.78	0.53	0.12	11.00	8.81	4.24	2.37	28.50
		SD	3.55	0.86	0.20	4.76	10.70	4.84	2.52	19.40
		min	0.00	0.02	0.00	0.50	0.14	0.31	0.77	8.36
		max	30.00	5.34	1.25	20.00	63.00	20.00	15.30	119.24
		count	93	101	58	54	83	84	44	42

Table C-4: Summary of water quality data collected in the Cape York region across the Sentinel-3 Reef water types (WT) as part of the wet season event sampling of the MMP. Multi-year samples were collected between December and April by CYWP since 2016–17 and up to 2022–23.

Cape York	Multi-year		TSS (mg L <sup>-1</sup> )	Chl- <i>a</i> (µg L <sup>-1</sup> )	CDOM (m <sup>-1</sup> )	Secchi depth (m)	DIN (µg L <sup>-1</sup> )	DIP (µg L <sup>-1</sup> )	PP (µg L <sup>-1</sup> )	PN (µg L <sup>-1</sup> )
	WT1									
		mean	20.36	1.10	1.65	2.21	13.37	3.49	10.66	85.03
		SD	44.13	1.13	1.59	2.23	14.64	2.58	12.64	87.21
		min	0.00	0.08	0.00	0.10	0.28	0.00	0.00	0.00
		max	320.00	8.82	6.17	12.00	73.14	13.32	68.00	563.78
		count	208	218	160	157	214	218	102	80
	WT2	mean	3.33	0.45	0.39	5.04	5.00	1.95	3.69	31.94
		SD	5.61	0.47	0.64	2.93	7.85	1.47	6.11	21.15
		min	0.00	0.04	0.00	0.20	0.28	0.31	0.00	8.84
		max	60.00	3.26	3.50	16.00	76.86	8.00	58.00	144.93
		count	295	301	180	225	301	301	188	170
	WT3	mean	1.53	0.27	0.08	8.54	3.64	1.67	2.47	22.18
		SD	2.00	0.31	0.24	3.81	6.99	1.46	5.83	12.89
		min	0.05	0.02	0.00	0.65	0.28	0.31	0.00	8.84
		max	14.00	1.95	2.00	17.50	68.00	7.14	49.00	76.71
		count	176	181	109	126	178	178	120	111
	WT4	mean	1.43	0.22	0.03	10.35	8.20	3.00	1.63	20.75
		SD	0.80	0.35	0.05	6.15	7.11	1.50	1.34	7.23
		min	0.41	0.02	0.00	2.50	0.28	0.31	0.77	13.84
		max	2.66	1.28	0.10	19.40	18.14	5.56	4.00	27.86
		count	13	13	4	8	13	13	5	4

Table C-5: Summary of water quality data collected in the Wet Tropics region across the Sentinel-3 Reef water types (WT) as part of the wet season event sampling of the MMP. Samples were collected between December and April by AIMS since 2016–17 and JCU since 2003–04 and up to and up to 2022–23.

Multi-year		TSS (mg L <sup>-1</sup> )	Chl- <i>a</i> (μg L <sup>-1</sup> )	CDOM (m <sup>-1</sup> )	Secchi depth (m)	DIN (μg L <sup>-1</sup> )	DIP (μg L <sup>-1</sup> )	PP (μg L <sup>-1</sup> )	PN (μg L <sup>-1</sup> )	
Wet Tropics	WT1	mean	8.66	1.27	0.64	2.05	45.66	7.13	8.86	53.22
		SD	9.71	1.84	0.51	2.13	57.32	5.12	8.50	29.84
		min	0.00	0.08	0.00	0.00	0.24	0.00	1.25	12.84
		max	92.00	30.90	3.56	14.00	338.00	21.00	40.86	173.71
		count	406	399	388	185	356	356	57	58
	WT2	mean	4.23	0.67	0.27	4.80	17.07	4.87	3.44	37.66
		SD	4.91	0.65	0.39	2.87	31.34	4.60	2.33	23.05
		min	0.00	0.02	0.00	0.50	0.12	0.00	0.53	7.74
		max	33.00	11.24	2.74	17.00	363.00	22.00	19.52	210.50
		count	623	637	574	400	611	615	228	229
	WT3	mean	2.84	0.42	0.11	8.77	9.23	3.28	2.34	28.46
		SD	4.79	0.49	0.16	4.53	13.20	3.60	1.72	13.68
		min	0.00	0.02	0.00	0.50	0.12	0.31	0.66	0.00
		max	31.00	5.34	1.38	22.00	77.00	21.00	13.53	100.19
		count	289	296	239	203	273	274	143	143
	WT4	mean	2.07	0.43	0.17	10.99	10.42	3.25	3.06	27.07
		SD	2.57	0.61	0.25	4.47	14.31	3.05	3.59	14.40
		min	0.01	0.10	0.00	1.30	0.14	0.31	0.93	8.36
		max	11.94	3.42	1.25	19.00	63.00	12.00	15.29	54.91
		count	33	35	29	25	33	33	19	19

Table C-6: Summary of water quality data collected in the Burdekin region across the Sentinel-3 Reef water types (WT) as part of the wet season event sampling of the MMP. Multi-years samples were collected between December and April by AIMS since 2016–17 and JCU since 2003–04 and up to 2022–23.

Burdekin	Multi-year		TSS (mg L <sup>-1</sup> )	Chl- <i>a</i> (µg L <sup>-1</sup> )	CDOM (m <sup>-1</sup> )	Secchi depth (m)	DIN (µg L <sup>-1</sup> )	DIP (µg L <sup>-1</sup> )	PP (µg L <sup>-1</sup> )	PN (µg L <sup>-1</sup> )
	WT1	mean	32.00	1.43	0.66	2.03	24.28	6.78	18.43	98.16
		SD	83.22	1.84	0.84	2.23	37.50	6.92	36.89	113.01
		min	0.05	0.10	0.00	0.00	0.28	0.31	1.27	15.23
		max	590.00	13.78	3.48	16.00	323.00	46.00	186.28	719.00
		count	157	156	113	102	151	155	63	73
	WS2	mean	3.99	0.66	0.13	3.98	9.22	3.34	3.92	46.03
		SD	8.47	0.82	0.25	2.30	17.98	3.64	3.33	29.68
		min	0.05	0.07	0.00	0.20	0.12	0.31	0.97	0.00
		max	130.00	8.69	2.03	16.00	223.38	27.90	26.35	182.88
		count	258	260	194	202	258	260	99	106
	WS3	mean	2.47	0.37	0.08	6.16	5.38	3.11	2.62	36.03
		SD	2.71	0.22	0.15	3.41	5.86	3.42	3.06	41.72
		min	0.05	0.08	0.00	1.00	0.14	0.31	0.91	0.00
		max	16.00	1.14	1.11	17.00	31.00	20.00	20.57	268.00
		count	97	96	71	61	81	82	40	40
	WS4	mean	1.77	0.34	0.06	11.21	4.38	2.45	1.90	31.46
		SD	5.35	0.38	0.09	4.76	5.37	2.96	0.95	24.87
		min	0.00	0.10	0.00	0.50	0.14	0.31	0.95	11.14
		max	30.00	2.20	0.33	20.00	19.00	9.00	4.64	119.24
count		33	39	23	21	28	29	20	19	

Table C-7: Summary of water quality data collected in the Mackay-Whitsunday region across the Sentinel-3 Reef water types (WT) as part of the wet season event sampling of the MMP. Multi-year samples were collected between December and April by AIMS since 2016–17 and JCU since 2003–04 and up to 2022–23.

Multi-year		TSS (mg L <sup>-1</sup> )	Chl- <i>a</i> (µg L <sup>-1</sup> )	CDOM (m <sup>-1</sup> )	Secchi depth (m)	DIN (µg L <sup>-1</sup> )	DIP (µg L <sup>-1</sup> )	PP (µg L <sup>-1</sup> )	PN (µg L <sup>-1</sup> )	
Mackay-Whitsundays	WT1	mean	12.44	1.26	0.45	1.88	21.12	10.89	6.85	62.20
		SD	20.59	1.22	0.56	1.41	16.50	7.43	2.56	26.91
		min	0.57	0.07	0.03	0.20	0.28	0.31	2.85	27.24
		max	110.00	6.78	2.60	6.00	68.04	36.23	12.35	115.28
		count	45	42	43	28	45	45	20	20
	WT2	mean	5.44	0.82	0.14	3.42	10.30	4.38	4.91	41.11
		SD	6.79	0.58	0.15	2.06	11.54	3.35	2.20	16.54
		min	0.10	0.03	0.01	0.40	0.14	0.00	1.83	16.43
		max	41.00	3.88	0.88	11.00	52.78	15.00	10.74	79.36
		count	134	129	98	73	127	132	74	75
	WT3	mean	1.56	0.49	0.02	6.48	2.57	1.69	2.98	44.30
		SD	2.10	0.24	0.01	1.97	4.02	1.28	0.97	29.15
		min	0.19	0.10	0.01	4.00	0.12	0.31	1.72	16.54
		max	12.00	1.19	0.05	10.50	22.00	7.00	5.57	136.91
		count	39	39	27	20	33	34	27	27

Table C-8: Summary of water quality data collected in the Fitzroy region across the Sentinel-3 Reef water types (WT) as part of the wet season event sampling of the MMP. Multi-year samples were collected between December and April by AIMS since 2021–22 and JCU since 2007–08 and up to 2022–23.

Fitzroy	Multi-year		TSS (mg L <sup>-1</sup> )	Chl- <i>a</i> (µg L <sup>-1</sup> )	CDOM (m <sup>-1</sup> )	Secchi depth (m)	DIN (µg L <sup>-1</sup> )	DIP (µg L <sup>-1</sup> )	PP (µg L <sup>-1</sup> )	PN (µg L <sup>-1</sup> )
	WT1	mean	7.63	1.30	0.29	2.52	15.10	10.27	6.62	45.53
		SD	10.41	1.88	0.36	1.39	25.42	12.46	4.73	21.59
		min	0.69	0.24	0.04	0.25	0.28	0.93	2.53	26.74
		max	39.88	8.73	1.41	6.00	116.83	57.76	22.64	97.76
		count	22	22	17	22	22	22	17	17
	WT2	mean	1.73	0.53	0.05	6.20	4.30	3.69	2.79	24.11
		SD	1.10	0.60	0.03	2.98	7.27	4.81	1.08	7.29
		min	0.17	0.11	0.01	2.50	0.28	0.31	1.28	14.04
		max	3.60	2.21	0.13	12.00	22.00	19.00	4.92	39.88
		count	27	27	22	27	27	27	22	22
	WT3	mean	0.49	0.22	0.02	8.91	0.77	0.87	1.95	17.92
		SD	0.38	0.09	0.02	2.22	0.21	0.55	0.59	5.09
		min	0.05	0.10	0.00	5.50	0.49	0.31	1.27	12.94
		max	1.70	0.35	0.05	11.50	0.98	1.70	3.21	29.44
		count	16	16	7	8	8	8	8	8

### C-5 Site-specific guideline values for MMP sites

Table C-9: Site-specific guideline values (GVs) used for comparison with water quality monitoring data. These GV's are used to calculate the annual condition version of the WQ Index for each water quality sampling location and were provided by the Reef Authority. GV's are derived from the Water Quality Guidelines for the Great Barrier Reef Marine Park (Great Barrier Reef Marine Park Authority, 2010, see Table B-1). Basin-level water quality objectives can be accessed online (Great Barrier Reef Marine Park Authority, Water quality guidelines for the Great Barrier Reef). Seasonal guideline values (i.e., wet vs. dry) are calculated as described in De'ath and Fabricius (2008). Guideline values for the Cape York region come from State of Queensland, (2020). See Appendix B for details on WQ Index calculation. DOF is direction of failure ('H' = high values fail, while 'L' = low values fail). Annual mean GV's are applied to annual mean values of monitoring data (and median GV's are applied to median data). Bold GV's are those applied to monitoring data.

GBRMPA group	GBRMPA sites	Measure	Water body	DOF	Annual Mean	Annual Median	Dry Median	Wet Median
30	ER01, AR01, PRN01, PRS01	Chl- <i>a</i> ( $\mu\text{g L}^{-1}$ )	Enclosed Coastal waters	H				<b>0.70</b>
		NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	Enclosed Coastal waters	H				<b>1.50</b>
		PN ( $\mu\text{g L}^{-1}$ )	Enclosed Coastal waters					
		PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	Enclosed Coastal waters	H				<b>3.00</b>
		PP ( $\mu\text{g L}^{-1}$ )	Enclosed Coastal waters					
		Secchi (m)	Enclosed Coastal waters	L				<b>3.00</b>
		TSS ( $\text{mg L}^{-1}$ )	Enclosed Coastal waters	H				<b>4.00</b>
		Turbidity (NTU)	Enclosed Coastal waters	H		<b>10.00</b>		
40	KR01, KR02, BR01, NR01, NR02	Chl- <i>a</i> ( $\mu\text{g L}^{-1}$ )	Enclosed Coastal waters	H				<b>0.70</b>
		NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	Enclosed Coastal waters	H				<b>1.00</b>
		PN ( $\mu\text{g L}^{-1}$ )	Enclosed Coastal waters					
		PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	Enclosed Coastal waters	H				<b>2.00</b>
		PP ( $\mu\text{g L}^{-1}$ )	Enclosed Coastal waters					
		Secchi (m)	Enclosed Coastal waters	L				<b>1.50</b>
		TSS ( $\text{mg L}^{-1}$ )	Enclosed Coastal waters	H				<b>6.00</b>
		Turbidity (NTU)	Enclosed Coastal waters	H		<b>11.00</b>		
50	ER02b, ER03, AR02b, AR03b, CI01, NR03, NR04, SR03, SR04, PRN02, PRN03, PRS02, PRS03, PRS2.5	Chl- <i>a</i> ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H		0.36	<b>0.25</b>	<b>0.46</b>
		NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H		0.35	<b>0.32</b>	<b>0.45</b>
		PN ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H		18.00		<b>20.00</b>
		PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H		1.40	<b>1.86</b>	<b>0.93</b>
		PP ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H		2.60		<b>3.00</b>
		Secchi (m)	Open Coastal waters	L	<b>10.00</b>			
		TSS ( $\text{mg L}^{-1}$ )	Open Coastal waters	H		1.90		<b>1.70</b>
		Turbidity (NTU)	Open Coastal waters	H			<b>0.90</b>	<b>0.80</b>
60	AE04, ER05, NR05, SR05, SR06, PRN04, PRN05, PRN06, PRBB, PRS05	Chl- <i>a</i> ( $\mu\text{g L}^{-1}$ )	Mid-shelf waters	H		<b>0.27</b>		
		NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	Mid-shelf waters	H		<b>0.35</b>		
		PN ( $\mu\text{g L}^{-1}$ )	Mid-shelf waters	H		<b>18.00</b>		

GBRMPA group	GBRMPA sites	Measure	Water body	DOF	Annual Mean	Annual Median	Dry Median	Wet Median
		PO <sub>4</sub> (µg L <sup>-1</sup> )	Mid-shelf waters	H		<b>0.62</b>		
		PP (µg L <sup>-1</sup> )	Mid-shelf waters	H		<b>2.00</b>		
		Secchi (m)	Mid-shelf waters	L	<b>10.00</b>			
		TSS (mg L <sup>-1</sup> )	Mid-shelf waters	H		<b>1.50</b>		
		Turbidity (NTU)	Mid-shelf waters	H		<b>0.50</b>		
70	NR06, ER06	Chl- <i>a</i> (µg L <sup>-1</sup> )	Offshore waters	H		<b>0.26</b>		
		NO <sub>x</sub> (µg L <sup>-1</sup> )	Offshore waters	H		<b>0.42</b>		
		PN (µg L <sup>-1</sup> )	Offshore waters	H		<b>16.00</b>		
		PO <sub>4</sub> (µg L <sup>-1</sup> )	Offshore waters	H		<b>0.39</b>		
		PP (µg L <sup>-1</sup> )	Offshore waters	H		<b>1.90</b>		
		Secchi (m)	Offshore waters	L	<b>17.00</b>			
		TSS (mg L <sup>-1</sup> )	Offshore waters	H		<b>0.50</b>		
		Turbidity (NTU)	Offshore waters	H		<b>0.50</b>		
1	C1, C6, C8, RM1, RM4, RM8, TUL1	Chl- <i>a</i> (µg L <sup>-1</sup> )	Open Coastal waters	H	0.45		<b>0.32</b>	<b>0.63</b>
		NO <sub>x</sub> (µg L <sup>-1</sup> )	Open Coastal waters	H		<b>0.35</b>		
		PN (µg L <sup>-1</sup> )	Open Coastal waters	H	20.00		<b>16.00</b>	<b>25.00</b>
		PO <sub>4</sub> (µg L <sup>-1</sup> )	Open Coastal waters	H		<b>2.00</b>		
		PP (µg L <sup>-1</sup> )	Open Coastal waters	H	2.80		<b>2.30</b>	<b>3.30</b>
		Secchi (m)	Open Coastal waters	L	<b>10.00</b>			
		TSS (mg L <sup>-1</sup> )	Open Coastal waters	H	2.00		<b>1.60</b>	<b>2.40</b>
		Turbidity (NTU)	Open Coastal waters	H		<b>1.00</b>		
2	RM9, RM10, TUL3, TUL4, TUL5, TUL6, TUL8, TUL9	Chl- <i>a</i> (µg L <sup>-1</sup> )	Open Coastal waters	H	0.45		<b>0.32</b>	<b>0.63</b>
		NO <sub>x</sub> (µg L <sup>-1</sup> )	Open Coastal waters	H		<b>0.35</b>		
		PN (µg L <sup>-1</sup> )	Open Coastal waters	H	20.00		<b>16.00</b>	<b>25.00</b>
		PO <sub>4</sub> (µg L <sup>-1</sup> )	Open Coastal waters	H		<b>2.00</b>		
		PP (µg L <sup>-1</sup> )	Open Coastal waters	H	2.80		<b>2.30</b>	<b>3.30</b>
		Secchi (m)	Open Coastal waters	L	<b>10.00</b>			
		TSS (mg L <sup>-1</sup> )	Open Coastal waters	H	2.00		<b>1.60</b>	<b>2.40</b>
		Turbidity (NTU)	Open Coastal waters	H		<b>1.00</b>		
3	C4, C5, C11, RM2, RM3, RM5, RM6, RM7, TUL2	Chl- <i>a</i> (µg L <sup>-1</sup> )	Mid-shelf waters	H		0.30	<b>0.32</b>	<b>0.63</b>
		NO <sub>x</sub> (µg L <sup>-1</sup> )	Mid-shelf waters	H		<b>0.31</b>		
		PN (µg L <sup>-1</sup> )	Mid-shelf waters	H		14.00	<b>16.00</b>	<b>25.00</b>
		PO <sub>4</sub> (µg L <sup>-1</sup> )	Mid-shelf waters	H		<b>2.00</b>		

GBRMPA group	GBRMPA sites	Measure	Water body	DOF	Annual Mean	Annual Median	Dry Median	Wet Median
		PP ( $\mu\text{g L}^{-1}$ )	Mid-shelf waters	H		2.00	<b>2.30</b>	<b>3.30</b>
		Secchi (m)	Mid-shelf waters	L		<b>13.00</b>		
		TSS ( $\text{mg L}^{-1}$ )	Mid-shelf waters	H		1.20	<b>1.60</b>	<b>2.40</b>
		Turbidity (NTU)	Mid-shelf waters	H		<b>0.60</b>		
4	RM12, TUL11	Chl- <i>a</i> ( $\mu\text{g L}^{-1}$ )	Mid-estuarine waters	H		<b>2.00</b>		
		NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	Mid-estuarine waters	H		<b>15.00</b>		
		PN ( $\mu\text{g L}^{-1}$ )	Mid-estuarine waters					
		PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	Mid-estuarine waters	H		<b>3.00</b>		
		PP ( $\mu\text{g L}^{-1}$ )	Mid-estuarine waters					
		Secchi (m)	Mid-estuarine waters	L		<b>1.50</b>		
		TSS ( $\text{mg L}^{-1}$ )	Mid-estuarine waters	H		<b>7.00</b>		
		Turbidity (NTU)	Mid-estuarine waters	H		<b>5.00</b>		
5	TUL7, TUL10	Chl- <i>a</i> ( $\mu\text{g L}^{-1}$ )	Lower estuarine waters	H		<b>1.10</b>		
		NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	Lower estuarine waters	H		<b>3.00</b>		
		PN ( $\mu\text{g L}^{-1}$ )	Lower estuarine waters					
		PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	Lower estuarine waters	H		<b>3.00</b>		
		PP ( $\mu\text{g L}^{-1}$ )	Lower estuarine waters					
		Secchi (m)	Lower estuarine waters	L		<b>1.60</b>		
		TSS ( $\text{mg L}^{-1}$ )	Lower estuarine waters	H		<b>5.00</b>		
		Turbidity (NTU)	Lower estuarine waters	H		<b>4.00</b>		
6	BUR1, BUR2	Chl- <i>a</i> ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H		0.35	<b>0.32</b>	<b>0.63</b>
		NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H		<b>0.28</b>		
		PN ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H		12.00	<b>16.00</b>	<b>25.00</b>
		PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H		<b>1.00</b>		
		PP ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H		2.20	<b>2.30</b>	<b>3.30</b>
		Secchi (m)	Open Coastal waters	L	<b>10.00</b>			
		TSS ( $\text{mg L}^{-1}$ )	Open Coastal waters	H		1.20	<b>1.60</b>	<b>2.40</b>
		Turbidity (NTU)	Open Coastal waters	H		<b>0.80</b>		
7	BUR3	Chl- <i>a</i> ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H	0.45		<b>0.32</b>	<b>0.63</b>
		NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H		<b>0.28</b>		
		PN ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H	20.00		<b>16.00</b>	<b>25.00</b>
		PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H		<b>1.00</b>		
		PP ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H	2.80		<b>2.30</b>	<b>3.30</b>

GBRMPA group	GBRMPA sites	Measure	Water body	DOF	Annual Mean	Annual Median	Dry Median	Wet Median
		Secchi (m)	Open Coastal waters	L	<b>10.00</b>			
		TSS (mg L <sup>-1</sup> )	Open Coastal waters	H	2.00		<b>1.60</b>	<b>2.40</b>
		Turbidity (NTU)	Open Coastal waters	H		<b>0.80</b>		
8	BUR4	Chl- <i>a</i> (µg L <sup>-1</sup> )	Open Coastal waters	H		0.59	<b>0.32</b>	<b>0.63</b>
		NO <sub>x</sub> (µg L <sup>-1</sup> )	Open Coastal waters	H		<b>0.28</b>		
		PN (µg L <sup>-1</sup> )	Open Coastal waters	H		17.00	<b>16.00</b>	<b>25.00</b>
		PO <sub>4</sub> (µg L <sup>-1</sup> )	Open Coastal waters	H		<b>1.00</b>		
		PP (µg L <sup>-1</sup> )	Open Coastal waters	H	2.80		<b>2.30</b>	<b>3.30</b>
		Secchi (m)	Open Coastal waters	L		<b>4.00</b>		
		TSS (mg L <sup>-1</sup> )	Open Coastal waters	H		1.90	<b>1.60</b>	<b>2.40</b>
		Turbidity (NTU)	Open Coastal waters	H		<b>1.30</b>		
9	BUR5	Chl- <i>a</i> (µg L <sup>-1</sup> )	Open Coastal waters	H		0.60	<b>0.32</b>	<b>0.63</b>
		NO <sub>x</sub> (µg L <sup>-1</sup> )	Open Coastal waters	H		<b>0.50</b>		
		PN (µg L <sup>-1</sup> )	Open Coastal waters	H	20.00		<b>16.00</b>	<b>25.00</b>
		PO <sub>4</sub> (µg L <sup>-1</sup> )	Open Coastal waters	H		<b>2.00</b>		
		PP (µg L <sup>-1</sup> )	Open Coastal waters	H	2.80		<b>2.30</b>	<b>3.30</b>
		Secchi (m)	Open Coastal waters	L		<b>3.00</b>		
		TSS (mg L <sup>-1</sup> )	Open Coastal waters	H		5.00	<b>1.60</b>	<b>2.40</b>
		Turbidity (NTU)	Open Coastal waters	H		<b>3.00</b>		
10	BUR6, BUR7	Chl- <i>a</i> (µg L <sup>-1</sup> )	Open Coastal waters	H	0.45		<b>0.32</b>	<b>0.63</b>
		NO <sub>x</sub> (µg L <sup>-1</sup> )	Open Coastal waters	H		<b>1.00</b>		
		PN (µg L <sup>-1</sup> )	Open Coastal waters	H		13.00	<b>16.00</b>	<b>25.00</b>
		PO <sub>4</sub> (µg L <sup>-1</sup> )	Open Coastal waters	H		<b>2.00</b>		
		PP (µg L <sup>-1</sup> )	Open Coastal waters	H		2.10	<b>2.30</b>	<b>3.30</b>
		Secchi (m)	Open Coastal waters	L	<b>10.00</b>			
		TSS (mg L <sup>-1</sup> )	Open Coastal waters	H		1.20	<b>1.60</b>	<b>2.40</b>
		Turbidity (NTU)	Open Coastal waters	H	<b>2.00</b>			
11	BUR8, BUR9	Chl- <i>a</i> (µg L <sup>-1</sup> )	Enclosed Coastal waters	H		<b>1.00</b>		
		NO <sub>x</sub> (µg L <sup>-1</sup> )	Enclosed Coastal waters	H		<b>4.00</b>		
		PN (µg L <sup>-1</sup> )	Enclosed Coastal waters					
		PO <sub>4</sub> (µg L <sup>-1</sup> )	Enclosed Coastal waters	H		<b>1.00</b>		
		PP (µg L <sup>-1</sup> )	Enclosed Coastal waters					
		Secchi (m)	Enclosed Coastal waters	L		<b>1.50</b>		

GBRMPA group	GBRMPA sites	Measure	Water body	DOF	Annual Mean	Annual Median	Dry Median	Wet Median
		TSS (mg L <sup>-1</sup> )	Enclosed Coastal waters	H		<b>2.00</b>		
		Turbidity (NTU)	Enclosed Coastal waters	H		<b>4.00</b>		
12	BUR10	Chl- <i>a</i> (µg L <sup>-1</sup> )	Mid-shelf waters	H		0.33	<b>0.32</b>	<b>0.63</b>
		NO <sub>x</sub> (µg L <sup>-1</sup> )	Mid-shelf waters	H		<b>0.28</b>		
		PN (µg L <sup>-1</sup> )	Mid-shelf waters	H		14.00	<b>16.00</b>	<b>25.00</b>
		PO <sub>4</sub> (µg L <sup>-1</sup> )	Mid-shelf waters	H		<b>1.00</b>		
		PP (µg L <sup>-1</sup> )	Mid-shelf waters	H		2.00	<b>2.30</b>	<b>3.30</b>
		Secchi (m)	Mid-shelf waters	L	<b>10.00</b>			
		TSS (mg L <sup>-1</sup> )	Mid-shelf waters	H		0.80	<b>1.60</b>	<b>2.40</b>
		Turbidity (NTU)	Mid-shelf waters	H		<b>0.50</b>		
13	BUR11, BUR12	Chl- <i>a</i> (µg L <sup>-1</sup> )	Open Coastal waters	H	0.45		<b>0.32</b>	<b>0.63</b>
		NO <sub>x</sub> (µg L <sup>-1</sup> )	Open Coastal waters	H		<b>1.00</b>		
		PN (µg L <sup>-1</sup> )	Open Coastal waters	H	20.00		<b>16.00</b>	<b>25.00</b>
		PO <sub>4</sub> (µg L <sup>-1</sup> )	Open Coastal waters	H		<b>2.00</b>		
		PP (µg L <sup>-1</sup> )	Open Coastal waters	H	2.80		<b>2.30</b>	<b>3.30</b>
		Secchi (m)	Open Coastal waters	L	<b>10.00</b>			
		TSS (mg L <sup>-1</sup> )	Open Coastal waters	H	2.00		<b>1.60</b>	<b>2.40</b>
		Turbidity (NTU)	Open Coastal waters	H		<b>2.00</b>		
14	BUR13, BUR14, BUR15	Chl- <i>a</i> (µg L <sup>-1</sup> )	Enclosed Coastal waters	H		<b>1.00</b>		
		NO <sub>x</sub> (µg L <sup>-1</sup> )	Enclosed Coastal waters	H		<b>4.00</b>		
		PN (µg L <sup>-1</sup> )	Enclosed Coastal waters					
		PO <sub>4</sub> (µg L <sup>-1</sup> )	Enclosed Coastal waters	H		<b>1.00</b>		
		PP (µg L <sup>-1</sup> )	Enclosed Coastal waters					
		Secchi (m)	Enclosed Coastal waters	L		<b>1.50</b>		
		TSS (mg L <sup>-1</sup> )	Enclosed Coastal waters	H		<b>2.00</b>		
		Turbidity (NTU)	Enclosed Coastal waters	H		<b>4.00</b>		
15	WHI1, WHI2, WHI3, WHI4, WHI5	Chl- <i>a</i> (µg L <sup>-1</sup> )	Open Coastal waters	H		0.36	<b>0.32</b>	<b>0.63</b>
		NO <sub>x</sub> (µg L <sup>-1</sup> )	Open Coastal waters	H		<b>1.00</b>		
		PN (µg L <sup>-1</sup> )	Open Coastal waters	H	14.00		<b>16.00</b>	<b>25.00</b>
		PO <sub>4</sub> (µg L <sup>-1</sup> )	Open Coastal waters	H		<b>1.00</b>		
		PP (µg L <sup>-1</sup> )	Open Coastal waters	H		2.30	<b>2.30</b>	<b>3.30</b>
		Secchi (m)	Open Coastal waters	L	<b>10.00</b>			
		TSS (mg L <sup>-1</sup> )	Open Coastal waters	H		1.40	<b>1.60</b>	<b>2.40</b>

GBRMPA group	GBRMPA sites	Measure	Water body	DOF	Annual Mean	Annual Median	Dry Median	Wet Median
		Turbidity (NTU)	Open Coastal waters	H		<b>1.10</b>		
16	WHI6	Chl- <i>a</i> ( $\mu\text{g L}^{-1}$ )	Enclosed Coastal waters	H		<b>1.30</b>		
		NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	Enclosed Coastal waters	H		<b>4.00</b>		
		PN ( $\mu\text{g L}^{-1}$ )	Enclosed Coastal waters					
		PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	Enclosed Coastal waters	H		<b>3.00</b>		
		PP ( $\mu\text{g L}^{-1}$ )	Enclosed Coastal waters					
		Secchi (m)	Enclosed Coastal waters	L		<b>1.60</b>		
		TSS ( $\text{mg L}^{-1}$ )	Enclosed Coastal waters	H		<b>5.00</b>		
		Turbidity (NTU)	Enclosed Coastal waters	H		<b>4.00</b>		
17	WHI7, WHI10	Chl- <i>a</i> ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H	0.45		<b>0.32</b>	<b>0.63</b>
		NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H		<b>0.25</b>		
		PN ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H		18.00	<b>16.00</b>	<b>25.00</b>
		PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H		<b>2.00</b>		
		PP ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H		2.10	<b>2.30</b>	<b>3.30</b>
		Secchi (m)	Open Coastal waters	L	<b>10.00</b>			
		TSS ( $\text{mg L}^{-1}$ )	Open Coastal waters	H		1.60	<b>1.60</b>	<b>2.40</b>
		Turbidity (NTU)	Open Coastal waters	H	<b>2.00</b>			
18	WHI8, WHI11	Chl- <i>a</i> ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H	0.45		<b>0.32</b>	<b>0.63</b>
		NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H		<b>1.00</b>		
		PN ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H	20.00		<b>16.00</b>	<b>25.00</b>
		PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H		<b>2.00</b>		
		PP ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H	2.80		<b>2.30</b>	<b>3.30</b>
		Secchi (m)	Open Coastal waters	L	<b>10.00</b>			
		TSS ( $\text{mg L}^{-1}$ )	Open Coastal waters	H	2.00		<b>1.60</b>	<b>2.40</b>
		Turbidity (NTU)	Open Coastal waters	H	<b>2.00</b>			
19	WHI9	Chl- <i>a</i> ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H	0.45		<b>0.32</b>	<b>0.63</b>
		NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H		<b>0.25</b>		
		PN ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H		18.00	<b>16.00</b>	<b>25.00</b>
		PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H		<b>2.00</b>		
		PP ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H		2.10	<b>2.30</b>	<b>3.30</b>
		Secchi (m)	Open Coastal waters	L	<b>10.00</b>			
		TSS ( $\text{mg L}^{-1}$ )	Open Coastal waters	H		1.60	<b>1.60</b>	<b>2.40</b>
		Turbidity (NTU)	Open Coastal waters	H	<b>1.00</b>			

GBRMPA group	GBRMPA sites	Measure	Water body	DOF	Annual Mean	Annual Median	Dry Median	Wet Median
20	WHI10.1, WHI10.2	Chl- <i>a</i> ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H	0.45		<b>0.32</b>	<b>0.63</b>
		NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H		<b>1.00</b>		
		PN ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H	20.00		<b>16.00</b>	<b>25.00</b>
		PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H		<b>2.00</b>		
		PP ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H	2.80		<b>2.30</b>	<b>3.30</b>
		Secchi (m)	Open Coastal waters	L	<b>10.00</b>			
		TSS ( $\text{mg L}^{-1}$ )	Open Coastal waters	H	2.00		<b>1.60</b>	<b>2.40</b>
		Turbidity (NTU)	Open Coastal waters	H			<b>2.00</b>	<b>12.00</b>
	FTZ1	Chl- <i>a</i> ( $\mu\text{g L}^{-1}$ )	Mid-shelf waters	H		0.27	<b>0.32</b>	<b>0.63</b>
		NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	Mid-shelf waters	H		<b>0.50</b>		
		PN ( $\mu\text{g L}^{-1}$ )	Mid-shelf waters	H		12.00	<b>16.00</b>	<b>25.00</b>
		PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	Mid-shelf waters	H		<b>2.00</b>		
		PP ( $\mu\text{g L}^{-1}$ )	Mid-shelf waters	H		1.90	<b>2.40</b>	<b>3.40</b>
		Secchi (m)	Mid-shelf waters	L		<b>12.00</b>		
		TSS ( $\text{mg L}^{-1}$ )	Mid-shelf waters	H		0.40	<b>1.70</b>	<b>2.50</b>
		Turbidity (NTU)	Mid-shelf waters	H		<b>0.30</b>		
	FTZ2, FTZ3	Chl- <i>a</i> ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H	0.45		<b>0.32</b>	<b>0.63</b>
		NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H		<b>0.50</b>		
		PN ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H	20.00		<b>16.00</b>	<b>25.00</b>
		PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H	<b>2.00</b>			
		PP ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H	2.80		<b>2.40</b>	<b>3.40</b>
		Secchi (m)	Open Coastal waters	L	<b>10.00</b>			
		TSS ( $\text{mg L}^{-1}$ )	Open Coastal waters	H	2.00		<b>1.70</b>	<b>2.50</b>
		Turbidity (NTU)	Open Coastal waters	H	<b>1.50</b>			
	FTZ4	Chl- <i>a</i> ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H	0.45		<b>0.32</b>	<b>0.63</b>
		NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H		<b>0.50</b>		
		PN ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H		15.00	<b>16.00</b>	<b>25.00</b>
		PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H		<b>2.00</b>		
		PP ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H		2.50	<b>2.30</b>	<b>3.30</b>
		Secchi (m)	Open Coastal waters	L		<b>10.00</b>		
		TSS ( $\text{mg L}^{-1}$ )	Open Coastal waters	H		1.00	<b>1.60</b>	<b>2.40</b>
		Turbidity (NTU)	Open Coastal waters	H		<b>0.50</b>		
	FTZ5	Chl- <i>a</i> ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H	0.45		<b>0.32</b>	<b>0.63</b>
		NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H		<b>0.50</b>		

GBRMPA group	GBRMPA sites	Measure	Water body	DOF	Annual Mean	Annual Median	Dry Median	Wet Median
		PN ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H	20.00		<b>16.00</b>	<b>25.00</b>
		PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H		<b>2.00</b>		
		PP ( $\mu\text{g L}^{-1}$ )	Open Coastal waters	H	2.80		<b>2.40</b>	<b>3.40</b>
		Secchi (m)	Open Coastal waters	L	<b>10.00</b>			
		TSS ( $\text{mg L}^{-1}$ )	Open Coastal waters	H	2.00		<b>1.70</b>	<b>2.50</b>
		Turbidity (NTU)	Open Coastal waters	H	<b>1.50</b>			
	FTZ6	Chl- <i>a</i> ( $\mu\text{g L}^{-1}$ )	Enclosed Coastal waters	H		<b>1.00</b>		
		NO <sub>x</sub> ( $\mu\text{g L}^{-1}$ )	Enclosed Coastal waters	H		<b>3.00</b>		
		PN ( $\mu\text{g L}^{-1}$ )	Enclosed Coastal waters	H				
		PO <sub>4</sub> ( $\mu\text{g L}^{-1}$ )	Enclosed Coastal waters	H		<b>3.00</b>		
		PP ( $\mu\text{g L}^{-1}$ )	Enclosed Coastal waters	H				
		Secchi (m)	Enclosed Coastal waters	L				
		TSS ( $\text{mg L}^{-1}$ )	Enclosed Coastal waters	H				
		Turbidity (NTU)	Enclosed Coastal waters	H			<b>7.00</b>	<b>15.00</b>

## C-6 Regional exposure assessments for waterbodies



Figure C-3: Areas (in km² and represented as horizontal bars) of seagrass (left) and coral reefs (right) in the a) Reef and regional waterbodies; b) Cape York, c) Wet Tropics, d) Burdekin, e) Mackay-Whitsunday, f) Fitzroy, g) Burnett-Mary. Percentages of total Reef habitats or total regional habitats are indicated in italics.

Table C-10: Areas (km<sup>2</sup>) (and percentages, %) of the Reef lagoon (total 348,839 km<sup>2</sup>) and division by water bodies (WB) (enclosed coastal, EC; open coastal, OC; mid-shelf, Mid; and offshore, Off) affected by the Reef WT1–2 combined, and the three Reef water types individually during the current wet season and for a range of reference periods.

Water type	Water body	Area of Reef affected in km <sup>2</sup> and %									
		2023–24 wet season		Long-term average		Average of coral recovery period (2012–2017)		Typical wet-year composite		Typical dry-year composite	
		km <sup>2</sup>	%	km <sup>2</sup>	%	km <sup>2</sup>	%	km <sup>2</sup>	%	km <sup>2</sup>	%
			(%WB)		(%WB)		(%WB)		(%WB)		(%WB)
WT1+2	Reef	68,303	20%	63,296	18%	58,870	17%	88,326	25%	42,047	12%
	EC	6,364	2%	6,366	2%	7,826	2%	6,367	2%	6,366	2%
			(77%)		(77%)		(95%)		(77%)		(77%)
	OC	30,672	9%	33,955	10%	32,085	9%	34,765	10%	28,672	8%
			(93%)		>99%		(97%)		>99%		(87%)
	Mid	21,426	6%	19,234	6%	16,296	5%	35,707	10%	6,566	2%
			(26%)		(23%)		(20%)		(44%)		(8%)
	Off	9,841	3%	3,741	1%	2,664	1%	11,488	3%	443	0%
			(4%)		(2%)		(1%)		(5%)		(0%)
WT1	Reef	13,974	4%	10,791	3%	10,140	3%	19,856	6%	7,363	2%
	EC	6,144	2%	6,018	2%	6,147	2%	6,199	2%	5,398	2%
			(75%)		(73%)		(75%)		(75%)		(65%)
	OC	7,241	2%	4,773	1%	3,989	1%	11,402	3%	1,965	1%
			(22%)		(14%)		(12%)		(35%)		(6%)
	Mid	574	0%	0	0%	4	0%	2,254	1%	0	0%
			(1%)		(0%)		(0%)		(3%)		(0%)
	Off	16	0%	0	0%	0	0%	0	0%	0	0%
			(0%)		(0%)		(0%)		(0%)		(0%)
WT2	Reef	65,966	19%	59,410	17%	55,074	16%	82,931	24%	39,382	11%
	EC	5,032	1%	5,118	1%	6,431	2%	5,490	2%	4,649	1%
			(61%)		(62%)		(78%)		(67%)		(56%)
	OC	30,637	9%	33,768	10%	31,894	9%	34,759	10%	28,461	8%
			(93%)		>99%		(97%)		>99%		(86%)

Water type	Water body	Area of Reef affected in km <sup>2</sup> and %									
		2023–24 wet season		Long-term average		Average of coral recovery period (2012–2017)		Typical wet-year composite		Typical dry-year composite	
		km <sup>2</sup>	%	km <sup>2</sup>	%	km <sup>2</sup>	%	km <sup>2</sup>	%	km <sup>2</sup>	%
			(%WB)		(%WB)		(%WB)		(%WB)		(%WB)
	Mid	20,550	6%	17,031	5%	14,387	4%	32,193	9%	5,840	2%
			(25%)		(21%)		(18%)		(39%)		(7%)
	Off	9,748	3%	3,493	1%	2,363	1%	10,489	3%	433	0%
			(4%)		(2%)		(1%)		(5%)		(0%)
WT3	Reef	213,491	61%	184,626	53%	165,582	47%	203,859	58%	135,874	39%
	EC	253	0%	36	0%	91	0%	168	0%	31	0%
			(3%)		(0%)		(1%)		(2%)		(0%)
	OC	28,114	8%	26,764	8%	25,620	7%	27,940	8%	25,874	7%
			(85%)		(81%)		(78%)		(85%)		(78%)
	Mid	79,112	23%	75,136	22%	71,728	21%	79,609	23%	53,525	15%
			(96%)		(92%)		(87%)		(97%)		(65%)
	Off	106,013	30%	82,691	24%	68,143	20%	96,142	28%	56,443	16%
			(48%)		(37%)		(31%)		(43%)		(25%)

## C-7 References

- Cooper TF, Uthicke S, Humphrey C, Fabricius KE (2007). Gradients in water column nutrients, sediment parameters, irradiance and coral reef development in the Whitsunday Region, central Great Barrier Reef. *Estuarine, Coastal and Shelf Science* 74:458-470.
- Cooper TF, Ridd PV, Ulstrup KE, Humphrey C, Slivkoff M, Fabricius KE (2008). Temporal dynamics in coral bioindicators for water quality on coastal coral reefs of the Great Barrier Reef. *Marine and Freshwater Research* 59:703-716.
- De'ath G and Fabricius KE (2008) Water quality of the Great Barrier Reef: distributions, effects on reef biota and trigger values for the protection of ecosystem health. Final Report to the Great Barrier Reef Marine Park Authority. Australian Institute of Marine Science, Townsville. 104 pp.
- Great Barrier Reef Marine Park Authority. Water quality guidelines for the Great Barrier Reef. Basin-level objectives for Wet Tropics, Townsville, and Mackay-Whitsundays regions. URL: <http://www.gbrmpa.gov.au/our-work/threats-to-the-reef/declining-water-quality?a=1394>. Accessed Jan 2019.
- Great Barrier Reef Marine Park Authority (2010). Water Quality Guidelines for the Great Barrier Reef Marine Park. Revised Edition 2010. Great Barrier Reef Marine Park Authority, Townsville. 100pp.
- State of Queensland (2020). Environmental Protection (Water and Wetland Biodiversity) Policy 2019: Environmental Values and Water Quality Objectives, Eastern Cape York Basins <https://environment.des.qld.gov.au/management/water/policy/cape-york-eastern-basins>.

## APPENDIX D: WATER QUALITY MONITORING IN THE FITZROY NRM REGION 2023–24



### D-1 Acknowledgements

The Fitzroy Basin Marine Monitoring Program for Inshore Water Quality is funded by the partnership between the Australian Government's Reef Trust and the Great Barrier Reef Foundation, and the Australian Institute of Marine Science.

We acknowledge Aboriginal and Torres Strait Islander Peoples as the Traditional Owners of the places where AIMS works, both on land and in the Sea Country of tropical Australia. We pay our respects to the Elders; past, present, and future; and their continuing culture, beliefs, and spiritual relationships and connection to the land and Sea Country. We would also like to thank the Woppaburra TUMRA Steering Committee and the Darumbal People Aboriginal Corporation RNTBC for consent to access sites located in Woppaburra and Darumbal Sea Country. We would also like to acknowledge the ongoing relationship with the Darumbal TUMRA steering committee.

We are grateful to everyone involved in field and labwork this year including Ulysse Bove, Bruce Brady, Kathy Connellan, Jack Massuger, Daniel Moran, Christian O'Dea, Keegan Vickers, Irena Zagorskis, the staff at Vision Environment, the AIMS Oceanographic Technician team, and the crews of the RV Cape Ferguson and RV Solander. Many thanks to the dedicated staff of the AIMS Analytical Technology Laboratory for analysis of water samples. Thanks also to Seanan Wild and the AIMS Indigenous Partnerships team for support with our engagement activities.

### D-2 Introduction and Background

The Fitzroy Natural Resource Management (NRM) Region extends from Carnarvon Gorge National Park to Rockhampton and out to the mouth of the Fitzroy River. Covering 15.7 million hectares, it has the largest catchment area draining into the Great Barrier Reef (the Reef) (Lewis *et al.*, 2021), equating to ~33% of all suspended sediment load from all the Great Barrier Reef Catchment Area (GBRCA) (Packett *et al.*, 2009). The region has a sub-tropical, semi-arid climate with high inter-annual variability in rainfall, high evaporation rates, and prolonged dry periods followed by infrequent major floods. The most consistent rain usually occurs during the wet season (November–March). Annual rainfall varies substantially across the region, from ~530 mm in the west to ~850 mm in the central and ~2,000 mm in the north-

east coastal area (Packett *et al.*, 2009). The region's rainfall has been decreasing by ~30 mm annually for the past 30 years (Yu *et al.*, 2013).

From 2005 to 2014, water quality monitoring by the Australian Institute of Marine Science (AIMS) occurred three times per year at three sites in the Fitzroy region under the Marine Monitoring Program (MMP) and results were published annually (Thompson *et al.*, 2014, previous reports cited therein). A program re-design of the MMP occurred in 2015 to increase the number of sites and the sampling frequency across the focus regions being monitored at that time. Due to funding constraints, this new program design could not be extended to the Fitzroy region, so monitoring in this region by AIMS ceased at the end of 2014.

A partnership between the Great Barrier Reef Foundation and AIMS began in 2020 to re-establish marine water quality monitoring in the Fitzroy region. The program design for this monitoring follows the same design principles as the MMP in other NRM regions (see [Appendix A](#)) and was funded until 2024.

The primary land use in The Fitzroy Region is cattle grazing (75%), followed by nature conservation (8.7%) and forestry (6.6%). This extensive grazing area supports 55% of the cattle in the GBRCA (Lewis *et al.*, 2021). Mining is another prominent industry in the area, accounting for 102,389 hectares of the catchment. The region holds 75% of Queensland's active coal mines and 47% of its gas mines (QLUMP, 2019). This type of land-use requires clearing of vegetation, leaving sediment susceptible to erosion. Erosion of hillslope and streambank soil ends up in local waterways and is transported into the Great Barrier Reef lagoon (Marwick *et al.*, 2014). Much of this sediment is extremely fine and remains suspended in the water, travelling onto coral reefs, seagrass beds and other sensitive marine ecosystems and reducing light availability (Bainbridge *et al.*, 2018). Catchment-derived sediments contain fertilizer and pesticides from agricultural sources. This additional nutrient input generates an increase in macroalgae and can cause algal blooms and eutrophication (Brodie *et al.*, 2011). Additionally, river discharge can increase susceptibility of coral to disease (Bruno *et al.*, 2003; Haapkylä *et al.*, 2011; Kline *et al.*, 2006; Kuntz *et al.*, 2005; Weber *et al.*, 2012; Vega Thurber *et al.*, 2013) and exacerbate coral bleaching (Wooldridge, 2009). This discharge often contains pesticides. Pesticide exposure can inhibit photosynthesis (Gallen *et al.*, 2014) and affect corals, seagrass, fish, and other marine organisms.

Large-volume floods, which occur once every few decades, can deliver catchment sediment and pollution to the mid and outer reef (Devlin *et al.*, 2001). However, small-scale local flooding adversely affects the inshore reef zone at a much higher frequency. The coastal area of the Fitzroy region comprises many inshore reefs and islands. This includes the 19 islands making up the Keppel group. Long-term monitoring of inshore coral reef sites in the Fitzroy region consistently classed these sites as 'very poor' to 'poor' and from 2005 to 2024 (Thompson *et al.*, 2024). Reefs closest to rivers had the lowest coral cover and improved with distance from the coast. Macroalgae abundance was 'high' at almost all reefs in the region.

### D-3 Methods

This Section provides a brief overview of the sampling methods and indicators that are monitored in the Fitzroy region. More details are presented in the main MMP report above and in a separate quality assurance/quality control (QA/QC) report (Great Barrier Reef Marine Park Authority, 2024).

#### Sampling design

This program is designed to measure the annual condition and long-term trends in coastal water quality rather than short-term episodic changes in water quality associated with periods of high river discharge. This type of monitoring is considered 'ambient', which refers to routine

sampling during the wet and dry seasons outside of major flood events. This program design is analogous to ambient monitoring conducted since 2005 under the MMP.

Monitoring site locations were selected along water quality gradients related to exposure to land-based runoff, with sites located with increasing distance from the Fitzroy River mouth and from the coast (Álvarez-Romero *et al.*, 2013). In order to maintain some continuity with the existing 10-year monitoring dataset (2005–2014), the three original sites (FTZ1–3) were re-instated in the current design (Table D-1, Figure D-1).

Table D-1: Description of the Fitzroy water quality sites monitored during 2023–24. Presence of data-logging instruments (turbidity/fluorescence or salinity loggers) is indicated by tick marks.

Site Name (Short Name)	Latitude	Longitude	Turbidity/fluorescence logger	Salinity logger	Number of times sampled (season)	Sampling
Barren Island (FTZ1)*	-23.152	151.069	✓		9 times per year (6 wet season and 3 dry season)	Duplicate samples surface and bottom
Humpy Island (FTZ2)*	-23.217	150.960	✓	✓	9 times per year (6 wet season and 3 dry season)	Duplicate samples surface and bottom
Pelican Island (FTZ3)*	-23.233	150.873			9 times per year (6 wet season and 3 dry season)	Duplicate samples surface and bottom
North Keppel (FTZ4)	-23.092	150.913			9 times per year (6 wet season and 3 dry season)	Duplicate samples surface and bottom
Peak West (FTZ5)	-23.341	150.905			9 times per year (6 wet season and 3 dry season)	Duplicate samples surface and bottom
Fitzroy River Mouth (FTZ6)	-23.475	150.938	✓	✓	9 times per year (6 wet season and 3 dry season)	Duplicate samples surface and bottom

\*Indicates sites that were monitored by AIMS from 2005–2014

From 2005 to 2014, monitoring occurred ~3 times per year at 3 sites in the various MMP monitoring regions including in the Fitzroy region (discontinued in 2015). These data are included in this report.

### Water quality sampling

At each of the sampling locations (Figure D-3, Table D-1), vertical profiles of water salinity and temperature were measured with a Conductivity Temperature Depth (CTD) profiler (Sea-Bird Electronics SBE19plus). CTD profiles are used to characterise the water column and to identify its state of vertical mixing. See the QA/QC report for a detailed description of CTD data processing (Great Barrier Reef Marine Park Authority, 2024).

Immediately following the CTD cast, discrete water samples were collected with Niskin bottles. Samples collected at ambient sites were from the surface (~0.5 m below water surface) and

bottom (~1 m above the seabed) of the water column, whereas for some event-based sampling only surface water samples were collected. Samples from the Niskin bottles were taken in duplicate and were analysed for a suite of water quality parameters (Table 2-1). Detailed descriptions of analytical chemistry techniques can be found in the QA/QC report (Great Barrier Reef Marine Park Authority, 2024). Values of water quality variables presented in this report are depth-weighted means calculated using surface and bottom samples.

Below is a brief description of each of the main water quality variables measured. These definitions are not all-encompassing but are meant to provide a short description of what aspects of water quality they measure and what processes influence the variables:

- **Total Suspended Solids (TSS)** is a measure of the suspended particulate material in the water column. These solids include suspended sediments (sand, silt, and clay), living plankton, and detrital (non-living organic) material. TSS concentrations are affected by oceanographic processes including primary production and resuspension, as well as inputs from other sources such as dredging and land-based runoff.
- **Secchi depth** is a visual measure of water clarity and proxy for light penetration, which is measured using a high-contrast black and white patterned disc called a Secchi disc. The Secchi depth is the average of the vertical disappearance and reappearance depths of the disc, where clarity increases with increasing Secchi depth. Secchi depth is a simple method that has been used for over 150 years, so is excellent for assessing Long-term change and for cross-system comparisons.
- **Turbidity** is a measure of light scattering caused by fine suspended particles, such as sediment, detritus, and plankton. Turbidity is affected by a wide range of factors including oceanographic processes such as resuspension of bottom sediments by wind, waves and currents; river discharge; and anthropogenic factors such as dredging.
- **Chlorophyll a (Chl-a)** concentration is a measure of phytoplankton biomass in a water body. Phytoplankton grow quickly in response to nutrient availability, so elevated values of Chl-a can indicate increased nutrient loading.
- **Dissolved inorganic nutrients (NH<sub>3</sub>, NO<sub>x</sub>, PO<sub>4</sub> and Si)** measure the amount of readily available nutrients for plankton growth in water samples. Inorganic nitrogen (NH<sub>3</sub>, NO<sub>x</sub>) and phosphate (PO<sub>4</sub>) represent around 1% of the nutrient pools in the Reef. The inorganic nutrient pools are affected by a complex range of biogeochemical processes including both natural (for example, plankton uptake, upwelling, nitrogen fixation, and remineralisation) and anthropogenic (for example, dredging and nutrient inputs from changed land use) processes.
- **Particulate nutrients (POC, PN and PP)** are a measure of the suspended material retained on a filter with a pore size of approximately 0.7 µm. This material consists of a minor fraction of living biomass (for example, bacteria, phytoplankton) and a major fraction of detritus (for example, dead cells, faecal pellets). Particulate nutrient concentrations are affected by oceanographic processes (primary production, bacterial production, resuspension, and remineralisation) as well as sources such as dredging and land-based runoff.
- **Dissolved organic carbon (DOC)** is a measure of organic carbon concentrations passing through a filter with a pore size of 0.45 µm. DOC has a complex chemical composition and is used by bacteria as a source of energy. The DOC pool is affected by a range of production and degradation pathways. The sources include primary production by phytoplankton, zooplankton grazing, resuspension events, river runoff, and abiotic breakdown of POC. DOC can be degraded by sunlight.

### ***In situ* loggers**

Continuous *in situ* chlorophyll fluorescence and turbidity were measured using WET Labs ECO FLNTUSB Combination Fluorometer and Turbidity Sensors located at three sites (Table D-1), which were deployed at 5 m below the surface and sampled at 10 min intervals. Water samples for analyses of Chl-*a* and TSS were collected three times per year to calibrate logger fluorescence and turbidity to *in situ* conditions. Diver-operated Niskin bottles were used to sample close to the moored loggers and samples were preserved and analysed in the same manner as ship-based water samples.

Daily averages of the chlorophyll and turbidity time-series are presented in Figure C-1. Annual means and medians of turbidity were also calculated for each site based on the 'water year' (1 October to 30 September) and compared with the water quality guideline value (GV) (Table C-2).

Salinity and temperature loggers (Sea-Bird Electronics SBE37) were deployed at two locations, with one of these being placed on a fixed mooring near the Fitzroy River mouth (Figure D-3, Table D-1). See the QA/QC report (Great Barrier Reef Marine Park Authority, 2024) for detailed descriptions of logger pre- and post-deployment procedures. Site-specific time-series of salinity and temperature can be found in Figure C-2.

### **Data analyses – Summary statistics and trends**

Concentrations of water quality parameters at each sampling occasion were calculated as depth-weighted means by trapezoidal integration of the data from all sampling depths. At most sites, only two vertical points are sampled (i.e., surface and bottom samples), and this method averages these values to derive the depth-weighted mean. Measurements falling below the instrumental detection limit were represented as half the detection limit. Summary statistics for all water quality variables are presented for all monitoring sites in Table C-1. Concentrations were compared to site-specific GVs (Table C-9), which are defined for Chl-*a*, PN, PP, TSS, Secchi depth, NO<sub>x</sub>, and PO<sub>4</sub>. Concentrations of water quality parameters are presented along the sampling transects for each focus region with distance from river mouths. Trends in water quality are represented with generalised additive models, fitted with a maximum of five knots and modelled with a gamma-distributed response and log-link function.

Temporal trends in key water quality variables (Chl-*a*, TSS, Secchi depth, turbidity, NO<sub>x</sub>, PN, PP, DOC, and POC) since 2005 are reported using only open coastal and mid-shelf sites, as GVs for enclosed coastal waters are derived differently and are not available for all variables, creating statistical imbalance.

Generalised additive mixed effects models (GAMMs) were used to decompose each irregularly spaced time-series into its trend cycles (long-term) and periodic (seasonal) components (Wood, 2006). GAMMs are an extension of additive models (which allow flexible modelling of non-linear relationships by incorporating penalised regression spline types of smoothing functions into the estimation process), where the degree of smoothing of each smooth term (and by extension, the estimated degrees of freedom of each smoother) is treated as a random effect and thus estimable via its variance as with other effects in a mixed modelling structure (Wood, 2006).

For each water quality variable within each focus region, the variable was modelled against a thin-plate smoother for date and a cyclical cubic regression spline (maximum of 5 knots) over months within the year. Spatial and temporal autocorrelation in the residuals was addressed by including sampling locations as a random effect and imposing a first-order continuous-time auto-regressive correlation structure (Pinheiro and Bates, 2000). All GAMMs were fitted using the *mgcv* (Wood, 2006, 2011) package in R 4.2.2 (R Core Team, 2022).

No monitoring occurred in the Fitzroy region from 2015–2020. GAMMs are therefore presented with no data through this period and should be interpreted with caution, until more data become available.

### Data analyses – Water Quality Index

The Water Quality Index (WQ Index) is an interpretation tool developed by AIMS to visualise trends in the suite of water quality variables measured, and to compare monitored water quality to existing Water Quality Guidelines (Department of Environment and Resource Management, 2009; Great Barrier Reef Marine Park Authority, 2010). The WQ Index uses a set of five key indicators:

- Water clarity
- Chl-*a* concentrations
- PN concentrations
- PP concentrations
- NO<sub>x</sub> concentrations.

These five indicators are a subset of the comprehensive suite of water quality variables measured in the MMP inshore water quality program. They have been selected because GVs are available for these measures, and they can be considered as relatively robust indicators that integrate a number of bio-physical processes in the coastal ocean.

Details of WQ Index calculation are given in Appendix B.

## D-4 Drivers and pressures influencing water quality in 2023–24

### Coastal development including agriculture

The Fitzroy Region is home to ~235,000 people, just 5% of Queensland's population (FBA, 2018). By area, cattle grazing is the primary land use in the catchment (Brodie *et al.*, 2003) and the initial clearing of vegetation for this purpose marked a significant change in the source and quantity of sediment exported by the Fitzroy River (Hughes *et al.*, 2009). Intensive cultivation of food crops and livestock feed production also contributes to the sediment load in the Fitzroy River (Hughes *et al.*, 2009). The region has over 472,000 ha of grain crops, largely for feeding livestock, and 25,000 ha of cotton (Thorburn and Wilkinson, 2013). Fluctuations in climate, cattle numbers and farming can greatly affect the state and nature of vegetation cover, and therefore, the susceptibility of soils to erosion, which leads to greater runoff of suspended sediments and nutrients. As Australia's cattle production has stayed relatively consistent since 2000 (MLA, 2021), the erosion of soil on cleared grazing and crop land will continue without the adoption of best management practice.

Annual TSS export from the Fitzroy Basin into the Reef lagoon is between 3 and 4.5 million tonnes per year, accounting for ~33% of all annual TSS reaching the Reef lagoon (Dougall *et al.*, 2005). The current best estimate of anthropogenically-derived TSS from the Fitzroy Region is 2.9 million tonnes per year (Kroon *et al.* 2012), which is 3.4 times greater than pre-European levels. The Fitzroy region has the second highest anthropogenically-derived TSS load in the Reef lagoon. TSS samples taken from the Fitzroy River during flood events were highest in areas with substantial intensive agriculture but were highly variable (Packet *et al.*, 2009). Inshore TSS and Chl-*a* annual mean values in the sector regularly exceed the water quality guideline values (Tracey *et al.*, 2017).

Fertilisers are lost from cropping systems and transported to nearby waterways, contributing to dissolved inorganic nitrogen (DIN) and phosphorus (DIP) concentrations in runoff (Brodie *et al.*, 2019). Pristine forested catchments export mostly organic forms of nitrogen, which are

largely unavailable to phytoplankton (Harris, 2001). Runoff enriched with anthropogenically-derived inorganic nutrients is linked to increased primary production and Chl-a concentrations in Reef inner-shelf waters (Wooldridge *et al.*, 2006). Estimated anthropogenic derived total nitrogen (TN) and total phosphorus (TP) in the Fitzroy Region is the highest in the Reef lagoon (TN = 14,000 tonnes  $y^{-1}$ , TP = 4,100 tonnes  $y^{-1}$ ) (Kroon *et al.*, 2012). This nutrient-enriched runoff means the Fitzroy produces 44% of the anthropogenic plume-based Chl-a content in coastal waters (Baird *et al.*, 2021).

The Fitzroy Region has 20% of the mapped seagrass beds in the Reef lagoon and many inshore reefs associated with the Keppel Islands. The main threats to seagrass are reduced light availability and smothering from suspended sediment and increased growth of epiphytic algae from excess nutrients. Seagrass monitoring in the region has rated seagrass sites as 'poor' (McKenzie *et al.*, 2024). Long-term monitoring of inshore coral reefs in the region shows distinct differences in benthic composition in relation to the water quality gradients (Thompson *et al.*, 2024). Coral cover in the Fitzroy has declined overall since 2005 but has seen some improvements following combined acute and chronic disturbances that influenced the region between 2006 and 2015 (Thompson *et al.*, 2024).

Given that benthic communities of the inshore reefs respond to gradients in water quality, especially sedimentation and nutrient availability, improved land management practices can decrease the stress caused by poor water quality on coastal marine ecosystems. Concerns about the water quality in the Reef lagoon led to the formulation of the Reef Plan for catchments adjacent to the Great Barrier Reef World Heritage Area by the Australian and Queensland governments in 2003 (The State of Queensland and Commonwealth of Australia 2003). The Reef Plan was revised and updated in 2009 and 2013 (The State of Queensland and Commonwealth of Australia, 2009; The State of Queensland, 2013), and further developed into The Reef 2050 Water Quality Improvement Plan (WQIP) (The State of Queensland, 2018). This plan set out ambitious targets to improve catchment and coastal water quality and has the aim of building resilience, improve ecosystem health and benefit communities. One of the main proposed actions is the establishment of the Paddock to Reef program, which aims to reduce threats to the health and resilience of the Great Barrier Reef by promoting the adoption of best management practices (Waterhouse *et al.*, 2018).

Best management practice has currently been implemented on 4% of grazing land, 14% of sugarcane land and 73% of banana production land (Great Barrier Reef Marine Park Authority, 2019); however, previous water quality targets were not met. The WQIP estimates the full adoption of class-B best management practice will reduce DIN load by 19%, and by 30% if class-A practices (innovative/aspirational practice) are fully adopted (Waltham *et al.* 2021). Complete adoption of best grazing practices could reduce TSS export from the Great Barrier Reef catchments by ~20% (Thorburn *et al.*, 2013; Thorburn and Wilkinson, 2013). However, there is limited data available on the direct efficacy of best practice strategies on water quality (Carroll *et al.*, 2012), and modelled values are used for improvement estimations (Baird *et al.*, 2021, Dougall *et al.*, 2014). Site surveys showed that practicing zero tillage on cropping land (wheat, sorghum and sunflower) in the Fitzroy basin reduced erosion rates by 75% compared to traditional cropping practices and that at a catchment level, using no tillage can reduce sediment yield by ~50% (Carroll *et al.*, 1997). Reducing nitrogen fertilizer application by 47% by increasing nitrogen use efficiency (Thorburn *et al.*, 2011) reduced nitrogen surplus (excess nitrogen left in the soil) by 60% and reduced DIN in runoff and deep drainage water (Webster *et al.*, 2012).

Several types of erosion occur due to heavy rainfall and poor land management. Hillslope erosion, the erosion of topsoil on hillslopes by water runoff, is primarily affected by cattle stocking rates. Increasing grazing pressure removes vegetation cover from the land, leaving the fertile topsoil vulnerable to erosion. Gully erosion is the incision of flowing surface water, creating deep, unstable channels, often leaving the land unsuitable for agriculture. Gully erosion is often the main source of sediment in floodwater (Wasson *et al.*, 2002; de Vente *et*

*al.*, 2005; Huon *et al.*, 2005). Streambank erosion arises when grazing occurs too close to vulnerable streambanks and riparian cover is not managed. These erosional processes cause fertile land to lose productive sediments, reducing vegetation, further destabilizing the area. Prolonged dry periods followed by intense rainfall cause massive gully erosion.

Sediment erosion can be managed by reducing stocking rates in the more vulnerable areas and slope stabilisation and sediment reinforcement by increasing root bonding (Shen *et al.*, 2017). Various types of restoration have been effective at reducing erosion rates. For example, Hillslope runoff in the Fitzroy Region was 50–90% less in sites with 40–50% ground cover when compared with sites with 10% cover (Owens *et al.*, 2003; Bartley *et al.*, 2006; Hawdon *et al.*, 2008; Silburn *et al.*, 2011). Streambank erosion is greatly reduced by increasing riparian forest buffers. This can reduce erosion rates by 59–91% (Zaimes *et al.*, 2008). This re-introduced vegetation stabilises existing sediments and reduces erosion, but also creates a sediment sink (Askey-Doran *et al.*, 1996; Furnas, 2003). This helps remove excess nutrients from groundwater and overland flooding (Apan *et al.*, 2002) and can capture up to 89% of nitrogen in runoff water (Thibault 1997). Sites along the Fitzroy River with higher riparian condition had better water quality (i.e., lower DOC, TN and dissolved metals), indicating the effectiveness of riparian restoration and management at improving water quality (Chua *et al.*, 2019). Fine sediments contribute 79% of the total nitrogen (TN) reaching the Great Barrier Reef, while DIN accounts for 17% (Kroon *et al.*, 2012). Decreasing erosion rates of agricultural land and the movement of both fine sediment and DIN to local waterways by riparian buffer restoration considerably decreases the TN load of river water (Thorburn and Wilkinson, 2013). However, the Fitzroy Region has only 1.3 million hectares of forested riparian areas, the lowest proportion within the GBRCA. From 2004 to 2008, 12,702 hectares of forested riparian areas were cleared. Sixty three percent of vegetation had been cleared in the region by 1999 and 0.5–0.75% of vegetation in the Fitzroy is currently cleared annually (CRC, 2003).

### **Climate and cyclone activity**

Climate is a major driver of the condition of water quality and ecosystems and can vary substantially between years. It is heavily driven by the El Niño Southern Oscillation (ENSO) cycle. Climate models predict continued warming (IPCC, 2021); increasing intensity of extreme rainfall events and acute disturbances; fewer but more intense tropical cyclones; and more frequent and extreme La Niña and El Niño events (Schaffelke *et al.*, 2017).

Cyclones create waves which resuspend sediment. Several cyclones have impacted the Reef lagoon in the past two decades; most notable are the intense category 4 and 5 systems: TC Larry (2006) and TC Yasi (2011) in the Wet Tropics and Burdekin Regions, and TC Debbie (2017) in the Mackay-Whitsunday Region. The Fitzroy Region was impacted by TC Tasha in 2010 and TC Marcia in 2015, a category 5 severe tropical cyclone. In recent years, there has been very limited cyclone activity in the Reef and none in the Fitzroy Region. The 2023–24 year also showed no cyclone activity directly impacting the Fitzroy region.

Freshwater discharge volumes into the Reef lagoon are closely related to rainfall during the wet season and have a significant influence on coastal water quality. The total annual freshwater discharge for all of the Reef basins relative to long-term medians (based on water year, calculated using the methods described in Section 2.7) is shown in Figure 3-4.

Due to its large catchment area, the Fitzroy Region is prone to flooding. Acute disturbances, such as heavy rainfall and storms, cause increased sediment load entering coastal waters and resuspension of particles already in the environment. In the Fitzroy flood associated with cyclone Joy (1991), extensive mortality of corals in shallow reefs of the Keppel Islands occurred (van Woessik and Done, 1996). This was primarily attributed to an extended period of low salinity at these sites (Brodie and Mitchell, 1992; O'Neill *et al.*, 1992). A period of compounding large flooding events and storms in the Fitzroy Region between 2006 and 2014 instigated a further decline in coral cover on the inshore reefs and the Keppel Islands (Thompson *et al.*, 2014a). Low salinity in the 2011 flooding triggered widespread mortality of

coral at 2 m on Peak, Pelican and Keppels South sites (Thompson *et al.*, 2013). Some sites experienced 100% mortality down to a depth of 8 m (Jones and Berkelmans, 2014). This coincided with high incidences of coral disease, supporting evidence that the confounding effects of low salinity (Haapkylä *et al.*, 2011) and increased organic matter and nutrients, most notably DOC (Brandt *et al.*, 2013; Kline *et al.*, 2006), can initiate coral disease outbreaks. The cumulative stress of high turbidity (low light) and an increase in algal growth from nutrients hinders coral recovery (Diaz-Pulido, 2009; Rogers and Miller, 2006; Roth *et al.*, 2018). Between 2005 and 2014, coral cover declined. Since 2014, coral cover and coral juvenile density have slightly improved but are still classified as 'poor' (Thompson *et al.*, 2024).

Flood plumes in the Reef lagoon rarely reach the mid and outer shelf, however, plumes from large flood events from cyclones can affect these reefs over short time periods. In the Wet Tropics region, the mid shelf reefs are closer to the coast (<30 km), and are more easily reached by flood plumes, for example cyclone Sadie (1994). However, the mid and outer shelf reefs of the Fitzroy Region are further from the coast (90 km to the Capricorn and Bunkers, 210 km to the Swains Reefs from Fitzroy River mouth). Physical conditions, such as wind are key factors in the spread of flood plumes. The prevailing winds in the Fitzroy Region are usually south-easterly, keeping flood plumes close to shore. However, the wind during cyclone Joy (1991) turned to blow offshore, moving the large plume out to the Capricorn and Bunker group, stretching 200 km offshore, lowering salinity on the Capricorn reefs to 27 ppt (Devlin *et al.*, 2001, Preker *et al.* 1992). Events such as this are seen on a scale of multiple decades, while the reefs associated with the Keppel Islands see extreme impacts from river plumes every 4–6 years, and Barren Island, Hummocky Island and Masthead Island every 10 years (Devlin *et al.*, 2001). Vertical distribution of flood plumes largely affects the salinity, turbidity and nutrient content of the top 3 to 4 m (Devlin *et al.*, 2001), but physical conditions, such as high winds can lead to deeper mixing during these extreme events.

Current patterns in the Reef lagoon are seasonal. South-easterly trade winds dominate most of the year, creating a strong north-westward alongshore current movement (Orpin *et al.*, 2010), especially from January to August. Some periods of southerly movement can be seen from August to December depending on prevailing oceanographic conditions (Luick *et al.*, 2007). This transport tends to move small, easily suspended particles discharged from the Fitzroy River northward. Water quality in the Whitsundays region has deteriorated since 2007 and it is speculated that this could be partly due to inputs of flood discharge from the Fitzroy River (Baird *et al.*, 2019, Cantin *et al.*, 2019). In more recent years, there has been a stabilisation and subsequent improvement in conditions observed in the Whitsundays region between 2017 to 2022 (Moran *et al.*, 2023). Analysis of coral cores in the Whitsunday Islands show that Barium/Calcium (Ba/Ca) ratios (which correlate well with flood events) of some sites increased significantly after large flood events in the Fitzroy Region. The Fitzroy River is thought to have the largest and longest-lasting influence of any river on the water quality of the Whitsunday region. However, the Ba/Ca ratios have not increased significantly from flood events before the period of poorer water quality in the Whitsunday region (2007–2017). Anomalous high Ba/Ca ratios at Scawfell do not seem to relate to Fitzroy River discharge. Hence, declining water quality in the area is more likely due to marine resuspension than flood plume inputs (Cantin *et al.* 2019). A high percentage of sediment in various sites around the Whitsunday Islands had a grain size of <63 µm (Thompson *et al.*, 2014b), which is easily resuspended during disturbance events.

The natural occurrence of the ENSO is closely linked to wet and dry periods on the east coast of Australia. Moderate La Niña conditions were observed in 2020–21 and continued through 2021–2023 and there has been minimal direct impact from acute disturbance associated with tropical cyclones. During La Niña, tropical cyclones are typically more common compared to intermediate years, and the first occurrences of cyclones is earlier in the season. This means these years have an increased likelihood of extensive flooding from rain and damage from high winds. Between 2010 and 2012, there was a strong La Niña event, increasing rainfall and

causing extensive flooding (NOAA, 2017). Discharge from the Fitzroy River was considerably higher than the long-term median discharge (2.8 million litres) in 2008 (Devlin, 2008) and 2010–2013 (Jones and Berkelmans, 2014). In 2010–11, it was nearly 38 million litres, reaching a peak mean daily discharge of 1.16 million mega-litres per day over a period of 18 days (Jones and Berkelmans, 2014).

Future changes to the ENSO due to anthropogenic warming are projected, but are difficult to quantify (Collins *et al.*, 2010). Strong ENSO events are predicted to become more common, especially extreme El Niño followed by La Niña events (Cai *et al.*, 2021). Tropical cyclones in the South Pacific are expected to become less frequent in future oceans (Murakami and Sugi, 2010; Sugi *et al.*, 2009; Zhao *et al.*, 2009; Gualdi *et al.*, 2008; Emanuel *et al.*, 2008), although by how much is disputed (Walsh *et al.*, 2012). Cyclone intensity, however, is predicted to increase marginally (Windlansky *et al.*, 2019). El Niño years bring drier weather to the Fitzroy Region, and with it, less vegetation growth to stabilise sediment in the catchment. La Niña years are wetter than normal years. If the frequency of intermediate years decreases, more extended dry years followed by major flood events may lead to an increase in erosion as vegetation will not have the time to recover between extreme dry and wet conditions, leaving more bare sediment that is vulnerable to erosion.

## D-5 Focus region water quality and Water Quality Index

Fringing reefs are formed around continental islands in Keppel Bay, many of which are used extensively for recreational and tourism activities. Six monitoring sites are sampled in this focus region ten times per year during ambient conditions in the dry and wet seasons. The monitoring sites are located in a transect from the river mouth in a north-easterly direction, representing a gradient in water quality. Four sites are located in the open coastal water body, one site is located in the mid-shelf water body, and one site is in enclosed coastal waters (Figure D-1).




Figure D-1: Sampling sites in the Fitzroy focus region, shown with the water body boundaries. Sites FTZ1–FTZ3 were monitored from 2005–2014 under the Marine Monitoring Program.

From 2008–2013, the Fitzroy NRM region experienced several years of intense flooding with annual discharge from the Fitzroy River exceeding the long-term median in 2008, 2010, 2011, 2012, and 2013 (Figure D-2, Figure 3-5). In four of these years, the freshwater discharge was greater than three times the long-term median, with the 2011 flood event being the largest on record (Figure D-3). Annual discharge of the Fitzroy River from 2014–2024 was generally close to or less than the long-term median (Figure D-2).

Annual discharge for the Fitzroy Basin in the 2023–24 water year was just below the long-term median (Figure 5-49; Table 3-1). The combined discharge and loads calculated for the 2023–24 water year were also below the long-term average. Over the 19-year period from 2005–06:

- discharge ranged from 437 GL (2020–21) to 41,736 GL (2010–11)
- TSS loads ranged from 16 kt (2013–14) to 7,000 kt (2010–11)
- DIN loads ranged from 74 t (2020–21) to 3,900 t (2010–11)
- PN loads ranged from 100 t (2020–21) to 17,000 t (2010–11).




Figure D-2: Loads of (A) TSS, DIN and PN and (B) discharge for the Fitzroy Basin from 2006 to 2024. The loads reported here are based on the monitoring data from the Fitzroy River as reported in the GBR Catchment Loads Monitoring Program with a long-term annual mean concentration of the existing data calculated to produce a load for the 2023–24 water year (where monitored load data have not yet been reported). Dotted line represents the long-term median for basin discharge. Note the different scales on the two y-axes.




Figure D-3: Total discharge from rivers in the Fitzroy region including the Fitzroy (gauge 130005A – The Gap) and Calliope (gauge 132001A – Castlehope) rivers and Waterpark creek (gauge 129001A – Byfield). Daily (blue) and water year (1 October to 30 September, red) discharge is shown. Red dashed line represents the long-term median annual discharge.

#### *Ambient water quality and the in situ Water Quality Index*

Water quality showed trends along the sampling transect (cross-shelf gradient in northerly direction). Sites located nearest to the river mouth (distance from river mouth = 0 km) had high concentrations of Chl-a,  $\text{NO}_x$ , TSS, and particulate nutrients (PN and PP), which declined with distance away from the river mouth, reaching low levels in mid-shelf waters (Figure D-4). Secchi depths were low at sites near the river mouth (water clarity was poor) and increased (water clarity improved) with distance from the river mouth. These spatial patterns are generally consistent with those that are typically observed in the region.

Seasonal differences in Chl-a,  $\text{NO}_x$ , PN, and PP were observed. Concentrations of Chl-a and  $\text{NO}_x$  were similar at inshore sites (enclosed coastal) between the wet and dry seasons but seasonal differences became more pronounced further offshore. PP and PN concentrations were higher in the wet season across the sampling transect. TSS and Secchi depth were generally similar along the transect between wet and dry seasons. (Figure D-4). These observed seasonal differences may be driven by strong physical forcing rather than by river discharge events given this was a year of near-median discharge.




Figure D-4: Water quality variables measured during ambient sampling in 2023–24 along the Fitzroy focus region transect. Chlorophyll *a* (Chl-*a*), nitrate/nitrite ( $\text{NO}_x$ ), total suspended solids (TSS), Secchi depth, particulate nitrogen (PN), and particulate phosphorus (PP) are shown with distance from the Fitzroy River mouth. Water body classifications are shown along the x-axes: Enclosed coastal (EC), open coastal (OC) and mid-shelf (MS). Note the y-axes are logarithmic scales. Fitted lines are generalised additive models.

Long-term trends in the water quality variables are shown in Figure D-5 and site-specific water quality results are presented in Table C-1.

During the 2023–24 water year, concentrations of Chl-*a* were below (met) the GV at most sites except Peak West (FTZ5). Concentrations of  $\text{PO}_4$  met GV at all sites except Fitzroy River Mouth (FTZ6) and concentrations of TSS and PN met GV at most sites. Concentrations of PP were just above (did not meet) the GV within the Fitzroy region. All other variables showed a mixed response relative to GV at most sites. Concentrations of  $\text{NO}_x$  exceeded (did not meet) the GV at most sites in the Fitzroy region (except Barren Island FTZ1) and Secchi depth did not meet GV at any sites in the region.

The gap in observational data between 2015 and 2020 limits the utility of the GAMMs in detecting long-term trends over this interval (Figure D-5). Despite this, trend analysis between

the previous end in monitoring (2015) and the current year suggest that Secchi depth, PN, PP, and PO<sub>4</sub> are similar between the time points while Chl-*a* may be showing early signs of improving and TSS may have deteriorated. There is a clearer trend emerging for NO<sub>x</sub>, which is showing signs of improvement (concentrations have decreased) in recent years. Although these are very preliminary findings; we expect long-term trends will continue to become clearer as additional data are collected. There is also evidence that DOC increased since 2015 (as it has in other focus regions monitored under the MMP), but may be declining again in recent years.




Figure D-5: Temporal trends in water quality variables for the Fitzroy focus region: a) chlorophyll a (Chl-a), b) Secchi depth, c) total suspended solids (TSS), d) turbidity, e) nitrate/nitrite ( $\text{NO}_x$ ), f) phosphate ( $\text{PO}_4$ ), g) particulate nitrogen (PN), h) particulate phosphorus (PP), i) particulate organic carbon (POC) and j) dissolved organic carbon (DOC). Generalised additive mixed effect models (trends) are represented by blue lines with shaded areas defining 95% confidence intervals of those trends and black dots represent observed data (depth weighted averages). These trends and data are accounting for the effects of wind, waves, tides, and seasons after applying x-z detrending. Trends of records from ECO FLNTUSB instruments are represented in red, and individual records can be found in Figure C-1. Dashed horizontal reference lines indicate annual guidelines. The apparent steep gradients in some variables during 2020–2023 are likely statistical artefacts of the lack of data from 2015–2020 and will improve as more monitoring data are collected for open coastal waters.

The WQ Index is calculated using two different formulations to communicate the a) long-term trend in water quality (based on the pre-2015 sampling design) and b) annual condition of water quality (based on the post-2015 sampling design, which increased the power to detect change). Section 2.5 and Appendix B contain details of the calculations for both Index formulations.

The long-term WQ Index showed a small (i.e., changing by a single grade) overall improvement in water quality from the period 2007–2015, driven by Chl-a, PN, and PP indicators (Figure D-6a). A trend of improvement in water quality has also been observed between 2020–2023, but the long-term index score stabilised in 2023–24 (Figure D-6a).

The annual condition WQ Index scored water quality as ‘good’ since its inception (2020–present), with the 2023–24 year also receiving a ‘good’ score (Figure D-6b). This version of the Index scores water quality parameters against GVs relevant to the season when samples are collected (wet versus dry GVs).

It is important to note that the two versions of the WQ Index are designed to answer separate questions and therefore differences in scores between the versions are expected.




Figure D-6: The Water Quality Index (WQ Index) for the Fitzroy focus region. The WQ Index uses two formulations to communicate: a) long-term trend (based on pre-2015 sampling design) and b) the annual condition (based on post-2015 sampling design). WQ Index colour coding: ● / ◆ – 'very good'; ○ / ◇ – 'good'; ● / ◇ – 'moderate'; ○ / ◇ – 'poor'; ● / ◇ – 'very poor'. Indicators or sub-indicators that are used to calculate the WQ Index are shown as coloured lines. Error bars (vertical black lines) on the WQ Index represent the 95% quantile intervals. Calculations for both formulations are described in Methods and Appendix B.

## D-6 Discussion and Conclusions

The discharge and loads calculated for the 2023–24 water year from the Fitzroy River were very close to the long-term median (Figure 3-5; Figure D-2; Table 3-1). The discharge has been near or below-median for the previous seven years in this region, and in the 2020–21 water year the discharge and loads calculated were amongst the lowest recorded over the past decade.

*Ambient water quality - Enclosed coastal, open coastal, and mid-shelf waters:*

- Secchi depth did not meet GVs at any sites in the region (Table D-2).
- Concentrations of NO<sub>x</sub> did not meet GVs at most sites in the Fitzroy region (with the exception of Barren Island FTZ1).
- PP met the local GVs at some sites but generally did not meet GVs for the region.
- TSS, PO<sub>4</sub>, and PN met the local GVs at most sites.
- Chl-*a* met GVs at all monitoring sites.
- The trend analysis conducted for other MMP focus regions (trends since the change of monitoring design in 2015) is not appropriate for this dataset as monitoring recommenced in 2020. However, preliminary trend analysis was conducted and as future monitoring occurs, the trends in conditions will become clearer. Trend analysis based on GAMMs suggests that over the last several years:
  - Secchi depth, PN, PP, and PO<sub>4</sub> concentrations show signs of stability since MMP monitoring ended in 2015
  - Chl-*a* may be showing early signs of improvement relative to 2015 while TSS may have deteriorated.
  - There is a clearer trend emerging for NO<sub>x</sub> which is showing signs of improvement (concentrations have decreased) in recent years.
  - More data are needed to confirm these conclusions.
- The long-term WQ Index showed a small (i.e., changing by a single grade) improvement in water quality over the period 2008 to 2015, which was driven by improvements in PN, PP, and Chl-*a* indicators. Over the period 2021–2023, a general trend of stability was seen. For the 2023–24 water year, the annual condition WQ Index score was 'good'.

Table D-2: Water quality indicator summary for Fitzroy region. Performance relative to guideline values is shown as: generally exceeding (✗) or meeting (✓) guideline values across all sites.

Water quality indicator	Mackay-Whitsunday
NO <sub>x</sub>	✗
PO <sub>4</sub>	✓
PN	✓
PP	✗
TSS	✓
Secchi depth	✗
Chl-a	✓

#### *Wet season and event water quality*

- There were no major flood events in the Fitzroy region during the 2023–24 wet season.

## D-7 References

- Álvarez-Romero JG, Devlin M, Teixeira da Silva E, Petus C, Ban NC, Pressey RL, Kool J, Roberts JJ, Cerdeira-Estrada S, Wenger AS Brodie J (2013) A novel approach to model exposure of coastal-marine ecosystems to riverine flood plumes based on remote sensing techniques. *Journal of Environmental Management* 119: 194-207.
- Apan AA, Raine SR, Paterson MS, (2002) Mapping and Analysis of Changes in the Riparian Landscape Structure of the Lockyer Valley Catchment, Queensland, Australia, 59(1): 43-57.
- Askey-Doran M, Bunn S, Hairsine P, Price P, Prosser I, Rutherford I (1996) *Riparian Management 3: Water Quality*, Land and Water Resources Research and Development Corporation, Canberra.
- Bainbridge Z, Lewis S, Bartley R, Fabricius K, Collier C, Waterhouse J, Garzon-Garcia A, Robson B, Burton J, Wenger A, Brodie J (2018) Fine sediment and particulate organic matter: A review and case study on ridge-to-reef transport, transformations, fates, and impacts on marine ecosystems. *Marine pollution bulletin*. 135:1205-20.
- Baird M, Margvelashvili N, Cantin N (2019) Historical context and causes of water quality decline in the Whitsunday region. CSIRO Oceans and Atmosphere. <https://www.environment.gov.au/system/files/resources/995dd4ca-bd44-407d-87e8-a323e67fef6c/files/historical-context-causes-water-quality-decline-whitsundays.pdf>, accessed 30 August 2021.
- Baird ME, Mongin M, Skerratt J, Margvelashvili N, Tickell S, Steven AD, Robillot C, Ellis R, Waters D, Kaniewska P, Brodie J (2021) Impact of catchment-derived nutrients and sediments on marine water quality on the Great Barrier Reef: An application of the eReefs marine modelling system. *Marine Pollution Bulletin*. 167:112297.
- Bartley R, Roth CH, Ludwig J, McJannet D, Liedloff A, Corfield J, Hawdon A, Abbott B (2006) Runoff and erosion from Australia's tropical semi-arid rangelands: Influence of ground cover for differing space and time scales. *Hydrological Processes: An International Journal*. 20(15):3317-33.
- Berkelmans R, Jones AM, Schaffelke B (2012) Salinity thresholds of *Acropora* spp. on the Great Barrier Reef. *Coral Reefs* 31(4): 1103-1110.
- BOM (Bureau of Meteorology) (2021) Fitzroy: Climate and water, accessed 11 October 2022. <http://www.bom.gov.au/water/nwa/2021/fitzroy/index.shtml>
- Brandt ME, Smith TB, Correa AM, Vega-Thurber R (2013) Disturbance driven colony fragmentation as a driver of a coral disease outbreak. *PLoS One*. 8(2):e57164.
- Brodie JE, Mitchell AW (1992) Nutrient composition of the January (1991) Fitzroy River flood plume, pp. 56–74 in *Workshop on the Impacts of Flooding*, ed. G.T. Byron, Workshop Series No. 17, Great Barrier Reef Marine Park Authority, Townsville.
- Brodie JE, McKergow LA, Prosser IP, Furnas M, Hughes AO, Hunter H. (2003) Sources of sediment and nutrient exports to the Great Barrier Reef World Heritage Area.
- Brodie J, Fabricius K, De'ath G, Okaji K (2005) Are increased nutrient inputs responsible for more outbreaks of crown-of-thorns starfish? An appraisal of the evidence. *Marine pollution bulletin*. 1;51(1-4):266-78.
- Brodie JE, Devlin M, Haynes D, Waterhouse J (2011) Assessment of the eutrophication status of the Great Barrier Reef lagoon (Australia). *Biogeochemistry*. 106(2):281-302.

- Brodie J, Grech A, Pressey B, Day J, Dale AP, Morrison T, Wenger A (2019) The future of the Great Barrier Reef: the water quality imperative. In *Coasts and Estuaries* (pp. 477-499). Elsevier.
- Bruno JF, Petes LE, Drew Harvell C, Hettinger A (2003) Nutrient enrichment can increase the severity of coral diseases. *Ecology letters*. 6(12):1056-61.
- Cai W, Santoso A, Collins M, Dewitte B, Karamperidou C, Kug JS, Lengaigne M, McPhaden MJ, Stuecker MF, Taschetto AS, Timmermann A (2021) Changing El Niño–Southern Oscillation in a warming climate. *Nature Reviews Earth & Environment*. 7:1-7.
- Cantin N, Wu Y, Fallon S, Lough J (2019) Historical records of terrestrial sediment and flood plume inputs to the Whitsunday Island region from coral skeletons: 1861-2017. Australian Institute of Marine Science, Townsville, Qld. 30p.
- Carroll C, Halpin M, Burger P, Bell K, Sallaway MM, Yule DF (1997) The effect of crop type, crop rotation, and tillage practice on runoff and soil loss on a Vertisol in central Queensland. *Soil Research*. 35(4):925-40.
- Chua EM, Wilson SP, Vink S, Flint N (2019) The influence of riparian vegetation on water quality in a mixed land use river basin. *River Research and Applications*. 35(3):259-67.
- Collins M, An SI, Cai W, Ganachaud A, Guilyardi E, Jin FF, Jochum M, Lengaigne M, Power S, Timmermann A, Vecchi G (2010) The impact of global warming on the tropical Pacific Ocean and El Niño. *Nature Geoscience*. 3(6):391-7.
- CRC for Coastal Zone, Estuary & Waterway Management (CRC) (2003) Central Queensland information paper. Volume 6: Fitzroy Catchment, Queensland.
- Department of Regional Development, Manufacturing and Water (DRMW) (2023). River discharge data. <https://water-monitoring.information.qld.gov.au/> Accessed October 2023.
- De Vente J, Poesen J, Verstraeten G (2005) The application of semi-quantitative methods and reservoir sedimentation rates for the prediction of basin sediment yield in Spain. *Journal of Hydrology*. 305(1-4):63-86.
- Devlin M, Waterhouse J, Taylor J, Brodie J (2001) Flood plumes in the Great Barrier Reef: spatial and temporal patterns in composition and distribution. GBRMPA research publication. 68.
- Devlin M (2008) The 2008 flood plume event on the Fitzroy River. Catchment to Reef Group, James Cook University. [https://www.qbrmpa.gov.au/\\_data/assets/pdf\\_file/0010/7687/JCU\\_Fitzroy\\_v1\\_2009.pdf](https://www.qbrmpa.gov.au/_data/assets/pdf_file/0010/7687/JCU_Fitzroy_v1_2009.pdf)
- Diaz-Pulido G, McCook LJ, Dove S, Berkelmans R, Roff G, Kline DI, Weeks S, Evans RD, Williamson DH, Hoegh-Guldberg O (2009) Doom and boom on a resilient reef: climate change, algal overgrowth and coral recovery. *PloS one*. 4(4):e5239.
- Dougall C, Packett R, Carroll C (2005) Application of the SedNet model in partnership with the Fitzroy Basin community. In MODSIM 2005 International Congress on Modelling and Simulation. Modelling and Simulation Society of Australia and New Zealand 2005 Dec (pp. 1119-1125).
- Dougall C, McCloskey GL, Ellis R, Shaw M, Waters D, Carroll C (2014) Modelling reductions of pollutant loads due to improved management practices in the Great Barrier Reef catchments – Fitzroy NRM region, Technical Report, Volume 6, Queensland Department of Natural Resources and Mines, Rockhampton, Queensland (ISBN: 978-0-7345- 0444-9).
- Emanuel K, Sundararajan R, Williams J (2008) Hurricanes and global warming: Results from downscaling IPCC AR4 simulations. *Bulletin of the American Meteorological Society*. 89(3):347-68.

- Fabricius KE (2005) Effects of terrestrial runoff on the ecology of corals and coral reefs: review and synthesis. *Marine pollution bulletin*. 50(2):125-46.
- FBA (Fitzroy Basin Association) (2018) Census of Population and Housing, 2016, accessed 30 August 2021, [Population demographics | Fitzroy Basin Association \(fba.org.au\)](https://www.fba.org.au/population-demographics)
- Furnas M, (2003) Catchments and Corals: Terrestrial Runoff to the Great Barrier Reef, Australian Institute of Marine Science and CRC Reef Research Centre, Townsville.
- Gallen C, Devlin M, Thompson K, Paxman C, Mueller J (2014) Pesticide monitoring in inshore waters of the Great Barrier Reef using both time-integrated and event monitoring techniques (2013 - 2014). The University of Queensland, The National Research Centre for Environmental Toxicology (Entox). <http://hdl.handle.net/11017/2930>, accessed 30 August 2021.
- Great Barrier Reef Marine Park Authority (2010). Water Quality Guidelines for the Great Barrier Reef Marine Park. Revised Edition 2010. Great Barrier Reef Marine Park Authority, Townsville. 100pp.
- Great Barrier Reef Marine Park Authority (2019) Great Barrier Reef Outlook Report 2019. Great Barrier Reef Marine Park Authority, <https://elibrary.gbrmpa.gov.au/jspui/bitstream/11017/3474/13/Outlook-Report-2019-Intro.pdf> , accessed 30 August 2021.
- Great Barrier Reef Marine Park Authority (2024), Great Barrier Reef Marine Monitoring Program Quality Assurance and Quality Control Manual 2022–23, Great Barrier Reef Marine Park Authority, Townsville, 189pp.
- Gualdi S, Scoccimarro E, Navarra A (2008) Changes in tropical cyclone activity due to global warming: Results from a high-resolution coupled general circulation model. *Journal of climate*. 21(20):5204-28.
- Haapkylä J, Unsworth RK, Flavell M, Bourne DG, Schaffelke B, Willis BL (2011) Seasonal rainfall and runoff promote coral disease on an inshore reef. *PloS one*. 10;6(2):e16893.
- Harris GP (2001) Biogeochemistry of nitrogen and phosphorus in Australian catchments, rivers and estuaries: effects of land use and flow regulation and comparisons with global patterns. *Marine and freshwater research*. 52(1):139-49.
- Hawdon AA, Keen RJ, Post DA, Wilkinson SN (2008) Hydrological recovery of rangeland following cattle exclusion. *IAHS publication*. 325:532.
- Hughes AO, Olley JM, Croke JC, McKergow LA (2009) Sediment source changes over the last 250 years in a dry-tropical catchment, central Queensland, Australia. *Geomorphology*. 104(3-4):262-75.
- Huon S, Bellanger B, Bonté P, Sogon S, Podwojewski P, Girardin C, Valentin C, de Rouw A, Velasquez F, Bricquet JP, Mariotti A (2005) Monitoring soil organic carbon erosion with isotopic tracers: two case studies on cultivated tropical catchments with steep slopes (Laos, Venezuela). *Soil Erosion and Carbon Dynamics* (pp. 301-328). CRC Press.
- IPCC (2021) Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S. L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M. I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J. B. R. Matthews, T. K. Maycock, T. Waterfield, O. Yelekçi, R. Yu and B. Zhou (eds.)]. Cambridge University Press. In Press.
- Jones AM, Berkelmans R (2014) Flood impacts in Keppel Bay, southern Great Barrier Reef in the aftermath of cyclonic rainfall. *PloS one*. 9(1):e84739.

- Kline DI, Kuntz NM, Breitbart M, Knowlton N, Rohwer F (2006) Role of elevated organic carbon levels and microbial activity in coral mortality. *Marine Ecology Progress Series*. 314:119-25.
- Kuntz NM, Kline DI, Sandin SA, Rohwer F (2005) Pathologies and mortality rates caused by organic carbon and nutrient stressors in three Caribbean coral species. *Marine Ecology Progress Series*. 294:173-80.
- Kroon FJ, Kuhnert PM, Henderson BL, Wilkinson SN, Kinsey-Henderson A, Abbott B, Brodie JE, Turner RD (2012) River loads of suspended solids, nitrogen, phosphorus and herbicides delivered to the Great Barrier Reef lagoon. *Marine pollution bulletin*. 65(4-9):167-81.
- Lewis SE, Bartley R, Wilkinson SN, Bainbridge ZT, Henderson AE, James CS, Irvine SA, Brodie JE (2021) Land use change in the river basins of the Great Barrier Reef, 1860 to 2019: A foundation for understanding environmental history across the catchment to reef continuum. *Marine Pollution Bulletin*. 166:112193.
- Luick JL, Mason L, Hardy T, Furnas MJ (2007) Circulation in the Great Barrier Reef Lagoon using numerical tracers and *in situ* data. *Continental Shelf Research*. 27(6):757-78.
- Marwick TR, Borges AV, Van Acker K, Darchambeau F, Bouillon S (2014) Disproportionate contribution of riparian inputs to organic carbon pools in freshwater systems. *Ecosystems*. 17(6):974-89.
- McKenzie, LJ, Collier, CJ, Langlois, LA, Brien, H and Yoshida, RL (2024). Marine Monitoring Program: Annual Report for Inshore Seagrass Monitoring 2022–23. Report for the Great Barrier Reef Marine Park Authority, Great Barrier Reef Marine Park Authority, Townsville. 178pp.
- Meat and Livestock Australia (MLA) (2021) Industry Projections 2021 - Australian cattle - February, [feb2021-mla-australian-cattle-industry-projections.pdf](#), accessed 30 August 2021.
- Moran, D., Robson, B., Gruber, R., Waterhouse, J., Logan, M., Petus, C., Howley, C., Lewis, S., James, C., Tracey, D., Mellors, J., O'Callaghan, M., Bove, U., Davidson, J., Glasson, K., Jaworski, S., Lefevre, C., Nordborg, M., Vasile, R., Zagorskis, I., Shellberg, J., 2023. Marine Monitoring Program: Annual Report for Inshore Water Quality Monitoring 2021–22. Report for the Great Barrier Reef Marine Park Authority, Great Barrier Reef Marine Park Authority, Townsville.
- Murakami H, Sugi M (2010) Effect of model resolution on tropical cyclone climate projections. *Sola*. 6:73-6.
- NOAA (2017) National Weather Service: Climate Prediction Center. [Climate Prediction Center - ONI \(noaa.gov\)](#), accessed 30 August 2021.
- O'Neill JP, Byron GT, Wright SC (1992) Some physical characteristics and movement of 1991 Fitzroy River flood plume. In Workshop Series (No. 17, pp. 36-51).
- Orpin AR, Ridd PV, Stewart LK (1999) Assessment of the relative importance of major sediment-transport mechanisms in the central Great Barrier Reef lagoon. *Australian Journal of Earth Sciences*. 46(6):883-96.
- Packett R, Dougall C, Rohde K, Noble R (2009) Agricultural lands are hot-spots for annual runoff polluting the southern Great Barrier Reef lagoon. *Marine pollution bulletin*. 58(7):976-86.
- Preker M, Byron GT (1992) The effects of the 1991 central Queensland floodwaters around Heron Island, Great Barrier Reef. In Workshop series. Great Barrier Reef Marine Park Authority.

- QLUMP (Queensland Land Use Mapping Program) (2019) Fitzroy and Burnett Mary natural resource management regions, <https://www.qld.gov.au/environment/land/management/mapping/statewide-monitoring/qlump/qlump-datasets> accessed 30 August, 2021.
- R Core Team (2022) R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing. Vienna, Austria. URL: <https://www.R-project.org/>
- Rogers CS, Miller J (2006) Permanent 'phase shifts' or reversible declines in coral cover? Lack of recovery of two coral reefs in St. John, US Virgin Islands. *Marine Ecology Progress Series*. 306:103-14.
- Roth F, Saalman F, Thomson T, Coker DJ, Villalobos R, Jones BH, Wild C, Carvalho S (2018) Coral reef degradation affects the potential for reef recovery after disturbance. *Marine Environmental Research*. 142:48-58.
- Shen P, Zhang LM, Chen HX, Gao L (2017) Role of vegetation restoration in mitigating hillslope erosion and debris flows. *Engineering Geology*. 216:122-33.
- Silburn DM, Carroll C, Ciesiolka CA, DeVoi RC, Burger P (2011) Hillslope runoff and erosion on duplex soils in grazing Lands in semi-arid central Queensland. I. Influences of cover, slope, and soil. *Soil Research*. 49(2):105-17.
- Sugi M, Murakami H, Yoshimura J (2009) A reduction in global tropical cyclone frequency due to global warming. *Sola*. 5:164-7.
- The State of Queensland and Commonwealth of Australia (2003) Reef water quality protection plan 2003: for the Great Barrier Reef World Heritage Area and adjacent catchments, [new reefplan.indd \(gbrmpa.gov.au\)](#), accessed 30 August 2021.
- The State of Queensland and Commonwealth of Australia (2009) Reef water quality protection plan 2009: for the Great Barrier Reef World Heritage Area and adjacent catchments, <http://hdl.handle.net/11017/1125>
- The State of Queensland (2013) Securing the health and resilience of the Great Barrier Reef World Heritage Area and adjacent catchment, [reef-plan-2013.pdf \(reefplan.qld.gov.au\)](#), accessed 30 August 2021.
- The State of Queensland (2018) Reef 2050 Water Quality Improvement Plan: 2017-2022. [Reef 2050 Water Quality Improvement Plan 2017-2022 \(reefplan.qld.gov.au\)](#), accessed 30 August 2021.
- Thibault PA. Ground cover patterns near streams for urban land use categories (1997) *Landscape and urban planning*. 39(1):37-45.
- Thompson A, Costello P, Davidson J, Schaffelke B, Uthicke S and Liddy M (2013) Reef Rescue Marine Monitoring Program. Report of AIMS Activities – Inshore coral reef monitoring 2012. Report for Great Barrier Reef Marine Park Authority. Australian Institute of Marine Science, Townsville. 120 p
- Thompson A., C. Lønborg, P. Costello, J. Davidson, M. Logan, M. Furnas, K. Gunn, M. Liddy, M. Skuza, S. Uthicke, M. Wright I. Zagorskis, and B. Schaffelke (2014a). Marine Monitoring Program. Annual Report of AIMS Activities 2013 to 2014– Inshore water quality and coral reef monitoring. Report for the Great Barrier Reef Marine Park Authority. Australian Institute of Marine Science, Townsville. 146 pp.
- Thompson A, Schroeder T, Brando VE, Schaffelke B (2014b) Coral community responses to declining water quality: Whitsunday Islands, Great Barrier Reef, Australia. *Coral Reefs*. 33(4):923-38.

- Thompson A, Davidson J, Logan M, Thompson C (2024). Marine Monitoring Program Annual Report for Inshore Coral Reef Monitoring: 2022–23. Report for the Great Barrier Reef Marine Park Authority, Great Barrier Reef Marine Park Authority, Townsville. 149 pp.
- Thorburn PJ, Wilkinson S, Meier E (2010) Prioritising practice changes in Reef Rescue. CSIRO Water for a Health Country, ISSN: 1835-095X, 56 pp.
- Thorburn PJ, Wilkinson SN (2013) Conceptual frameworks for estimating the water quality benefits of improved agricultural management practices in large catchments. *Agriculture, ecosystems & environment*. 180:192-209.
- Thorburn PJ, Wilkinson SN, Silburn DM (2013) Water quality in agricultural lands draining to the Great Barrier Reef: a review of causes, management and priorities. *Agriculture, ecosystems & environment*. 180:4-20.
- Tracey D, Waterhouse J, Da Silva E (2017) Preliminary investigation of alternative approaches for the Reef Plan Report Card Water Quality Metric report. Great Barrier Reef Marine Park Authority.
- van Woessik R, Done TJ (1997) Coral communities and reef growth in the southern Great Barrier Reef. *Coral Reefs*. 16(2):103-15.
- Vega-Thurber RL, Burkepile DE, Fuchs C, Shantz AA, McMinds R, Zaneveld JR (2014) Chronic nutrient enrichment increases prevalence and severity of coral disease and bleaching. *Global change biology*. 20(2):544-54.
- Walsh KJ, McInnes KL, McBride JL (2012) Climate change impacts on tropical cyclones and extreme sea levels in the South Pacific—A regional assessment. *Global and Planetary Change*. 80:149-64.
- Waltham NJ, Wegscheidl C, Volders A, Smart JC, Hasan S, Lédée E, Waterhouse J. Land use conversion to improve water quality in high DIN risk, low-lying sugarcane areas of the Great Barrier Reef catchments (2021) *Marine Pollution Bulletin*. 167:112373.
- Wasson RJ, Caitcheon G, Murray AS, McCulloch MA, Quade JA (2002) Sourcing sediment using multiple tracers in the catchment of Lake Argyle, Northwestern Australia. *Environmental Management*. 29(5):634-46.
- Waterhouse J, Henry N, Mitchell C, Smith R, Thomson B, Carruthers C, Bennett J, Brodie J, McCosker K, Northey A, Poggio M, Moravek T, Gordon B, Orr G, Silburn M, Shaw M, Bickle M, Ronan M, Turner R, Waters D, Tindall D, Trevithick R, Ryan T, VanderGragt M, Houlden B, Robillot C (2018) Paddock to Reef Integrated Monitoring, Modelling and Reporting Program: Program Design 2018-2022, [Revised P2R draft for input \(reefplan.qld.gov.au\)](#), accessed 30 August 2021.
- Weber M, De Beer D, Lott C, Polerecky L, Kohls K, Abed RM, Ferdelman TG, Fabricius KE (2012) Mechanisms of damage to corals exposed to sedimentation. *Proceedings of the National Academy of Sciences*. 109(24):E1558-67.
- Webster AJ, Bartley R, Armour JD, Brodie JE, Thorburn PJ (2012) Reducing dissolved inorganic nitrogen in surface runoff water from sugarcane production systems. *Marine Pollution Bulletin*. 65(4-9):128-35.
- Wooldridge S, Brodie J, Furnas M (2006) Exposure of inner-shelf reefs to nutrient enriched runoff entering the Great Barrier Reef Lagoon: Post-European changes and the design of water quality targets. *Marine pollution bulletin*. 52(11):1467-79.
- Wooldridge SA (2009) Water quality and coral bleaching thresholds: Formalising the linkage for the inshore reefs of the Great Barrier Reef, Australia. *Marine Pollution Bulletin*. 58(5):745-51.

- Yu B, Joo M, Carroll C (2013) Land use and water quality trends of the Fitzroy River, Australia. Understanding Freshwater Quality Problems in a Changing World Proceedings of H. 4:1-8.
- Zaimes GN, Schultz RC, Isenhardt TM (2008) Streambank Soil and Phosphorus Losses Under Different Riparian Land-Uses in Iowa 1. JAWRA Journal of the American Water Resources Association. 44(4):935-47.
- Zhao M, Held IM, Lin SJ, Vecchi GA (2009) Simulations of global hurricane climatology, interannual variability, and response to global warming using a 50-km resolution GCM. Journal of Climate. 22(24):6653-78.

## APPENDIX E. SCIENTIFIC PUBLICATIONS AND PRESENTATIONS ASSOCIATED WITH THE PROGRAM, 2023–24

### E-1 Publications

#### *Reports and scientific publications*

Great Barrier Reef Marine Park Authority (2024) Great Barrier Reef Marine Monitoring Program Quality Assurance and Quality Control Manual 2022-23, Great Barrier Reef Marine Park Authority, Townsville, 189 pp.

Great Barrier Reef Marine Park Authority (2024) Great Barrier Reef Marine Monitoring Program: Monitoring of the Inshore Reef Regions 2022-23, Great Barrier Reef Marine Park Authority, Townsville.

Gruber, R., Waterhouse, J., Petus, C., Howley, C., Lewis, S., Moran, D., James, C., Logan, M., Bove, U., Brady, B., Choukroun, S., Connellan, K., Davidson, J., Mellors, J., O'Callaghan, M., O'Dea, C., Shellberg, J., Tracey, D., Zagorskis, I., (2024). Great Barrier Reef Marine Monitoring Program: Annual Report for Inshore Water Quality Monitoring 2022–23. Report for the Great Barrier Reef Marine Park Authority, Great Barrier Reef Marine Park Authority, Townsville. 284 pp.

Prazeres, M., Thompson, A., Gruber, R., Waterhouse, J., McKenzie, L., Howley, C., Thompson, C., Lewis, S., Walker, K. (2024) Great Barrier Reef Marine Monitoring Program: Synthesis Report 2022–23, Great Barrier Reef Marine Park Authority, Townsville.

#### *Data used for model validation and external investigations*

During the 2023–24 year, MMP WQ data has been used by several external groups, including:

- Validation of the eReefs marine models for the Great Barrier Reef, led by Jenny Skerratt at CSIRO. An extensive list of resulting publications is available from: <https://research.csiro.au/ereefs/models/further-reading/>
- Incorporation of data into the IMOS Bio-Optical Database for validation of satellite imagery, led by Thomas Schroeder at CSIRO.
- Water quality data was provided to NRM technical officers at the Mackay-Whitsunday-Isaac Healthy Rivers to Reef Partnership, the Dry Tropics Partnership for Healthy Waters, and the Wet Tropics Healthy Waterways Partnership to be used in preparation of the latest Regional Report Cards.

#### *Related papers and reports – linking to MMP data/methods:*

Lewis S, Bainbridge Z, Smithers S (2024) Question 2.3 What evidence is there for changes in land-based runoff from pre-development estimates in the Great Barrier Reef? In Waterhouse J, Pineda M-C, Sambrook K (Eds) 2022 Scientific Consensus Statement on land-based impacts on Great Barrier Reef water quality and ecosystem condition. Commonwealth of Australia and Queensland Government.

Lewis S, Bainbridge Z, Smithers S (2024) Question 3.1 What are the spatial and temporal distributions of terrigenous sediments and associated indicators within the Great Barrier Reef? In Waterhouse J, Pineda M-C, Sambrook K (Eds) 2022 Scientific Consensus Statement on land-based impacts on Great Barrier Reef water quality and ecosystem condition. Commonwealth of Australia and Queensland Government.

- Flom, M. I. (2023). Towards establishing spatial and temporal patterns in inshore pelagic microbial communities: a baseline study in the Wet Tropics, Great Barrier Reef, Australia. Honours thesis, James Cook University.
- Puignou Lopez, O. Lewis, S. James, C. Davis, A. Mackay, S. (in review). Hydrology of the Great Barrier Reef catchment area along a latitudinal gradient: Upscaling discharge to reflect catchment inputs. *Journal of Hydrology*.
- Robson B, Brown A, Uthicke S (2024) Question 4.1 What is the spatial and temporal distribution of nutrients and associated indicators within the Great Barrier Reef? In Waterhouse J, Pineda M-C, Sambrook K (Eds) 2022 Scientific Consensus Statement on land-based impacts on Great Barrier Reef water quality and ecosystem condition. Commonwealth of Australia and Queensland Government.
- Terzin M., Robbins S.J., Bell S.C., Cao K.L., Gruber R.K., Frade P.R., Webster N.S., Yeoh Y.K., Bourne D.G., Laffy P.W. (in review) Gene Content of Seawater Microbes is a Strong Predictor of Water Chemistry Across the Great Barrier Reef, 05 September 2024, PREPRINT (Version 1) available at Research Square [<https://doi.org/10.21203/rs.3.rs-4900069/v1>]

## E-2 Presentations

- Davidson J., Grant C., Evans-Illidge E., Forrester M. Meeting with Manbarra TOs regarding FPIC for AIMS / MMP sampling. Meeting held at Mingga Mingga Ranger HQ on Palm Island. 4 July 2024
- Gruber R., GBR MMP: Inshore water quality results 2022-23. AIMS Program 2 staff webinar, online via Teams, 5 February 2024
- Gruber R., Monitoring water quality in the GBR lagoon presented to ~30 Burdekin graziers at a visit to AIMS organised via NQ Dry Tropics. AIMS, Townsville, QLD, 17 April 2024.
- Gruber R. Inshore water quality: 2023-24 condition and long-term trends. Presentation at MMP Symposium to ~100 stakeholder (50 in person, 50 online) from MMP teams, GBRMPA, RRCs, OGBR, TOs. Symposium held at Townsville Yacht Club and online, Townsville QLD. 13 November 2024.
- Howley, Shellberg, Albert-Mitchell. Environmental Impacts of Cyclone Jasper in the Cape York region. March-July 2024. This was part of a series of presentations funded by Commonwealth & DESI DRFA, including overview of details collected via MMP Inshore WQ monitoring team.
- Howley et al., Inshore Water Quality in the Cape York Region 2023-24 (MERI Workshop)
- Howley C, Herkess S, Shellberg J. Reducing Land Use Impacts & Sediment Pollution to the Great Barrier Reef - SE Cape York Peninsula. Presented to GBRF Regional Sediment Program Coordinator Forum. Online 8 May 2023
- Howley et al., Inshore Water Quality in the Cape York Region 2023-(Yinting Aboriginal Corp TUMRA Meeting- June 2024)
- Howley, C. and Polglase, L. Inshore Water Quality in the Cape York Region, Pascoe sub-region focus. Presented to Kuku Ya'u Aboriginal Corporation AGM. Lockhart River and online- 15 November 2024.
- Howley C. Inshore water quality monitoring – Cape York condition, trends and Cyclone Jasper. Presentation at MMP Symposium to ~100 stakeholder (50 in person, 50 online) from MMP teams, GBRMPA, RRCs, OGBR, TOs. Symposium held at Townsville Yacht Club and online, Townsville QLD. 13 November 2024.

- Lewis S. Inshore water quality monitoring – Event sampling in the Wet Tropics. Presentation at MMP Symposium to ~100 stakeholder (50 in person, 50 online) from MMP teams, GBRMPA, RRCs, OGBR, TOs. Symposium held at Townsville Yacht Club and online, Townsville QLD. 13 November 2024.
- Moran et al. Inshore water quality ambient monitoring for the Inshore Reef 2023-24 (MERI Workshop)
- Moran D. Inshore water quality monitoring at AIMS. Presentation to MH Premium Farms Management and Executives group visit hosted at AIMS, Townsville, 30 August 2023.
- Moran D, Connellan KL., and CYWP team. MMP WQ Annual QA/Training Workshop. Workshop with AIMS and CYWP teams, at CYWP Water Shed', Cooktown, QLD, 27 Sept 2023.
- Moran D. and JCU team. MMP WQ Annual QA/Training Workshop. Workshop with AIMS and JCU teams, JCU ATSIP Building, Townsville, QLD, 9 Nov 2023.
- Moran D., Introduction to MMP Inshore Water Quality Monitoring presented to Reef Authority staff at Reef Authority Internal Seminar. Online via Teams, 3 October 2023
- Moran D. Introduction to Water Quality Monitoring presented to 92 Indigenous Women Rangers from Traditional Owner groups throughout Queensland for the Queensland Indigenous Womens Rangers Network open day at AIMS, Townsville, QLD, 23 November 2023.
- Moran D., Introduction to MMP Inshore Water Quality Monitoring presented to ~20 marine science students at a visit to AIMS organised via the JCU winter school program. AIMS, Townsville, QLD, 26 June 2024
- Moran D. Presentation to primary school students relating to sustainability and water cycle themes and linking to MMP inshore water quality, coral and seagrass monitoring. Presentation at St Anthony's Primary School. Townsville, QLD. 27 August 2024.
- Moran D., Grant C. Meeting with Jabalbina PBC to introduced the MMP coral and water Quality programs ahead of future meetings and discussions regarding FPIC. Meeting held online via Teams. 21 October 2024.
- Robson B. Fitzroy Basin Marine Monitoring Program for Inshore Water Quality results: 2022-23. Presented at Fitzroy Partnership for River Health Water Forum, Rockhampton, QLD. 24 October 2024.
- Streten C., Wachenfeld D. The science of long term inshore water quality and coral monitoring and links to broader government ecosystem around The Reef. Presentation to AIMS Council. Townsville, QLD. 29 October 2024.
- Thompson A., Grant C. Meeting with Gunggaandji PBC to introduced the MMP coral and water Quality programs ahead of future meetings and discussions regarding FPIC. Meeting held online via Teams. 13 August 2024.
- Waterhouse J., Howley C., Moran D. IMPACTS OF TROPICAL CYCLONE JASPER - How a 1-1000 year event transformed Great Barrier Reef catchments and impacted inshore marine ecosystems. Presentation of MMP Inshore Water Quality observations from 2023-24 to Reef Authority Science Seminar Series. Townsville, QLD. 5 September 2024.
- Waterhouse J., Howley C., Moran D., Robson B. Presentations and discussion of 2023-24 inshore water quality observations at MMP Synthesis Workshop 2024. Townsville QLD. 23 October 2024.
- Waterhouse, Howley, Petus, Lewis, Mellors, James, O'Callaghan. Marine Monitoring Program Inshore WQ monitoring, October 2023

Waterhouse et al., Inshore water quality wet season monitoring in the GBR 2023-24 (MERI Workshop)

Waterhouse J. The 2023-24 wet season: water quality drivers and conditions. Presentation at MMP Symposium to ~100 stakeholder (50 in person, 50 online) from MMP teams, GBRMPA, RRCs, OGBR, TOs. Symposium held at Townsville Yacht Club and online, Townsville QLD. 13 November 2024.